

The spatial distribution of ammonia, methane and nitrous oxide emissions from agriculture in the UK 2009

Annual Report to Defra (Project AC0112)

Dragosits U. and Sutton M.A.

*Centre for Ecology & Hydrology (Edinburgh Research Station)
Bush Estate, Penicuik, Midlothian, EH26 0QB, UK.*

April 2011

Client: North Wyke Research	
Client Project number: Defra AC0112	CEH Project number: C03642
Project Title: Inventories of Ammonia and Greenhouse Gasses from UK Agriculture	
Start date: 1-May-2008	Completion date: 31/03/2013
Client Project Officers: Tom Misselbrook (NWR) for Defra	
CEH Project Officer: Ulli Dragosits	
Main authors: Dragosits U. and Sutton M.A.	
Reporting period: 05/10 – 04/11	
Report date: 15-04-2011	

EXECUTIVE SUMMARY

Modelling and mapping UK ammonia and greenhouse gas emissions from agriculture Defra project AC0112

- Agricultural emissions of ammonia, methane and nitrous oxide for 2009 were spatially distributed across the UK, and maps produced.
- Agricultural emission sources were spatially distributed using the CEH/University of Edinburgh AENEID model. The model incorporates detailed agricultural census data, landcover data, agricultural practice information (e.g. fertiliser application rates, stocking densities) and emission source strength data from the UK emissions inventories for agriculture 2009 (Misselbrook *et al.* 2010, Cardenas *et al.* 2010).
- All emission maps correspond to the totals reported by North Wyke Research (NWR) for 2009 (Misselbrook *et al.* 2010, Cardenas *et al.* 2010).

CONTENTS

EXECUTIVE SUMMARY	3
CONTENTS	4
1. INTRODUCTION	5
1.1. BACKGROUND	5
1.2. ANNUAL WORK SCHEDULE/DELIVERABLES	5
2. METHODS - SPATIAL DISTRIBUTION OF NH₃, CH₄ AND N₂O EMISSIONS FROM AGRICULTURAL SOURCES	6
3. RESULTS.....	7
3.1. SPATIALLY DISTRIBUTED EMISSIONS OF NH ₃ , CH ₄ AND N ₂ O FOR 2009.....	7
3.2. MAJOR CHANGES AND CONSEQUENCES.....	11
3.2.1. <i>Changes in emissions from agricultural NH₃ sources</i>	11
3.2.2. <i>Changes in emissions from agricultural CH₄ and N₂O sources</i>	11
4. CONCLUSIONS.....	12
ACKNOWLEDGEMENTS.....	12
REFERENCES	12
APPENDIX A - NARSES CATEGORIES.....	14

1. INTRODUCTION

1.1. Background

Emissions of ammonia, methane and nitrous oxide for 2009 were spatially distributed using the AENEID model (Dragosits *et al.* 1998, Hellsten *et al.* 2008) and mapped for the UK. This report briefly describes the methodology used for the sources listed above, including any changes in the methodology and the consequences of these changes.

The agricultural emission estimates for ammonia (NH₃), methane (CH₄) and nitrous oxide (N₂O) are derived annually under Defra project AC0112 (inventories by Misselbrook *et al.* and Cardenas at North Wyke Research (NWR); see Table 1). The current contract exploits the expertise of CEH in spatially distributing emissions from agricultural sources, and complements the expertise of North Wyke Research in producing UK emission estimates from experimental data, peer-reviewed literature and agricultural management practices, including mitigation options.

Due to data licensing restrictions in relation with the Data Protection Act, the detailed 5 km model output can currently only be shown as “emissions from livestock” rather than for individual livestock sectors.

Table 1: UK emissions of ammonia (NH₃), methane (CH₄) and nitrous oxide (N₂O) for 2009, as collated by NWR and mapped by CEH (in kt yr⁻¹). Totals may not add up exactly due to rounding.

Gas	Source	UK emission (kt) 2009
NH ₃	Livestock manure	195.8 kt NH ₃
	Fertiliser application	36.0 kt NH ₃
	Total agriculture	231.8 kt NH₃
CH ₄	Enteric fermentation	718.7 kt CH ₄
	Livestock manure	133.3 kt CH ₄
	Total agriculture	852.1 kt CH₄
N ₂ O	Crops & soils	80.3 kt N ₂ O
	Livestock manure	6.4 kt N ₂ O
	Total agriculture	86.8 kt N₂O

1.2. Annual work schedule/deliverables

- Task 1: To acquire source data (agricultural census) from the devolved authorities for spatially distributing agricultural ammonia emissions from livestock manures and fertiliser application.
- Task 2: To model NH₃, CH₄ and N₂O emissions from agricultural sources at a 5 km grid resolution using the AENEID model for the UK, including conversion of results for Northern Ireland from the Irish Ordnance Survey Grid to the Great Britain Ordnance Survey Grid (OSGB).
- Task 3: To provide a short report describing the methodology and results, highlighting any changes and their consequences.
- Task 4: To streamline the inventory process jointly between CEH and NWR. This includes updating the CEH AENEID model to match inventory requirements, e.g., for dealing with new livestock categories supplied by the devolved authorities.

- Task 5: To submit the spatial datasets to AEA for inclusion in the National Atmospheric Emission Inventory (NAEI) and Greenhouse Gas Inventory (GHGI).

2. METHODS - SPATIAL DISTRIBUTION OF NH_3 , CH_4 AND N_2O EMISSIONS FROM AGRICULTURAL SOURCES

Agricultural census/survey data for 2009 were acquired at the finest available spatial resolution from the devolved authorities in the UK, i.e. Defra (England), the Scottish Executive (Scotland), Welsh Assembly (Wales) and DARD NI (Northern Ireland). The census data for the different countries were aggregated to a common set of categories, referred to as the “NARSES categories” (see Appendix A), to ensure compatibility between the different countries’ systems.

The agricultural emission inventory for NH_3 was mapped using output from the NARSES model at NWR (Misselbrook *et al.* 2010, Defra project AC0112). As in previous years, detailed emission source strength estimates were derived for the main livestock emission components (housing, manure storage, landspreading of manures, grazing) for each NARSES category. Average fertiliser N application rates to different crops were taken from the British Survey of Fertiliser Practice for 2009 (BSFP 2010).

These detailed data were applied in the AENEID model (Dragosits *et al.* 1998, Hellsten *et al.* 2008), which distributes livestock and fertiliser emissions to different land cover types (e.g. arable land, improved grass, part-improved grass, rough grazing etc.) derived from the CEH landcover map (LCM2000).

Methane emissions from agriculture were distributed using the greenhouse gas version of the AENEID model (Sutton *et al.* 2004, 2006), which takes account of the spatial location of the CH_4 sources (i.e. animals and manures) specifically. Nitrous oxide emissions from livestock manures were distributed using the N_2O version of AENEID, while the spatial distribution of N_2O emissions summarised under “soils” includes a number of different sources, which were modelled mapped as follows:

- Fertiliser application (19.0 kt N_2O): Emissions from this source were spread via the AENEID output for all crops and grassland, combined with detailed data on fertiliser application rates from the British Survey of Fertiliser Practice for 2000 (BSFP 2010).
- Grazing and manure spreading (21.2 kt N_2O): Emissions from this source were calculated via N excretion rates of grazed livestock combined with the spatial distribution of grazed livestock from AENEID and the manure spreading calculations used in AENEID.
- Leaching (24.8 kt N_2O): Emissions from this source were spread via the AENEID output for all crops and grassland.
- Crop residues (8.5 kt N_2O): Emissions from this source were spread evenly over all NARSES crop categories, excluding grassland, fruit and nursery stock etc.
- Nitrogen deposition to agricultural land (4.9 kt N_2O): This source originates from re-emission of N deposited to agricultural land as N_2O , and has been distributed using a combination of the spatial distribution of agricultural land from AENEID and the current CEH estimates of N deposition for the UK.
- Legumes (0.9 kt N_2O): Emissions from this source were spread via the spatial distribution of the NARSES category “other crops”, which contains legumes such as peas and beans from the agricultural census.
- Biological fixation of nitrogen (N) by improved grassland (0.6 kt N_2O): This source mainly originates from clover on organic farms, which were assumed to be distributed

evenly among all farms in the UK. The total emissions from this source were distributed using the spatial distribution of agricultural grassland from the AENEID model.

- Histosols (0.5 kt N₂O): Histosols are agricultural soils with a high organic carbon content. A dataset was derived by combining a map of % organic carbon and the CEH landcover map (LCM2000). N₂O emissions from this source were distributed over the resulting map of suitable agricultural land.

The resulting spatially distributed N₂O emission estimates were individually checked for consistency with the BBSRC inventory and then aggregated to the categories listed in Table 1 (above). It should be noted that N₂O emissions from the application of livestock manures are included in the “soils” category, rather than with livestock emissions, as for NH₃.

The resulting spatially distributed emission estimates were then aggregated as follows:

- NH₃: emissions from **livestock manures** and **fertilisers**,
- CH₄: emissions from **enteric fermentation** and **livestock manures**
- N₂O: emissions from **livestock manures** and **soils**

All output data were checked for consistency with the NARSES inventory.

3. RESULTS

3.1. Spatially distributed emissions of NH₃, CH₄ and N₂O for 2009

All UK maps were produced on the Ordnance Survey GB Grid at a resolution of 5 km x 5 km. The units for all GIS datasets submitted are kg ammonia (NH₃), methane (CH₄) and nitrous oxide (N₂O), respectively, per grid square. All spatial datasets were submitted to NWR (Defra Contract AC0112) and to AEAT (for use in the National Atmospheric Emission Inventory (NAEI, see www.naei.org.uk) and the Greenhouse Gas Inventory (GHGI, see www.ghgi.org.uk)). Figures 1, 2 and 3 show the 2009 maps resulting from the spatial modelling of emissions for NH₃, CH₄ and N₂O, respectively (units: kg ha⁻¹ year⁻¹).




Figure 1: UK ammonia emissions from a) livestock manures, b) fertilisers and c) total agriculture (c = a + b) for 2009 (Units: kg NH₃ ha⁻¹ year⁻¹).




Figure 2: UK methane emissions from a) enteric fermentation, b) livestock manure management and c) total livestock (c = a + b) for 2009 (Units: kg CH₄ ha⁻¹ year⁻¹).




Figure 3: UK nitrous oxide emissions from a) soils, b) livestock manure management and c) total agriculture (c = a + b) for 2009 (Units: kg N₂O ha⁻¹ year⁻¹).

3.2. Major changes and Consequences

3.2.1. CHANGES IN EMISSIONS FROM AGRICULTURAL NH₃ SOURCES

Overall, the estimate of NH₃ emissions from UK agriculture increased by 2.3 kt NH₃ between 2008 and 2009, with 229.5 and 231.8 kt NH₃ emitted, respectively (Misselbrook *et al.* 2010). This includes decreases in livestock emissions by 1.9 kt NH₃ and increases in fertiliser emissions by 4.2 kt NH₃.

Revisions to the calculations during the current inventory year include updates to housing data for cattle, pigs and poultry and manure management data for sheep, goats and farmed deer. The decrease in estimated livestock emissions between 2008 and 2009 is mainly due to a decline in livestock numbers for most sectors (apart from pigs and horses). Overall, emissions from cattle showed a small increase due to new housing data, despite a small decrease in the UK cattle population. Conversely, emissions from pigs increased slightly, with housing emissions reduced due to new data, offset by a small increase in the UK pig population.

Fertiliser emissions increased due to an overall increase (by 1.3%) in fertiliser N application (BSFP 2010) as well as an increase in the use of urea (by 28%). As urea is associated with a much larger NH₃ volatilisation rate than other N fertilisers, this resulted in a substantial increase in fertiliser NH₃ emissions between 2008 and 2009.

Table 2: Differences between the 2008 and 2009 inventories for NH₃ emissions from UK agriculture (adapted from Misselbrook *et al.* 2010). Totals may not add up exactly due to rounding.

	2008 kt NH ₃	2009 Kt NH ₃	difference kt NH ₃	difference %
All cattle	132.0	132.5	+0.5	+0.4%
All Sheep, Goats & Deer	10.7	10.1	-0.6	-6%
Pigs	19.8	18.8	-1.0	-5%
All Poultry	30.7	29.5	-1.2	-4%
Horses	4.7	4.8	+0.1	+2%
Livestock total	197.7	195.8	-1.9	-1%
N fertilisers	31.8	36.0	+4.2	+13%
Agriculture total	229.5	231.8	+2.3	+1%

3.2.2. CHANGES IN EMISSIONS FROM AGRICULTURAL CH₄ AND N₂O SOURCES

Between 2008 and 2009, the inventory total for CH₄ emissions from UK livestock has decreased by 14.1 kt CH₄, mainly due to the real general decline in livestock numbers for most sectors. This decrease has been partly offset by new, slightly higher, Tier 2 emission factors for dairy cows in milk.

Table 3: Differences between the 2008 and 2009 inventories for CH₄ emissions from UK agriculture (adapted from Cardenas *et al.* 2008, 2009). Totals may not add up exactly due to rounding.

	2008 kt CH ₄	2009 kt CH ₄	difference kt CH ₄	difference %
Enteric fermentation	731.0	718.7	-12.3	-2%
Livestock manure	135.2	133.3	-1.9	-1%
Total agriculture	866.2	852.1	-14.1	-2%

Overall, emissions of N₂O increased by 5 kt between 2008 and 2009, according to the inventories of Cardenas *et al.* (2009, 2010). In the 2009 inventory, land spreading of sewage

sludge to agricultural land was included in the calculations for the first time, resulting in an additional 1.4 kt N₂O, split over direct and indirect soil emissions. Further main reasons for the increase in N₂O emissions between 2008 and 2009 are the increase in fertiliser N application rates and a revision in the calculation method of indirect emissions from leaching (previously, volatilised N and N₂O emissions were subtracted before calculating leaching rather than from the total N applied). These increases are partly offset by the continuing decrease in livestock numbers.

Table 3: Differences between the 2008 and 2009 inventories for N₂O emissions from UK agriculture (adapted from Cardenas *et al.* 2009, 2010). Totals may not add up exactly due to rounding.

	2008	2009	difference	difference
	kt N ₂ O	kt N ₂ O	kt N ₂ O	%
Direct soil emissions	50.2	50.7	+0.5	+1%
Indirect soil emissions	25.0	29.7	+4.7	+19%
All crops & soils	75.2	80.3	+5.1	+7%
Manure management	6.6	6.4	-0.2	-3%
Agriculture total	81.8	86.8	+5.0	+6%

4. CONCLUSIONS

New ammonia emission maps were derived for the UK (Defra project AC0112), and submitted for inclusion in the 2009 version of the NAEI and GHGI for agriculture for the UK.

Agricultural emission sources were distributed using the CEH/University of Edinburgh AENEID model, which incorporates agricultural census data, landcover data, agricultural practice information (e.g. fertiliser application rates, stocking densities). The latest source strength estimates from the UK NH₃ Emissions Inventory (Misselbrook *et al.* 2010) and the UK greenhouse gas inventory (CH₄ and N₂O; Cardenas *et al.* 2010), produced at NWR, were also applied in AENEID.

ACKNOWLEDGEMENTS

The authors are grateful to Defra, who are funding this work under project AC0112, as a contribution to the National Ammonia Emissions Inventory (NAEI) and Greenhouse Gas Inventory (GHGI), and the Devolved Authorities of the UK for providing agricultural census/survey data.

REFERENCES

- BSFP (2010): *The British Survey of Fertiliser Practice. Fertiliser use on farm crops for crop year 2009*. Defra, London. 100pp.
- Cardenas L., Gilhespy S. and Misselbrook T. (2009) Inventory of UK emissions of methane and nitrous oxide from agricultural sources for the year 2008. MS EXCEL spreadsheet. North Wyke Research, Devon, UK.
- Cardenas L., Gilhespy S. and Misselbrook T. (2010) Inventory of UK emissions of methane and nitrous oxide from agricultural sources for the year 2009. MS EXCEL spreadsheet. North Wyke Research, Devon, UK.
- Dragosits U., Sutton M.A., Place C.J. and Bayley A.A. (1998) Modelling the spatial distribution of agricultural ammonia emissions in the UK. *Environmental Pollution* **102** (S1), 195-203
- Hellsten S., Dragosits U., Place C.J., Vieno M., Dore A.J., Misselbrook T.H., Tang Y.S. and Sutton M.A. (2008) Modelling the spatial distribution of ammonia emissions in the UK. *Environmental Pollution* **154**, 370-379.

- Misselbrook T.H., Chadwick D.R., Gilhespy S.L., Chambers B.J., Smith K.A., Williams J. and Dragosits U. (2009) Inventory of ammonia emissions from UK agriculture 2008. Defra Contract Report (AC0112). North Wyke Research, Devon. 31 pp.
- Misselbrook T.H., Chadwick D.R., Gilhespy S.L., Chambers B.J., Smith K.A., Williams J. and Dragosits U. (2010) Inventory of ammonia emissions from UK agriculture 2009. Defra Contract Report (AC0112). North Wyke Research, Devon. 34 pp.
- Sutton M.A., Dragosits U., Simmons I., Tang Y.S., Hellsten S., Love L., Viero M., Skiba U., di Marco C., Storeton-West R.L., Fowler D., Williams J., North P., Hobbs P. and Misselbrook T. (2006) Monitoring & modelling trace-gas changes following the 2001 outbreak of Foot & Mouth Disease to reduce the uncertainties in agricultural emissions abatement. *Environmental Science & Policy* 9, 407-422.
- Sutton M.A., Dragosits U., Dore A.J., McDonald A.G., Tang Y.S., van Dijk N., Bantock T., Hargreaves K.J., Skiba U., Fowler D., Misselbrook T., Brown L. and Hobbs P. (2004) The potential to use trace-gas changes following the 2001 outbreak of Foot and Mouth Disease in Great Britain to reduce the uncertainties in agricultural emissions abatement. *J. Environ. Science and Policy* 7, p. 177-194.
- Webb J., Anthony S., Fawcett C., Sutton M.A., Hellsten S., Dragosits U., Place C.J., Misselbrook T., Scholefield D., Sneath R. and ApSimon H.M. (2004) National Ammonia Reduction Strategy Evaluation System (NARSES) Final Report to Defra, Project AM0101. ADAS, CEH, University of Edinburgh, IGER, SRI and ICON. 25 pp.

APPENDIX A - NARSES CATEGORIES

Cattle

- 1 Dairy cows & heifers
- 2 Dairy heifers in calf, 2 years +
- 3 Dairy heifers in calf, <2 years
- 4 Beef cows & heifers
- 5 Beef heifers in calf, 2yrs +
- 6 Beef heifers in calf, <2 years
- 7 Bulls >2yrs
- 8 Bulls 1-2yrs
- 9 Other cattle, 2yrs +
- 10 Other cattle, 1-2yrs
- 11 Other cattle, <1yr

Sheep

- 12 Sheep
- 13 Lambs, under 1 year old

Pigs

- 14 Sows in pig & other sows
- 15 Gilts in pig & barren sows
- 16 Gilts > 50kg not yet in pig
- 17 Boars
- 18 Other pigs, 110kg and over
- 19 Other pigs, 80-110kg
- 20 Other pigs, 50-80kg
- 21 Other pigs, 20-50kg
- 22 Other pigs, under 20kg

Poultry

- 23 Layers
- 24 Breeding birds
- 25 Broilers
- 26 Pullets
- 27 Turkeys
- 28 Other poultry

Other livestock

- 29 Horses
- 30 Goats
- 31 Deer

Crops

- 32 Set-aside land
- 33 Wheat
- 34 Winter Barley
- 35 Spring Barley
- 36 Sugar beet
- 37 Oilseed rape
- 38 Potatoes
- 39 Other cereals
- 40 Other root crops
- 41 Other crops
- 42 Vegetables for human consumption
- 43 (Soft) Fruit
- 44 Bulbs, flowers and nursery stock
- 45 Grassland less than 5 years old
- 46 Permanent grassland