

BRITISH GEOLOGICAL SURVEY


King Edward Point

Observatory

Monthly Magnetic Bulletin

August 2012

12/08/KE



**British
Geological Survey**

NATIONAL ENVIRONMENT RESEARCH COUNCIL

KING EDWARD POINT OBSERVATORY MAGNETIC DATA

1. Introduction

King Edward Point observatory was installed by the British Geological Survey (BGS), with the permission of the Government of South Georgia and the South Sandwich Islands (GSGSSI), in collaboration with British Antarctic Survey (BAS), and became operational in February 2011.

This bulletin is published to provide rapid access to the provisional geomagnetic observatory results. The information is freely available for personal, academic, educational and non-commercial research or use. Magnetic observatory data are presented as a series of plots of one-minute, hourly and daily values, followed by tabulations of monthly values. The operation of the observatory and presentation of data are described in the rest of this section.

Enquiries about the data should be addressed to:

Geomagnetism Team
Earth Hazards and Systems
British Geological Survey
Murchison House, West Mains Road
Edinburgh EH9 3LA
Scotland, UK

Tel: +44 (0) 131 667 1000
Fax: +44 (0) 131 650 0265
E-mail: enquiries@bgs.ac.uk
Internet: www.geomag.bgs.ac.uk

2. Position

King Edward Point Observatory, one of the geomagnetic observatories maintained and operated by the British Geological Survey (BGS), is situated on a site adjacent to the Government of South Georgia and the South Sandwich Islands (GSGSSI) research station at King Edward Point, Cumberland East Bay, South Georgia. The observatory co-ordinates are:

Geographic: $54^{\circ} 16' 55''S$ $323^{\circ} 30' 25''E$
Geomagnetic: $45^{\circ} 59' 02''S$ $029^{\circ} 32' 06''E$
Height above mean sea level: 7 m

The geomagnetic co-ordinates are approximations, calculated using the 11th generation International Geomagnetic Reference Field (IGRF) at epoch 2012.5. On-line access to models (including IGRF), charts and navigational data are available at http://www.geomag.bgs.ac.uk/data_service/models_compass/home

3. The Observatory Operation

3.1 GDAS

The observatory operates under the control of the Geomagnetic Data Acquisition System (GDAS), which was developed by BGS staff, installed and became operational in February 2011. The data acquisition software, running on QNX operated computers, controls the data logging and the communications.

There are two sets of sensors used for making magnetic measurements. A tri-axial linear-core fluxgate magnetometer, manufactured by the Danish Meteorological Institute, is used to measure the variations in the horizontal (H) and vertical (Z) components of the field. The third sensor is oriented perpendicular to these, and measures variations, which are proportional to the changes in declination (D). Measurements are made at a rate of 1 Hz.

In addition to the fluxgate sensors there is a proton precession magnetometer (PPM) making measurements of the absolute total field intensity (F) at a rate of 0.1Hz.

The raw unfiltered data are retrieved automatically via Internet connections to the BGS office in Edinburgh in near real-time. The fluxgate data are filtered to produce one-minute values using a 61-point cosine filter and the total field intensity samples are filtered using a 7-point cosine filter. The one-minute values provide input for various data products, available on-line at www.geomag.bgs.ac.uk/data_service/home

3.2 Absolute Observations

The GDAS fluxgate magnetometers accurately measure variations in the components of the geomagnetic field, but not the absolute magnitudes. One set of absolute measurements of the field are made manually once per month. A fluxgate sensor mounted on a theodolite is used to determine D and inclination (I); the GDAS PPM measurements, with a site difference correction applied, are used for F . The absolute observations are used in conjunction with the GDAS variometer measurements to produce a continuous record of the absolute values of the geomagnetic field elements as if they had been measured at the observatory reference pillar.

4. Observatory Results

The data presented in the bulletin are in the form of plots and tabulations described in the following sections.

4.1 Absolute Observations

The absolute observation measurements made during the month are tabulated. Also included are the corresponding baseline values, which are the differences between the absolute measurements and the variometer measurements of D , H and Z (in the sense absolute–variometer). These are also plotted (markers) along with the derived preliminary daily baseline values (line) throughout the year. Daily mean differences between the measured absolute F and the F computed from the baseline corrected H and Z values are plotted in the fourth panel (in the sense measured–derived). The bottom panel shows the daily mean temperature in the fluxgate chamber.

4.2 Summary magnetograms

Small-scale magnetograms are plotted which allow the month's data to be viewed at a glance. They are plotted 16 days to a page and show the one-minute variations in D , H and Z . The scales are shown on the right-hand side of the page. On disturbed days the scales are multiplied by a factor, which is indicated above the panel for that day. The variations are centred on the monthly mean value, shown on the left side of the page.

4.3 Magnetograms

The daily magnetograms are plotted using one-minute values of D , H and Z from the fluxgate sensors, with any gaps filled using back-up data. The magnetograms are plotted to a variable scale; scale bars are shown to the right of each plot. The absolute level (the monthly mean value) is indicated on the left side of the plots.

4.4 Hourly Mean Value Plots

Hourly mean values of D , H and Z for the past 12 months are plotted in 27-day segments corresponding to the Bartels solar rotation number. Magnetic disturbances associated with active regions and/or coronal holes on the Sun may recur after 27 days: the same is true for geomagnetically

quiet intervals. Plotting the data in this way highlights this recurrence. Diurnal variations are also clear in these plots and the amplitude changes throughout the year highlight the seasonal changes. Longer term secular variation is also illustrated.

4.5 Daily and Monthly Mean Values

Daily mean values of D , H , Z and F are plotted throughout the year. In addition, a table of monthly mean values of all the geomagnetic elements is provided. These values depend on accurate specification of the fluxgate sensor baselines. It is anticipated that these provisional values will not be altered by more than a few nT or tenths of arcminutes before being made definitive at the end of the year.

5. Conditions of Use

The data presented in this bulletin are provided for personal, academic, educational, non-commercial research or other non-commercial use and are not for sale or distribution to third parties without written permission from BGS.


Reproduction of any part of this bulletin should be accompanied by the statement: 'Reproduced with the permission of the British Geological Survey ©NERC. All rights Reserved'. Publications making use of the data should include an acknowledgment statement of the form: 'The results presented in this paper rely on the data collected at King Edward Point magnetic observatory, South Georgia operated by the British Geological Survey in co-operation with the British Antarctic Survey and the Government of South Georgia and the South Sandwich Islands (GSGSSI).'

Commercial users can contact the geomagnetism team for information on the range of applications and services offered. Full contact details are available at www.geomag.bgs.ac.uk/contactus/staff

KING EDWARD POINT OBSERVATORY

ABSOLUTE OBSERVATIONS


King Edward Point 2012




King Edward Point

August

2012






Date: 01-08-2012


King Edward Point

Day number: 214



Date: 02-08-2012


Day number: 215



Date: 03-08-2012


King Edward Point

Day number: 216



Date: 04-08-2012


Day number: 217



Date: 05-08-2012


King Edward Point

Day number: 218



Date: 06-08-2012


Day number: 219



Date: 07-08-2012


King Edward Point

Day number: 220



Date: 08-08-2012


Day number: 221



Date: 09-08-2012


King Edward Point

Day number: 222



Date: 10-08-2012


Day number: 223



Date: 11-08-2012


King Edward Point

Day number: 224



Date: 12-08-2012


Day number: 225



Date: 13-08-2012


King Edward Point

Day number: 226



Date: 14-08-2012


Day number: 227



Date: 15-08-2012


King Edward Point

Day number: 228



Date: 16-08-2012


Day number: 229



Date: 17-08-2012


King Edward Point

Day number: 230



Date: 18-08-2012


Day number: 231



Date: 19-08-2012


King Edward Point

Day number: 232



Date: 20-08-2012


Day number: 233



Date: 21-08-2012


King Edward Point

Day number: 234



Date: 22-08-2012


Day number: 235



Date: 23-08-2012


King Edward Point

Day number: 236



Date: 24-08-2012


Day number: 237



Date: 25-08-2012


King Edward Point

Day number: 238



Date: 26-08-2012


Day number: 239



Date: 27-08-2012


King Edward Point

Day number: 240



Date: 28-08-2012


Day number: 241



Date: 29-08-2012


King Edward Point

Day number: 242



Date: 30-08-2012


Day number: 243




Date: 31-08-2012

King Edward Point


Day number: 244




King Edward Point Observatory: Declination (degrees)




King Edward Point Observatory: Horizontal Intensity (nT)



King Edward Point Observatory: Vertical Intensity (nT)



King Edward Point Observatory 2012



Monthly Mean Values for King Edward Point Observatory 2012

Month	<i>D</i>	<i>H</i>	<i>I</i>	<i>X</i>	<i>Y</i>	<i>Z</i>	<i>F</i>
January	-7° 10.4'	15959 nT	-55° 27.4'	15834 nT	-1993 nT	-23182 nT	28144 nT
February	-7° 10.4'	15944 nT	-55° 29.0'	15819 nT	-1991 nT	-23183 nT	28136 nT
March	-7° 11.0'	15919 nT	-55° 31.7'	15794 nT	-1991 nT	-23187 nT	28125 nT
April	-7° 12.2'	15918 nT	-55° 31.7'	15793 nT	-1996 nT	-23185 nT	28124 nT
May	-7° 12.0'	15921 nT	-55° 31.0'	15795 nT	-1995 nT	-23179 nT	28120 nT
June	-7° 12.2'	15913 nT	-55° 31.5'	15787 nT	-1995 nT	-23175 nT	28112 nT
July	-7° 11.8'	15903 nT	-55° 32.4'	15778 nT	-1992 nT	-23173 nT	28105 nT
August	-7° 11.9'	15907 nT	-55° 31.5'	15782 nT	-1993 nT	-23167 nT	28102 nT

Note

- i. The values shown here are provisional.