




Structural considerations suggest that by end-Precambrian times the Charnwood Terrane had become tectonically merged with chemically different volcanic arc rocks (Wrekin Terrane) seen at the Wrekin and Long Mynd. This juxtaposition occurred along a major northerly trending structure, named as the 'Malvern lineament' by Lee *et al.* (1991), which in this region broadly coincides with the faults defining the Knowle and Needwood Triassic basins (Fig. 4). This 'Malvernian' tectonic influence persisted long afterwards, and will be discussed later.

In Charnwood Forest, the Precambrian rocks form a distinctive landscape of rolling hills crowned by craggy knolls, with intervening valleys excavated in the much softer, unconformable Triassic strata. They are divided (Moseley & Ford, 1985) into two lower groups of volcanoclastic rocks, of which the younger Maplewell Group contains primary volcanic components in the form of tuffs and extremely coarse, bouldery fragmental rocks (Fig. 3). The latter are



**Figure 3.** Precambrian volcanic breccia at the 'Bomb Rocks', in the Charnwood Lodge Nature Reserve.



**Figure 2.** Simplified geology of the Trent catchment basin

interpreted as the products of pyroclastic block flows, similar to the recent eruptions on Montserrat in the Caribbean island arc (Carney, 1999; 2000). Their presence is due to the close proximity of local Precambrian volcanic centres, which were situated in the Bardon Hill and Whitwick-Sharpley areas (Carney, 2000). The Caldecote Volcanic Formation of Nuneaton differs in containing tuffaceous beds, up to 60 m thick, characterised by abundant whole or fragmentary quartz and plagioclase crystals (Bridge *et al.*, 1998). As in Charnwood Forest, these rocks are cut by two sets of quartz diorite intrusions. The youngest of these has a distinctive granophyric texture and at Nuneaton it has yielded zircons giving a Late Neoproterozoic U/Pb age of  $603 \pm 2$  Ma (Tucker & Pharaoh, 1991). The stratigraphically higher volcanoclastic strata of the Maplewell Group in Charnwood Forest have, however, given younger U/Pb zircon ages of around 566-560 Ma (Compston *et al.*, 2002). That part of the succession is famous for its fossil fauna (Boynton & Ford, 1995), which includes *Charnia*, a major index fossil of the newly-established Ediacaran Stage – the final division of Precambrian time, which is considered to have ended at c 543 Ma.

By the close of the Precambrian, the various volcanic arc terranes had been tectonically amalgamated to form the elongate microcontinent of Eastern Avalonia (Gibbons & Horák, 1996; Pharaoh & Carney, 2000). The sea then invaded this eroded landmass, depositing a transgressive sedimentary sequence, the fullest development of which is exposed within the Nuneaton inlier (Fig. 2). It commences with the Hartshill Sandstone Formation, deposited in nearshore, tidally influenced environments (Brasier *et al.*, 1978; Bridge *et al.*, 1998), which rests with erosional unconformity on deeply weathered Precambrian rocks (Carney, 1995). Near the top, this formation contains a minor depositional hiatus represented by the Home Farm Member ('*Hyolithes* Limestone'), a

condensed sequence of Lower Cambrian age (Tommotian-Attabanian) hosting the earliest shelly fossils to be found in Britain (Brasier, 1984). Trilobite-bearing mudrocks of the overlying Stockingford Shale Group are at least 700 m thick at Nuneaton where the topmost unit, the Merevale Shale Formation, has fossils indicative of a lowermost Ordovician (Tremadoc) age (Taylor & Rushton, 1971). Remarkably, Tremadocian mudrocks are also encountered in deep boreholes beneath Leicester (Molyneux, 1991), 33 km farther east. As borehole cores indicate that these rocks commonly dip steeply, the most likely explanation for their regional extent, without invoking extraordinary thicknesses, is that the Stockingford Shale Group has been tectonically repeated across faults and folds in a structurally complex basement.

In Charnwood Forest the suggestion of a Lower Cambrian age for the youngest, Brand Group rocks is a recent major development that has followed from the discovery of *Teichichnus*, a Phanerozoic trace fossil, on local headstones carved from quarries in the Swithland Formation (Bland & Goldring, 1995). The Brand Group may thus be a close contemporary of the Stockingford Shale Group, although there is no other faunal evidence to corroborate this.

Further rock sequences of probable early Ordovician (Tremadoc) age to the west and north of the Birmingham conurbation (Fig. 2) are represented by the Barnt Green Volcanic Formation, which includes water-laid tuffs, and the overlying Lickey Quartzite Formation, the latter probably deposited in nearshore, tidally influenced environments (Molyneux in Old *et al.*, 1991; Powell *et al.*, 2000). There are possible links between these isolated exposures and the more complete successions of the Welsh Basin, which includes igneous rocks generated by the subduction of Iapetus oceanic crust beneath Avalonia.

Silurian rocks are preserved only in the far west of the region, their most extensive outcrop being the inlier centred on Walsall, north of Birmingham (Fig. 2). They locally rest unconformably on the Lickey Quartzite Formation and their deposition is attributed to a marine transgression that occurred in Llandovery (Telychian) times (Powell *et al.*, 2000). Silurian strata mainly consist of mudstones interbedded with limestone-dominant units, the most famous of which is the Much Wenlock Formation, exposed at the Wren's Nest Nature Reserve. The overlying mudstones of the Lower Ludlow Shales and Ledbury Formation (Pridoli age) are the youngest preserved elements of this transgressive sequence, the deposition of which would have been terminated, in earliest Devonian times, by the onset of the late Caledonian earth movements.


East of Birmingham, no strata between Tremadoc and late Devonian age have been found. However, igneous intrusions emplaced within the Precambrian, Cambrian and Tremadoc rock sequences have been radiometrically dated to Ordovician age (Caradoc to Ashgill), by Noble *et al.* (1993). Their calc-alkaline

chemistry is compatible with magma generation during subduction of the Iapetus/Tornquist plate system beneath the Midlands, which then formed part of the northwards-migrating Avalonia microcontinent (Pharaoh, 1999). In the Trent region, these igneous intrusions are major sources of hard-rock aggregate and are well known from their exposures in large quarries, such as those currently operating at Croft and Mountsorrel (see article by A. McGrath, this issue). They fall into two chemically and mineralogically distinct 'clans': the Midlands Minor Intrusive Suite, of olivine-bearing lamprophyres and hornblende diorites, is exposed in quarries around Nuneaton (Bridge *et al.*, 1998). Farther east are the granodiorites and quartz-diorites of the Mountsorrel Complex and South Leicestershire Diorites (Le Bas, 1972). The Mountsorrel and South Leicestershire plutonic rocks are chemically comparable with the contemporary Caradocian intrusions of Snowdonia and the Lake District, confirming the extension of the Caledonian magmatic system - the 'concealed Caledonides' of Pharaoh *et al.* (1987b) - down the eastern side of England.

## Forming the structural template: late Caledonian orogenesis

This important tectonic episode is here divided into two parts a) movements that accompanied the late Silurian docking of Avalonia with the Laurentian plate along the Iapetus and Tornquist suture zones, and b) the Acadian orogeny (*sensu* McKerrow *et al.*, 2000), which occurred some 20 Ma later, in Devonian (Emsian) times (Soper & Woodcock, 2003). The deformation created a structural template for much of the basement of southern Britain. In the Trent region, however, its effects have mainly been deciphered by considering the movement histories of the late Caledonian, and in some cases Precambrian, faults that have been rejuvenated through an extensive cover of younger (Upper Palaeozoic to Mesozoic) rocks. The orientations of these fundamental basement structures show significant variation across the Trent catchment, outlining the three tectonic domains shown in the inset of Figure 4 (Smith *et al.*, 2005). The least deformed domain is represented by the Midlands Microcraton, where northerly fault systems were ultimately inherited from the latest Precambrian phase of volcanic arc amalgamation along the 'Malvern lineament', discussed above. Those fault systems are truncated to the west by north-easterly structures of the Iapetus domain, representing Acadian deformation within the Welsh Basin. In the east, they are beheaded by the structures of the Tornquist domain, reflecting displacements within the concealed Caledonides basement of eastern England.

Charnwood Forest provides an important window on local Tornquist deformation, which here was particularly intense and accompanied by upper



**Figure 4.** Triassic outcrop (shaded) and named extensional basins, overlaid with major Variscan faults. For explanation of pre-Triassic rocks (blank areas) see Fig. 2. Inset shows the Precambrian to Palaeozoic tectonic domains that influence the underlying structure of the Trent region. Data from Smith *et al.* (2005). Urban conurbations shown are: B, Birmingham; C, Chesterfield; D, Derby; S, Stoke.

greenschist metamorphism (Merriman & Kemp, 1997). The structures that resulted included north-west trending displacements, such as the Thringstone Fault and the adjacent Charnwood anticline, as well as a west-north-westerly trending, penetrative cleavage fabric (Carney *et al.*, 2001). Argon isotope dating of mica cleavage fabrics suggest that in this part of Britain the cleavage, and associated folding and faulting, was actually a pre-Acadian event, which occurred in late Silurian times, about 425-416 Ma (Unpublished BGS data). The importance of these structures to the subsequent geological evolution of the region cannot be overstated; they exerted a tectonic control that persisted ‘posthumously’ long afterwards, and the Charnwood cleavage direction had a particular influence. It is seen in the orientation of Variscan structures such as the Mackworth-Hoton Fault System and other parallel Tornquist domain faults (Fig. 4), some of which remained periodically active into post-Jurassic times.

### Late Devonian to end-Carboniferous: sedimentary and structural events

The sheer variety of sedimentary rocks produced during this period is a major feature of Trent Basin geology, and an important landscape agency. It also reflects the underlying influence of the Variscan

tectonic cycle that was developing throughout the Carboniferous Period in response to stresses generated by movements within the Variscan suture and associated fold belt, which lay across southern Britain. This orogenic system marked the final stage in the tectonic amalgamation of the Pangaea supercontinent.

Following the fifty million years or so of erosion after the end-Silurian and Acadian uplifts, a change to at least localised subsidence in this region is detected in latest Devonian (Frasnian-Famennian) times, with the accumulation of mainly continental, fluvial deposits. These are only preserved along the western margin of the Nuneaton inlier, as the Oldbury Farm Sandstone Formation (Bridge *et al.*, 1998). Progressive crustal extension subsequently affected the north-east of the region, where Eastern Caledonide, ‘Tornquist’ structures predominate (inset, Fig. 4). Deep, sediment-filled, asymmetric grabens were formed, controlling the syn-rift phase of Carboniferous deposition (Fraser & Gawthorpe, 1990; 2003). Their bounding faults have west-north-westerly orientations suggesting an underlying ‘basement’ structural control that is related to the tectonic ‘grain’ produced by the Charnwood Forest cleavage direction. In the Trent area the deepest of these troughs was the Widmerpool half-graben (or ‘Gulf’), in which about 5.5 km of turbiditic, mud-dominated sediment accumulated during the Early Carboniferous (Dinantian) Period (Carney *et al.*, 2001) along the northern, hangingwall side of the Mackworth-Hoton Fault System (Fig. 4). Coral reefs and carbonate shelves were established in the shallower marine environments created in parts of this tilted block and graben topography (Miller & Grayson, 1982). They belong to the fossiliferous Peak Limestone Group (formerly the Carboniferous Limestone Series or Supergroup), a major landscape-forming sequence exposed within the core of the Pennine Anticline (Fig. 2).

By Namurian times crustal extension had largely ceased, heralding the commencement of the ‘post-rift’ tectonic phase, characterised by regional thermal subsidence (Fraser & Gawthorpe, 1990). Sediments filled in the remaining basins, eventually expanding outwards across the bounding faults. Turbiditic mudstones, siltstones and sandstones of the Edale Shales (now the Bowland Shale Formation) were the initial products of this cycle. They were followed by the southwards encroachment of deltas that deposited the thick, feldspathic sandstones of the Millstone Grit Group. The resistance of these sandstones to erosion, compared with the intervening mudstone beds, produces the spectacular ‘edges’ that dominate the landscape of the Dark Peak (Fig. 5). Subsequently, during the Westphalian Carboniferous Epoch, a vast, featureless, equatorial delta plain occupied the gradually subsiding Pennine Basin (Fig. 8A). The strata deposited, belonging to the Pennine Coal Measures Group, mostly comprise repeated sedimentary cycles (Guion *et al.*, 1995), commencing with dark grey to black, lacustrine or marine



**Figure 5.** Burbage Edge, Derbyshire; typical upland Carboniferous scenery developed on tilted sandstone beds of the Millstone Grit Group. The slope below the sandstone exposure is veneered by a periglacial waste-mantle of Late Devensian age.

mudstones passing upwards into sandy siltstones of overbank or lacustrine delta facies, then into channel sandstones that are commonly surmounted by a seatearth (palaeosol horizon) and coal seam (swamps and mires). This lithological diversity, when combined with later erosion, has produced a strongly featured terrain that is typical of all Coal Measures outcrops.

It is tempting to attribute this essentially quiescent geological interval to the absence of local tectonism; however, ‘growth’ faults have been recognized in Westphalian strata, and to the east of the region, in the Vale of Belvoir area, boreholes show that virtually the whole of the concealed Lower Coal Measures sequence was replaced by low-angled shield volcanoes. From these were erupted ‘within-plate’ - type alkali olivine basalt lavas and peperitic breccias (Kirton, 1984; Carney *et al.*, 2004), a style of volcanism that is commonly associated with fissure activity, implying at least localized extension. Coal Measures deposition was terminated by tectonic movements that ushered in better-drained, alluvial environments in which were deposited the predominantly red-coloured mudstones and sandstones of the Warwickshire Group (formerly ‘Barren Measures’). These Bolsovian to Stephanian sequences are exemplified by the exposures in the South Staffordshire and Warwickshire coalfields, west of Birmingham and Nuneaton respectively (Fig. 2). The reddened, ferruginous palaeosol horizons distinctive to many parts of this group signify deep weathering associated with emergence. Uplift was probably in part fault-controlled, and was a prelude to widespread inversion of the Pennine Basin during the culmination of the Variscan Orogeny in latest Carboniferous to earliest Permian times (Besly, 1988).

The end-Variscan uplifts are most obviously manifested by the fold that formed the limestone-cored Pennine Anticline in the north (Fig. 2). Different structural styles prevailed farther south, however, in

the area occupied by the Midlands Microcraton basement block. There, the Pennine Basin Coal Measures were inverted as a series of synclinal structures, the margins of which are both defined and controlled by faults which, with predominant northerly trends (Fig. 4), reflect the underlying but persistent influence of structures associated with the ‘Malvern lineament’ Precambrian terrane boundary. Intervening between the inverted ‘coalfield synclines’ were uplifted massifs composed of Precambrian and Lower Palaeozoic basement rocks (Fig. 8B). In the Tornquist structural domain the associated faults and fractures acted as conduits for the expulsion of hot, metal-rich basinal fluids that gave rise to the Derbyshire lead and fluorspar mineralisation (e.g. Ford, 2001), and many faults were bordered by inversion anticlines that favoured oil migration and accumulation, with important economic consequences for the East Midlands (Fraser & Gawthorpe, 2003).

### **Permian to end-Triassic: sedimentation and structural development**

Throughout much of the Permian Period, of almost 40 million years duration, the land surface of eastern England was undergoing erosion within an arid, rock-desert located just to the north of the Equator, in the heart of the Pangaea supercontinent. Late in Permian times, however, marginal marine sedimentation occurred as the Southern North Sea Basin encroached across the northern parts of the Trent region. Strata of the Zechstein Group were deposited (Fig. 2), their main representative being the Cadeby Formation (Lower Magnesian Limestone), which forms the escarpment overlooking the Nottinghamshire-Derbyshire coalfield at places such as Bolsover.

By earliest Triassic times, crustal extension associated with the lead-up to Atlantic opening triggered widespread subsidence across the northern margin of Pangaea (Chadwick *et al.*, 1989). In the west of the Trent region this subsidence was greatly accentuated by the development of deep, fault-bounded extensional basins (Fig. 8C). Figure 4 shows that the distribution of these basins was in large part controlled by the rejuvenation of pre-existing Variscan or earlier structures within the Midlands Microcraton, particularly those with inherited northerly, ‘Malvernian’ orientations; for example, the Hopton, Sandon, Burton and Western Boundary faults.

Three phases of sedimentation deposited the Triassic strata that dominate the Trent valley catchment geology (Fig. 6). Initially, major rivers flowed from the south (Warrington & Ivimey-Cook, 1992), exploiting the developing extensional basins and depositing sandstones and conglomerates of the Sherwood Sandstone Group. These strata, which are major aquifers, host the famous caves of Nottingham (Waltham, 1996), and form the many exposures around Nottingham University campus (Howard, 2003). The magnitude of differential subsidence during this

earliest part of the Triassic is exemplified by the 760 m of sandstone present in the Knowle Basin (Powell *et al.*, 2000), as opposed to the 50-150 m thickness range that is typical outside such basins.

Later in the Triassic Period, intensely arid climatic environments characterised deposition of the Mercia Mudstone Group. The widespread distribution and thickness of these strata is attributed to regional crustal downwarping that created a basin in which the Triassic sediments were confined and preserved, allowing them to thicken and eventually to completely cover remaining topographical elements, such as the Precambrian mountainland of Charnwood Forest. In the latter area, the basal Triassic unconformity is spectacularly displayed in Bardon Hill Quarry (Fig. 7), and at Buddon Wood Quarry, Mountsorrel (*this issue*). It is the locus of sporadic mineralization that includes base metals (Pb-Cu-V-Mo) and, more rarely, gold and silver (King, 1968). The red-coloured Mercia Mudstone strata that are so distinctive to the landscapes of the Trent catchment have been compared with loess-type deposits, and latterly (Jefferson *et al.*, 2002) with the modern 'parna' of the south-eastern Australian desert. A complex of mainly continental environments is represented, albeit with occasional marine influences, in which were accumulated thick sequences of red-brown or rarely green-grey mudstone of aeolian to lacustrine origin, punctuated by fluvial episodes that deposited beds of green-grey dolomitic siltstone and sandstone, commonly referred to as 'skerries' because of their relative hardness and resistance to erosion. Higher in the group, evaporitic

conditions are indicated by the incoming of gypsum, of local commercial importance.


The Penarth Group, of Rhaetian (latest Triassic) age represents the final phase of sedimentation. These predominantly argillaceous strata are of marginal marine facies (Swift & Martill, 1999) and represent the initial deposits of a major transgression. They form a small but conspicuous escarpment feature throughout the Trent region.

## Jurassic to Cretaceous: submergence of the Pangaea margin


Marine conditions persisted throughout the Trent region during this 140 million year interval. Jurassic strata of the Lias and Inferior Oolite groups are the main survivors of later Cenozoic erosion. They are disposed within a 'wolds'- type landscape of cuestas and dip-slopes on the eastern margin of the Trent catchment (Fig. 2). In part, their outcrop limit determines the course of the Trent as it approaches the Humber estuary (Fig. 1). The grey mudstones of the Lias Group accumulated in the warm, shallow, sub-tropical sea that was now established across the East Midlands Shelf. The waters deepened with time, leading to better oxygenation and a transition into hemipelagic shelf environments (Weedon, 1986) that supported a diverse fauna of ammonites and bivalves. The Marlstone Rock Formation gives rise to a particularly dramatic escarpment overlooking the Vale of Belvoir, and has been a major source of ironstone and building stone. With its locally prominent cross-bedding, the unit represents one of the shallow water, regressive episodes on the East Midlands Shelf. A later regression is recorded by the Northampton Sand Formation, which is basal to the Inferior Oolite Group (Hallam, 2001).

## Cenozoic uplift and erosion: the modern landscape emerges

Cretaceous strata probably accumulated across the whole of the Trent region during the final stages of the Pangaea shelf sea; however, little is known of their final extent or age because they were largely removed during 60 million years of Cenozoic erosion (Green *et al.*, 2001). The latter study suggested at least two episodes of uplift, which are usually attributed to a combination of tectonic events: the opening of the Atlantic Ocean and compression transmitted from the Alpine Orogeny, which developed from the middle Cenozoic onwards. Recent work suggests that the Cenozoic tectonic regime was asymmetric, involving a principal axis of uplift along the western seaboard of England (Bott & Bott, 2004). This produced eastward tilting, about one degree, or less, on average, that allowed erosion to etch out the scarp and dip-slope topography that characterises the 'wolds' landscape on the Jurassic and Cretaceous outcrops in the far east of



**Figure 6.** Outcrop of mainly Permo-Triassic strata (shaded), with named Triassic basins (see Fig. 4), in relation to the distribution of Trent floodplain deposits.



*Figure 7. Early Triassic palaeovalleys excavated on a mountainous Precambrian landsurface and filled with Mercia Mudstone strata, revealed on the east face of Bardon Hill Quarry, Charnwood Forest.*


the Trent region. The tilting initiated systems of east-flowing trunk streams, which were the main agencies for dissecting and removing the Jurassic and younger sequences (Gibbard & Lewin, 2003). One of these systems was the Bytham (or ‘proto-Soar’) River, the sandy deposits of which indicate that it originally flowed north-eastwards through Leicester (Fig. 1), along the present Soar valley and thence eastwards, along what is now the Wreake valley (Rice, 1991).

The geomorphological process of drainage superimposition, acting on uplifted and tilted Cretaceous strata, explains the eastward-draining river systems proposed by Gibbard and Lewin (2003), and elements of this direction are indeed represented in central England; for example the upper and middle Trent and Dove rivers. This pattern is, however, disrupted in parts where the trunk streams follow northerly courses, as shown in Figure 6. The most obvious control over this deflected drainage pattern is geological structure, with the northerly flowing streams favoured by the former sites of early Triassic rifting; for example, along the Knowle and Hinckley basins, Bratch Graben and parts of the Needwood Basin. This control was most probably facilitated by reactivation of the Triassic structures with Variscan inheritance that originally delineated these basins (Fig. 4). Thus as it was being uplifted and tilted, the Jurassic to Cretaceous cover strata were in places subsiding along fault-controlled troughs (Fig. 8D), the formation of which would have interfered with and locally deflected the easterly-flowing, superimposed river courses. There is abundant evidence in the Trent region

for such post-Triassic fault reactivation (Smith *et al.*, 2005; figure 44), including displacement of the youngest-preserved (Lower Jurassic) strata; for example by the Princethorpe and Whitnash faults north of Warwick (Old *et al.*, 1987).

### **Quaternary drainage development**

The progressive northwards drift of the Eurasian Plate throughout the Cenozoic Period, acting in combination with other factors, culminated in the onset of colder climatic conditions early in the Pleistocene Period. In terms of the deposits left behind in the Trent region, the most significant glaciation occurred during the Anglian Quaternary Stage, about 440 000 years ago, when ice sheets traversed the whole area, depositing locally thick ‘superficial’ sequences of glacial material dominated by till (boulder clay). BGS mapping has shown that the glacial deposits mantle a pre-existing topography, which includes pre-glacial valley systems such as that of the Bytham River (Rice, 1991). Thus the topography revealed following the partial erosional removal of the Anglian deposits is largely that of the Cenozoic landscape. Since ice withdrawal, however, there have been many minor, and some significant, drainage reorganisations to the pre-Anglian



**Figure 8.** Schematic rejuvenation history of a typical late Caledonian structure in Central England, using as an example the record of past movements documented for the Polesworth Fault (Bridge *et al.*, 1998). A, possible role as a growth fault during Coal Measures sedimentation; B, end-Variscan fault rejuvenation and basin inversion; C, extensional relaxation and reversal of previous throw, with probable associated flexuring, to form early Triassic rift basins; D, further fault rejuvenation during Cenozoic regional uplift and initiation of the modern drainage pattern.

systems. For example, the drainage in the Wreake valley (Fig. 1) was reversed to its modern westwards flowing direction.

Regional isostatic rebound and superimposed glacioeustatic fluctuations, dating from the Anglian ice withdrawal, have further influenced not only landscape development, but also the nature and distribution of fluvial deposits throughout the later part of the Pleistocene and into Holocene times. Successive aggradations and incisions over this period have resulted in a 'flight' of five Trent river terraces (e.g. Posnansky, 1960; Carney *et al.*, 2001), each separated

by a 4-7 m vertical interval. The highest and oldest terraces (Eagle Moor and Balderton terraces) have been radiometrically age-dated by Brandon & Sumbler (1991); their outcrops indicate that in pre-Ipswichian ('Wolstonian') times at least, the Trent must have flowed eastwards through the gap in the Jurassic escarpment at Lincoln (Fig. 1). Its subsequent diversion northwards to the Humber estuary may be a result of the younger, Late Devensian glaciation that occurred about 30 000 to 12 000 years ago, the ice front of which would have presented a barrier to drainage around the eastern, northern and western fringes of the Trent Basin. The youngest Trent terrace, the 'floodplain terrace' of Posnansky (1960), represents the valley-confined glacial outwash deposits of this latest cold stage; it is commonly thickly developed beneath the modern alluvium and is a major producer of sand and gravel. In Figures 1 and 6 its outcrops (named as either the Syston or Holme Pierrepont terraces) have been combined with those of the modern alluvium to provide a geology-based model of the Trent catchment in the form of its active floodplain network. This is perhaps a more realistic depiction of a river system than more conventional portrayals that are simply based on distribution of the main river channels and tributary streams. The mid-Pleistocene through to Holocene geomorphological and archaeological development of the Trent valley is summarised by Knight & Howard (2004).

## Conclusions

The protracted geological history of the Trent region has played an important, albeit subtle role in determining its modern physiography. This article has documented the effects of major plate tectonic changes that have underpinned such a role, generating varied rock sequences and perpetuating structures controlling geological and geomorphological processes. The most obvious legacy of this structural evolution is a plethora of 'weak' crustal zones in the form of faults, folds and cleavage belts. Many of these were initiated hundreds of millions of years ago but they have persisted through time as a result of their repeated, 'posthumous' reactivations, a process recognised by Turner (1949). By extrapolation into Cenozoic time, it is likely that inherited structure continued to be an important geomorphological influence, imparting a differential component to uplift and tilting and contributing to the wide variety of rocks, landscapes and drainage patterns seen today in the Trent catchment.

## Acknowledgements

This paper, less minor modifications, was first published in the Open University Geological Society Journal (Symposium Edition: 2006). The author and editor are grateful to the Open University for permission to reproduce large parts of it here. A S Howard and P J Strange are thanked for their comments on an earlier draft. Published with the permission of the Executive Director, British Geological Survey (NERC).

## References

- Besly, B.M. 1988. Palaeogeographic implications of late Westphalian to early Permian red-beds, Central England. 200-221 in *Sedimentation in a Synorogenic Basin Complex: the Upper Carboniferous of Northwest Europe*. Besly B M & Kelling J (editors). (Blackie, Glasgow and London).
- Bland, B.H. & Goldring R. 1995. *Teichichnus* Seilacher 1955 and other trace fossils (Cambrian?) From the Charnian of Central England. *Neues Jb. Palaeont. Abh*, **195**, 5-23.
- Bott, M.H. P. & Bott J.D.J. 2004. The Cenozoic uplift and earthquake belt of mainland Britain as a response to an underlying hot, low-density upper mantle. *Journal of the Geological Society of London*, **161**, 19-31.
- Boynton, H.E. & FORD T.D. 1995. Ediacaran fossils from the Precambrian (Charnian Supergroup) of Charnwood Forest, Leicestershire. *Mercian Geologist*, **13**, 165-183.
- Brasier, M.D. 1984. Microfossils and small shelly fossils from the Lower Cambrian *Hyalolithes* Limestone at Nuneaton, English Midlands. *Geological Magazine*, **121**, 229-253.
- Brasier, M.D., Hewitt, R.A. & Brasier C.J. 1978. On the late Precambrian-early Cambrian Hartshill Formation of Warwickshire. *Geological Magazine*, **115**, 21-36.
- Brandon, A. & Sumbler, M.G. 1991. The Balderton Sand and Gravel: pre-Ipswichian cold stage fluvial deposits near Lincoln, England. *Journal of Quaternary Science*, **6**, 117-138.
- Bridge, D. McC., Carney, J.N., Lawley R.S. & Rushton, A.W.A. 1998. Geology of the country around Coventry and Nuneaton. *Memoir of the British Geological Survey*, Sheet 169 (England and Wales).
- Carney, J.N. 1995. Precambrian and Lower Cambrian rocks of the Nuneaton Inlier: A field excursion to Boon's and Hartshill quarries. *Mercian Geologist*, **13**, 189-198.
- Carney, J.N. 1999. Revisiting the Charnian Supergroup: new advances in understanding old rocks. *Geology Today*, **15**, 221-229.
- Carney, J. N. 2000. Igneous processes within late proterozoic volcanic centres near Whitwick, northwestern Charnwood Forest. *Mercian Geologist*, **15**, 7-28.
- Carney, J.N., Ambrose, K., Brandon, A., Royles, C P, Cornwell J D & Lewis M. A. 2001. Geology of the country between Loughborough, Burton and Derby. *Sheet Description of the British Geological Survey*, 1:50 000 Series Sheet 141 Loughborough (England and Wales).
- Carney, J.N., Ambrose, K., Brandon, A., Lewis, M.A., Royles, C. P. & Sheppard, T. H. 2004. Geology of the country around Melton Mowbray. *Sheet Description of the British Geological Survey*, 1:50 000 Series Sheet 142 Melton Mowbray (England and Wales).
- Chadwick, R.A., Livermore, R.A. & Penn, I.E. 1989. Continental extension in Southern Britain and surrounding areas and its relationship to the opening of the North Atlantic Ocean. *American Association Petroleum Geologists Memoir* **46**, 411-424.
- Compston, W., Wright, A.E. & Toghil, P. 2002. Dating the Late Precambrian volcanicity of England and Wales. *Journal of the Geological Society, London*. **159**, 323-339.
- Ford, T.D. 2001. The geology of the Matlock mines: a review. *Bulletin of the Peak District Mines Historical Society*, **14**, 34 pp
- Fraser, A.J. & Gawthorpe, R.L. 1990. Tectono-stratigraphic development and hydrocarbon habitat of the Carboniferous in northern England. 49-86 in *Tectonic Events Responsible for Britain's Oil and Gas Reserves*. Hardman R F P & Brooks J (editors). *Geological Society Special Publication*, **55**.
- Fraser, A.J. & Gawthorpe, R.L. 2003. *An Atlas of Carboniferous Basin evolution in Northern England*. Geological Society of London Memoir, **28**.
- Gibbard, P.L & Lewin, J. 2003. The history of the major rivers of southern Britain during the Tertiary. *Journal of the Geological Society of London*, **160**, 829-847.
- Gibbons, W. & Horák, J.M. 1996. The evolution of the Neoproterozoic Avalonian subduction system: Evidence from the British Isles, 269-280 in *Avalonian and Related Peri-Gondwana Terranes of the Circum-Atlantic*. Nance, R D, & Thompson, M D (editors). *Geological Society of America Special Paper*, **304**.
- Green, P.F., Thomson, K. & Hudson, J. 2001. Recognition of tectonic events in undeformed regions: contrasting results from the Midland Platform and East Midlands Shelf, Central England. *Journal of the Geological Society of London*, **158**, 59-73.
- Guion, P.D., Fulton, I.M. & Jones N.S. 1995. Sedimentary facies of the coal-bearing Westphalian A and B north of the Wales-Brabant High. *Geological Society of London Special Publication*, **82**, 45 78.
- Hallam, A. 2001. A review of the broad pattern of Jurassic sea-level changes and their possible causes in the light of current knowledge. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **167**, 23-37.
- Howard, A.S. 2003. Excursion 9. The Permo-Triassic rocks of Nottingham. *Geologists' Association Guide* 63, 80-91.
- Jefferson, I.F., Rosenbaum, M.R. & Smalley, I.J. 2002. Mercia Mudstone as a Triassic aeolian desert sediment. *Mercian Geologist*, **15**, 157-162.
- King, R.J. 1968. Mineralization. 112-137 in Sylvester-Bradley P C & Ford T D (Editors). *The Geology of the East Midlands*. (Leicester University Press.)
- Knight, D. & Howard, A.J. 2004. *Trent Valley Landscapes: the archaeology of 500, 000 years of change* (King's Lynn, Norfolk: Heritage Marketing and Publications Ltd), 202pp.
- Kirton, S.R. 1984. Carboniferous volcanicity in England with special reference to the Westphalian of the E. and W. Midlands. *Journal of the Geological Society of London* **141**, 161-170.
- Le Bas, M.J. 1972. Caledonian igneous rocks beneath Central and Eastern England. *Proc. Yorkshire Geological Society*, 39, 71-86.
- Lee, M. K., Pharaoh, T.C. & Soper N J, 1990. Structural trends in central Britain from images of gravity and aeromagnetic fields. *Journal of the Geological Society of London*, **147**, 241-258.
- McKerrow, W.S., Mac Niocaill, C. & Dewey, J.F. 2000. The Caledonian Orogeny redefined. *Journal of the Geological Society, London*, **157**, 1149-1155.
- Merriman R.J. & Kemp, S.J. 1997. Metamorphism of the Charnian Supergroup in the Loughborough District, 1:50K Sheet 141. *British Geological Survey Technical Report WG/97/7*.
- Miller, J. & Grayson, R.F. 1982. The regional context of Waulsortian facies in northern England. 17-33 in *Symposium on the palaeoenvironmental setting and distribution of the Waulsortian Facies*. Bolton K, Lane R H, & Le Mone D V (editors). (El Paso: El Paso Geological Society and the University of Texas).
- Molyneux, S.G. 1991. The contribution of palaeontological data to an understanding of the early Palaeozoic framework of eastern England. *Annales de la Societe de Belgique*, **114**, 93-195.
- Moseley, J. & Ford, T.D. 1985. A stratigraphic revision of the Late Precambrian rocks of the Charnwood Forest, Leicestershire. *Mercian Geologist*, **10**, 1-18.
- Noble, S.R., Tucker, R.D. & Pharaoh, T.C. 1993. Lower Palaeozoic and Precambrian igneous rocks from eastern England, and their bearing on late Ordovician closure of the Tornquist Sea: constraints from U-Pb and Nd isotopes. *Geological Magazine*, **130**, 835-846.
- Old R A, Sumbler M G & Ambrose K, 1987, Geology of the country around Warwick. *Memoir of the British Geological Survey*, Sheet 184 (England and Wales).
- Old, R.A., Hamblin, R.J.O., Ambrose, K. & Warrington, G. 1991. Geology of the country around Redditch. *Memoir of the British Geological Survey*, Sheet 183 (England and Wales).
- Pharaoh, T.C. 1999. Palaeozoic terranes and their lithospheric boundaries within the Trans-European Suture Zone (TESZ): a review. *Tectonophysics*, **314**, 17-41.
- Pharaoh, T.C., Webb, P.C., Thorpe, R.S. & Beckinsale R D, 1987a. Geochemical evidence for the tectonic setting of late Proterozoic volcanic suites in central England. *Geological Society of London Special Publication*, **33**, 541-552.

- Pharaoh, T.C., Merriman, R.J., Webb, P.C. & Beckinsale, R.D. 1987b. The concealed Caledonides of eastern England: preliminary results of a multidisciplinary study. *Proceedings of the Yorkshire Geological Society*, **46**, 355-369.
- Pharaoh, T.C. & Carney, J.N. 2000. Introduction to the Precambrian rocks of England and Wales. 3-17 in: Precambrian Rocks of England and Wales. *Geological Conservation Review Series*, **20**.
- Posnansky, M. 1960. The Pleistocene succession in the middle Trent basin. *Proceedings Geologist's Association*, **71**, 285-311.
- Powell, J.H., Glover, B.W. & Waters, C. N. 2000. Geology of the Birmingham area. *Memoir of the British Geological Survey*, Sheet 168 (England and Wales).
- Rice, R.J. 1991. Distribution and provenance of the Baginton Sand and Gravel in the Wreake valley, northern Leicestershire, England: implications for inter-regional correlation. *Journal of Quaternary Science*, **6**, 39-54.
- Smith, N.J.P., Kirby, G.A. & Pharaoh, T.C. 2005. Structure and evolution of the south-west Pennine Basin and adjacent area. *Subsurface memoir of the British Geological Survey*.
- Soper, N.J. & Woodcock, N.H. 2003. The lost Old Red Sandstone of England and Wales: a record of post-Iapetan flexure of Early Devonian transtension? *Geological Magazine*, **140**, 627-647.
- Swift, A. & Martill, D.M. 1999. Fossils of the Rhaetian Penarth Group. *Paleontological Association Field Guides to Fossils*, **9**.
- Taylor, K. & Rushton, A.W.A. 1971. The pre-Westphalian geology of the Warwickshire Coalfield, with a description of three boreholes in the Merevale area. *Bulletin Geological Survey of Great Britain*, **35** (issued in 1972).
- Tucker, R. D. & Pharaoh, T.C. 1991. U-Pb zircon ages for Late Precambrian igneous rocks in southern Britain. *Journal of the Geological Society of London*, **148**, 435-443.
- Turner, J.S. 1949. The deeper structure of central and northern England. *Proceedings Yorkshire Geological Society*, **44**, 59-88.
- Waltham, T. 1996. *Sandstone Caves of Nottingham*. East Midlands Geological Society, Nottingham, 56pp.
- Warrington, G. & Ivimey-Cook, H.C. 1992. Triassic. 97-106 in Atlas of palaeogeography and lithofacies. Cope J C W, Ingham J K & Rawson P F (editors). *Geological Society of London Memoir*, **13**.
- Weedon, G.P. 1986. Hemipelagic shelf sedimentation and climatic cycles: the basal Jurassic (Blue Lias) of southern Britain. *Earth and Planetary Science Letters*, **76**, 321-335.

J. N. Carney  
 British Geological Survey  
 Keyworth, Nottingham NG12 5GG  
 jnca@bgs.ac.uk