



Contents lists available at ScienceDirect

## Journal of Insect Physiology

journal homepage: [www.elsevier.com/locate/jinsphys](http://www.elsevier.com/locate/jinsphys)

## Corrigendum

## Corrigendum to “The effect of acclimation temperature on thermal activity thresholds in polar terrestrial invertebrates” [J. Insect Biol. 59 (2013) 1057–1064]


Q1 M.J. Everatt <sup>a,\*</sup>, J.S. Bale <sup>a</sup>, P. Convey <sup>b,c</sup>, M.R. Worland <sup>b</sup>, S.A.L. Hayward <sup>a</sup><sup>a</sup>School of Biosciences, University of Birmingham, Edgbaston, Birmingham B15 2TT, UK<sup>b</sup>British Antarctic Survey, Natural Environment Research Council, High Cross, Madingley Road, Cambridge CB3 0ET, UK<sup>c</sup>National Antarctic Research Center, IPS Building, University Malaya, 50603 Kuala Lumpur, Malaysia

The authors regret that a significance asterisk was mistakenly added to Fig. 2 on plot B (*A. antarcticus*, summer acclimatised) and that significance asterisks were accidentally missed out from Fig. 3 in the first version of the paper. These have now been corrected. The authors would like to apologise for any inconvenience caused.


DOI of original article: <http://dx.doi.org/10.1016/j.jinsphys.2013.08.003>

\* Corresponding author. Tel.: +44 789 620 1770.

E-mail address: [mxe746@bham.ac.uk](mailto:mxe746@bham.ac.uk) (M.J. Everatt).



**Fig. 2.** CTmax (A) and heat coma (B) of *C. antarcticus*, *A. antarcticus* and *M. arctica*, following acclimation at 4, 9 and  $-2^{\circ}\text{C}$ , and individuals acclimatised to the Antarctic summer (*C. antarcticus* and *A. antarcticus* only). Means  $\pm$  S.E.M. are presented for approximately 30 individuals. Asterisks indicate a treatment significantly different from  $4^{\circ}\text{C}$  acclimated individuals for each species at  $P < 0.05$  (Kruskal-Wallis test).



**Fig. 3.** Locomotion analysis (distance travelled in 5 min) of *M. arctica* (A), *C. antarcticus* (B) and *A. antarcticus* (C), following acclimation at 4, 9, and -2 °C (0 °C for *M. arctica*). Means ± S.E.M. are presented for approximately 10 individuals. Asterisks indicate a treatment significantly different from 4°C acclimated individuals for each species at  $P < 0.05$  (Kruskal-Wallis test; Tukey's multiple range test). Movement speeds at 25°C were not analysed for *M. arctica* and *C. antarcticus*.