

1 **Can an Anthropocene Series be defined and recognised?**

2

3 JAN ZALASIEWICZ¹, MARK WILLIAMS¹, COLIN N. WATERS²

4

5 *1. Department of Geology, University of Leicester, University Road, Leicester*
6 *LE1 7RH, UK.*

7 *2. British Geological Survey, Keyworth, Nottingham, NG12 5DP, UK*

8

9

10 **Abstract:** We consider the Anthropocene as a physical, chronostratigraphic unit
11 across terrestrial and marine sedimentary facies, from both a present and a far
12 future perspective, provisionally using a ~1950 CE base that approximates with
13 the 'Great Acceleration' of Steffen *et al.* (2007), worldwide sedimentary
14 incorporation of A-bomb-derived radionuclides and light nitrogen isotopes
15 linked to the growth in fertilizer use, and other markers. More or less effective
16 recognition of such a unit today (with annual/decadal resolution) is facies-
17 dependent and variably compromised by the disturbance of stratigraphic
18 superposition that commonly occurs at geologically brief temporal scales, and
19 that particularly affects soils, deep marine deposits and the pre-1950 parts of
20 current urban areas. The Anthropocene, thus, more than any other geological
21 time unit, is locally affected by such blurring of its chronostratigraphic boundary
22 with Holocene strata. Nevertheless, clearly separable representatives of an
23 Anthropocene Series may be found in lakes, land ice, certain river/delta systems,
24 in the widespread dredged parts of shallow marine systems on continental
25 shelves and slopes and in those parts of deepwater systems where human-raftered
26 debris is common. From a far future perspective, the boundary is likely to appear
27 geologically instantaneous and stratigraphically significant.

28

29 Keywords: Anthropocene, chronostratigraphy, sedimentary facies

30

31

32

33 The concept that we may be living in an Anthropocene geological time interval
34 has attracted considerable interest and scrutiny since its latest restatement by
35 Crutzen & Stoermer (2000) and Crutzen (2002) (see also Revkin 1992). These
36 authors effectively regarded the Holocene as having terminated because of the
37 scale and significance of human impact upon the Earth System. In this view, a
38 new and distinct phase of Earth history has already begun, and Crutzen in 2002
39 regarded the beginning of the Industrial Revolution as marking the beginning of
40 profound global change.

41

42 Formalizing this concept within the Geological Time Scale (Zalasiewicz *et al.*
43 2008, 2011, 2012) would result in the creation of an Anthropocene Epoch.
44 Higher levels (e.g. Period, Era) might be considered because of the lack of
45 precedent in Earth history for some of the component stratigraphical signals,
46 such as the lithostratigraphical signal in urban regions (Price *et al.* 2011, Ford *et
al.* this volume) and the scale and character of the biotic change (Barnosky 2008,
47 this volume; Barnosky *et al.* 2011, 2012). Lower hierarchical levels are possible
48 too (e.g. an Anthropocene Age as subdivision of the Holocene Epoch), and this
49 would result in less **modification of** the Geological Time Scale. However, we
50 continue to discuss the Anthropocene in terms of the hierarchical level of Epoch,
51 not least because it brings clear focus on the important scientific question of
52 whether or not the Earth system now lies outside of the ‘Holocene envelope’ of
53 stratigraphically significant environmental conditions (cf. Röckstrom *et al.* 2009;
54 Steffen *et al.* 2011).

56

57

58 **Anthropocene boundary level**

59

60 To carry out the analysis below, we must provisionally select a start date for the
61 Anthropocene. Potential dates for the beginning of this phenomenon have fallen
62 into three categories. Firstly, dates a few to several millennia back within the
63 Holocene (Certini & Scalenghe; Ruddiman 2013) have been suggested, reflecting
64 the growing evidence for widespread, low-intensity human modification of the
65 terrestrial environment (Ellis 2011, Kaplan *et al.* 2011) and, more

66 controversially (Ruddiman 2003, 2013; cf. Elsig *et al.* 2009) resultant release of
67 sufficient greenhouse gases to maintain the Holocene within stable conditions of
68 climate and sea level. Secondly, the beginning of the Anthropocene at ~1800 CE,
69 as originally suggested by Crutzen (2002): that is, around the beginning of the
70 Industrial Revolution when the rapid increase in human numbers, energy use
71 and atmospheric carbon dioxide levels began (Zalasiewicz *et al.* 2008, Fig. 1).
72 And thirdly ~1950 CE, the beginning of the post-war ‘Great Acceleration’ of
73 economic activity (Steffen *et al.* 2007).

74

75 We regard the latter two as the more suitable candidates, because of the clear
76 break between Holocene global stability (or very slow change) and the more
77 rapid and geologically striking changes of the last two centuries (e.g. Zalasiewicz
78 *et al.* 2008, Fig. 1; Steffen *et al.*, 2011, Fig. 1). The Anthropocene does not
79 represent the detectable incoming of human influence (which in any case is
80 clearly diachronous: e.g. Kaplan *et al.* 2011) but major change to the Earth
81 system, that happens to be currently driven by human forcing, but that may
82 geologically soon be more significantly controlled by a number of secondary
83 positive feedbacks such as methane release from permafrost and ice-albedo
84 changes (e.g. Hay 2013, pp. 897-939).

85

86 For the purposes of this exercise we choose the later, ~1950 CE date. This level
87 coincides with changes to lacustrine dynamics and sedimentation worldwide
88 (expressly linked to a potential Holocene/Anthropocene boundary by Wolfe *et*
89 *al.* 2013, and partly reflecting worldwide shift in nitrogen isotopes associated
90 with increase in global fertilizer use: Holtgrieve *et al.* 2011). It also coincides
91 with the beginning of the nuclear age and the spread of artificial radionuclides
92 into contemporary sediments worldwide, and both biotic (Barnosky, this
93 volume, Wilkinson *et al.*, this volume) and physical (Ford *et al.*, this volume)
94 stratigraphical signals that seem to be both stratigraphically sharp (to ~decadal
95 level) and globally widespread. These changes are traceable by scientists living
96 today, and not just by hypothetical ‘far-future’ geologists. They represent a
97 significant and permanent shift in the Earth system, though likely not the

98 greatest changes, that will almost certainly take place in the coming centuries
99 and millennia (Barnosky *et al.* 2011, 2012; New *et al.* 2011).

100

101 Thus, while it is still too early to make a formal recommendation, the ~1950
102 level currently seems to provide sharper stratigraphic definition than the
103 relatively more diffuse and diachronous signals associated with the Industrial
104 Revolution (e.g. the shift in carbon isotopes from the increase in fossil fuel
105 burning: Al-Rousan *et al.* 2004).

106

107 We do not here examine the question of whether the boundary should be defined
108 by a Global Standard Stratigraphic Age (GSSA or more simply a numerical age) or
109 Global Stratigraphic Section and Point (GSSP = 'golden spike'). For practical
110 purposes in current use, we consider that either would be effective. By
111 whichever means defined, this ~1950 CE level might be regarded as
112 stratigraphically challenging, in encompassing (to date) the geologically almost
113 infinitesimally brief interval of ~65 years: over three orders of magnitude
114 shorter than the Holocene and over five orders of magnitude shorter than the
115 average epoch in the Cenozoic (Fig. 1).

116

117

118 **The Anthropocene in geochronology and chronostratigraphy**

119

120 Given current stratigraphic practice, we must consider the Anthropocene as a
121 potential formal stratigraphic unit in not one but two meanings.

122

123 Firstly, it is a potential *geochronological* unit, that is, one of geological time, over
124 which a variety of events have taken place on Earth. An Anthropocene Epoch, as
125 an Earth-based time unit, would (as with the Holocene Epoch and all other
126 geochronological units) hence be used as temporal reference for events in the
127 Earth's deep interior as much as those at the surface.

128

129 Separate geological time scales have been set up for other bodies such as the
130 Moon and Mars (Tanaka & Hartmann 2012) and so an Anthropocene Epoch

131 would be limited to this planet, as holds currently true for other terrestrial
132 geochronological units. The limits on Earth extend from the core to the
133 atmosphere and arguably to the region of space immediately dominated by the
134 Earth's gravitational field, though excluding the Moon, that has a separate
135 stratigraphic scheme (Tanaka & Hartmann 2012). However, we note that it is
136 now beginning to be possible to correlate the Anthropocene across space, in
137 what might be regarded as the first interplanetary stratigraphic marker since the
138 products of the Late Heavy Bombardment of the late Archaean. Infinitesimally
139 smaller in bulk though very much more synchronously distributed, human-
140 projected spacecraft and associated debris have now left physical traces on and
141 around several planets and moons of this Solar System.

142

143 For all past geological units, with the exception of the later part of the Holocene
144 Epoch (that we still, formally, live in) all of our knowledge of the history of the
145 Earth is derived from the rock record. From the beginning of a human written
146 record, this proxy record began to be augmented by human observations of
147 terrestrial events. This human observation has developed, today, to the extent
148 that many terrestrial processes are now routinely monitored, recorded and
149 analysed; this means that geological proxy data of the Anthropocene, being
150 captured within rock currently forming, can now be directly compared with the
151 geological events themselves.

152

153 The Anthropocene, in tandem with other geological units, should also be
154 considered as a material rock unit of chronostratigraphy (commonly referred to
155 as 'time-rock'). Chronostratigraphic units are commonly regarded as the
156 material 'rock' record of geological time, and thus the physical embodiment of
157 (and evidence for) the passage of time. Thus, the Jurassic System comprises all
158 of the rock formed during the Jurassic Period, while the Pleistocene Series is the
159 equivalent rock record of the Pleistocene Epoch. There is hence a hierarchical
160 system of chronostratigraphical terms, exactly parallel to those of
161 geochronology. The Anthropocene, if considered as an Epoch, should also be
162 considered as a Series.

163

164

165 **Chronostratigraphy, scale-dependence and the Anthropocene**

166

167 Not all geologists consider chronostratigraphy to be a necessary and
168 fundamental part of the Geological Time Scale (e.g. Carter 2007; Zalasiewicz *et*
169 *al.* 2004, 2007). In such an interpretation there need not be both a Pleistocene
170 Epoch and a Pleistocene Series, but simply an Epoch, to which the material
171 record is referred descriptively (thus: strata formed during the Pleistocene
172 Epoch, or more simply ‘Pleistocene strata’). Currently, though, most
173 stratigraphers, as represented by voting members of the International
174 Commission on Stratigraphy, prefer to use the dual hierarchy of geochronology +
175 chronostratigraphy (Zalasiewicz *et al.* 2013), and so we here regard
176 consideration of an Anthropocene Series as an integral part of the analysis of the
177 Anthropocene concept.

178

179 Chronostratigraphy in practice only effectively applies to stratified rocks, where
180 superposition applies and hence ‘lower’ equals ‘older’ and ‘upper’ equals
181 ‘younger’ (Zalasiewicz *et al.* 2013). Single hand specimens of igneous and
182 (especially) metamorphic rocks commonly include a number of intermeshing
183 fabrics of distinctly different ages (that can be dated and placed within a
184 geochronological framework), and so ‘upper’ and ‘lower’ have no meaning and
185 the rock itself cannot be regarded as having ‘formed’ at a particular moment in
186 time. Thus, a putative Anthropocene Series encompasses only stratified deposits
187 currently accumulating and not (say) mineral assemblages now crystallizing (i.e.
188 during the Anthropocene Epoch) in the roots of current mountain belts.

189

190 Chronostratigraphy is also scale-dependent (Zalasiewicz *et al.* 2007). That is, on
191 short time-scales, the superpositional fabrics of sedimentary stratification may
192 be disrupted by such processes as bioturbation (in marine deposits especially:
193 Anderson 2001) or by soil-forming processes (Bacon *et al.* 2012), giving
194 disrupted sedimentary fabrics in which temporal information has been mixed or
195 homogenized. This process commonly affects time units of durations of some
196 thousands of years (Anderson 2001) but it can also act over time scales of

197 millions of years (e.g. Bacon et al. 2012) and length scales of kilometres in the
198 case of subsurface sedimentary diapirism (e.g. Shoulders & Cartright 2004).
199
200 For most stratigraphic units in the deep time record, this scale-dependence effect
201 may be neglected, given that currently achievable levels of time resolution are
202 typically measured **in fractions of millions of years**. However, the duration of
203 epochs, both actual (Holocene) and potential (Anthropocene) becomes much
204 shorter towards the present day (Fig. 1). Thus, for Pleistocene and (especially)
205 for Holocene strata the scale dependence effect becomes significant, and for the
206 Anthropocene (where decadal time resolution may reasonably be sought) it
207 becomes an important factor in chronostratigraphic definition.
208
209

210 **Components of an Anthropocene Series**

211
212 Despite the complications noted above, an attempt to define an Anthropocene
213 Series is both part of formal stratigraphic analysis and, independently of this, is
214 useful in helping to understand the Anthropocene phenomenon (formal or
215 informal) as a part of Earth history.
216
217 What might an Anthropocene Series, and its various material stratigraphic
218 components, comprise? We consider the strata that accumulate in a range of
219 geographic settings, from terrestrial (**in the sense of 'land-based'**) to deep
220 marine, and discuss how they might be recognised and characterised. We
221 reiterate that the Anthropocene here is a time boundary, and not a boundary
222 between anthropogenic 'artificial' and 'natural' sedimentary facies. Hence an
223 Anthropocene Series (and, indeed, pre-Anthropocene deposits) will include both
224 of these facies, the boundary between them being diachronous. Nevertheless,
225 the extent of facies diachroneity will vary, both geographically and between
226 different types of stratigraphic signal, and this might offer the possibility of
227 effective discrimination of an Anthropocene Series.
228

229 These strata include a number of proxies for time – not least fossils, a form of
230 evidence that remains key to the subdivision of Phanerozoic strata (Gradstein *et*
231 *al.* 2012) and that has the potential to help characterise an Anthropocene
232 interval (Barnosky, this volume; Wilkinson *et al.*, this volume), when used in
233 combination with other stratigraphic indicators (Waters *et al.*, this volume).

234

235

236 **Terrestrial settings**

237

238 Geologically, the terrestrial realm may be divided into areas of erosion,
239 particularly of older rock, and areas of deposition. The former in stratigraphy
240 may be considered as unconformity surfaces, only to be preserved at the
241 transition between phases of erosion and subsequent sedimentation. Although
242 such erosion surfaces may be studied by techniques such as Terrestrial
243 Cosmogenic Nuclide (TCN) dating (e.g. Gosse & Phillips 2001), we will not
244 consider them further here, except via the indirect record they leave via the
245 sedimentary deposits eroded from them. These may be broadly categorized as
246 the following.

247

248 **Soils:** Soils are perhaps the most widespread terrestrial sedimentary facies,
249 forming on both erosional and depositional surfaces, and having deep time
250 equivalents, palaeosols, when preserved upon depositional surfaces.

251

252 The alteration of soils by anthropogenic activities is widespread, striking and
253 increasingly well documented (Richter, 2007). But, the spread of anthropogenic
254 soils has been strongly diachronous through the Holocene, and reflects the
255 spread of agriculture across the globe (Ellis *et al.* 2012). At present, therefore:
256 which soils are Holocene and which are Anthropocene?

257

258 One approach here has been to take a major phase of soil expansion two
259 thousand years ago across northern Europe (Certini & Scalenghe 2011) and
260 suggest that the base of that may be taken as a ‘golden spike’ to mark the base of
261 the Anthropocene. This is an intriguing and imaginative suggestion, but is not

262 without problems (Gale & Hoare 2012). Firstly, the base of a soil upon older
263 regolith is gradational and cannot capture a boundary with the resolution
264 required for the Anthropocene. Secondly, and more generally, soils exemplify
265 the ‘scale-dependence’ phenomenon noted above, being continually reworked by
266 both natural and anthropogenic processes as long as they are at the Earth’s
267 surface. Hence, it may in some ways be more appropriate to place *all* surface
268 soils in the Anthropocene, because they are continually being modified, even
269 though many of them have fabrics and components which range back for
270 thousands and, in some cases (Bacon *et al.* 2012) for millions of years. This
271 ongoing modification is arguably greatest for agricultural soils, because of the
272 intensive nature of human reworking. Because of the breakdown of
273 superposition, soils are generally problematic to classify chronostratigraphically
274 at the very high levels of temporal resolution required for the Anthropocene.
275 Thirdly, the criteria for definition of a ‘golden spike’ recommends that a section
276 be used in which there is a continuous succession, where observed gaps in
277 deposition are absent or at a minimum. In existing chronostratigraphical units,
278 palaeosols are considered to represent time-gaps and would be avoided as a
279 basis on which to define a chronostratigraphical boundary (Remane *et al.* 1996).

280

281 **Lacustrine deposits:** Lake deposits are perhaps the most straightforward to
282 deal with stratigraphically. Their deposits commonly form ordered strata, which
283 - especially in those lakes with low-oxygen bottom waters – tend not to be
284 seriously disrupted by bioturbation. The resulting high-resolution stratigraphic
285 archives can show a clear signal of the environmental changes that may
286 potentially characterise a ~1950 CE Anthropocene Series base, such as
287 widespread, marked N isotope (Holtgrieve *et al.* 2011) and palaeontological
288 (Wolfe *et al.* 2013) signals in northern lakes far from urban centres, while the
289 incoming of A-bomb test-related radionuclides provides another marker
290 (Appleby 2008; Hancock *et al.* 2011; Yan *et al.* 2002; Hancock, this volume). If it
291 was decided to define the Anthropocene boundary via a physical reference level
292 or GSSP (‘golden spike’) rather than a designated numerical date GSSA (see
293 discussion below), then lake deposits will figure strongly as settings for
294 candidate stratotypes. Lacustrine sediments, though, include anthropogenic

295 signals of other ages too, some markedly diachronous, such as sediment influxes
296 associated with land use changes (Edwards & Whittington 2001).

297

298 **Fluvial deposits:** The human management of rivers, and consequent alteration
299 of their patterns of sedimentation and erosion, has a long history, and the
300 consequent spread of indirect anthropogenic deposits has been marked (e.g.
301 Syvitski & Kettner 2011; Merritts *et al.* 2011; Brown *et al.* 2013), multi-faceted
302 (e.g. the 19th century modification of fluvial sedimentation in north America, as
303 numbers of beavers – and hence beaver dams – fell sharply as a result of hunting:
304 Kramer *et al.* 2011) and strikingly diachronous across the world, and even in
305 part on a regional scale within the UK (Lewin 2012). Indeed, the difficulty of
306 consistently recognising an Anthropocene boundary in modern fluvial deposits
307 was regarded by Autin & Holbrook (2012) as one reason to reject the concept of
308 a formalised Anthropocene.

309

310 However, globally, the rate of fluvial transformation saw significant rises that
311 coincided with the two main inflections in human economic activity at ~1800
312 and at ~1950 (Syvitski & Kettner 2011) both of which are candidate dates for
313 the beginning of the Anthropocene. To what extent these may be generally
314 ‘traceable’ within the sedimentary record seems still to be an open question.
315 Locally, at least, major, distinct Anthropocene bodies of sediment are building up
316 behind the major dams that in recent decades have been constructed on nearly
317 all major rivers of the world (Syvitski & Kettner 2011), with rates of sediment
318 supply commonly increased by deforestation and related processes (Wilkinson,
319 2005). For instance, most sediment that used to be transported down to the Nile
320 Delta is now trapped behind the Aswan Dam (producing a substantial, and
321 rapidly growing Anthropocene sediment body) or held within artificially
322 multiplied (for irrigation) distributaries within a system that has been
323 completely altered by human activity (Stanley 1996).

324

325 **Significant future rise in sea-level** would be expected to result in development of
326 transgressive estuarine to marine deposits in the distal parts of river systems.

327 **However, the interplay of associated changes in precipitation, vegetation and**

328 human forcing would certainly be complex, making patterns of sedimentation
329 hard to predict.

330

331 **Aeolian deposits** Windblown deposits occur both within the major sand seas of
332 the world, such as the ergs of the Sahara desert, as more localised dune fields,
333 such as those associated with coastal areas, and also as far-travelled loess and
334 related deposits. All are sensitive to local climate and to vegetation cover, and
335 human activity, in particular through over-grazing, over-cultivation,
336 unsustainable irrigation techniques and deforestation, which has strongly
337 influenced the generation of loess through desertification, and whose effects
338 include increases in dust flux (Goudie 2009). There is evidence of an increase of
339 a factor of two in background dust loads over the Atlantic since the mid-1960's,
340 the likely product of desertification caused by the doubling of the population in
341 the Sahel region over the past 40 years (Moulin & Chiapello 2006). The extent to
342 which these might translate into an Anthropocene Series boundary is uncertain.
343 It seems likely that within contemporary large, long-lived dune fields, at least,
344 the shifting sands will render a boundary difficult to locate and trace precisely –
345 though in this the Anthropocene is not alone in facing difficulties of
346 chronostratigraphic classification (see below).

347

348 **Glacial deposits:** Glacial deposits are sensitive recorders of changes in ice
349 volume and extent, and many present-day glacial valleys in Europe include
350 terminal moraines reflecting the greater extent of ice during the Little Ice Age of
351 the 16th to mid19th centuries (Mann 2002). Similarly, the shrinking of most
352 mountain glaciers since the 1850's, with regional variations in both retreat and
353 advance during the mid-20th Century and large-scale retreats since the 1980's
354 (IPCC 2001, figure 2.18.), linked to global temperature increases, has exposed
355 morainic deposits that may be clearly identified and mapped as of Anthropocene
356 age, particularly where detailed cartographic and photographic records occur of
357 glacier extents earlier in the 20th century (e.g. Kulkarni *et al.* 2007). Associated
358 deposits include those laid down catastrophically by dam-bursts, as increased
359 volumes of meltwater have accumulated behind and destabilized morainic dams
360 and wasting morainic ice cores (Nayar 2009).

361

362 **Ice:** This is also a terrestrial sedimentary deposit that is found on all the
363 continents (except in Australia, and probably not for much longer in Africa,
364 where it is represented only by rapidly-thawing Kilimanjaro). Ice sheets record
365 snow layers extending back many thousands of years, and encapsulating (in the
366 Arctic and Antarctic) the entire interval of human history, including levels that
367 can be identified for 1800 and 1950, and which provide data on rising CO₂
368 intervals. Snow layers record human pollutants from the atmosphere back to
369 classical times (e.g. lead aerosols derived from Roman smelting). Following this,
370 there is a succession of recorded events that might provide geochemical criteria
371 to identify either a ~1800 CE or a ~1950 CE level. This includes the CO₂ levels
372 preserved in air pockets (though this is compromised by the 'lock-in' time for air
373 post-dating the deposition of the snow). However, events such as the
374 appearance of nitrogen derived from the Haber-Bosch process (cf. Holtgrieve *et*
375 *al.* 2011), the change in lead isotopes reflecting the use and then abandonment of
376 lead additives in petrol (Bollhöfer & Rosman 2000), and the incorporation of
377 artificial radionuclides provide useful global stratigraphic markers. The range of
378 palaeoenvironmental proxies recorded in this medium and the annual resolution
379 make selection of a GSSP within a snow/ice core a potential option, as for the
380 Pleistocene/Holocene boundary (cf. Walker *et al.* 2009).

381

382 **Artificial deposits** The transformation of primary raw materials (sand and
383 gravel, limestone, mudrock, metal ores) into the fabric of urban areas represents
384 the creation of a novel and substantial type of stratum in which the buildings
385 themselves and the associated landscape changes (the latter mapped as various
386 types of Artificial Ground on British Geological Survey maps, for instance: Price
387 *et al.* 2011; Ford *et al.* this volume) provide something that combines features of
388 a lithostratigraphic unit and of an extraordinarily large trace fossil system.

389

390 The resulting deposit is clearly anthropogenic but, because towns and cities have
391 been a feature of human civilization since the Epi-palaeolithic (Mesolithic) about
392 9000 BC (see Edgeworth this volume), also clearly diachronous. We may discuss
393 two features of relevance here.

394

395 Firstly, the extraordinary post-war growth of cities and megacities allows, by
396 simply mapping the historical growth of urban areas, a distinction between post-
397 1950 CE artificial deposits and those that predate them (Fig. 2). Prior to the
398 1950's, large cities tended to be located close to natural resources or be suitable
399 coastal locations for the import/export of these resources. The post-1950s
400 evolution of megacities has relied upon the contained population of the megacity
401 to be the key resource, and these cities have been a centre for the inward influx
402 of natural resources sourced from rural areas and transported to the cities to
403 fuel industry and construction. This change can be seen as a product of
404 improvement of transport networks and greater efficiencies in the mass-
405 transport of bulk materials during the late 20th century (Haff, this volume; see
406 also Williams *et al.*, this volume). This creation of laterally continuous but
407 temporally distinct deposits may be compared with, say, those created naturally
408 during the progradation (outgrowth) of a delta system.

409

410 Secondly, even within the older parts of existing cities, the continuous
411 replacement of the urban fabric, both above and below ground, means that these
412 artificial deposits comprise complex mixtures of pre-Anthropocene and
413 Anthropocene rocks and minerals (and, locally, indeed fossils). The presence of
414 novel materials and minerals in both direct and to a lesser extent indirect
415 anthropogenic deposits (Ford *et al.* this volume, Zalasiewicz *et al.* this volume b)
416 provides an approach to dating these deposits to decadal level, a resolution far
417 beyond that applicable for previous epochs. This is a rather coarser-grained
418 equivalent of the situation noted above with soils, and again underscores the
419 awkwardness of chronostratigraphy in dealing with short time scales and
420 complex sedimentary processes and geometries. It is only towns and cities
421 abandoned pre-1950 that may be said to comprise wholly pre-Anthropocene
422 representatives of this deposit type.

423

424 Below ground, artificial ground locally deeply extends into underlying strata via
425 the many mineshafts and boreholes sunk to extract resources, with considerable
426 'halo' effects via such as hydrocarbon extraction (and now, injection of fluids and

427 sand for shale gas extraction). Geologically, this is something of a hybrid,
428 combining features of burrowing, albeit on an enormous scale, with those of
429 intrusive bodies, showing cross-cutting relationships, and even of diagenetic
430 alteration. Neither of the last two phenomena are generally classified within
431 chronostratigraphic units (as they do not show superpositional relationships), as
432 their history may be protracted and only related in general terms to processes
433 acting at the Earth's surface (Ford *et al.* this volume). The subsurface
434 anthropogenic phenomena, by contrast, are very much related to surface
435 activities [and can also impact on the surface, as for instance with aquaculture-
436 related subsidence on the Yellow River delta in China now reaches 250
437 mm/year: Higgins *et al.* 2013]. They clearly form a pronounced and temporally
438 constrained event, given the post-war surge in drilling and mining (Ford *et al.*
439 this volume).

440

441

442 **Marine settings**

443

444 **Coastal systems** These systems include beaches, tidal flats and deltas.
445 Throughout much of the latter half of the Holocene, these have been commonly
446 progradational, as sediment eroded from the land, has accumulated around a
447 coastline more or less fixed as sea level stabilized following its post-glacial rise.
448 Where sediment has built up and built out in this way, then distinct stratal
449 packets that relate to industrialization and land use change have been
450 recognised and suggested as Anthropocene markers (e.g. Poirier *et al.* 2011).
451 Some are distinctive through their content of heavy metals, organic chemicals
452 and so on (e.g. Allen 1988; Marshall *et al.* 2007; Vane *et al.* 2011; Galuszka *et al.*,
453 this volume), with eutrophication of coastal environments due to influx of excess
454 nitrogen, and these may also be used to help identify an Anthropocene/Holocene
455 boundary. Globally, the overall facies changes are diachronous, but within them
456 some signals (such as distinct chemical markers related to particular industrial
457 processes: Kruse 1999) may provide more or less synchronous marker levels.

458

459 Within the last couple of centuries – and particularly the last several decades,
460 many coastal systems have seen large-scale change that is clearly relevant to the
461 historical characterisation of the Anthropocene, but that complicates the simple
462 progradational picture. For instance, as rivers have been dammed, sediment is
463 temporally stored behind the dams (see above) and does not nourish growing
464 deltas, some of which have as a consequence shrunk back (e.g. Nile, Mississippi
465 etc – Törnqvist *et al.* 2008). Related phenomena include the draining of coastal
466 wetlands for farmland, resulting in the large-scale loss of such strata as surface
467 peat deposits through desiccation, deflation and oxidation. For instance, some
468 2000 km² of peat up to 4 m thick in the English Fenland alone has disappeared
469 since the 18th century, resulting in the exposure of the underlying geology, now
470 itself compacted and oxidised (Smith *et al.* 2011). Attempts at stabilization of
471 coastal erosion rates through construction of coastal defences produces artificial
472 deposits, while impacting upon sediment flux and erosion rates adjacent to
473 protected regions. The Anthropocene boundary here approximates to a regional
474 sedimentary hiatus and disconformity - likely to be buried beneath new coastal
475 sedimentary deposits, a century or two hence, as only a geologically trivial sea
476 level rise will suffice to trigger marine transgression across such areas.

477

478 The expression of the Anthropocene in the environmentally sensitive coastal
479 systems, therefore, represents a diverse patchwork of deposits and lacunae that
480 reflect local interplays of natural and anthropogenic forces.

481

482 **Shelf/slope marine systems** Human impact on open marine systems has in
483 general substantially lagged those on land. The marine fisheries in northern
484 Europe began in earnest in Medieval times, perhaps as a result of technological
485 improvements (e.g. effective drift nets) and their spread across the world has
486 been charted by Roberts (2007). The concomitant, diachronous decline in fish
487 stocks through overfishing changed the structure of marine ecosystems, though
488 impacts on the kind of organisms (e.g. foraminifera, dinoflagellates – much lower
489 in the food chain) used in biostratigraphy have likely been small, even with the
490 dramatic fish declines reported (e.g. Myers & Worm 2003).

491

492 More profound physical and chemical impacts on recent marine strata are
493 associated with the industrial age, from ~1800 CE. The greatest physical impact
494 on sediments has been the physical disruption caused by sea bottom trawling.
495 This is not a modern technique: the 14th century saw a petition to regulate the
496 use of the ‘wondyrechaun’ – essentially a wooden beam trawl used in shallow
497 coastal waters (Roberts 2007) – but open sea trawling came with steam-
498 powered ships, and has continued to expand markedly in recent decades, moving
499 into slope settings in waters approaching a kilometre deep.

500

501 Sea bottom trawling now affects some 15 million km² each year (Gattuso *et al.*
502 2009) – representing most of the world’s continental shelf area and also
503 including significant areas of deepwater slope (Puig *et al.* 2012) and seamount
504 surface. The process in effect ploughs the sea floor, producing a coarsening-
505 upwards sedimentary signature (Palanques *et al.* 2001; M. Coughlan, pers.
506 comm.), with mud swept up into an expanded nepheloid layer and transported
507 more distally, and nutrients redistributed (Dounas *et al.* 2007). Benthic
508 assemblages are altered (Malakoff, 2002) and some sensitive ones (e.g.
509 deepwater coral systems) effectively destroyed (Sheppard 2006). Topographic
510 effects may be substantial, with evident smoothing of topographic contours (Puig
511 *et al.* 2012).

512

513 More recent extension of ‘Worked Ground’ into a marine setting can be seen with
514 increased extraction of mineral resources including hydrocarbons and
515 aggregates. It is only since the 1940’s that technology and economics has made
516 offshore extraction of hydrocarbons feasible and it has grown to the point where
517 it currently accounts for about 30% of total global output. Aggregate extraction
518 significantly modifies the marine landscape, causes habitat modification and
519 impacts on benthic communities both within, and downcurrent, of extraction
520 sites and can significantly change sediment fluxes, potentially starving supplies
521 of sand to coastal areas. Similar concerns are being raised about offshore wind
522 turbine construction, an even more recent and expanding innovation.

523

524 Within tropical waters, bleaching of coral reefs in response to rising sea
525 temperatures, in addition to other stressors, such as increased turbidity of
526 marine waters due to runoff, the fishing process of dynamiting reefs and
527 ultimately decreasing ocean pH (Tyrrell 2011) may lead to the extinction of
528 whole reef systems, resulting in a drowned reef horizon.

529

530 In aggrading sedimentary systems, the resultant facies should have considerable
531 preservation potential. It is of limited diachroneity, given the marked post-1950
532 expansion of many of the processes involved.

533

534 **Deep sea:** This is usually considered as those areas where water is >200 m deep
535 (i.e. largely below wave base and off the continental shelf edge) and might be
536 simplified into two main systems: the clastic wedges of turbidite fans and
537 contourite drifts that fringe the continental masses, and the slowly accumulating
538 deep-sea oozes that lie beyond. Both systems have been and continue to be
539 affected by physical disturbance (e.g. by trawling, offshore mineral extraction),
540 by input of particulate material ('litter') varying from micron to metre scale in
541 size (and locally indeed larger, in the case of shipwrecks), by chemical
542 contamination with both organic and inorganic substances, by effects associated
543 with atmospheric CO₂ increase and warming (such as variations in pH and
544 dissolved oxygen content) and by biological changes driven by all of the above
545 processes, either directly or indirectly. The extent of these effects – all of which
546 can affect the nature of sediments being deposited - have been qualitatively
547 described but not yet rigorously mapped (Ramirez-Llodra *et al.* 2011). The
548 stratigraphic signal is patchy but locally may be striking. As with the effects of
549 urbanization, local signals go back millennia. Major expansions of activity and
550 hence extent of stratigraphic imprint were associated with the Industrial
551 Revolution at ~1800 CE and with the ongoing 'Great Acceleration' that started
552 ~1950 CE.

553

554 The accumulation of litter – material dropped overboard - has reached the level
555 where it rivals the extent of ice rafted debris (IRD) in scale (Ramirez-Llodra *et al.*
556 2011), and is now seen in most surveys of the sea floor, where it is easily distinct

557 from the surrounding (mostly very fine-grained) sediment. We suggest hence
558 terming this material, sedimentologically, as human-rafted debris (HRD) to help
559 characterize a deep-water facies of a putative Anthropocene Series. Given
560 technical progress, it shows the kind of extremely high-resolution
561 'biostratigraphy' of human artefacts and products also seen on land (Ford *et al.*,
562 this volume). Hence, spreads of clinker (combustion products from the coal that
563 powered steam-ships) were universally dumped on the sea floor in the period
564 ~1800 CE to ~1950 CE, - now colonised by a specific biota - might be regarded
565 as immediately pre-Anthropocene in our provisional definition, while those with
566 plastics, aluminium and other such more modern materials largely date from
567 after 1950 CE (Ramirez-Llodra *et al.* 2011). In the distal, naturally slow-
568 accumulating parts of the sea floor, such HRD from different centuries will in
569 effect fall within and contribute to the same physical layer.

570

571 More broadly, within the clastic wedges, the pattern of turbidite/contourite
572 deposition seems not yet to have been substantially affected by human activity; it
573 is not clear that changes in sediment supply caused by large-scale anthropogenic
574 modification of river systems (e.g. Syvitski & Kettner 2011) have yet filtered
575 down to cause substantial change to deep-sea clastic systems, though we regard
576 significant longer-term change as likely (see below). However, local effects
577 include the triggering of turbidity currents (that may also rework HRD into
578 concentrations: Ramirez-Llodra *et al.* 2011) by bottom trawling (Puig *et al.*
579 2012). In the longer-term, clastic shut-off caused by sea-level rise may be
580 envisaged.

581

582 The slowly accumulating deep-ocean oozes beyond will, in addition to such
583 accumulations, be influenced by anthropogenic change, via such signals as a
584 lighter carbon isotopic content in foraminifera shells, from the burning of fossil
585 fuels. Additional chemical signals such as those from anthropogenic organic
586 pollutants or artificial radionuclides are rapidly (e.g. Robison *et al.* 2005) if
587 unevenly (Buesseler *et al.* 2007) transported to the sea via aggregated sinking
588 planktonic debris. However, the very slow accumulation rate over most of this
589 realm means that this material is thoroughly intermixed, by bioturbation, with

590 pre-Anthropocene sediment, precluding recognition at such scale of a distinct
591 Anthropocene Series. Only in regions of significantly more rapid deposition (e.g.
592 Al-Rousan *et al.* 2004) does such a potential Series emerge as a distinct entity
593 with coherent upper (sedimenting) and lower surfaces. However, potential
594 changes to ocean chemistry may result in more extensive anoxia, with eutrophic
595 bottom conditions limiting bioturbation, and changes to the elevation of the
596 Calcite Compensation Depth in response to reduced oceanic pH (Tyrrell 2011),
597 producing a carbonate dissolution layer. In addition, the types of deep-sea
598 mineral extraction planned, if put into practice (of manganese nodules, for
599 instance), will cause widespread and distinct physical and biological
600 modification.

601

602

603 **Duration of the Anthropocene: the long-term perspective**

604

605 The complexities of diachronous event and process boundaries and scale
606 dependence effects, visible today, will largely or wholly disappear in any
607 consideration of far future perspective.

608

609 We do not consider the Anthropocene as a short transitional phase to some kind
610 of post-Anthropocene interval, even were there to be a catastrophic decrease in
611 the global human population in the near future. Rather, we consider that the
612 future course of geological evolution, with both natural and human feedbacks,
613 will inevitably be shaped by the anthropogenic perturbations that have taken
614 place to date. Thus the Anthropocene has only just begun and will play out over
615 geological rather than human timescales. The Toarcian and Paleocene-Eocene
616 Thermal Maximum (PETM) (Cohen *et al.* 2007; Zachos *et al.* 2005) events may be
617 regarded as comparable, with an initial perturbation of the carbon cycle,
618 amplification by natural feedbacks including massive carbon release from
619 ground to air, modulated by astronomical pacing (Kemp *et al.* 2005), and slow
620 recovery over the order of 0.1-0.2 Myr. Although each of these events in detail
621 represents a succession of distinct phases, each may also be (and are, in practice)
622 regarded as a whole.

623

624 In detail, the Anthropocene departs from the Toarcian and PETM models in a
625 number of ways. It is an incipient 'hyperthermal' in an icehouse rather than
626 greenhouse world, and so the ultimate sea level rise (barely begun) should give a
627 stronger transgressive signal (Rahmstorf, 2007) than that in an essentially ice-
628 free world. Indeed, if the glacial-interglacial cycle is significantly perturbed
629 (Tyrrell 2011) with ice loss that exceeds Quaternary norms, then the geologically
630 rapid transgression that followed the collapse of the end-Ordovician glaciation
631 (Brenchley et al. 1994) might be considered as a closer analogue (Zalasiewicz &
632 Williams, in press).

633

634 The Anthropocene also has a biotic pattern where perturbations (habitat
635 clearance, predation, trans-global rather than local species invasions) are not
636 simply forced by climate and ocean chemistry; as with previous biotic
637 revolutions, these will be geologically long-lasting quantitatively (i.e. regarding
638 diversity measures) and effectively permanent qualitatively (with new lineages
639 arising from survivors and invaders) (see also Barnosky, this volume). This
640 pattern is also unique in modification by unpredictable but likely important
641 feedbacks, both planned and unplanned, within the perturbatory human system
642 (Kellie-Smith & Cox 2011).

643

644 One might compare the scale of effects with those recently proposed (the 8.2 and
645 4.2 kyr events) to subdivide the Holocene Epoch into Ages (Walker et al. 2012;
646 see also Gibbard & Walker, this volume). As regards global climate, current
647 effects (a <1 degree C global temperature rise since the beginning of the 20th
648 century) might not be regarded as yet comparing with the 8.2 and 4.2 kyr events
649 in magnitude. However, near-future temperature rises are projected to
650 considerably exceed these (IPCC 2001, 2007), given the unprecedented and
651 ongoing rise in greenhouse gas levels. Other signals, though (lithostratigraphic,
652 biostratigraphic, chemostratigraphic) are already pronounced and, as an
653 ensemble, have no parallel in Earth's stratigraphic history. Debate over the
654 current formal significance of the Anthropocene will need to assess the relevant
655 importance of all the relevant signals, and this is not a trivial task.

656

657 Nevertheless, the unprecedented rate of change in its early stages (within a small
658 part of a single interglacial phase) means that the lower boundary to deposits of
659 Anthropocene facies will appear synchronous globally. One may develop the
660 'superinterglacial' concept of Broecker (1987) by envisaging a variety of
661 stratigraphic signals that vary from 'event beds' (e.g. the urban lithostratigraphic
662 signal), to longer-lasting perturbations of chemical cycles and related effects on
663 global temperature and sea level, to the effectively permanent changes to the
664 course of the Earth's biotic evolution.

665

666

667 **Discussion**

668

669 How might the Anthropocene be characterized? Clearly, it is not simply by the
670 appearance of anthropogenic signals in the stratigraphic record, as these are
671 diachronous, locally dating back to earlier parts of the Holocene and indeed into
672 pre-Holocene deposits. Such early records have been used in favour of an 'early
673 Anthropocene' hypothesis that encompasses much of the Holocene (e.g.
674 Ruddiman 2003, 2013) and also in criticism of the attempt to define an
675 Anthropocene unit in stratigraphy at all (e.g. Gale & Hoare 2012; Gibbard &
676 Walker, this volume).

677

678 The key question seems to be whether the present-day Earth system now has
679 been changed (by whatever agent) sufficiently in scale and permanence to justify
680 a new geological time interval. If that is the case, one also may accept that the
681 change from a putative pre-Anthropocene to an Anthropocene state has taken
682 place non-instantaneously and diachronously. Most changes to the Earth system
683 in our planet's history have been neither instantaneous nor globally synchronous
684 (e.g. Williams *et al.* this volume), and most established geological time
685 boundaries have been compromises – generally vigorously debated - of one sort
686 or another.

687

688 The task then becomes one of finding the most effective – or, if one prefers, the
689 least worst – criteria for defining a boundary. Then, one has to decide whether a
690 boundary so defined can function effectively to define both a unit of time (an
691 Anthropocene Epoch) and a body of strata (an Anthropocene strata). This is the
692 question we examine here. We note that the further test for a formal
693 Anthropocene – its use to both geological and arguably wider (Vidas, 2011;
694 Nature, 2012; Zalasiewicz 2013) communities – falls outside the scope of this
695 paper, as does the question – see above and Zalasiewicz *et al.* 2008, 2011, 2012,
696 Wolfe *et al.* 2013 – over whether a boundary, if agreed, is best defined by GSSP
697 ('golden spike') or GSSA (numerical date).

698

699 It is clear that the material record of a putative Anthropocene Series, even
700 considered with a ~1950 CE boundary, is locally distinctive and substantial – a
701 feature reflecting the globally enhanced rates of erosion and sedimentation
702 caused by humans (Hooke, 2000; Wilkinson 2005; Syvitski & Kettner 2011;
703 Price *et al.* 2011). It is in many places also effectively distinguishable from pre-
704 Anthropocene strata, on a decadal or even annual scale of resolution.

705

706 Elsewhere, though, the distinction of Anthropocene from pre-Anthropocene
707 strata is less obvious. This may be because there are no significant markers or
708 facies changes (as in desert dune strata, for instance). Or, it might reflect
709 widespread irresolvable mixing of Anthropocene and pre-Anthropocene strata,
710 through non-human bioturbation and other mixing processes (as in the deep
711 ocean). Or there may have been protracted, complex human reworking of the
712 ground (as in long-inhabited cities). Such phenomena prevent the clear,
713 unambiguous and consistent delineation of a laterally continuous 'Anthropocene
714 Series'. We may discuss them in turn as regards comparison with older
715 chronostratigraphic units.

716

717 The local inability to unambiguously assign particular units of strata to
718 chronostratigraphic units is a problem as old as is geology. One may take the
719 case of the 'Permo-Triassic', long used as a descriptive bucket label given the
720 difficulty of locating a boundary between Permian and Triassic deposits in 'red

721 bed' deposits that lack fossils, even if it is as sharp and catastrophically founded
722 as that between the Permian and Triassic systems (and between the Palaeozoic
723 and Mesozoic Erathems). Even in less stratigraphically opaque strata,
724 chronostratigraphic boundaries, away from the reference 'golden spike' section,
725 can rarely be located within an error bar of less than a few hundred thousand
726 years (Zalasiewicz *et al.* 2013). Most stratigraphic research is based upon the
727 most informative and correlatable sections, but between these there are many
728 stratal units within which major chronostratigraphic boundaries are located only
729 approximately.

730

731 Similar uncertainty will certainly apply to an 'Anthropocene Series', with
732 boundaries (now being placed at a decadal/annual scale) being effectively
733 locatable in some places and more uncertainly placed in others. Hence, at least
734 qualitatively, the Anthropocene shares the correlation problems attached to
735 chronostratigraphic units generally, and it is not yet clear whether it possesses
736 these kinds of uncertainties in greater measure than do the established units of
737 the Geological Time Scale.

738

739 The problem of the disruption of superposition is rather different. This arises in
740 part out of the exceedingly short timescale of the Anthropocene (to date) and in
741 part out of complex, intermingled sedimentary geometries commonly created by
742 human activity, where clear principles of superposition cannot be applied. This
743 creates situations that archaeologists, for instance, are more familiar with, in
744 discriminating numerous successive historical events within geometrically
745 complex deposits (Edgeworth this volume) and on palimpsest surfaces (where
746 the evidence from different phases of human history is preserved upon
747 essentially two-dimensional surfaces).

748

749 The practice of basing chronostratigraphic subdivision upon the principle of
750 superposition reflects the tendency on Earth for thick successions of strata to
751 have built up, virtually since the origin of the planet. For most of the geological
752 record it is an effective means to build and operate the geological time scale, and
753 in older rocks, where stratigraphic uncertainties are measured in millions of

754 years, then superpositional blurring through bioturbation and allied processes
755 may be regarded as negligible. In such circumstances, chronostratigraphy and
756 geochronology have operated in parallel, in their long-established 'dual
757 hierarchy'.

758

759 However, at brief geological time scales and/or when extremely fine temporal
760 resolution is sought, disruption of superpositional relationships may become a
761 practical, rather than theoretical problem. This is already the case in the
762 discrimination of high-resolution climate histories from deep sea floor deposits,
763 where those strata with the highest sedimentation rates (and therefore least
764 prone to bioturbational mixing) are actively sought. This phenomenon is, hence,
765 most acutely expressed in the Anthropocene, with its extremely short timescale
766 exacerbated by its peculiarly human-made complex stratigraphic geometries. It might
767 be regarded as a problem as much inherent of chronostratigraphic practice as it is
768 of the Anthropocene.

769

770 Nevertheless, despite the complicating effects of these various processes, we
771 propose that a reasonably consistent Holocene-Anthropocene boundary placed
772 at ~1950 CE might be effectively traceable over large areas in both marine and
773 non-marine settings. Attempts to consistently trace and delineate such a unit
774 would reveal the extent to which this proposal is true. They would also help in
775 the understanding of the extraordinary episode of history – whether formalised
776 in stratigraphy or not – which the Earth is currently experiencing.

777

778

779 **Conclusions**

780

- 781 • A material 'Anthropocene Series' might be defined with a historically
782 recent boundary at ~1950 CE, characterised by time proxies such as artificial
783 radionuclides, biostratigraphic changes and human-made novel materials such
784 as plastics and uncombined aluminium. It locally forms substantial, distinct and
785 correlatable sediment bodies in both terrestrial and marine realms.

786

787 • Locally, too, Anthropocene deposits so defined are difficult to recognise
788 and correlate for want of appropriate time markers to fix the boundary. These
789 are analogous to stratigraphically indeterminate deposits in the older
790 stratigraphic record.

791

792 • Commonly, also, Anthropocene deposits are difficult to separately
793 recognise as distinct units because of intermixing, for instance by human or no-
794 human bioturbation, reflecting the very short duration of the Anthropocene.
795 This may be regarded as a problem inherent in very high-resolution
796 chronostratigraphy as much as one of the Anthropocene.

797

798 • Attempts to better delineate and analyse the material expression of the
799 Anthropocene will increase our understanding of the phenomenon as a whole.

800

801

802 **Acknowledgements:** Colin Waters publishes with the permission of the
803 Executive Director, British Geological Survey, Natural Environment Research
804 Council. We thank Will Steffen and Mike Ellis for thorough and helpful reviews.

805

806

807 References

808

- 809 ALLEN, J.R.L. 1988. Modern-period muddy sediments in the Severn Estuary
810 (southwestern U.K.): a pollutant-based model for dating and correlation.
811 *Sedimentary Geology*, **58**, 1-21.
- 812 AL-ROUSAN, S., PÄTZOLD, J., AL-MOGHRABI, S., & WEFER, G., 2004, Invasion of
813 anthropogenic CO₂ recorded in planktonic foraminifera from the northern Gulf
814 of Aquaba. *International Journal of Earth Sciences*, **93**, 1066–1076, doi:
815 10.1007/s00531-004-0433-4.
- 816 ANDERSON, D.M. 2001. Attenuation of millenial-scale events by bioturbation in
817 marine sediments. *Palaeoceanography*, **16**, 352-357.
- 818 APPLEBY, P.G. 2008. Three decades of dating recent sediments by fallout
819 radionuclides: a review. *The Holocene*, **18 (1)**, 83-93.

- 820 AUTIN, W.J. & HOLBROOK, J.M. 2012. Is the Anthropocene an issue of stratigraphy or
821 pop culture? *GSA Today*, **22** (7), 60-61.
- 822 BACON, A.R., RICHTER, D. DEB, BIERMAN, P.R. & ROOD, D.H. 2012. Coupling meteoric
823 ^{10}Be with pedogenic losses of ^9Be to improve soil residence times on an
824 ancient North American interfluve. *Geology*, **40**, 847-850.
- 825 BARNOSKY, A.D. 2008. Megafauna biomass tradeoff as a driver of Quaternary and
826 future extinctions. *Proceedings of the National Academy of Sciences USA*,
827 **105**(suppl. 1), 11543-11548.
- 828 BARNOSKY, A.D. 2013. Palaeontological evidence for defining the Anthropocene.
829 In: *A Stratigraphical Basis for the Anthropocene*. Waters *et al.* (eds)
- 830 BARNOSKY, A.D., MATZKE, N., TOMIYA, S., WOGAN, G.O.U., SWARTZ, B., QUENTAL, T.,
831 MARSHALL, C., MCGUIRE, J.L., LINDSEY, E.L., MAGUIRE, K.C., MERSEY, B. & FERRER,
832 E.A. 2011. Has the Earth's sixth mass extinction already arrived? *Nature*,
833 **471**, 51-57.
- 834 BARNOSKY, A.D., HADLY, E.A., BASCOMPTE, J., BERLOW, E.L., BROWN, J.H., FORTELIUS, M.,
835 GETZ, W.M., HARTE, J., HASTINGS, A., MARQUET, P.A., MARTINEZ, N.D., MOOERS, A.,
836 ROOPNARINE, P., VERMEEIJ, G., WILLIAMS, J.W., GILLESPIE, R., KITZES, J., MARSHALL,
837 C., MATZKE, N., MINDELL, D.P., REVILLA, R. & SMITH, A.B. 2012. Approaching a
838 state-shift in the biosphere. *Nature*, **486**, 52-56.
- 839 BOLLHÖFER, A. & ROSMAN, K.J.R. 2000. Isotopic source signatures for atmospheric
840 lead: The Southern Hemisphere. *Geochimica Cosmochimica Acta*, **64**, 3251-
841 3262.
- 842 BRENCHLEY, P.J., MARSHALL, J.D., CARDEN, G.A.F., ROBERTSON, D.B.R., LONG,
843 D.G.F., MEIDLÀ, T., HINTS, L. & ANDERSON, T.F. 1994. Bathymetric and
844 isotopic evidence for a short-lived Late Ordovician glaciation in a greenhouse
845 period. *Geology*, **22**, 295-298.
- 846 BROWN, A.G., TOMS, P., CAREY, C. & RHODES, E. 2013. Geomorphology of the
847 Anthropocene: time-transgressive discontinuities of human-induced alluviation.
848 *Anthropocene* **1**, doi: 10.1016/j.ancene.2013.06.002.
- 849 BROECKER, W.S. 1987. *How to build a habitable planet*. New York, Eldigio Press,
850 291 pp.
- 851 BUESSELER, K.O. and 18 others. 2007. Revisiting carbon flux through the ocean's
852 twilight zone. *Science*, **316**, 567-570.

- 853 CARTER, R.M. 2007, Stratigraphy into the 21st century. *Stratigraphy*, **4**, 187-194.
- 854 CERTINI, G. & SCALENGHE, R. 2011. Anthropogenic soils are the golden spikes for the
855 Anthropocene. *The Holocene*, **21**, 1269-1274
- 856 COHEN, A.S., COE, A.L. & KEMP, D.B. 2007. The Late Palaeocene-Early Eocene and
857 Toarcian (Early Jurassic) carbon isotope excursions: a comparison of their time
858 scales, associated environmental changes, causes and consequences. *Journal of*
859 *the Geological Society, London*, **164**, 1093-1108.
- 860 CRUTZEN, P.J. 2002, Geology of Mankind. *Nature*, **415**, 23.
- 861 CRUTZEN, P.J. & STOERMER, E.F. 2000, The ‘Anthropocene’. *Global Change*
862 *Newsletter*, **41**, 17-18.
- 863 DOUNAS, C., DAVIES, I., TRIANTAFYLLOU, G., KOULOURI, P., PETIHAKIS, G.,
864 ARVANITIDIS, C., SOURLATZIS, G. & ELEFTHERIOU, A. 2007. Large-scale
865 impacts of bottom trawling on shelf primary productivity. *Continental Shelf*
866 *Research*, **27**, 2198-2210.
- 867 EDGEWORTH, M. 2013. Archaeological stratigraphy and artificial ground. In: A
868 Stratigraphical Basis for the Anthropocene. Waters *et al.* (eds).
- 869 EDWARDS, K.J. & WHITTINGTON, G. 2001. Lake sediments, erosion and landscape
870 change during the Holocene in Britain and Ireland. *Catena*, **42**, 143-173.
- 871 ELLIS, E.C. 2011. Anthropogenic transformation of the terrestrial biosphere.
872 *Philosophical Transactions of the Royal Society*, **A369**, 1010-1035.
- 873 ELLIS, E.C., ANTILL, E.C. & KREFT, H. 2012. All Is Not Loss: Plant Biodiversity in
874 the Anthropocene. *PLoS ONE*, **7**, e30535.
- 875 ELSIG, J., SCHMITT, J., LEUENBERGER, D., SCHNEIDER, R., EYER, M., LEUENBERGER,
876 M., JOOS, F., FISCHER, H. & STOCKER, T.F. 2009. Stable isotope constraints on
877 Holocene carbon cycle changes from an Antarctic ice core. *Nature*, **461**, 507-
878 510.
- 879 FINNEY 2013 In: A Stratigraphical Basis for the Anthropocene. Waters *et al.* (eds).
- 880 FORD, J.R., PRICE, S.J., COOPER, A.H. & WATERS, C.N. An assessment of
881 lithostratigraphy for anthropogenic deposits. In: A Stratigraphical Basis for
882 the Anthropocene. Waters *et al.* (eds).
- 883 GALE, S.J. & HOARE, P.G. 2012. The stratigraphic status of the Anthropocene. *The*
884 *Holocene*, DOI:10.1177/0959683612449764.

- 885 GALUSZKA et al. this volume. Assessing the Anthropocene with geochemical
886 methods. In: A Stratigraphical Basis for the Anthropocene. Waters *et al.*
887 (eds).
- 888 GATTUSO, J. P., SMITH, S. V., HOGAN, C. M. & DUFFY, J. E. 2009 'Coastal zone'. In
889 Encyclopedia of Earth (ed. C. J. Cleveland). Washington, DC: Environmental
890 Information Coalition, National Council for Science and the Environment. (First
891 published in the Encyclopedia of Earth. See
892 http://www.eoearth.org/article/Coastal_zone.)
- 893 GIBBARD, P.L. & WALKER, M.J.C. 2013. The term 'Anthropocene' in the context of
894 formal geological classification. In: A Stratigraphical Basis for the
895 Anthropocene. Waters *et al.* (eds)
- 896 GOSSE, J.C. & PHILLIPS, F.M. 2001. Terrestrial in situ cosmogenic nuclides: theory
897 and application. *Quaternary Science Reviews*, **20**, 1475-1560.
- 898 GOUDIE, A.S. 2009. Dust storms: Recent developments. *Journal of Environmental
899 Management*, **90**, 89-94.
- 900 GRADSTEIN, F., OGG, G., SCHMITZ, M. & OGG, G. (eds) *A Geological Time Scale
901 2012*. Elsevier, 1144 pp.
- 902 HAFF, P. this volume. Technology as a geological phenomenon: Implications for
903 human well-being. In: A Stratigraphical Basis for the Anthropocene. Waters
904 *et al.* (eds).
- 905 HANCOCK, G.J., LESLIE, C., EVERETT, S.E., TIMS, S.G., BRUNSKILL, G.J. & HAESE, R. 2011.
906 Plutonium as a chronomarker in Australian and New Zealand sediments: a
907 comparison with ¹³⁷Cs. *Journal of Environmental Radioactivity*, **102**, 919-
908 929.
- 909 HAY, W.W. 2013. *Experimenting on a small planet: a scholarly entertainment*.
910 Springer-Verlag: Berlin Heidelberg, 983 pp.
- 911 HIGGINS, S., OVEREEM, I., TANAKA, A. & SYVITSKI, J.P.M. 2013. Land subsidence at
912 aquaculture facilities in the Yellow River delta, China. *Journal of Geophysical
913 Research* **40**, 3898-3902.
- 914 HOLTGRIEVE, G.W., SCHINDLER, D.E., HOBBS, W.O., LEAVITT, P.R., WARD, E.J.,
915 BUNTING, L., CHEN, G., FINNEY, B.P., GREGORY-EAVES, I., HOLMGREN, S.,
916 LISAC, M.J., LISI, P.J., NYDICK, K., ROGERS, L.A., SAROS, J.E., SELBIE, D.T.,
917 SHAPLEY, M.D., WALSH, P.B. & WOLFE, A.P. 2011. A coherent signature of

- 918 anthropogenic nitrogen deposition to remote watersheds of the northern
919 hemisphere. *Science*, **334**, 1546-1548.
- 920 HOOKE, R. LEB., 2000, On the history of humans as geomorphic agents. *Geology* **28**:
921 843-846.
- 922 IPCC (Intergovernmental Panel on Climate Change) 2001. *Climate Change 2001: The*
923 *Scientific Basis. Contribution of Working Group I to the Third Assessment*
924 *Report of the Intergovernmental Panel on Climate Change*. Houghton, J.T., Y.,
925 Ding, D.J., Griggs, M., Noguer, P.J., van der Linden, X., Dai, K., Maskell &
926 Johnson, C.A. (eds). Cambridge University Press, Cambridge, United Kingdom
927 and New York, NY, USA, 881pp.
- 928 IPCC (Intergovernmental Panel on Climate Change). 2007. *Climate change 2007:*
929 *synthesis report. Summary for policy makers*. Available at:
930 <http://www.ipcc.ch/SPM2feb07.pdf>.
- 931 KAPLAN, J.O., KRUMHARDT, K., ELLIS, E.C. RUDDIMAN, W.F., LEMMEN, C. &
932 GOLDEWIJK, K.K. 2011. Holocene carbon emissions as a result of anthropogenic
933 land cover change. *The Holocene*, **21** (5), 775-791.
- 934 KELLIE-SMITH, O. & COX, P.M. 2011. Emergent dynamics of the climate-economy
935 system in the Anthropocene. *Philosophical Transactions of the Royal Society*,
936 **A369**, 868-886.
- 937 KEMP, D.B., COE, A.L., COHEN, A.S. & SCHWARK, L. 2005. Astronomical pacing of
938 methane release in the Early Jurassic period. *Nature*, **437**, 396-399.
- 939 KRAMER, N., WOHL, E.E. & HARRY, D.L. 2011. Using ground penetrating radar to
940 ‘unearth’ buried beaver dams. *Geology*, **40**, 43-46.
- 941 KRUGE, M. A. 1999. Molecular organic geochemistry of New York Bight sediments.
942 Sources of biogenic organic matter and polycyclic aromatic hydrocarbons.
943 *Northeastern Geology and Environmental Sciences*, **21**, 121-128.
- 944 KULKARNI, A.V., BAHUGUNA, I.M., RATHORE, B.P., SINGH, S.K., RANDHAWA, S.S.,
945 SOOD, R.K. & DHAR, S. 2007. Glacial retreat in Himalaya using Indian Remote
946 Sensing satellite data. *Current Science*, **92**, 69-74.
- 947 LARMER, B. 2010. Shanghai Dreams. *National Geographic*, March 2010, 124-141
- 948 LEWIN, J. 2012. Enlightenment and the GM floodplain. *Earth Surface Processes and*
949 *Landforms*. DOI:10.1002/esp.3230
- 950 MALAKOFF, D. 2002. Trawling’s a drag for marine life, say studies. *Science* **298**,
951 2123.

- 952 MANN, M.E. 2002. Little Ice Age. In: *The Earth system: physical and chemical*
953 *dimensions of global environmental change*. MacCracken, M.C. & Perry, J.S.
954 (eds). *Encyclopedia of Global Environmental Change*, Volume 1, 504–509.
955 (Chichester: John Wiley & Sons, Ltd.) ISBN 0-471-97796-9.
- 956 MARSHALL, W.A., GEHRELS, W.R., GARNETT, M.H., FREEMAN, S.P.H.T., MADEN, C.
957 & XU, SHENG, 2007, The use of ‘bomb spike’ calibration and high-precision
958 AMS 14C analyses to date salt marsh sediments deposited during the last three
959 centuries. *Quaternary Research*, **68**, 325-337.
- 960 MERRITS, D., WALTER, R., RAHNIS, M., HARTRANFT, J., COX, S., GELLIS, A., POTTER,
961 N., HILGARTNER, W., LANGLAND, M., MANION, L., LIPPINCOTT, C., SIDDIQUI, S.,
962 REHMAN, Z., SCHEID, C., KRATZ, L., SHILLING, A., JENSCHKE, M., DATIN, C.,
963 CRANMER, E., REED, A., MATUSZEWSKI, D., VOLI, M., OHLSON, E.,
964 NEUGEBAUER, A., AHAMED, A., NEAL, C., WINTER, A. & BECKER, C. 2011.
965 Anthropocene streams and base-level controls from historic dams in the
966 unglaciated mid-Atlantic region, USA. *Philosophical Transactions of the Royal*
967 *Society, A***369**, 976-1009.
- 968 MOULIN, C. & CHIAPELLO, I. 2006. Impact of human-induced desertification on the
969 intensification of Sahel dust emission and export over the last decades.
970 *Geophysical Research Letters*, **33**, L18808.
- 971 MYERS, R.A. & WORM, B., 2003, Rapid worldwide depletion of predatory fish
972 communities. *Nature*, **423**, 280-283.
- 973 NATURE (editorial) 2011. The human epoch. *Nature*, **473**, 254.
- 974 NAYAR, A. 2009. When the ice melts. *Nature*, **461**, 1042-1046.
- 975 NEW, M.G., LIVERMAN, D.M., BETTS, R.A., ANDERSON, K.L. & WEST, C.C. (eds)
976 2011. Four degrees and beyond: the potential of a global temperature increase
977 of four degrees and its implications. *Philosophical Transactions of the Royal*
978 *Society, A***369**.
- 979 PALANQUES, A., GUILLÉN, J. & PUIG, P. 2001. Impact of bottom trawling on water
980 turbidity and muddy sediment of an unfished continental shelf. *Limnology and*
981 *Oceanography*, **46**, 1100-1110..
- 982 POIRIER, C., CHAUMILLON, E. & ARNAUD, F. 2011. Siltation of river-influenced
983 coastal environments: Respective impact of late Holocene land use and high-
984 frequency climate changes. *Marine Geology*, **290**, 51-62.

- 985 PRICE, S.J., FORD, J.R., COOPER, A.H. & NEAL, C. 2011. Humans as major geological
986 and geomorphological agents in the Anthropocene: the significance of artificial
987 ground in Great Britain. *Philosophical Transactions of the Royal Society*, **A369**,
988 1056-1084.
- 989 PUIG, P., CANALS, M., COMPANY, J.B., MARTIN, J., AMBLAS, D., LASTRAS, G.,
990 PALANQUES, A. & CALAFAT, A.M. 2012. Ploughing the deep sea floor. *Nature*,
991 **489**, 286-289.
- 992 RAHMSTORF, S., 2007, A semi-empirical approach to projecting future sea-level rise.
993 *Science*, **315**, 368-370.
- 994 RAMIREZ-LLODRA, E., TYLER, P.A., BAKER, M.C., BERGSTAD, O.A., CLARK, M.R.,
995 ESCOBAR, E., LEVIN, L.A. MENOT, L., ROWDEN, A.A., SMITH, C.R. & VAN
996 DOVER, C.L. 2011. Man and the Last Great Wilderness: Human Impact on the
997 Deep Sea. *Plos One* **6** (8), e22588, 1-25.
- 998 REMANE, J., BASSETT, M.G., COWIE, J.W., GOHRBANDT, K.H., LANE, H.R.,
999 MICHELSON, O. & NAIWEN, W. 1996. Revised guidelines for the establishment
1000 of global chronostratigraphic standards by the International Commission on
1001 Stratigraphy (ICS). *Episodes* **19**, 77-81.
- 1002 REVKIN, A. 1992. *Global warming: understanding the forecast*. New York, NY:
1003 American Museumn of Natural History, Environmental Defense Fund,
1004 Abbeville Press.
- 1005 RICHTER, D. 2007. Humanity's transformation of Earth's soil: pedology's new
1006 frontier. *Soil Science* **172**, 957-967.
- 1007 ROBERTS, C. 2007. *The unnatural history of the sea*. Island Press.
- 1008 ROBISON, B.H., REISENBICHLER, K.R. & SHERLOCK, R.E. 2005. Giant larvacean
1009 houses: rapid carbon transport to the deep sea floor. *Science*, **308**, 1609-1611.
- 1010 ROCKSTRÖM, J., STEFFEN, W., NOONE, K., PERSSON, Å, CHAPIN, III, F.S., LAMBIN,
1011 E.F., LENTON, T.M., SCHEFFER, M., FOLKE, C., SCHELLNHUBER, H.J., NYKVIST,
1012 B., DE WIT, C.A., HUGHES, T., VAN DER LEEUW, S., RODHE, H., SÖRLIN, S.,
1013 SNYDER, P.K., COSTANZA, R., SVEDIN, U., FALKENMARK, M., KARLBERG, L.,
1014 CORELL, R.W., FABRY, V.J., HANSEN, J., WALKER, B., LIVERMAN, D.,
1015 RICHARDSON, K., CRUTZEN, P. & FOLEY, J.A. 2009. A safe operating space for
1016 humanity. *Nature*, **461**, 472-475.
- 1017 RUDDIMAN, W F. 2003. The Anthropogenic Greenhouse Era began thousands of
1018 Years Ago. *Climatic Change*, **61**, 261-293.

- 1019 RUDDIMAN, W.F. 2013. Anthropocene. *Annual Review of Earth and Planetary*
1020 *Sciences*. DOI: 10.1146/annurev-earth-050212-123944
- 1021 SHEPPARD, C. 2006. Trawling the sea bed. *Marine Pollution Bulletin*, **52**, 831-835.
- 1022 SHOULDERS, S.J. & CARTWRIGHT, J.A., 2004, Constraining the depth and timing of
1023 large-scale conical sandstone intrusions. *Geology*, **32**, 661-664.
- 1024 SMITH, D.M., ZALASIEWICZ, J.A., WILLIAMS, M., WILKINSON, I., REDDING, M. &
1025 BEGG, C. 2010. Holocene drainage of the English Fenland: roddons and their
1026 environmental significance. *Proceedings of the Geologists' Association*, **121**,
1027 256-269.
- 1028 STANLEY, D.J. 1996. Nile delta: extreme case of sediment entrapment on a delta plain
1029 and consequent coastal land loss. *Marine Geology*, **129**, 189-195.
- 1030 STEFFEN, W., CRUTZEN, P.J. & MCNEILL, J.R., 2007, The Anthropocene: are humans
1031 now overwhelming the great forces of Nature? *Ambio*, **36**, 614-621.
- 1032 STEFFEN, W.Å.P, DEUTSCH, L., ZALASIEWICZ, J., WILLIAMS, M., RICHARDSON, K.,
1033 CRUMLEY, C., CRUTZEN, P., FOLKE, C., GORDON, L., MOLINA, M.,
1034 RAMANATHAN, V., ROCKSTRÖM, J., SCHEFFER, M., SCHELLNHUBER, J. &
1035 SVEDIN, U. 2011. The Anthropocene: from global change to planetary
1036 stewardship. *Ambio*, **40**, 739-761.
- 1037 SYVITSKI, J.P.M. & KETTNER, A. 2011. Sediment flux and the Anthropocene.
1038 *Philosophical Transactions of the Royal Society*, **A369**, 957-975.
- 1039 TANAKA, K.L. & HARTMANN, W.K. 2012. Ch. 15: The Planetary Time Scale. In: The
1040 Geological Time Scale 2012. GRADSTEIN, F.M., OGG, J., SCHMITZ, M. & OGG,
1041 G. (eds), 275-298. Elsevier.
- 1042 TÖRNQVIST, T E, WALLACE, D J, STORMS, J E A, WALLINGA, J, VAN DAM, R L,
1043 BLAAUW, M, DERKSEN, M S, KLERKS, C J W, MEIJNEKEN, C, & SNIJDERS, E M
1044 A. 2008. Mississippi Delta subsidence primarily caused by compaction of
1045 Holocene strata. *Nature Geoscience*, **1**, 173-176.
- 1046 TYRRELL, T. 2011. Anthropogenic modification of the oceans. *Philosophical*
1047 *Transactions of the Royal Society*, **A369**, 887-908.
- 1048 VANE, C.H., CHENERY, S.R., HARRISON, I., KIM, A.W., MOSS-HAYES, V. & JONES,
1049 D.G. 2011. Chemical signatures of the Anthropocene in the Clyde estuary, UK:
1050 sediment-hosted Pb, 207/206 Pb, total petroleum hydrocarbon and
1051 polychlorinated biphenyl pollution records. *Philosophical Transactions of the*
1052 *Royal Society*, **A369**, 1085-1111.

- 1053 VIDAS, D. 2011. The Anthropocene and the international law of the sea. *Philosophical
1054 Transactions of the Royal Society, A***369**, 909-925.
- 1055 WALKER, M., JOHNSEN, S., RASMUSSEN, S.O., STEFFENSEN, J.P., POPP, T., GIBBARD,
1056 P., HOEK, W., LOWE, J., BJÖRCK, S., CWYNAR, L., HUGHEN, K., KERSHAW, P.,
1057 KROMER, B., LITT, T., LOWE, D.J., NAKAGAWA, T., NEWNHAM, R. &
1058 SCHWANDER, J. 2009. Formal definition and dating of the GSSP (Global
1059 Stratotype Section and Point) for the base of the Holocene using the Greenland
1060 NGRIP ice core, and selected auxiliary records. *Journal of Quaternary Science,*
1061 **24**, 3-17.
- 1062 WALKER, M.J.C., BERKELHAMMER, M., BJÖRK, S., CWYNAR, L.C., FISHER, D.A.,
1063 LONG, A.J. & LOWE, J.J. 2012. Formal subdivision of the Holocene
1064 Series/Epoch: a discussion paper by a Working Group of INTIMATE
1065 (Integration of ice-core, marine and terrestrial records) and the Subcommission
1066 on Quaternary Stratigraphy (International Commission on Stratigraphy).
1067 *Journal of Quaternary Studies* doi: 10.1002/jqs.2565.
- 1068 WILKINSON, B.H., 2005, Humans as geologic agents: A deep-time perspective.
1069 *Geology*, **33**, 161-164.
- 1070 WILKINSON, I.P. et al. this volume. Micropalaeontological signatures of the
1071 Anthropocene. In: *A Stratigraphical Basis for the Anthropocene*. Waters *et*
1072 *al.* (eds).
- 1073 WILLIAMS, M., et al. This volume.
- 1074 WOLFE, A.P., HOBBS, W.O., BIRKS, H.W., BRINER, J.P., HOLMGREN, S.W.,
1075 INGOLFSSÓN, Ó., KAUSHAL, S.J., MILLER, G.H., PAGANI, M., SAROS, J.E. &
1076 VINEBROOK, R.D. 2013. Stratigraphic expressions of the Holocene-
1077 Anthropocene transition revealed in sediments from remote lakes. *Earth-
1078 Science Reviews*, **116**, 17-34.
- 1079 YAN, P., SHI, P., GAO, S., CHEN, L., ZHANG, X. & BAI, L. 2002. ^{137}Cs dating of
1080 lacustrine sediments and human impacts on Dalian Lake, Qinghai Province,
1081 China. *Catena*, **47**, 91-99.
- 1082 ZACHOS, J.C., RÖHL, U., SCHELLENBERG, S.A., SLUIJS, A., HODELL, D.A., KELLY,
1083 D.C., THOMAS, E., NICOLO, M., RAFFI, I., LOURENS, L.J., MCCARREN, H. &
1084 KROON, D., 2005, Rapid acidification of the ocean during the Paleocene-
1085 Eocene thermal maximum. *Science*, **308**, 1611-1615.
- 1086 ZALASIEWICZ, J. 2013. The epoch of humans. *Nature Geoscience* **6**, 8-9.

- 1087 ZALASIEWICZ, J.A., SMITH, A, BRENCHLEY, P., EVANS, J., KNOX, R., RILEY, N., GALE,
1088 A., RUSHTON, A., GIBBARD, P., HESSELBO, S., MARSHALL, J., OATES, M.,
1089 RAWSON, P. & TREWIN, N. 2004. Simplifying the stratigraphy of time.
1090 *Geology*, **32**, 1-4.
- 1091 ZALASIEWICZ, J., WILLIAMS, M, SMITH, A., BARRY, T.L., BOWN, P.R., RAWSON, P.,
1092 BRENCHLEY, P., CANTRILL, D., COE, A.E., COPE, J.C.W., GALE, A., GIBBARD,
1093 P.L., GREGORY, F.J., HOUNSLOW, M., KNOX, R., POWELL, P., WATERS, C.,
1094 MARSHALL, J., OATES & STONE, P. 2008. Are we now living in the
1095 Anthropocene? *GSA Today*, **18** (2), 4-8.
- 1096 ZALASIEWICZ, J., WILLIAMS M, FORTEY, R.A., SMITH, A.G., BARRY, T.L. COE, A.L.,
1097 BOWN, P.R., GALE, A., GIBBARD, P.L., GREGORY, F.J., HOUNSLOW, M.W.,
1098 KERR, A.C., PEARSON, P., KNOX, R., POWELL J., WATERS, C., MARSHALL, J.,
1099 OATES, M., RAWSON P. & STONE, P. 2011. Stratigraphy of the Anthropocene.
1100 *Philosophical Transactions of the Royal Society*, **A369**, 1036-1055.
- 1101 ZALASIEWICZ, J., CRUTZEN, P.J. & STEFFEN, W. 2012. Anthropocene. Chapter 32 (pp.
1102 1033-1040) in *A Geological Time Scale 2012* (ed. GRADSTEIN, F., OGG, G.,
1103 SCHMITZ, M. & OGG, G.). Elsevier, 1144 pp.
- 1104 ZALASIEWICZ, J., WILLIAMS, M, SMITH, A., BARRY, T.L., BOWN, P.R., RAWSON, P.,
1105 BRENCHLEY, P., CANTRILL, D., GALE, A., GIBBARD, P.L., GREGORY, F.J.,
1106 HOUNSLOW, M., KNOX, R., POWELL, P., WATERS, C., MARSHALL, J., OATES,
1107 STONE, P. & TREWIN, N. 2007. The scale-dependence of strata-time relations:
1108 implications for stratigraphic classification. *Stratigraphy*, **4**, 139-144.
- 1109 ZALASIEWICZ, J., CITA, M.B., HILGEN, F., PRATT, B.R., STRASSER, A. THIERRY, J.,
1110 WEISSERT, H. 2013. Chronostratigraphy and geochronology: a proposed
1111 realignment. *GSA Today*, **23** (3), 4-8.
- 1112 ZALASIEWICZ, J., KRYZA, R. & WILLIAMS, M. this volume. The mineral signature of
1113 the Anthropocene. In: A Stratigraphical Basis for the Anthropocene. Waters *et*
1114 *al.* (eds).
- 1115 ZALASIEWICZ, J. & WILLIAMS, M. In press. The Anthropocene: a comparison with the
1116 Ordovician-Silurian boundary. *Rendiconti Lincei*.
- 1117
- 1118
- 1119 **List of figures**
- 1120

1121

1122

1123 Figure 1. Comparison of lengths of epochs from the mid-Cenozoic to the present,
1124 showing progressive shortening in time span. Dates from Gradstein et al. 2012.

1125

1126 Fig. 2. The rapid mid-twentieth century growth of Shanghai, as an example of the
1127 formation of a distinct, extensive sedimentary facies that may be referred to a putative
1128 Anthropocene Series. Information from Larmer (2010) and Map of Central Shanghai,
1129 printed by the British War Office/US Army Map Service in 1935.

1130







