

Article (refereed) - postprint

Grez, Audrey A.; Zaviezo, Tania; Roy, Helen E.; Brown, Peter M.J.; Segura, Bernardo. 2017. **In the shadow of the condor: invasive *Harmonia axyridis* found at very high altitude in the Chilean Andes.** *Insect Conservation and Diversity*, 10 (6). 483-487. <https://doi.org/10.1111/icad.12258>

© 2017 The Royal Entomological Society

This version available <http://nora.nerc.ac.uk/id/eprint/518682/>

NERC has developed NORA to enable users to access research outputs wholly or partially funded by NERC. Copyright and other rights for material on this site are retained by the rights owners. Users should read the terms and conditions of use of this material at <http://nora.nerc.ac.uk/policies.html#access>

This document is the author's final manuscript version of the journal article, incorporating any revisions agreed during the peer review process. Some differences between this and the publisher's version remain. You are advised to consult the publisher's version if you wish to cite from this article.

The definitive version is available at <http://dx.doi.org/10.1111/icad.12258>

Contact CEH NORA team at
noraceh@ceh.ac.uk

1 **In the shadow of the condor: Invasive *Harmonia axyridis* found at very high**
2 **altitude in the Chilean Andes**

3

4 Audrey A Grez¹

5 Tania Zaviezo²

6 Helen E Roy³

7 Peter MJ Brown⁴

8 Bernardo Segura¹

9

10 Running Head: *Harmonia axyridis* at high altitude in the Andes

11

12

13

14 ¹Facultad de Ciencias Veterinarias y Pecuarias, Universidad de Chile, Casilla 2-15,
15 Santiago, Chile

16 ²Facultad de Agronomía e Ingeniería Forestal, Pontificia Universidad Católica de Chile,
17 Casilla 306-22, Santiago, Chile

18 ³Centre for Ecology & Hydrology, Wallingford, Oxfordshire, OX10 8BB, UK

19 ⁴Animal & Environment Research Group, Anglia Ruskin University, Cambridge, CB1
20 1PT, UK

21

22 Corresponding author: Audrey Grez; agrez@uchile.cl; telephone number: +56-
23 29785549

24

25

26 **Abstract.** 1. *Harmonia axyridis* has invaded many regions of the world, with negative
27 effects on local biodiversity, and thus it is of global concern for biological conservation.
28 Recently it has invaded central Chile, one of the world's biodiversity hotspots, where
29 the abundance and richness of ladybird species, particularly native species in
30 agroecosystems, have declined following its arrival.

31 2. *Harmonia axyridis* is particularly abundant in spring in the valleys of central Chile,
32 but there is a dramatic decline in its abundance during the hot summer months.

33 3. This study reports the occurrence of this invasive alien species in the summer at
34 high altitudes (3578 m asl) in the Andes, which is the highest record worldwide.
35 Individuals were observed on native cushion plants, in a unique environment rich in
36 endemic species. *Harmonia axyridis* were active, reproducing and co-occurring with
37 three other species of coccinellids.

38 4. The dispersal of *H. axyridis* and other coccinellids from the valleys to high altitudes
39 could represent a mechanism to escape the adverse high temperatures during the
40 summer, returning to the valleys in autumn. Our study highlights the need to study the
41 impacts of invasive alien species across a range of habitats, including not only
42 agricultural landscapes, but also extreme ecosystems.

43

44

45 **Key words.** Altitude, Coccinellidae, extreme ecosystems, harlequin ladybird, multi-
46 coloured Asian ladybeetle, non-native species.

47

48 **Introduction**

49

50 The rate of translocation of species by humans beyond their native ranges is increasing
51 and, for most taxa, there appears to be no indication of deceleration (Seebens *et al.*,
52 2017). The effects of alien species on biodiversity and ecosystem functioning are
53 widely recognised (Simberloff *et al.*, 2013). Invasive alien species, the subset of alien
54 species that threaten biodiversity, society or the economy, are in part defined by their
55 ability to spread rapidly within invaded regions. Some studies have suggested that
56 natural habitats are resistant to invasion and evoke mechanisms related to the resident
57 community structure and specifically species diversity (Lyons & Schwartz, 2001; Shea
58 & Chesson, 2002).

59 One invasive alien species of global concern is *Harmonia axyridis* (Pallas)
60 (Coleoptera: Coccinellidae), the harlequin or multicoloured Asian ladybeetle, that has
61 invaded many regions of the world, with negative effects on local biodiversity,
62 particularly native coccinellids (Alyokhin & Sewell, 2004; Harmon *et al.*, 2007; Brown *et*
63 *al.*, 2011; Roy *et al.*, 2012). In the last 25 years it has expanded its distribution to five
64 continents, with Oceania (New Zealand) being added in 2016. Notably, this species
65 reaches high population numbers in temperate regions of the northern and southern
66 hemispheres, particularly in anthropogenic habitats (Roy *et al.*, 2016).

67 In Chile, the first wild populations of *H. axyridis* were recorded in 2003 in the
68 central zone, 100 km north of Santiago, but populations started to rise in 2010-2011,
69 spreading rapidly throughout the country. More recently *H. axyridis* has reached very
70 high abundance in crops and has also invaded natural habitats, such as the
71 sclerophyllous matorral (Greze *et al.*, 2016). This is of concern because central Chile is
72 one of the world's 35 biodiversity hotspots, an important reservoir of biodiversity
73 (Mittermeier *et al.*, 2011). Also, this region has the highest number of endemic
74 coccinellids in the country (Alaniz & Greze, unpublished data) which could be negatively
75 affected by *H. axyridis*. Indeed there has been a decline in abundance and richness of

76 ladybird species, particularly native species, within alfalfa crops following the arrival of
77 *H. axyridis* (Grez *et al.*, 2016).

78 In Chile it has been observed that *H. axyridis* reaches sustained high
79 abundance in alfalfa crops during early spring. Noticeably, in summer its populations
80 decline and it becomes almost absent within these crops until autumn, when it
81 recolonizes for a short period of time before the onset of winter (Grez & Zaviezo,
82 unpublished data). The phenology of *H. axyridis* within central Chile could be explained
83 by the Mediterranean climate of this region, with wet and cold winters and hot and dry
84 summers (Di Castri & Hajek, 1976), and by the low tolerance of *H. axyridis* to high
85 temperatures (Benelli *et al.*, 2015; Barahona-Segovia *et al.*, 2016). The behavioural
86 responses of *H. axyridis* under these unfavourable field conditions is unknown. One
87 hypothesis is that during the summer it migrates towards places where temperatures
88 are lower, for example high altitudes. This note reports the colonization by *H. axyridis*
89 in the high Andes of central Chile, an extreme native habitat, during summer.

90

91 **Study site**

92 Two surveys were conducted during summer 2017 (29 January and 12 March), in the
93 high Andes of central Chile, in the surroundings of Valle Nevado ski resort (UTM
94 WGS84 19S 384386 E; 6313663 S), approximately 50 km east of Santiago (Appendix
95 S1). This area has an alpine climate, with very cold winters (mean air temperature ~
96 1.7°C, minimum of -15°C) and mild summers (mean air temperature ~ 6.8°C, and
97 maximum 17°C) (Molina-Montenegro *et al.*, 2006). In these two surveys, coccinellids
98 were searched in an altitudinal gradient, from 2700 to 3600 m asl. In this area there are
99 no trees or bushes, and the vegetation is very patchy, dominated by two native cushion
100 plants (Apiaceae): *Azorella madreporica* Clos and *Laretia acaulis* (Cav.) Gill. et Hook
101 (Fig. 1A). *Azorella madreporica* is a very flat and tightly knit cushion species, extending
102 from 33°S to 50°S, and growing from above 3200 m asl in the Andes of central Chile
103 (33°S) and close to sea level in its southern distribution (Hoffmann *et al.*, 1998). *Laretia*

104 *acaulis* is a hard resinous cushion plant native of the high Andes of Chile and
105 Argentina, extending from 28°S to 35°S and most abundant between 2100 to 3100 m
106 asl (Hoffmann *et al.*, 1998). Cushion plants are one of the best-adapted growth forms in
107 this habitat, generating more suitable micro-habitats for other plants and insects
108 (Cavieres *et al.*, 1998; Molina-Montenegro *et al.*, 2009).

109

110 **Results and discussion**

111 In total, 37 adult *H. axyridis* (succinea form, the only recorded form in Chile so
112 far; Grez & Zaviezo 2015) were found distributed from 2790 to 3578 m asl, all of them
113 on patches of the cushion plants (Table 1). In lower altitude samples *H. axyridis* was
114 absent, and the highest altitude where it was found coincides with the highest places
115 with cushion plants in this area. Previous to this survey, from a citizen science
116 observation reported through <http://www.chinita-arlequin.uchile.cl/> (Grez & Zaviezo,
117 2015), one record at high altitudes was known in Chile (3086 m asl, 150 km north of
118 the current records, Cordillera de Cuncumén; 372642 E 6460876 S) on *Eleocharis*
119 *pseudoalbibracteata* Nes & Meyen ex Kunth. (Cyperaceae), another native plant.
120 These records are the highest of *H. axyridis* worldwide. Other records of this species in
121 high altitudes are from around 2500 m asl (Lesotho, southern Africa), and 1800 m asl
122 (South Africa) (Stals 2010) in autumn-winter. In Europe it has been found in a range of
123 altitudes up to 2280 m asl in Carinthia (Austria) (Roy *et al.*, 2016). Additionally, *H.*
124 *axyridis* is known to overwinter at moderate altitudes (e.g. on rocky mountains) in its
125 native range (Wang *et al.*, 2011). In contrast to these other high altitude records from
126 around the world, data of this study were collected in summer. It has been reported that
127 some ladybirds estivate at high altitudes (Stewart *et al.*, 1967), but in our observations
128 the majority of individuals were active and reproducing. Therefore, the high Andes
129 would represent a refuge for *H. axyridis* during the summer, escaping from the high
130 temperatures in the valley.

131 Together with *H. axyridis*, three other ladybirds were present: the native (and
132 endemic) *Eriopsis chilensis* Hofmann and two other non-native species, *Hippodamia*
133 *convergens* (Guerin-Meneville) (a North American species first introduced to Chile in
134 1903, but only established in the early 90's) and *Hippodamia variegata* (Goeze) (a
135 Eurasian species introduced from South Africa in 1967) (Fig. 1B - D). Additionally, the
136 native *Adalia deficiens* Mulsant was found, but only at the lowest altitude samples
137 (Table 1). Mainly larvae and adults of these species were observed, but also a few
138 eggs and pupae. Previously, in summer 2003 and 2006, the first three species were
139 also found in the same area of the present survey, associated with the same species of
140 cushion plants, which were suggested to provide more suitable microclimatic conditions
141 for the ladybirds in this habitat (Molina-Montenegro *et al.*, 2006; 2009). Notably, in the
142 previous studies at these high altitude sites *H. axyridis* was not observed, suggesting
143 that this invasive alien species has only recently colonized the high Andes, considering
144 that it had arrived in the adjacent valleys near Santiago in 2008.

145 Co-occurrence of *H. axyridis* with these other species may represent a threat to
146 biodiversity as it has been documented in other regions, including the nearby valleys
147 (Grez *et al.*, 2016; Roy *et al.*, 2016). It has been reported that *H. axyridis* is a strong
148 competitor and intraguild predator of other coccinellids due to its larger size (*H.*
149 *axyridis*: up to 8 mm; *H. convergens*: up to 6.5 mm; *H. variegata* and *E. chilensis* up to
150 6 mm; González, 2006) and strong physical and chemical defences (reviewed within
151 Roy *et al.*, 2016). The asymmetry of these interactions in favour of *H. axyridis*, coupled
152 with its recent colonization of this unique habitat, could result in the displacement of the
153 other species. Additionally, *H. axyridis* could contain high numbers of obligate parasitic
154 microsporidia and Laboulbeniales fungi (Vilcinskas *et al.*, 2013; Roy *et al.*, 2016),
155 potentially impacting the health of the other species they interact with. Observations
156 during the survey of mating attempts between a male *H. variegata* and a female *H.*
157 *axyridis* may indicate that parasite transmission between species is plausible. The

158 potential of *H. axyridis* of negatively impacting biodiversity in the high Andes is worthy
159 of further study.

160 Apart from the species reported here, only two other ladybirds have been
161 recorded at altitudes in excess of 3000 m: *Coccinella septempunctata* L. at up to 3475
162 m (Rice, 1992) and *Hippodamia quinquesignata* (Kirby) at up to 3354 m (Edwards,
163 1957), both in the USA. Also, in invertebrate surveys at the summit of Mauna Kea,
164 Hawaii, *H. convergens* was found at a number of very high altitude sites up to 4226 m
165 and *C. septempunctata* at a single site at the same altitude (Englund *et al.*, 2010).
166 Nevertheless, these insects were considered aeolian, i.e. not resident of the area
167 where they were collected, but rather blown up from lower elevations (Englund *et al.*,
168 2010).

169 In summary, this note provides evidence of the occurrence of the invasive alien
170 species *H. axyridis* at high altitudes in the Andes near Santiago, in a unique
171 environment rich in endemic species (Arroyo & Cavieres, 2013), which constitutes the
172 highest report worldwide. This is unique not only because of the high altitude and
173 associated extreme environment, but also because it highlights the way in which *H.*
174 *axyridis* can survive adverse high temperatures during the summer within an invaded
175 region through dispersal to elevated positions, a further mechanism explaining the
176 success of this invading species. This report also highlights the importance of including
177 landscape scale factors (i.e. altitude) into models predicting establishment, spread and
178 ultimately impacts of invasive alien species to ensure that the diversity of habitats and
179 associated niche opportunities for invading species are considered.

180

181 **Conflict of interest**

182 Authors have no conflict of interest.

183

184 **Acknowledgements**

185 This study was supported by the Chilean government through FONDECYT 1140662.
186 HER receives co-funding from the Natural Environment Research Council (NERC) and
187 the Joint Nature Conservation Committee (JNCC). PMJ Brown also thanks Anglia
188 Ruskin University.

189

190 **Supporting Information**

267 Appendix S1. Map of central Chile and surveyed location.

268

269 **References**

270

271 Alyokhin, A. & Sewell, G. (2004) Changes in a lady beetle community following the
272 establishment of three alien species. *Biological Invasions*, **6**, 463-471.

273

274 Arroyo, M.T.K. & Cavieres, L.A. (2013) High-Elevation Andean Ecosystems.
275 *Encyclopedia of Biodiversity* (Second Edition) (ed by Levin, S.A.) pp. 96-110. Academic
276 Press, San Diego.

277

278 Barahona-Segovia, R.M., Grez, A.A. & Bozinovic, F. (2016) Testing the hypothesis of
279 greater invasive eurythermality in ladybugs: from physiological performance to life
280 history strategies. *Ecological Entomology*, **41**, 182-191.

281

282 Benelli, M., Leather, S.R., Francati, S., Marchetti, E. & Dindo, M.L. (2015) Effect of two
283 temperatures on biological traits and susceptibility to a pyrethroid insecticide in an
284 exotic and native coccinellid species. *Bulletin of Insectology*, **68**, 23-29.

285

286 Brown, P. M., Frost, R., Doberski, J., Sparks, T. I. M., Harrington, R. & Roy, H. E.
287 (2011) Decline in native ladybirds in response to the arrival of *Harmonia axyridis*: early
288 evidence from England. *Ecological Entomology*, **36**, 231-240.

289

290 Cavieres, L.A., Peñalosa, A., Papic, C. & Tambutti, M. (1998) Efecto nodriza del cojín
291 *Laretia acaulis* (Umbelliferae) en la zona alto-andina de Chile central. *Revista Chilena*
292 *de Historia Natural*, **71**, 337-347.

293

294 Di Castri, F. & Hajek, E.R. (1976) Bioclimatografía de Chile. Ediciones Universidad
295 Católica de Chile, Santiago, Chile.

296

297 Edwards, J.G. (1957) Entomology above timberline: II. The attraction of ladybird
298 beetles to mountain tops. *Coleopterists Bulletin*, **11**, 41-46.

299

300 Englund, R.A., Preston, D.J., Myers, S., Englund, L.L., Imada, C. & Evenhuis, N.L.
301 (2010) Results of the 2009 Alien Species and Wēkiu Bug (*Nysius wekiuicola*) Surveys
302 on the Summit of Mauna Kea, Hawaii Island. Final Report, Honolulu.

303

304 Grez, A.A. & Zaviezo, T. (2015) Chinita arlequín: *Harmonia axyridis* en Chile.
305 www.chinita-arlequin.uchile.cl.

306

307 Grez, A.A., Zaviezo, T., Roy, H., Brown, P.M.J. & Bizama, G. (2016) Rapid spread of
308 *Harmonia axyridis* in Chile and its effects on ladybeetle biodiversity. *Diversity and*
309 *Distributions*, **22**, 982-994.

310

311 Harmon, J.P., Stephens, E. & Losey, J. (2007) The decline of native ladybirds
312 (Coleoptera: Coccinellidae) in the United States and Canada. *Journal of Insect*
313 *Conservation*, **11**, 85–94.

314

315 Hoffmann, A., Kalin Arroyo, M., Liberona, F., Muñoz, M. & Watson, J. (1998) *Plantas*
316 *Altoandinas en la Flora Silvestre de Chile*. Ediciones Fundación Claudio Gay,
317 Santiago, Chile.
318

319 Lyons, K.G. & Schwartz, M.W. (2001) Rare species loss alters ecosystem function –
320 invasion resistance. *Ecology Letters*, **4**, 358-365.
321

322 Mittermeier, R.A., Turner, W.R., Larsen, F.W., Brooks, T.M. & Gascon, C. (2011)
323 Global biodiversity conservation: the critical role of hotspots. *Biodiversity hotspots:*
324 *Distribution and Protection of Conservation priority Areas* (ed by Zachos, F.E. &
325 Habekl, J.C.). pp. 3–22. Springer Publishers, London.
326

327 Molina-Montenegro, M.A., Briones, R. & Cavieres, L.A. (2009) Does global warming
328 induce segregation among alien and native beetle species in a mountain-top?
329 *Ecological Research*, **24**, 31-36.
330

331 Molina-Montenegro, M.A., Badano, E.I. & Cavieres, L.A. (2006) Cushion plants as
332 microclimatic shelters for two ladybird beetles species in Alpine zone of Central Chile.
333 *Arctic, Antarctic, and Alpine Research*, **38**, 224-227.
334

335 Rice, M.E. (1992) High altitude occurrence and westward expansion of the seven-
336 spotted lady beetle, *Coccinella septempunctata* (Coleoptera: Coccinellidae), in the
337 Rocky Mountains. *Coleopterists Bulletin*, **46**, 142-143.
338

339 Roy, H. E., Adriaens, T., Isaac, N. J., Kenis, M., Onkelinx, T., Martin, G. S., Brown,
340 P.M.J., Hautier, L., Poland, R., Roy, D.B., Comont, R., Eschen, R., Frost, R., Zindel, R.,
341 Van Vlaenderen, J., Nedved, O., Ravn, H.P., Gregoire, J-C., de Biseau, J-C. & Maes,

342 D. (2012) Invasive alien predator causes rapid declines of native European
343 ladybirds. *Diversity and Distributions*, **18**, 717-725.

344

345 Roy, H.E., Brown, P.M.J., Adriaens, T., Berkvens, N., Borges, I., Clusella-Trullas, S.,
346 De Clercq, P., Eschen, R., Estoup, A., Evans, E.W., Facon, B., Gardiner, M.M., Gil, A.,
347 Grez, A., Guillemaud, T., Haelewaters, D., Herz, A., Honek, A., Howe, A.G., Hui, C.,
348 Hutchinson, W.D., Kenis, M., Koch, R.L., Kulfan, J., Lawson Handley, L., Lombaert, E.,
349 Loomans, A., Losey, J., Lukashuk, A.O., Maes, D., Magro, A., Murray, K.M., San
350 Martin, G., Martinkova, Z., Minnaar, I., Nedved, O., Orlova-Bienkowskaja, M.J.,
351 Rabitsch, W., Ravn, H.P., Rondoni, G., Rorke, S.L., Ryndevich, S.K., Saethre, M-G.,
352 Sloggett, J.J., Soares, A.O., Stals, R., Tinsley, M.C., Vandereycken, A., van Wielink,
353 P., Vigišová, S., Zach, P., Zaviezo, T. & Zhao, Z. (2016) The harlequin ladybird,
354 *Harmonia axyridis*: an inspiration for global collaborations on invasion biology.
355 *Biological Invasions*, **18**, 997-1044.

356

357 Seebens, H., Essl, F. & Blasius, B. (2017) The intermediate distance hypothesis of
358 biological invasions. *Ecology Letters*, **20**, 158-165.

359

360 Shea, K. & Chesson, P. (2002) Community ecology theory as a framework for
361 biological invasions. *Trends in Ecology and Evolution*, **17**, 170-176.

362

363 Simberloff, D., Martin, J.L., Genovesi, P., Maris, V., Wardle, D.A., Aronson, J.,
364 Courchamp, F., Galil, B., García-Berthou, E., Pascal, M., Pyšek, P., Sousa, R.,
365 Tabacchi, E. & Vilà, M. (2013) Impacts of biological invasions: what's what and the way
366 forward. *Trends in Ecology and Evolution*, **28**, 58-66.

367

368 Stals, R. (2010) The establishment and rapid spread of an alien invasive lady beetle:
369 *Harmonia axyridis* (Coleoptera: Coccinellidae) in southern Africa, 2001–2009.
370 *IOBC/WPRS Bulletin*, **58**,125–132.
371
372 Stewart, J.W., Whitcomb, V.H. & Bell, K.O. (1967) Estivation studies of the convergent
373 lady beetle in Arkansas. *Journal of Economic Entomology*, **60**, 1730-1735.
374
375 Vilcinskis, A., Stoecker, K., Schmidtberg, H., Röhrich, C. R. & Vogel, H. (2013)
376 Invasive harlequin ladybird carries biological weapons against native competitors.
377 *Science*, **340**, 862-863.
378
379 Wang, S., Michaud, J.P., Tan, X.L., Zhang, F. & Guo, X.J. (2011) The aggregation
380 behavior of *Harmonia axyridis* in its native range in Northeast China. *BioControl*, **56**,
381 193-206.
382

383 **FIGURE CAPTIONS**

384

385 **Fig. 1** A) Surveyed habitat at Valle Nevado, Andes Mountains; B) *Harmonia axyridis*
386 with a larva of *Hippodamia convergens*; C) male *Hippodamia variegata* mating with a
387 female *Harmonia axyridis*; D) the native *Eriopsis chilensis*;

388



Table 1. *Harmonia axyridis* records from surveys on summer 2017 on an altitudinal gradient in the Andes Mountains, east of Santiago.

Date	Grid ref		<i>H. axyridis</i> abundance	Other coccinellid species
	(UTM WGS84 19S)	Altitude (m asl)		
29 January 2017	376341 E;	1952	0	> 10 <i>Adalia deficiens</i> , 1 <i>Eriopis chilensis</i>
	6308900 S			
	376721 E;	2200	0	1 <i>Hippodamia convergens</i>
	6308240 S			
	381665 E;	2467	0	>10 <i>Hippodamia convergens</i>
	6307303 S			
	382954 E;	2790	2	3 <i>Eriopis chilensis</i> , 2 <i>Hippodamia variegata</i> , 5 <i>Hippodamia convergens</i> .
	6308428 S			
	383788 E;	3025	1	
	6308770 S			
	384127 E;	3230	21	5-10 <i>Eriopis chilensis</i> , > 30 <i>Hippodamia variegata</i> , > 30 <i>Hippodamia convergens</i>
	6311547 S			
12 March 2017	383664 E;	3340	1	> 30 <i>Hippodamia convergens</i> ; > 10 <i>Hippodamia variegata</i> ; 5 <i>Eriopis chilensis</i>
	6311320 S			
	383574 E;	33508	9	> 30 <i>Hippodamia convergens</i> , > 30 <i>Hippodamia variegata</i> and 5 <i>Eriopis chilensis</i>
	6311097 S			

385411 E; 3578 3 > 10 *H. convergens*, > 10 *H. variegata*
6313115 S

