

ENVIRONMENTAL RESEARCH
LETTERS

LETTER

Global importance of methane emissions from drainage ditches and canals

OPEN ACCESS

RECEIVED
22 January 2021REVISED
25 February 2021ACCEPTED FOR PUBLICATION
2 March 2021PUBLISHED
15 March 2021

Original content from this work may be used under the terms of the [Creative Commons Attribution 4.0 licence](#).

Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.



M Peacock^{1,*}, J Audet², D Bastviken³, M N Futter¹, V Gauci^{4,11}, A Grinham⁵, J A Harrison⁶, M S Kent⁷, S Kosten⁸, C E Lovelock⁹, A J Veraart⁸ and C D Evans^{1,10}

¹ Department of Aquatic Sciences and Assessment, Swedish University of Agricultural Sciences, Uppsala 756 51, Sweden

² Department of Bioscience, Aarhus University, Vejløvej 25, Silkeborg 8600, Denmark

³ Department of Thematic Studies—Environmental Change, Linköping University, Linköping 581 83, Sweden

⁴ School of Geography, Earth and Environmental Sciences, University of Birmingham, Birmingham B15 2TT, United Kingdom

⁵ School of Civil Engineering, The University of Queensland, Brisbane 4072, Australia

⁶ School of the Environment, Washington State University, Vancouver Campus, Vancouver, WA 98686, United States of America

⁷ Environmental Science, School of Biosciences, University of Nottingham, Sutton Bonington Campus, Loughborough LE12 5RD, United Kingdom

⁸ Department of Aquatic Ecology and Environmental Biology, Institute for Water and Wetland Research, Radboud University, Nijmegen, The Netherlands

⁹ School of Biological Sciences, The University of Queensland, Brisbane 4072, Australia

¹⁰ Centre for Ecology & Hydrology, Environment Centre Wales, Bangor, Gwynedd LL57 2UW, United Kingdom

¹¹ Birmingham Institute of Forest Research, University of Birmingham, Birmingham, United Kingdom

* Author to whom any correspondence should be addressed.

E-mail: michael.peacock@slu.se and mikepeacocknin@yahoo.co.uk

Keywords: methane, ditch, greenhouse gas, drainage, land use

Supplementary material for this article is available [online](#)

Abstract

Globally, there are millions of kilometres of drainage ditches which have the potential to emit the powerful greenhouse gas methane (CH₄), but these emissions are not reported in budgets of inland waters or drained lands. Here, we synthesise data to show that ditches spanning a global latitudinal gradient and across different land uses emit large quantities of CH₄ to the atmosphere. Area-specific emissions are comparable to those from lakes, streams, reservoirs, and wetlands. While it is generally assumed that drainage negates terrestrial CH₄ emissions, we find that CH₄ emissions from ditches can, on average, offset ~10% of this reduction. Using global areas of drained land we show that ditches contribute 3.5 Tg CH₄ yr⁻¹ (0.6–10.5 Tg CH₄ yr⁻¹); equivalent to 0.2%–3% of global anthropogenic CH₄ emissions. A positive relationship between CH₄ emissions and temperature was found, and emissions were highest from eutrophic ditches. We advocate the inclusion of ditch emissions in national GHG inventories, as neglecting them can lead to incorrect conclusions concerning the impact of drainage-based land management on CH₄ budgets.

1. Introduction

Inland waters including lakes, rivers, wetlands and ponds play an important role in the global carbon cycle and climate system because they emit large amounts of greenhouse gases (GHGs) including methane (CH₄), carbon dioxide (CO₂), and nitrous oxide (N₂O) to the atmosphere [1–5]. CH₄ is an important GHG emitted from inland waters; this is because many waterbodies have high fluxes and because CH₄ has a 100 year global warming potential (GWP) of 28 [6] resulting in a substantial radiative forcing effect [4, 7–9].

Total inland water CH₄ emissions have been estimated as 100–200 Tg CH₄ yr⁻¹ [10, 11], equivalent to ~30% of global CH₄ emissions [11]. Globally and regionally important emissions of CH₄ have been documented from reservoirs [4], lakes [12], streams and rivers [13], small natural ponds [8], and artificial ponds [14]. It would therefore appear that CH₄ emissions from the major categories of inland waters have been quantified. However, emissions from another, potentially important category of inland waters have yet to be quantified at a global scale: ditches. Here, we present a synthesis of new and existing flux data from ditches showing that ditches release large amounts

of CH₄ and that these emissions are significant on national and global scales.

The Ramsar Convention includes ditches as human-made wetlands under the category ‘canals and drainage channels or ditches’ [15]. Artificial irrigation channels are also classified as ‘irrigated land’. However, there are no strict Ramsar definitions for these categories, which are a ‘broad framework to aid rapid identification of the main wetland habitats’ [15]. Ditches, canals and channels are created to: (a) improve the productivity of wet soils through drainage; (b) reclaim land from flooded areas; (c) move water through a landscape for agricultural or industrial use; (d) reduce agricultural soil erosion, or (e) remove stormwater in urban areas. For our purposes, we define ditches as constructed linear waterways, whilst acknowledging that their physical characteristics and function may vary widely (figure 1). Thus, irrigation channels and water supply canals in arid regions fall within our definition. We recognise that not all linear waterways included in our definition can strictly be described as drainage ditches in the traditional sense (i.e. they do not all drain land/lower the water table), however we use this term for simplicity, and because drainage ditches comprise the large majority of our dataset, as well as the majority of mapped linear water features globally (e.g. global ditch length is several orders of magnitude larger than global canal length; see SI text 9 (available online at stacks.iop.org/ERL/16/044010/mmedia)). Finally, it is worth noting that terminology is inconsistent and the words ‘ditch’ and ‘canal’ may even be used interchangeably; for example, larger drainage channels in SE Asian peatlands are navigable and often referred to locally as ‘canals’.

The spatial extent of ditches has been recognised on both small and large landscape scales. For instance, in a boreal catchment ditches doubled the length of the stream network [16], whilst in Great Britain the total length of ditches is estimated at ≈604 000 km which is more than twice that of streams and rivers combined (≈267 000 km) [17]. Data are scarce, but other estimates of drainage networks exist, and show that ditches can occupy a significant proportion of the waterscape in some countries and regions across the world [18–22].

Ditches possess particular characteristics which means that they cannot be assumed to function identically to most streams and rivers (although similarities may exist with streams in low-relief agricultural landscapes). Firstly, ditches do not in general follow natural topographic gradients and do not have natural catchments. They are often, but not always, situated in low-lying, low-relief landscapes which result in low flow rates. This creates a set of conditions favourable to CH₄ production and emission, namely: (a) the accumulation of sediment; (b) the development of anoxia; and (c) the growth of emergent

plants. Ditches within agricultural and urban landscapes may also receive high inputs of labile organic matter and nutrients, providing a substrate for methanogenesis [21]. In such environments, CH₄ is produced autochthonously in anaerobic bottom waters or sediments by methanogenesis at rates dependent on an array of factors including temperature and labile carbon content [23]. In addition, ditches may act as conduits for the emission of CH₄ which is produced under anaerobic conditions in adjacent terrestrial environments [24]. This process could be particularly important because, unlike other inland waters, ditches are specifically created *to drain* (i.e. to receive lateral transfers of water from adjacent land), and at an extremely high drainage density [20, 25] when compared to natural streams. In summary, ditches have a particular set of hydrological, chemical and morphological characteristics that differentiate them from natural streams in ways that favour CH₄ production and emission. It has been suggested that ditches may have higher CH₄:CO₂ ratios than streams, albeit based on a small dataset (see SI text 1) [13].

Given that CH₄ emissions from streams have been recognized for over 100 years [26, 27], there have been surprisingly few quantitative assessments of CH₄ fluxes from ditches until the last two decades. Roulet and Moore [24] performed one of the first studies, in which they suggested that peatland ditches ‘depending on the flow rate, depth, and morphology, could provide an ideal environment for the transport and *in situ* production of CH₄’. They found that drainage resulted in CH₄ uptake by the terrestrial peat surface, but that the ditches themselves emitted large amounts of CH₄. Thus, when scaled across the entire peatland they suggested that, depending on the spacing between ditches, drainage could result in a net increase in landscape CH₄ emissions. The potential of peatland ditches to emit CH₄ was confirmed by other studies [28, 29]. Ditches were subsequently incorporated into IPCC guidance for assessment of GHG fluxes from drained organic soils as part of the 2013 Wetlands Supplement [30]. This assessment identified a limited dataset of 19 publications reporting ditch CH₄ emissions from drained peat soils [31]. Because of the growing appreciation of the important role of CH₄ in regulating the global climate, the number of relevant studies has increased since 2013. However, to date no comprehensive, global-scale analysis of CH₄ emissions from ditches has been undertaken. Furthermore, whereas stream (and lake) emissions are a natural component of the global CH₄ budget (albeit subject to potential anthropogenic influences) ditches are constructed, therefore all associated CH₄ emissions must be considered anthropogenic. Our aim here is to draw attention to the potential importance of ditch fluxes in the global CH₄ cycle, and to highlight the lack of data from ditches which are



Figure 1. Photographs of ditches differing in design and function. Clockwise from top left: (1) ditch in a UK upland blanket bog; (2) an urban canal in the Netherlands; (3) a ditch in a felled peatland forest in Sweden; (4) a water management channel in an Indonesian Acacia plantation; (5) a rice paddy irrigation channel in Malaysia; (6) a freshly dug ditch on ex-Mega Rice Project land in Central Kalimantan; (7) a UK urban ditch under flood conditions.

a near ubiquitous landscape feature across climate zones and land use classes.

Here, we update and extend estimates of CH_4 emissions from ditches by including new literature and previously unpublished data. In total, we were able to calculate annual emissions for 64 unique ditch sites encompassing a wide range of ditch morphologies, latitudes and land uses. The majority of studies used the floating chamber method, and so we focused on these, and excluded a small number of flux

estimates based on dissolved CH_4 concentrations. For all studies we extracted data on land-use, eutrophication status, and climate.

2. CH_4 emissions from ditches: a synthesis

Reported CH_4 fluxes ranged from 0.1 to $386 \text{ g CH}_4 \text{ m}^{-2} \text{ yr}^{-1}$ with a mean (\pm standard error) of $64.6 \pm 11.1 \text{ g CH}_4 \text{ m}^{-2} \text{ yr}^{-1}$. Emissions from ditches in tropical and sub-tropical zones are




Figure 2. Mean annual CH₄ emissions from ditches (table SI1) grouped according to climate zone (panel A), eutrophication status (panel B) and in relation to mean annual temperature and nutrient status (panel C). For panel A, lowercase letters show significant differences: between boreal and sub/tropical (test statistic = -19.8 , $p = 0.004$) and temperate and sub/tropical (test statistic = -20.0 , $p = 0.001$). For panel B, Kruskal–Wallis shows no significance ($p = 0.089$). For panel C, Spearman correlation coefficient = 0.48 ($p < 0.001$). Orange circles = oligotrophic, yellow = mesotrophic, green = eutrophic. Note that CH₄ flux is presented on a log scale for panels A and B, but not C, and all statistical tests were performed on non-transformed data.

significantly higher than those from temperate and boreal zones (figure 2(A)) and higher fluxes are associated with higher temperatures, as shown by a significant correlation between CH₄ flux and temperature (figure 2(C)). Fluxes were significantly higher from continuously inundated ditches compared to intermittently dry ditches which can sometimes show negative fluxes (SI figure 1, SI text 8). There was no significant ($p = 0.09$) effect of nutrient status, but at sites where mean annual temperature (MAT) was above 15 °C eutrophic fluxes were noticeably higher (figure 2). Fluxes from eutrophic ditches also spanned

the greatest range, with some eutrophic ditches exhibiting very low fluxes.

Despite variation within each land-use category, mean fluxes tended to increase with increasing land use intensity in the order: natural land and forest \approx peat extraction < low-intensity grassland < urban land < cropland \approx high-intensity grassland (figure 3). Differences were significant between natural land and forest, and both cropland and high-intensity grassland.

More flux estimates were obtained from ditches draining peat soils ($n = 50$) compared to mineral

soils ($n = 14$). There was no significant difference between the two groups, with means of $58 \pm 11 \text{ g CH}_4 \text{ m}^{-2} \text{ yr}^{-1}$ and $87 \pm 33 \text{ g CH}_4 \text{ m}^{-2} \text{ yr}^{-1}$ for ditches draining peat and mineral soils, respectively (Mann–Whitney U test, $p = 0.91$). We compared these ditch emissions against the reduction in terrestrial emissions following drainage, assuming that ditch surface areas occupy, on average, 3% of the landscape (so called $\text{Frac}_{\text{ditch}} = 0.03$), and find that, on average, ditch emissions equate to 4% (error = 2%–8%) and 18% (15%–22%) of the terrestrial CH_4 reduction in mineral and peat soils respectively. The discrepancy in percentages between soil types arises primarily due to the large IPCC emission factors given for wetlands on mineral soils, which are substantially larger than fluxes from undrained peatlands.

3. Ditch emissions and upscaling

A comparison shows that, on an areal basis, ditches emit similar amounts of CH_4 to reservoirs, streams and tropical wetlands (figure 4).

Ditches have been mapped and total lengths calculated for some countries [18, 22], but for many regions of the world there is a lack of data on ditch densities. Additionally, although our synthesised CH_4 data covers a reasonable geographic spread, it is biased towards European and North American locations. Considering this, a detailed upscaling of global CH_4 emissions (i.e. aggregating by land use, soil type, climate zone, etc) from ditches would be flawed. Instead, and with the aim of calling attention to the possible global magnitude of these emissions, we present a rough ‘back of the envelope’ upscaling. For this, we took the average annual flux from our dataset and a global estimate of drained land area, and calculated total ditch surface areas using values of drainage ditch areas from the IPCC [30] and literature searches. This gives a mean flux of 3.46 Tg yr^{-1} ($0.61\text{--}10.5 \text{ Tg yr}^{-1}$) (table 1).

4. Discussion

Our analysis shows that mean areal ditch CH_4 emissions are as large as those from other well characterised systems (inland waters and wetlands) that are considered to be important at national or global scales. Our global estimate is 3.5 Tg yr^{-1} ($0.6\text{--}10.5 \text{ Tg yr}^{-1}$) which is 0.1%–1.8% of global CH_4 emissions or 1.0% (0.2%–2.9%) of global anthropogenic CH_4 [11]. For comparison, our estimate is greater than emissions from permafrost soils (1 Tg yr^{-1}) [11] which receive considerably more attention compared to ‘dull as ditchwater’ drainage channels. On a 20 year and 100 year time frame using a sustained-flux GWP [32], global emissions respectively equate to 332 and 156 $\text{Tg CO}_2 \text{ eq yr}^{-1}$. Thus, ditch emissions shift the balance and reduce

the expected CH_4 sink from drained soils, which is estimated globally as $38 \text{ Tg CH}_4 \text{ yr}^{-1}$ for all unsaturated oxic soils [11]. Ditch emissions could be even more important on a national scale. For example, in densely drained countries such as the Netherlands and Finland ditches could emit the equivalent of 7% and 9% of national anthropogenic CH_4 emissions (see SI text 10).

Multiple studies now show that ditch emissions can dominate the CH_4 budget of drained landscapes and, in some situations (e.g. if ditch density is high, or within-field water table is low) can outweigh any terrestrial CH_4 uptake [33–35]. Evidence of high emissions from ditches within drained landscapes counters the commonly held view, previously embedded in IPCC land-use emissions reporting guidance [36], that drained landscapes have zero CH_4 emissions (see SI text 11). Furthermore, a recent review of cropland drainage ignored the emissions from ditches and concluded that drainage could lead to a net reduction in CO_2 equivalents, and thus have a beneficial impact on climatic warming [37] (although wetland drainage can enhance terrestrial CO_2 and N_2O emissions [30]). We acknowledge that ditch emissions can be implicitly included in landscape budgets measured by eddy covariance towers, assuming that the ditch network within the tower footprint is representative of that in the wider landscape, but assume this is an uncommon occurrence (see SI text 11). Instead, we have shown that omission of ditch CH_4 emissions from landscape-scale GHG budgets will lead to underestimation of the overall global warming impact of drainage-based land management.

We found that ditch emissions increased with rising MAT, and that emissions were significantly higher in sub-tropical/tropical climate zones, reflecting the temperature-dependence of CH_4 production [38]. Higher fluxes were also associated with more intensive land uses, such as cropland. Annual fluxes were lower from ditches that periodically dried out, and this is likely due to sediments become aerobic and inhibiting methanogenesis [23]. For sites with MAT above 15°C , eutrophic ditches had higher fluxes but, surprisingly, no significant difference ($p = 0.09$) was found between mean fluxes from different nutrient statuses (although this may be due to a relatively small sample size resulting in an underpowered test). Taken together, we infer that CH_4 emissions will tend to increase with temperature, in the absence of other biogeochemical constraints such as intermittent drying, low substrate supply, or a low nutrient status.

We consider our upscaled estimate to be conservative for the following reasons. Firstly, we did not quantify channel area in irrigated land or ditches in urban land. Secondly, our estimate is based on chamber measurements which typically do not include plant mediated fluxes, and may not capture ebullitive fluxes (see SI text 2) which are temporally and spatially highly variable [39]. We also note that ditches




Table 1. Estimated lower, middle and upper global CH₄ emissions from drainage ditches. Lower and upper CH₄ bounds are based on 95% CIs for flux. $Frac_{ditch}$ is a measure of the proportion of the landscape area occupied by ditches, effectively a function of ditch spacing, configuration and width [31]. $Frac_{ditch}$ estimates and areas of drained land are taken from the literature. See section 5 for further detail, including references.

	Lower	Middle	Upper
Ditch flux (g CH ₄ m ⁻² y ⁻¹)	42.7	64.6	97.7
Total drained land (ha)	142 102 750	178 447 500	214 792 250
$Frac_{ditch}$	0.01	0.03	0.05
Total ditch area (ha)	1 421 028	5 353 425	10 739 613
Total ditch CH ₄ emission (Tg yr ⁻¹)	0.6	3.5	10.5
Global CH ₄ emission (Tg yr ⁻¹)	572	572	572
% global flux from ditches	0.1	0.6	1.8

can emit CO₂ and N₂O [40] which further increase GHG forcing associated with drained landscapes [41]. N₂O emissions may be particularly large from ditches in drained wetlands that have been converted to croplands and subjected to extensive nitrogen fertilization. However, at this stage, there are insufficient data to enable these emissions of CO₂ and N₂O to be upscaled [42].

Our work also carries limitations. Firstly, our synthesised data is biased towards ditches draining peat soils ($n = 50$) compared to mineral soils ($n = 14$). Thus, if there is a difference in ditch fluxes

between the two soil groups our upscaling could also be biased. There was no significant difference in mean fluxes between the two groups and we also found studies where, although annual fluxes could not be calculated, there was evidence for large CH₄ emissions from ditches in mineral soils. For example, an extremely high flux of 250 g CH₄ m⁻² d⁻¹ was reported from a ditch on a dairy farm in the Netherlands [43] and a UK survey detected a biogenic CH₄ hotspot near an urban canal [44]. There is therefore no reason to assume that emissions from mineral soils are smaller than those from peat soils,

but future measurements of ditch fluxes on mineral soils should be a key priority to robustly test this assumption. It is feasible, however, that CH₄ production pathways may vary: *in situ* production may dominate in well-drained mineral soils, whilst lateral inputs of CH₄ into ditches are likely to also contribute in poorly-drained soils. A second limitation is the lack of information on ditch lengths and maps of drained land, which leads to a large uncertainty when upscaling. Note that the largest uncertainty in our calculations comes from ditch area: our upper estimate is ≈ 7.5 times larger than the lower, whilst for ditch CH₄ flux the upper estimate is ≈ 2.3 times larger than the lower. Drained agricultural land, the majority of which is on mineral soils, occupies the largest cumulative area of drained land, but few measurements of $\text{Frac}_{\text{ditch}}$ exist for this land (see SI text 7). Mapping ditch networks within this land should therefore be another key priority for future work. Thirdly, there is an absence of nighttime ditch CH₄ measurements. It has been shown that CH₄ concentrations in channelized eutrophic streams [45] and ditches [46] can be approximately double those of daytime concentrations, due to diel fluctuations in oxygen concentrations. If this is a widespread phenomenon in ditches then our calculated annual fluxes may be low. Fourthly, ditch CH₄ measurements in boreal regions are often restricted to the growing season (SI table 1). Non-growing season emissions from shallow aquatic systems, marshes and peatlands in the boreal comprise, on average, 16% of annual emissions [47], and thus annual CH₄ emissions from ditches may be underestimated in our synthesis. Finally, the effect of aquatic plants is uncertain. Emergent vegetation in streams can act as a pathway for CH₄ emission and enhance sediment CH₄ concentrations, whilst floating plant species can act to increase or decrease net aquatic CH₄ emission depending on local conditions [48].

Increasing global populations and associated food and fibre demands are likely to lead to increased land drainage, ditch density and nutrient loadings [37]. Our results suggest that this will lead to significant increases in global ditch CH₄ emissions. Ditch emissions may also increase under a warmer climate. However, there are opportunities to mitigate these emissions. Nutrient enrichment increases CH₄ fluxes in other aquatic ecosystems such as lakes and reservoirs [4, 7], and there is no reason to assume that this effect does not occur in ditches. Reducing nutrient runoff into ditches may offer a pathway to mitigate CH₄ emissions to the atmosphere, whilst also improving water quality, thereby enhancing human and aquatic ecosystem health. Adaptive design and maintenance of ditch networks might also be an option to limit emissions. For example, clearing vegetation would prevent plant-mediated emissions and additionally remove a source of labile carbon for methanogens, although we acknowledge a

conflict here with the biodiversity value provided by ditch vegetation. Maintaining deep, stable water levels may also limit emissions by minimising the warming of sediments and thereby reducing CH₄ production.

In some regions rewetting of drained wetlands is being pursued as a climate change mitigation measure. Rewetted sites often possess ditch networks that may be partially blocked or infilled, or that are still used for active hydrological management, and high CH₄ fluxes have been measured from such ditches [49, 50]. If CH₄ emissions from relict ditches are of a similar magnitude to emissions from the adjacent wetland area, as assumed by the IPCC [30], then this is not a cause for concern. However, for intermediately rewetted systems such as wet meadows and paludiculture, it is feasible that ditch emissions may make a significant contribution to the overall GHG balance of the restored landscape. At present, a lack of data means that no conclusions can be drawn on this topic, but it is a highly relevant area of scientific and policy uncertainty.

5. Materials and methods

5.1. Experimental design

Data on emissions of CH₄ from ditches were collected by searching the published literature (including grey literature) and by collating unpublished data. Some studies did not use the term 'ditches' but maps or written descriptions suggested that this term was appropriate. For example, some studies measured 'streams' [51] or 'rivers' [52]. Contact with the authors confirmed which of their waterbodies could be regarded as ditches. In total, we gathered 52 studies and from these we extracted 75 flux estimates disaggregated by site. Different studies took different approaches in the number and type of sites measured. For instance, if one study reported data from two sites under different land-uses then these were assigned to their two respective categories. However, other studies sometimes measured more than one site under the same land-use category, but reported a single annual flux. For these studies, only one annual flux could be extracted and used in the analysis. For our synthesis, we adopted and expanded the categories of land-use from Evans [31]: natural land and forest ($n = 21$), peat extraction ($n = 8$), low-intensity grassland ($n = 9$), high-intensity grassland ($n = 9$), cropland ($n = 11$), and urban ($n = 6$). In many cases studies reported an annual flux, or an average flux that was representative for the annual period, which we then extracted, but for some sites we had to calculate annual fluxes using other methods (see SI text 3 and 5). Routine sampling can miss periods of high discharge which have been shown to be important hot moments for CH₄ emissions from streams, and more so the greater the stream channel slope [53], but we assume that ditches generally have lower slopes than streams, due to the fact that waterlogged land (the

typical target area for drainage ditches) occupies flatter landscapes. We categorized sites by trophic status using a published method [54] (see SI text 4) and obtained MAT from each study. If MAT was not given, we located the field site in Google Earth, and searched for the nearest town/city where we could find proxy MAT data online.

5.2. Comparison of terrestrial CH₄ reduction and ditch CH₄ emissions

We took the mean ditch emissions from our synthesis of 58 ± 11 g CH₄ m⁻² yr⁻¹ and 87 ± 33 g CH₄ m⁻² yr⁻¹ for ditches draining peat and mineral soils. We then calculated the mean drainage-induced decrease in terrestrial CH₄ emissions. For undrained peat soils we assumed mean terrestrial emissions of 16 g CH₄ m⁻² yr⁻¹ and 5.5 g CH₄ m⁻² yr⁻¹ for northern [55] and tropical peatlands [30], respectively, giving an overall undrained peat mean of 10.7 g CH₄ m⁻² yr⁻¹. From this we subtracted the mean emission from drained organic soils (excluding rice), given by the IPCC as 0.8 g CH₄ m⁻² yr⁻¹, resulting in a drainage-induced mean terrestrial decrease of 9.9 g CH₄ m⁻² yr⁻¹. For mineral soils we used the IPCC [30] data for rewetted inland mineral soils across all climate zones as a proxy for mean emission from undrained wetlands on mineral soils (note that the majority, $n = 32$, of these data are actually from natural wetlands, with $n = 7$ from created/restored wetlands), but we split the data into continuously inundated wetlands (mean = 96.5 g CH₄ m⁻² yr⁻¹) and intermittently flooded wetlands (mean = 48.7 g CH₄ m⁻² yr⁻¹). The IPCC [30] assumption is that these soils no longer emit CH₄ when drained, therefore we assume that drainage leads to a complete cessation of terrestrial CH₄ emissions. We then applied an average Frac_{ditch} value of 0.03 (see 'Global and National Ditch Emissions' discussion on Frac_{ditch}), with corresponding terrestrial land fraction as 0.97. The reduction in terrestrial flux, and ditch emission, are then weighted according to their respective fractions. This gives weighted ditch emissions of 1.4–2.1 g CH₄ m⁻² yr⁻¹ and 1.6–3.6 g CH₄ m⁻² yr⁻¹ for peat and mineral soils, respectively. When expressed as percentages offsetting the drainage-induced terrestrial reduction, the means and errors are: peat soil = 18% (15%–22%); drained intermittently flooded wetland on mineral soil = 5.5% (3.4%–7.6%); drained continuously inundated wetland on mineral soil = 2.8% (1.7%–3.8%).

5.3. Global and national ditch emissions

We used Finland and the Netherlands as case studies to estimate the importance of ditch emissions in densely drained countries. These countries were chosen as information is available on ditch length/drainage densities and CH₄ fluxes. We upscaled by multiplying total ditch surface area, calculated using

literature values [22, 56], by the mean ditch flux measured in each respective country (see SI text 10). Upscaled emissions were compared against national anthropogenic CH₄ emissions [57].


For our global upscaling we combined estimates of CH₄ emission, drained land area, and ditch surface area. We calculated total drained land as the sum of drained high-latitude peatland forestry (13 447 500 ha [58, 59]) and drained agricultural land (165 000 000 ha [37, 60]; smaller than other estimates, e.g. 190 000 000 ha [61]). Drained tropical peatland covers a moderately large area (7750 000 ha [62]) but the majority of this is used for oil palm (3055 000 ha) and smallholder farmland (3321 000 ha) [62]. To avoid double-counting we thus assume drained tropical peatland area is captured under total drained agricultural land. We used a 10% error margin on the area of drained high-latitude peatland forest land for upper and lower errors. For drained agricultural land we took lower and upper values of 130 000 000 ha and 200 000 000 from a recent study [37].

The fraction of ditches (Frac_{ditch}) is a measure of the proportion of the landscape area occupied by ditches/canals, effectively a function of ditch spacing, configuration and width [31]. Values of Frac_{ditch} were collated for peatland soils by the IPCC [30] in 2014. We searched the literature for more peatland values of Frac_{ditch} published since then, or that had been overlooked, and found 14 more values which we combined with the IPCC dataset, giving a total of 37 Frac_{ditch} values (SI table 4). Data for mineral soils are scarce and depend on studies reporting both drainage densities and ditch widths. We searched the literature and managed to collate 11 Frac_{ditch} values for mineral soils from nine studies (SI text 7, SI table 2). Frac_{ditch} ranged from 0.004 to 0.1, with most values being 0.01–0.03. For our upscaling we took 0.03 as an average Frac_{ditch}, and used 0.01 and 0.05 as lower and upper bounds.

For our emission estimate we took the mean ditch CH₄ flux from the studies we collated (SI table 1). For our upper and lower ditch flux estimates, we used 95% confidence intervals calculated on the log-transformed CH₄ flux data. We compared the global ditch emission against a global CH₄ emission of 572 Tg yr⁻¹ and a global anthropogenic emission of 357 Tg yr⁻¹ [11].

5.4. Statistical analysis

Statistical analysis was performed using SPSS. We used the nonparametric Kruskal–Wallis test (2 sided), with inbuilt post hoc test, to test for differences in ditch flux between climate zone, land use, and nutrient status. For the land use test (figure 3) there were many potential pair wise comparisons. If all these were performed then any post-hoc correction would be overly conservative. We therefore used figure 3 to guide which comparisons to make, by looking at categories where visual differences (i.e. no/minimal



overlap between fluxes) were apparent. Following this, we performed post-hoc tests for forest/semi natural vs cropland, and forest vs high intensity, with no correction. For the correlation between MAT and flux (figure 2(C)) we used Spearman correlation coefficients to test for monotonic relationships between variables.

Data availability statement

All data that support the findings of this study are included within the article (and any supplementary files).

Acknowledgments

No specific grant funded this work. MP acknowledges funding from Formas (Grant 2020-00950). DB contributions were supported by European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (Grant Agreement No. 725546). JA acknowledges funding from Carl Tryggers Stiftelse. VG is grateful for support from the AXA Research Fund. We thank Marcus Klaus and Dongqi Wang for providing information on the sampling sites in their respective papers, and Frances Manning for providing data ahead of publication. We thank Jesper Rydén and Gareth Harvey for their advice on different approaches to statistically analysing the data. We thank Andy Baird and Sophie Green for providing ebullition data that were used in

an earlier draft of this paper. We thank one anonymous reviewer and Tim Moore for their constructive review comments that helped to improve the manuscript.

Author contributions

The initial collation of ditch emission data was undertaken by CE as part of his contribution to the 2013 IPCC Wetland Supplement (Inland drained organic soils chapter), which was expanded by CM and MP for the Flooded Lands chapter of the Intergovernmental Panel on Climate Change 2019 Refinement to the 2006 Guidelines for National Greenhouse Gas Inventories (43), for which CL, CE, JH and DB were Lead Authors, and MP, SK and AG were Contributing Authors. MP, JA, AG, MK, SK, AJV contributed data. AG and MP created the figures. All authors searched for literature data and provided ideas. MP wrote the paper with input from all co-authors. The authors declare no competing interests.

ORCID iDs

M Peacock <https://orcid.org/0000-0002-3086-2854>

J Audet <https://orcid.org/0000-0001-5839-8793>

D Bastviken <https://orcid.org/0000-0003-0038-2152>

V Gauci <https://orcid.org/0000-0002-2452-7291>

J A Harrison  <https://orcid.org/0000-0002-0677-5478>

S Kosten  <https://orcid.org/0000-0003-2031-0965>

References

- [1] Battin T J, Luysaert S, Kaplan L A, Aufdenkampe A K, Richter A and Tranvik L J 2009 The boundless carbon cycle *Nat. Geosci.* **2** 598–600
- [2] Ciais P et al 2013 Carbon and other biogeochemical cycles *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* ed T F Stocker, D Qin, G-K Plattner, M Tignor, S K Allen, J Boschung, A Nauels, Y Xia, V Bex and P M Midgley (Cambridge: Cambridge University Press) 465–570
- [3] Cole J J et al 2007 Plumbing the global carbon cycle: integrating inland waters into the terrestrial carbon budget *Ecosystems* **10** 172–85
- [4] Deemer B R, Harrison J A, Li S, Beaulieu J J, DelSontro T, Barros N, Bezerra-Neto J F, Powers S M, Dos Santos M A and Vonk J A 2016 Greenhouse gas emissions from reservoir water surfaces: a new global synthesis *Bioscience* **66** 949–64
- [5] Maavara T, Lauerwald R, Laruelle G G, Akbarzadeh Z, Bouskill N J, Van Cappellen P and Regnier P 2019 Nitrous oxide emissions from inland waters: are IPCC estimates too high? *Glob. Change Biol.* **25** 473–88
- [6] Myhre G et al 2013 Anthropogenic and natural radiative forcing *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* ed T Stocker, D Qin, G-K Plattner, M Tignor, S Allen, J Boschung, A Nauels, Y Xia, V Bex and P Midgley (Cambridge: Cambridge University Press) 659–740
- [7] DelSontro T, Beaulieu J J and Downing J A 2018 Greenhouse gas emissions from lakes and impoundments: upscaling in the face of global change *Limnol. Oceanogr. Lett.* **3** 64–75
- [8] Holgerson M A and Raymond P A 2016 Large contribution to inland water CO₂ and CH₄ emissions from very small ponds *Nat. Geosci.* **9** 222–6
- [9] Selvam B P, Natchimuthu S, Arunachalam L and Bastviken D 2014 Methane and carbon dioxide emissions from inland waters in India—implications for large scale greenhouse gas balances *Glob. Change Biol.* **20** 3397–407
- [10] Bastviken D, Tranvik L J, Downing J A, Crill P M and Enrich-Prast A 2011 Freshwater methane emissions offset the continental carbon sink *Science* **331** 50
- [11] Saunio M et al 2020 The global methane budget 2000–2017 *Earth Syst. Sci. Data* **12** 1561–623 (<https://essd.copernicus.org/articles/12/1561/2020/>)
- [12] Bastviken D, Cole J, Pace M and Tranvik L 2004 Methane emissions from lakes: dependence of lake characteristics, two regional assessments, and a global estimate *Glob. Biogeochem. Cycles* **18** GB4009
- [13] Stanley E H, Casson N J, Christel S T, Crawford J T, Loken L C and Oliver S K 2016 The ecology of methane in streams and rivers: patterns, controls, and global significance *Ecol. Monogr.* **86** 146–71
- [14] Grinham A, Albert S, Deering N, Dunbabin M, Bastviken D, Sherman B, Lovelock C E and Evans C D 2018 The importance of small artificial water bodies as sources of methane emissions in Queensland, Australia *Hydrol. Earth Syst. Sci.* **22** 5281–98
- [15] Ramsar 2009 Ramsar classification system for wetland type. Information sheet on Ramsar wetlands, Annex 1 (available at: http://archive.ramsar.org/pdf/ris/key_ris_e.pdf)
- [16] Hasselquist E M, Lidberg W, Sponseller R A, Ågren A and Laudon H 2018 Identifying and assessing the potential hydrological function of past artificial forest drainage *Ambio* **47** 546–56
- [17] Brown C D, Turner N, Hollis J, Bellamy P, Biggs J, Williams P, Arnold D, Pepper T and Maund S 2006 Morphological and physico-chemical properties of British aquatic habitats potentially exposed to pesticides *Agric. Ecosyst. Environ.* **113** 307–19
- [18] Ågren A M and Lidberg W 2019 The importance of better mapping of stream networks using high resolution DEMs *Hydrol. Earth Syst. Sci. Discuss.* 1–20 (<https://hess.copernicus.org/preprints/hess-2019-34/hess-2019-34>)
- [19] Fang J, Yingjie M and Zili F 2002 A study on salt pollution cycle of cultivated land between drainage and irrigation in the main stream of Tarim River, Xinjiang, China *Prospects for Saline Agriculture (Tasks for Vegetation Science vol 37)* ed R Ahmad and K A Malik (Berlin: Springer) 37–42 (https://link.springer.com/chapter/10.1007/978-94-017-0067-2_5)
- [20] Herzon I and Helenius J 2008 Agricultural drainage ditches, their biological importance and functioning *Biol. Conserv.* **141** 1171–83
- [21] Needelman B A, Kleinman P J, Strock J S and Allen A L 2007 Drainage ditches improved management of agricultural drainage ditches for water quality protection: an overview *J. Soil Water Conserv.* **62** 171–8
- [22] Verdonschot R C, Keizer-Vlek H E and Verdonschot P F 2011 Biodiversity value of agricultural drainage ditches: a comparative analysis of the aquatic invertebrate fauna of ditches and small lakes *Aquat. Conserv. Mar. Freshwater Ecosyst.* **21** 715–27
- [23] Segers R 1998 Methane production and methane consumption: a review of processes underlying wetland methane fluxes *Biogeochemistry* **41** 23–51
- [24] Roulet N T and Moore T R 1995 The effect of forestry drainage practices on the emission of methane from northern peatlands *Can. J. For. Res.* **25** 491–9
- [25] Levvasseur F, Lagacherie P, Bailly J S, Biarnès A and Colin F 2015 Spatial modeling of man-made drainage density of agricultural landscapes *J. Land Use Sci.* **10** 256–76
- [26] Browne C A 1925 Thomas Paine's theory of atmospheric contagion and his account of an experiment performed by George Washington upon the production of marsh gas *J. Chem. Educ.* **2** 99–101
- [27] Phillips F C 1985 On the possibility of the occurrence of hydrogen and methane in the atmosphere *J. Am. Chem. Soc.* **107** 801–9
- [28] Best E P and Jacobs F H 1997 The influence of raised water table levels on carbon dioxide and methane production in ditch-dissected peat grasslands in the Netherlands *Ecol. Eng.* **8** 129–44
- [29] Van Den Pol-van Dasselaar A, Van Beusichem M L and Oenema O 1999 Methane emissions from wet grasslands on peat soil in a nature preserve *Biogeochemistry* **44** 205–20
- [30] IPCC 2014 2013 Supplement to the 2006 IPCC Guidelines for National Greenhouse Gas Inventories: Wetlands ed T Hiraishi, T Krug, K Tanabe, N Srivastava, J Baasansuren, M Fukuda and T G Troxler (Geneva: Intergovernmental Panel on Climate Change)
- [31] Evans C D, Renou-Wilson F and Strack M 2016 The role of waterborne carbon in the greenhouse gas balance of drained and re-wetted peatlands *Aquat. Sci.* **78** 573–90
- [32] Neubauer S C and Megonigal J P 2015 Moving beyond global warming potentials to quantify the climatic role of ecosystems *Ecosystems* **18** 1000–13
- [33] Korhonen M, Tuovinen J P, Aurela M, Koskinen M, Minkkinen K, Ojanen P, Penttilä T, Rainne J, Laurila T and Lohila A 2017 Methane exchange at the peatland forest floor—automatic chamber system exposes the dynamics of small fluxes *Biogeosciences* **14** 1947–67
- [34] Peacock M, Ridley L M, Evans C D and Gauci V 2017 Management effects on greenhouse gas dynamics in fen ditches *Sci. Total Environ.* **578** 601–12
- [35] Teh Y A, Silver W L, Sonnentag O, Detto M, Kelly M and Baldocchi D D 2011 Large greenhouse gas emissions from a temperate peatland pasture *Ecosystems* **14** 311–25

- [36] IPCC 2006 *2006 IPCC Guidelines for National Greenhouse Gas Inventories* ed H S Eggleston, L Buendia, K Miwa, T Ngara and K Tanabe (Tokyo: Intergovernmental Panel on Climate Change, IGES)
- [37] Castellano M J, Archontoulis S V, Helmers M J, Poffenbarger H J and Six J 2019 Sustainable intensification of agricultural drainage *Nat. Sustain.* **2** 914–21
- [38] Yvon-Durocher G, Allen A P, Bastviken D, Conrad R, Gudasz C, St-Pierre A, Thanh-Duc N and Del Giorgio P A 2014 Methane fluxes show consistent temperature dependence across microbial to ecosystem scales *Nature* **507** 488–91
- [39] Wik M, Thornton B F, Bastviken D, Uhlbäck J and Crill P M 2016 Biased sampling of methane release from northern lakes: a problem for extrapolation *Geophys. Res. Lett.* **43** 1256–62
- [40] McPhillips L E, Groffman P M, Schneider R L and Walter M T 2016 Nutrient cycling in grassed roadside ditches and lawns in a suburban watershed *J. Environ. Qual.* **45** 1901–9
- [41] Audet J, Wallin M B, Kyllmar K, Andersson S and Bishop K 2017 Nitrous oxide emissions from streams in a Swedish agricultural catchment *Agric. Ecosyst. Environ.* **236** 295–303
- [42] IPCC 2019 Agriculture, forestry and other land use (AFOLU) *2019 Refinement to the 2006 IPCC Guidelines for National Greenhouse Gas Inventories* ed C E Lovelock et al vol 4 (Geneva: Intergovernmental Panel on Climate Change)
- [43] Hensen A, Groot T T, Van Den Bulk W C, Vermeulen A T, Olesen J E and Schelde K 2006 Dairy farm CH₄ and N₂O emissions, from one square metre to the full farm scale *Agric. Ecosyst. Environ.* **112** 146–52
- [44] Zazzeri G, Lowry D, Fisher R E, France J L, Lanouisellé M, Grimmond C S and Nisbet E G 2017 Evaluating methane inventories by isotopic analysis in the London region *Sci. Rep.* **7** 4854
- [45] Harrison J A, Matson P A and Fendorf S E 2005 Effects of a diel oxygen cycle on nitrogen transformations and greenhouse gas emissions in a eutrophied subtropical stream *Aquat. Sci.* **67** 308–15
- [46] Deng O et al 2020 Emission of CO₂ and CH₄ from a multi-ditches system in rice cultivation region: fluxtemporal-spatial variation and effect factors *J. Environ. Manage.* **270** 110918
- [47] Treat C C, Bloom A A and Marushchak M E 2018 Nongrowing season methane emissions—a significant component of annual emissions across northern ecosystems *Glob. Change Biol.* **24** 3331–43
- [48] Kosten S, Piñeiro M, De Goede E, De Klein J, Lamers L P and Ettwig K 2016 Fate of methane in aquatic systems dominated by free-floating plants *Water Res.* **104** 200–7
- [49] Chamberlain S D, Boughton E H and Sparks J P 2015 Underlying ecosystem emissions exceed cattle-emitted methane from subtropical lowland pastures *Ecosystems* **18** 933–45
- [50] Cooper M D, Evans C D, Zielinski P, Levy P E, Gray A, Peacock M, Norris D, Fenner N and Freeman C 2014 Infilled ditches are hotspots of landscape methane flux following peatland re-wetting *Ecosystems* **17** 1227–41
- [51] Klaus M, Geibrink E, Jonsson A, Bergstrom A K, Bastviken D, Laudon H, Klaminder J and Karlsson J 2018 Greenhouse gas emissions from boreal inland waters unchanged after forest harvesting *Biogeosciences* **15** 5575–94
- [52] Yu Z, Wang D, Li Y, Deng H, Hu B, Ye M, Zhou X, Da L, Chen Z and Xu S 2017 Carbon dioxide and methane dynamics in a human-dominated lowland coastal river network (Shanghai, China) *J. Geophys. Res.: Biogeosci.* **122** 1738–58
- [53] Natchimuthu S, Wallin M B, Klemetsson L and Bastviken D 2017 Spatio-temporal patterns of stream methane and carbon dioxide emissions in a hemiboreal catchment in Southwest Sweden *Sci. Rep.* **7** 39729
- [54] Evans C D, Futter M N, Moldan F, Valinia S, Frogbrook Z and Kothawala D N 2017 Variability in organic carbon reactivity across lake residence time and trophic gradients *Nat. Geosci.* **10** 832–5
- [55] Abdalla M, Hastings A, Truu J, Espenberg M, Mander Ü and Smith P 2016 Emissions of methane from northern peatlands: a review of management impacts and implications for future management options *Ecol. Evol.* **6** 7080–102
- [56] Peltomaa R 2007 Drainage of forests in Finland *Irrig. Drain.* **56** S151–9
- [57] Janssens-Maenhout G et al 2019 EDGAR v4.3.2 Global Atlas of the three major greenhouse gas emissions for the period 1970–2012 *Earth Syst. Sci. Data* **11** 959–1002
- [58] Paavilainen E and Päivänen J 1995 *Peatland Forestry: Ecology and Principles. Ecological Studies* vol 111 (Berlin: Springer)
- [59] Biancalani R and Avagyan A 2014 Towards climate-responsible peat-lands management *Mitigation of Climate Change in Agriculture* Series 9 (Rome: Food and Agriculture Organization of the United Nations) 100
- [60] Feick S, Siebert S, Döll P and Digital Global A 2005 Map of artificially drained agricultural areas Frankfurt Hydrology Paper 04 (Frankfurt am Main: Institute of Physical Geography, Frankfurt University)
- [61] Schultz B, Thatte C D and Labhsetwar V K 2005 Irrigation and drainage. Main contributors to global food production *Irrig. Drain.* **54** 263–78
- [62] Wijedasa L S, Sloan S, Page S E, Clements G R, Lupascu M and Evans T A 2018 Carbon emissions from South-East Asian peatlands will increase despite emission-reduction schemes *Glob. Change Biol.* **24** 4598–613
- [63] Sjögersten S, Black C R, Evers S, Hoyos-Santillan J, Wright E L and Turner B L 2014 Tropical wetlands: a missing link in the global carbon cycle? *Glob. Biogeochem. Cycles* **28** 1371–86