

OVERVIEW

Rewilding and restoring nature in a changing world

Benis N. Egoh¹*, Charity Nyelele¹, Karen D. Holl²‡, James M. Bullock³‡, Steve Carver⁴‡, Christopher J. Sandom⁵‡

1 Department of Earth System Science, University of California Irvine, Irvine, California, United States of America, **2** Environmental Studies Department, University of California, Santa Cruz, California, United States of America, **3** UK Centre for Ecology & Hydrology, Wallingford, Oxfordshire, United Kingdom, **4** School of Geography, University of Leeds, Leeds, United Kingdom, **5** School of Life Sciences, University of Sussex, Brighton, United Kingdom

☉ These authors contributed equally to this work.


✉ Current address: Sussex Sustainability Research Programme, Brighton, United Kingdom

‡ These authors also contributed equally to this work.

* ebenis@gmail.com, begoh@uci.edu

Increased anthropogenic pressure, invasive alien species and climate change, among other factors, continue to negatively impact and degrade the planet's ecosystems and natural environment. As nature declines at alarming rates, the loss of biodiversity is not only a huge concern, but it also undermines the many ecological, social, human health and wellbeing benefits nature provides us. Numerous reports, including those from the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services (IPBES, <https://www.ipbes.net/>), have documented this unprecedented decline in nature across space and time. For example, the 2019 IPBES global assessment report on biodiversity and ecosystem services shows that 75% of the global land surface has been significantly altered, 66% of the ocean area is experiencing increasing cumulative impacts, and over 85% of wetland area has been lost (Brondizio et al. [1]). All the recent IPBES reports from global to regional scales and the Millennium Ecosystem Assessment of 2005 (Reid et al. [2]), point to one thing: the urgency for us to act to save nature and humankind. Ecological restoration has emerged as a powerful approach to combat degradation in land and water, mitigate climate change, and restore lost biodiversity and key ecosystem functions and services. In June this year (2021), the United Nations (UN) is launching the Decade on Ecosystem Restoration (<https://www.decadeonrestoration.org/>), an ambitious program to trigger a global movement for restoring the world's ecosystems. In line with that, PLOS ONE commissioned this Collection on Rewilding and Restoration. This is consistent with the year's Earth Day theme, "Restore Our Earth", which calls on everyone to be a part of the change and to focus on natural processes, emerging green technologies and innovative thinking that can restore the world's ecosystems.

When PLOS ONE launched this Rewilding and Restoration collection, we were asked to identify exciting advances and emerging trends observed recently in the areas of rewilding and restoration. We highlight: 1) increasing recognition of the value of restoration in ecosystems worldwide, particularly in a time of rapid global environmental change; 2) understanding and incorporating benefits and beneficiaries in supporting and financing restoration initiatives; 3) exploring the theoretical underpinning for the importance of 'megabiota'—the largest plants and animals—for driving biosphere scale processes such as ecosystem total biomass, resource flows and fertility; and 4) showcasing success stories on how rewilding nature in the developing world is reversing the impact of invasive species (<https://everyone.plos.org/2020/08/28/taking-a-walk-on-the-wild-side/>). The broad range of publications in this Collection cover all



OPEN ACCESS

Citation: Egoh BN, Nyelele C, Holl KD, Bullock JM, Carver S, Sandom CJ (2021) Rewilding and restoring nature in a changing world. *PLoS ONE* 16(7): e0254249. <https://doi.org/10.1371/journal.pone.0254249>

Editor: RunGuo Zang, Chinese Academy of Forestry, CHINA

Published: July 14, 2021

Copyright: This is an open access article, free of all copyright, and may be freely reproduced, distributed, transmitted, modified, built upon, or otherwise used by anyone for any lawful purpose. The work is made available under the [Creative Commons CC0](https://creativecommons.org/licenses/by/4.0/) public domain dedication.

Data Availability Statement: All relevant data are within the paper.

Funding: NA

Competing interests: The authors have declared that no competing interests exist.

these areas and much more, making it one of the most exciting collections on rewilding and restoring nature in recent times. The two main themes that emerge from the collection are related to restoration success stories (>40%) and best practices in restoration around the globe (>30%). The selected studies in this Collection, which cover six continents and at least 13 countries, were carried out in diverse settings and contexts, such as marine, fresh water and terrestrial habitats including forests and grasslands, rivers and coastal areas, woodlands, wetlands, and mountains (e.g., Sansupa et al. [3], Broughton et al. [4], Schulz et al. [5], Ndangalasi et al. [6]). Features of interest included in this Collection span from bacteria through large vertebrates (e.g., wild dogs, elephants) to ecosystems and their functions. These articles also showcase a range of methodological approaches from a series of small-scale field experiments (Wasson et al. [7]), wildlife tracking and remote sensing (Mata et al. [8]), and large-scale models to predict restoration outcomes (D'Acunto et al. [9]).

This rich collection from PLOS ONE addresses a range of related and interesting issues: 1) Different restoration approaches, from passive rewilding to active target driven restoration, are needed to achieve different restoration goals in different circumstances. 2) Nature is complex and context dependent and so diverse approaches to restoration will help ensure different taxonomic groups and ecosystem functions and services are supported. 3) Developing and recording best practice for different restoration approaches will greatly aid the achievement of restoration aims. 4) Measuring restoration success needs comprehensive, multi-dimensional, and quantifiable metrics to account for potentially complex trade-offs. 5) Arguments for restoration based on ecocentric and nature's contribution to people both have merit and appeal to different audiences, but it should not be assumed goals derived from these different ways of thinking will be aligned. This is a diverse collection of restoration and rewilding research, and that diversity neatly reflects the diverse approaches and goals needed for restoration to be successful.

The articles in this issue discuss case studies that span a continuum of restoration interventions from removing anthropogenic disturbance and allowing the ecosystem to regenerate naturally (i.e., passive restoration or rewilding) to intensive interventions with ongoing management. For example, Broughton et al. [4] found that secondary woodlands in England that were adjacent to ancient woodlands recovered naturally over a period of a few decades. Díaz-García et al. [10] compared recovery of amphibians, ants, and dung beetles in naturally regenerating and actively planted tropical forests in Mexico; they found that passive and active restoration approaches were similarly effective in restoring species richness of all guilds, but that forest specialists were enhanced through active planting. In contrast, other studies show that intensive anthropogenic interventions such as transplanting corals (Ferse et al. [11]), or controlling invasive species and reintroducing fauna (Roberts et al. [12]) are necessary to facilitate recovery. The diversity of responses reported highlights the need to tailor restoration strategies to the local ecosystem type, the species of interest, and the level of prior disturbance.

Similarly, studies in this collection demonstrate complex interactions between wild and domestic herbivory, controls on grazing intensity and spatial ecological variables, making generalizations difficult and stressing the need for context-specific studies and understanding to guide management of disturbance regimes. One study in African savanna (Young et al. [13]) explores the impact of grazing on biodiversity and shows that plots protected from herbivory by large wild herbivores for the past 25 years have developed a rich diversity of woody vegetation species which could disappear upon rewilding depending on level of predation and associated behavioral patterns. However, they also show that individuals of the dominant tree species in this system, *Acacia drepanolobium*, greatly reduce their defense in the absence of browsers; hence the sudden arrival of these herbivores resulted in far greater elephant damage than for conspecifics in adjacent plots that had been continually exposed to herbivory.

Similarly, Peacock et al. [14] suggests that cattle negatively impact regeneration of gallery forests in Bolivia and alter both the structure and composition of the shrub and ground layers with potential consequences for the diversity and abundance of wildlife. Previous studies including Hanke et al. [15] have shown increases in species diversity and ecological functioning with grazing. These results suggest that the impact of grazing on ecosystems, species and ecosystem functioning depends on the system, the grazing species, and their numbers, and overall carrying capacity.

Several best practices are highlighted in the Collection. Larson et al. [16] created a model to determine an “optimal maximum distance” that would maximize availability of native prairie seed in the midwestern United States (US) from commercial sources while minimizing the risk of novel invasive weeds via contamination. Pedrini et al. [17] test seed pretreatment methods to enhance vegetation establishment from direct seeding and illustrate how a range of life stage transitions including germination, emergence and survival of native grass species used in restoration programs can be improved by seed coating with salicylic acid. Roon et al. [18] used a before-after-control-impact experiment across three stream networks in the northwestern to provide guidance on riparian thinning to provide optimal stream habitat. These best practices are key in our ability to replicate in different places and achieving restoration success.

Determining the success of ongoing restoration efforts is crucial to assessing management actions but requires comprehensive, multi-dimensional, and quantifiable metrics and approaches consistent with restoration goals. Despite the plethora of restoration projects around the world, it is only now that we are beginning to understand whether the restoration goals have been met and what trade-offs exist (Mugwedi et al. [19]). The importance of measuring restoration outcomes against clearly specified goals and objectives cannot be overemphasized, as shown in a recent restoration study in China that aimed to improve carbon storage through tree planting but has severely depleted water resources (Zhao et al. [20]). Similarly, Valach et al. [21] show that productive wetlands restored for carbon sequestration quickly become net carbon dioxide (CO₂) sinks although the trade-offs need to be further assessed. In their study exploring restoration success in South Africa, del Río et al. [22] improve our understanding on how techniques such as remote sensing can be used to measure restoration success.

As shown in this Collection and in other studies, trade-offs in restoration efforts are not uncommon and ultimately, restoration is successful when we can achieve restoration goals while minimizing trade-offs. The successful stories from the restoration interventions across different habitats and species showcased in the Collection (e.g., Sansupa et al. [3], Roon et al. [18], Valach et al. [21], Bouley et al. [23]) are a valuable addition to the science needed to advocate for restoration as a pathway to the recovery of previously degraded, damaged, or destroyed ecosystems. Reporting successful restoration outcomes can help increase buy-in for further restoration projects and increase funding availability for such projects. However, such buy-in can only occur if stakeholders are interested in the set restoration goals. For example, the need for climate mitigation has been used to justify several restoration programs around the world (Alexander et al. [24], Griscom et al. [25]). In this Collection, Matzek et al. [26] ask whether including ecosystem services as a restoration goal will engage a different set of values and attitudes than biodiversity protection alone. They found that support for habitat restoration is generally based on ecocentric values and attitudes, but that positive associations between pro-environmental behavior and egoistic values emerge when emphasis is placed on ecosystem service outcomes. They emphasize the notion that the ecosystem services concept garners non-traditional backers and broadens the appeal of ecological restoration as it is seen as a means of improving human well-being. Nevertheless, several studies (Bullock et al. [27], Egoh et al. [28], Newton et al. [29]) have shown that there can be trade-offs between

biodiversity and services during restoration and among different services, so restoration aims need to be clear rather than assuming win-win outcomes. Indeed, previous studies including Berry et al. [30] have suggested that a broad spectrum of perspectives on biodiversity conservation exist and should be used as arguments for conservation actions, from intrinsic to utilitarian values. In their analysis, the main differences between types of arguments appeared to result from the espousal of ecocentric or anthropocentric viewpoints, rather than from differences between the various stakeholder groups. This suggests that to promote restoration goals, a broad range of restoration goals are needed, including those that are more anthropocentric such as economic development.

While the positive impacts of ecological restoration on biodiversity are well established, less evidence is available regarding its impacts on economic development and employment. Although restoration efforts centered around economic development in Africa, such as the Working for Water Project and eThekweni forest restoration project in South Africa have generated strong support from government, not many such initiatives exist in other parts of the world (Mugwedi et al. [19]). In this collection, Newton et al. [29] examine the impacts of restoration on economic development and employment. They conclude that landscape-scale restoration or rewilding of agricultural land can potentially increase the contribution of farmland to economic development and employment, by increasing flows of multiple ecosystem services to the many economic sectors that depend on them. Indeed, restoration has contributed to the economy in many parts of the world leading to the framing of the term “restoration economy” or “green economy” which is now commonly used in the restoration literature (Bek et al. [31], Formosa et al. [32]). A recent report by Dasgupta [33] states that “our economies are embedded within Nature, not external to it”. While we are all looking forward to the UN Decade on Ecosystem Restoration launching this year, the uptake of restoration projects will depend on financing. Generating funds to support and sustain restoration projects is one of the biggest challenges facing restoration activities worldwide (FAO and Global Mechanism of the UNCCD [34]). The inclusion of a broad range of goals that span from biodiversity to anthropocentric goals such as those related to benefits of nature’s contribution to people to those that are purely development such as job creation may be the way forward.

References

1. Brondizio ES, Settele J, Díaz S, Ngo HT. Global assessment report on biodiversity and ecosystem services of the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services. Bonn: IPBES Secretariat; 2019.
2. Reid WV, Mooney HA, Cropper A, Capistrano D, Carpenter SR, Chopra K, et al. Ecosystems and human well-being-Synthesis: A report of the Millennium Ecosystem Assessment. Washington, DC: Island Press; 2005.
3. Sansupa C, Purahong W, Wubet T, Tiansawat P, Pathom-Aree W, Teamroong N, et al. Soil bacterial communities and their associated functions for forest restoration on a limestone mine in northern Thailand. *PLoS One*. 2021 Apr 8; 16(4):e0248806. <https://doi.org/10.1371/journal.pone.0248806> PMID: 33831034
4. Broughton RK, Bullock JM, George C, Hill RA, Hinsley SA, Maziarz M, et al. Long-term woodland restoration on lowland farmland through passive rewilding. *PLoS One*. June 16, 2021; e0252466 <https://doi.org/10.1371/journal.pone.0252466> PMID: 34133452
5. Schulz K, Stevens PW, Hill JE, Trotter AA, Ritch JL, Tuckett QM, et al. Coastal restoration evaluated using dominant habitat characteristics and associated fish communities. *PLoS one*. 2020 Oct 22; 15(10):e0240623. <https://doi.org/10.1371/journal.pone.0240623> PMID: 33091008
6. Ndangalasi HJ, Martínez-Garza C, Harjo TC, Pedigo CA, Wilson RJ, Cordeiro NJ. Seedling recruitment under isolated trees in a tea plantation provides a template for forest restoration in eastern Africa. *PLoS One*. 2021 May 6; 16(5):e0250859. <https://doi.org/10.1371/journal.pone.0250859> PMID: 33956837
7. Wasson K, Tanner KE, Woolfolk A, McCain S, Suraci JP. Top-down and sideways: Herbivory and cross-ecosystem connectivity shape restoration success at the salt marsh-upland ecotone. *PLoS One*. 2021 Feb 22; 16(2):e0247374. <https://doi.org/10.1371/journal.pone.0247374> PMID: 33617558

8. Mata JC, Buitenwerf R, Svenning JC. Enhancing monitoring of rewilding progress through wildlife tracking and remote sensing. *PLoS One*. 2021; Forthcoming.
9. D'Acunto LE, Pearlstine L, Romañach SS. Joint species distribution models of Everglades wading birds to inform restoration planning. *PLoS One*. 2021 Jan 28; 16(1):e0245973. <https://doi.org/10.1371/journal.pone.0245973> PMID: 33508032
10. Díaz-García JM, López-Barrera F, Pineda E, Toledo-Aceves T, Andresen E. Comparing the success of active and passive restoration in a tropical cloud forest landscape: A multi-taxa fauna approach. *PLoS One*. 2020 Nov 10; 15(11):e0242020. <https://doi.org/10.1371/journal.pone.0242020> PMID: 33170890
11. Ferse SC, Hein MY, Rölfer L. A survey of current trends and suggested future directions in coral transplantation for reef restoration. *PLoS One*. 2021 May 3; 16(5):e0249966. <https://doi.org/10.1371/journal.pone.0249966> PMID: 33939716
12. Roberts JL, Cooper WJ, Luther D. Global assessment of forest quality for threatened terrestrial vertebrate species in need of conservation translocation programs. *PLoS One*. 2021 Apr 14; 16(4):e0249378. <https://doi.org/10.1371/journal.pone.0249378> PMID: 33852601
13. Young TP, Kimuyu DM, Odadi WO, Wells HB, Wolf AA. Naïve plant communities and individuals may initially suffer in the face of reintroduced megafauna: An experimental exploration of rewilding from an African savanna rangeland. *PLoS One*. 2021 Apr 6; 16(4):e0248855. <https://doi.org/10.1371/journal.pone.0248855> PMID: 33822786
14. Peacock J, Tonra CM, King J, Davies GM. Restoration of gallery forest patches improves recruitment of motacú palms (*Attalea princeps*) while diversifying and increasing wildlife populations. *PLoS One*. 2021 Apr 29; 16(4):e0250183. <https://doi.org/10.1371/journal.pone.0250183> PMID: 33914765
15. Hanke W, Böhner J, Dreber N, Jürgens N, Schmiedel U, Wesuls D, et al. The impact of livestock grazing on plant diversity: an analysis across dryland ecosystems and scales in southern Africa. *Ecol Appl*. 2014 Jul; 24(5):1188–203. <https://doi.org/10.1890/13-0377.1> PMID: 25154106
16. Larson JL, Larson DL, Venette RC. Balancing the need for seed against invasive species risks in prairie habitat restorations. *PLoS One*. 2021 Apr 7; 16(4):e0248583. <https://doi.org/10.1371/journal.pone.0248583> PMID: 33826620
17. Pedrini S, Stevens JC, Cross AT, Dixon KW. Seed encrusting with salicylic acid: a novel approach to improve establishment of grass species in ecological restoration. *PLoS One*. 2021 June 9; 16(6), p. e0242035 <https://doi.org/10.1371/journal.pone.0242035> PMID: 34106919
18. Roon DA, Dunham JB, Groom JD. Shade, light, and stream temperature responses to riparian thinning in second-growth redwood forests of northern California. *PLoS One*. 2021 Feb 16; 16(2):e0246822. <https://doi.org/10.1371/journal.pone.0246822> PMID: 33592001
19. Mugwedi LF, Rouget M, Egoh B, Ramdhani S, Slotow R, Rentería JL. An assessment of a community-based, forest restoration programme in Durban (eThekweni), South Africa. *Forests*. 2017 Aug; 8(8):255.
20. Zhao M, Geruo A, Zhang J, Velicogna I, Liang C, Li Z. Ecological restoration impact on total terrestrial water storage. *Nat Sustain*. 2021 Jan; 4(1):56–62.
21. Valach AC, Kasak K, Hemes KS, Anthony TL, Dronova I, Taddeo S, et al. Productive wetlands restored for carbon sequestration quickly become net CO₂ sinks with site-level factors driving uptake variability. *PLoS One*. 2021 Mar 25; 16(3):e0248398. <https://doi.org/10.1371/journal.pone.0248398> PMID: 33765085
22. del Río T, Willemen L, Vrieling A, Snoeys A, Nelson A. Long-term assessment of ecosystem services at ecological restoration sites using Landsat time series. *PLoS One*. 2021; In Press.
23. Bouley P, Paulo A, Angela M, Du Plessis C, Marneweck DG. The successful reintroduction of African wild dogs (*Lycaon pictus*) to Gorongosa National Park, Mozambique. *PLoS One*. 2021 Apr 22; 16(4):e0249860. <https://doi.org/10.1371/journal.pone.0249860> PMID: 33886594
24. Alexander S, Nelson CR, Aronson J, Lamb D, Cliquet A, Erwin KL, et al. Opportunities and challenges for ecological restoration within REDD+. *Restor Ecol*. 2011 Nov; 19(6):683–9.
25. Griscom BW, Adams J, Ellis PW, Houghton RA, Lomax G, Miteva DA, et al. Natural climate solutions. *Proc Natl Acad Sci*. 2017 Oct 31; 114(44):11645–50. <https://doi.org/10.1073/pnas.1710465114> PMID: 29078344
26. Matzek V, Wilson KA. Public support for restoration: Does including ecosystem services as a goal engage a different set of values and attitudes than biodiversity protection alone? *PLoS One*. 2021 Jan 19; 16(1):e0245074. <https://doi.org/10.1371/journal.pone.0245074> PMID: 33465097
27. Bullock JM, Aronson J, Newton AC, Pywell RF, Rey-Benayas JM. Restoration of ecosystem services and biodiversity: conflicts and opportunities. *Trends Ecol Evol*. 2011 Oct 1; 26(10):541–9. <https://doi.org/10.1016/j.tree.2011.06.011> PMID: 21782273

28. Ego B.N., Paracchini M.L., Zulian G., Schägner J.P. and Bidoglio G., 2014. Exploring restoration options for habitats, species and ecosystem services in the European Union. *Journal of Applied Ecology*, 51(4), pp.899–908.
29. Newton AC, Evans PM, Watson SC, Ridding LE, Brand S, McCracken M, et al. Ecological restoration of agricultural land can improve its contribution to economic development. *PLoS One*. 2021 Mar 5; 16(3): e0247850. <https://doi.org/10.1371/journal.pone.0247850> PMID: 33667265
30. Berry PM, Fabók V, Blicharska M, Bredin YK, Llorente MG, Kovács E, et al. Why conserve biodiversity? A multi-national exploration of stakeholders' views on the arguments for biodiversity conservation. *Biodivers Conserv*. 2018 Jun; 27(7):1741–62.
31. Bek D, Nel E, Binns T. Jobs, water or conservation? Deconstructing the green economy in South Africa's Working For Water Programme. *Environ Dev*. 2017 Dec 1; 24:136–45.
32. Formosa ML, Kelly EC. Socioeconomic Benefits of a Restoration Economy in the Mattole River Watershed, USA. *Soc Nat Resour*. 2020 Sep 1; 33(9):1111–28.
33. Dasgupta P. The economics of biodiversity: the Dasgupta review. London: HM Treasury; 2021.
34. FAO and Global Mechanism of the UNCCD. Sustainable Financing for Forest and Landscape Restoration-Opportunities, Challenges and the Way Forward. Liagre L, Lara Almuedo P, Besacier C, Conigliaro M, editors. Rome: Food and Agricultural Organization; 2015.