

Distribution of conodonts in the type Ludlow Series

Series and stages		Formation	<i>Panderodus equicostatus</i>	<i>Ozarkodina excavata</i>	<i>Panderodus</i> sp.	<i>Ozarkodina</i>	<i>Kockelella</i>	<i>Kockelella variabilis</i>	<i>Panderodus recurvatus</i>	<i>Panderodus unicosatus</i>	<i>Oulodus</i> sp.	<i>Pelekyognathus dubius</i>	<i>Coryssognathus dubius</i>	<i>Ozarkodina confluens</i>	<i>Pseudooneotodus beckmanni</i>	<i>Dapsilodus obliquicostatus</i>	<i>Decoriconus</i> sp.	<i>Ozarkodina snajdri</i>	<i>Ozarkodina wimani</i>	<i>Panderodus serratus</i>	<i>Ozarkodina remscheidensis eosteinhorrensis</i>	<i>Ozarkodina remscheidensis</i> subsp.	<i>Ozarkodina cf. crispa</i>
Pridoli ?		Downton Castle Sandstone	x									x	x								x		
Ludlow	Ludfordian	Upper Whitcliffe	x	x								x	x	x				x	x	x	x	x	x
		Lower Whitcliffe	x										x	x				x	x	x	x		
		Upper Leintwardine	x							x	x		x	x	x			x					
		Lower Leintwardine	x	x	x			x		x	x	x	x	x	x	x	x	x					
	Gorstian	Upper Bringewood	x	x				x		x	x	x	x	x	x								
		Lower Bringewood			x	x	x	?															
		Upper Elton	x	x																			
		Middle Elton			x																		
		Lower Elton	x	x																			
	Wenlock		Much Wenlock Limestone	x	x																		

See 'The type Ludlow Series: Biostratigraphy - conodonts' for sources of information.

[Return to 'The Ludlow Series \(Upper Silurian\) of the type area - introductory page'.](#)

(Author: SGM)