

British Geological Survey
NATURAL ENVIRONMENT RESEARCH COUNCIL

Office of the Deputy Prime Minister
Central Government

OXFORDSHIRE

Mineral Resource Information in Support of National, Regional and Local Planning

Mineral Resources

Scale 1:100 000

Compiled by G.E. Norton, D.G. Cameron, A.J. Bloodworth, D.J. Evans, G.K. Lott, K.A. Arton and E.E. Highway
Project Leader: D.E. Highway
Digital cartography by N.A. Spencer, British Geological Survey
Published 2004

This map comprises part of a summary of the Mineral Resources of the South-East of England Region. For further information see www.britain.gov.uk

BIBLIOGRAPHIC REFERENCE

Norton, G.E. and others. 2004. Mineral Resource Information in Support of National, Regional and Local Planning. Oxfordshire. British Geological Survey Commissioned Report CRO406/04

Production of this map was commissioned and funded by the Office of the Deputy Prime Minister (Contract MP0677).

SAND & GRAVEL

Superficial deposits

- Sub-alluvial: Indicated resources
- Sub-alluvial: Indicated resources (only in area assessed by BGS)
- River terrace deposits
- River terrace deposits: Concealed (only in area assessed by BGS)
- Glaciofluvial sand and gravel deposits
- Boundary of area assessed for sand and gravel at the indicated resource level

Bedrock deposits

- Horseshay Sand Formation, Kingston Formation, Fairford Stone, Gravelformation and Lower Greensand } Jurassic and Cretaceous
- Lower Greensand (Baulking - Fernham area only) } Lower Cretaceous

FULLER'S EARTH

- White Chalk Subgroup: Upper Cretaceous
- Grey Chalk Subgroup: Upper Cretaceous

CHALK

- Chalk: Higher purity (> 93% CaCO₃) } White Chalk Subgroup: Upper Cretaceous
- Chalk: Low purity (< 93% CaCO₃) } Grey Chalk Subgroup: Upper Cretaceous

LIMESTONE

- Chipping Norton Limestone, White Limestone, Highworth Limestone, Stamford Limestone, Wheatley Limestone and Coral Rag } Jurassic
- Ironstone (Marlstone Rock Formation) } Jurassic

MINERAL PLANNING PERMISSION (as at 31/12/03)

Source: Oxfordshire County Council

- Surface planning permission (valid and expired)

MINERAL WORKINGS

- Burford Active site
- Inactive (including yet to be worked), worked-out and/or restored site
- Rail aggregate terminal
- Coal-fired Power Station

Ch	Chalk	Fi	Flint	Sg	Sand & gravel
Ci	Common clay & shale	Istn	Ironstone	Ss	Silica Sand
CR	Crushed Rock	Lst	Limestone	Sst	Sandstone
FE	Fuller's Earth	Sec	Secondary aggregate		

ENVIRONMENTAL DESIGNATIONS (as at 30/06/03)

- National nature conservation designations (SSSIs and NNrS)
- International nature conservation designations (SACs, SPA and Ramsar sites)
- Area of Outstanding Natural Beauty (AONB): Chilterns (part) and Cotswolds (part)
- Scheduled Monument

ADMINISTRATIVE AREAS

- Mineral Planning Authority
- District

Topography reproduced from the OS map by British Geological Survey with the permission of Ordnance Survey on behalf of the Controller of Her Majesty's Stationery Office. © Crown Copyright. All rights reserved. Unauthorised reproduction infringes Crown copyright and may lead to prosecution or civil proceedings. Licence number: 10007672/2004.

Digital ISBN: NBN, SPA, SAC and Ramsar boundaries © English Nature 2003
Contact address: English Nature, Northminster House, Northminster, Peterborough, PE1 1UA, UK. Tel: 01753 652000. Fax: 01753 652015. Web page: www.english-nature.gov.uk

Portions of Scheduled Monuments at 20th September 2003 as published by English Heritage.

The majority of monuments are protected against a control NNrS symbol. Consequently the actual area and/or length of a monument protected by the legal constraints of a NNrS symbol are those shown. Monuments scheduled as part of a NNrS are not covered by the NNrS symbol.

Contact address: English Heritage, 23 Savile Row, London, W1P 2ET, UK. Tel: 020 7973 3123. Web page: www.english-heritage.org.uk

Digital AONB boundaries © Countryside Commission 1984 (now Countryside Agency)

Contact address: Countryside Agency, John Dower House, Clozemont Place, Chatterham, Gloucestershire, GL50 3JA, UK. Tel: 0142 521381. Fax: 0142 546270. Web page: www.countryside.gov.uk

Published by the Office of the Deputy Prime Minister © Queen's Printer and Controller of Her Majesty's Stationery Office 2004.

This publication (including logos) may be reproduced free of charge in any format or medium for research, private study or circulation within an organisation. This subject is not to be reproduced, stored in a retrieval system, or used in a promotional context. The material must be acknowledged as Crown copyright and the title of the publication specified.

Applications for reproduction should be made in writing to: The Copyright Unit, Her Majesty's Stationery Office, St Clements House, 1-15 Colston, Norwich NR3 1BQ. Fax: 01603 723000 or e-mail: copyright@hmo.gov.uk

Aims and Limitations

The purpose of this map is to show the broad distribution of mineral resources which may be of current or potential economic interest and to state those that are subject to national planning restrictions. The maps are intended to assist in the consideration and preparation of development plans in respect of mineral extraction and the production of important mineral resources against development. They bring together a wide range of information, much of which is scattered and not always available in a convenient form.

The maps have been produced by collation and interpretation of mineral resource data primarily held by the British Geological Survey. Information on the extent of mineral planning permissions has been obtained from the relevant Mineral Planning Authority (MPA). Some of these permissions may have expired. The status of individual areas can be ascertained from the appropriate MPA. Location information on national planning permissions has been obtained from the appropriate statutory body (Countryside Agency, English Nature and English Heritage). For further information the relevant body should be contacted.

The mineral resource data presented are based on the best available information, but are not comprehensive and their quality is variable. The information shown above, therefore, approximate. Mineral resources defined on the map delineate areas within which potentially extractable mineral resources may occur. These areas are not of uniform potential and also take account of planning constraints which may limit their working. The economic potential of specific sites can only be proved by a detailed evaluation programme. Such an investigation is an essential prerequisite to submitting a planning application for mineral extraction. Extensive areas are shown as having no mineral resource potential, but some limited mineral resources may occur in these areas. The presence of these resources generally reflects very local or specific situations.

The maps are intended for general consideration of mineral issues and are not a source of detailed information on specific sites. The maps should not be used to determine individual planning applications or to taking other decisions on the acquisition or use of a particular piece of land, although they may give useful background information which will help in such decisions.

PLANNING PERMISSIONS FOR MINERAL EXTRACTION

The extent of known extent and former planning permissions for mineral extraction is shown on the map, irrespective of their current planning or operational status. The origins were digitised by BGS from Planning Orders and other documents supplied by Oxfordshire County Council and queries regarding the sites shown should be directed to the authority at the address shown below. The polygons cover active, former and restored mineral workings and occasionally, unworked deposits.

Planning Permissions represent areas where a commercial decision to work mineral has been made. A successful application has been made through the provisions of the Town and Country Planning legislation and the permitted resources will have been adapted to a greater or lesser extent. Current planning status is not qualified on the map but is available in the underlying database.

Contact address:
Oxfordshire County Council, Environmental Services, County Hall, New Road, Oxford OX1 1TD, UK. Tel: 01865 618178. Fax: 01865 246110. Web page: www.oxfordshire.gov.uk

CHALK

Chalk is a soft, white, crystalline, sedimentary rock, consisting mostly of the mineral calcite, which is composed of calcium carbonate (CaCO₃). It is formed in a broad band running from east to west. On this resource map, the chalk is subdivided into two categories: the pure and high purity, low purity chalk (generally near the EPR CaCO₃ boundary) and the low purity chalk (generally in the area of the Upper Cretaceous). These resources tend to have a relatively high clay content. The Grey Chalk Subgroup (Ironstone Lower Chalk) is about 60 m thick in Oxfordshire. High purity chalk, which is also known as white chalk, occurs in the White Chalk Subgroup of the Upper Cretaceous. The White Chalk Subgroup is composed of two main units, the upper Chalk and the lower Chalk, and has sometimes been worked as a decorative stone. The White Chalk Subgroup formerly Middle and Upper Chalk is about 160 m thick in Oxfordshire.

The White Chalk Subgroup is quarried for agricultural lime at one location in south Oxfordshire.

Chalk is also an important aquifer and a major source of groundwater in the south of the county.

BUILDING STONE

The Jurassic and Cretaceous rocks of the county have provided a wide variety of stone for building purposes. In the Lower Jurassic succession, the limonitic limestones, sandstones and ironstones (Marlstone Rock Formation) were extensively worked for building stone around Alton, Horton and Oxton.

The Middle Jurassic limestones were the most important building stone resource and were quarried over much of the county, including operations at Taynton, Barton, Milton, Bladen, Cotbury and Hatherwood. Also locally important were extensive quarries working the beds limestones of the Great Oolite Group at Blonfield, Fines, Westwell and Fiddlers for roofing slates which dated back to the early 1700s. The Tertiary Forest Marble Formation limestones were worked for roofing slates and paving around Birta Norton, Hallow and Bradwell.

Upper Jurassic limestones and ironstones were worked around Oxford at Headington and Wheatley (Cornwall Group) and a Great Milton and Great Henny Forests Sandstone Formation.

Elsewhere, in the south and east of the county, chalk from the Cretaceous and cemented sandstones from the Palaeogene (Tertiary) were worked locally for building stone.

There are currently three quarries working the Marlstone Rock Formation for aggregate at Alton, Horton and Wroxton, which are also worked for building stone and a building stone quarry at Great Tew.

FULLER'S EARTH

The term 'Fuller's Earth' is used to describe clay composed essentially of the clay mineral Ca-zeolite, which exhibits a unique combination of properties on which its industrial applications are based. Ca-zeolite can be widely converted to a simple sodium-exchange process and most Fuller's Earth is used in this form in papermaking and laundry powder applications. Fuller's earth deposits were formed as a result of the interaction of volcanic ash with the oceanic crust. The accumulation and preservation of volcanic ash into thick beds involved a complex set of geological processes. Consequently, Fuller's Earth deposits of potential economic interest have a very restricted distribution in Britain and Oxfordshire is one of the very few countries where it occurs.

Larger-scale beds of Fuller's Earth occur within the Lower Cretaceous Lower Greensand in Oxfordshire. A number of small, isolated nodules of Fuller's Earth occur unconformably overlying Jurassic strata in Oxfordshire. Fuller's Earth occurs where the Lower Greensand interfingers with the underlying Jurassic sandstones and is confined to the Baulking/Fernham area to the south of Farnham. Only the outlier at Farnham is worked.

In 1884 a smaller, stable deposit of Fuller's Earth was discovered at Moor Mill Farm about 2 km west of the Baulking Deposit. This deposit was granted planning permission for extraction in 1988 and is due to be opened up in 2004. Reserves are reported to be some 300,000 tonnes in a single bed up to 3.8 m thick.

Work carried out by the BGS in 1989 indicated that Fuller's Earth also occurs in the Fernham area. However, beds were only 0.3 to 0.6 m thick and are rare and only one bed was proved to be thicker than 1 m. The remaining parts of the Baulking/Fernham Lower Greensand outlier do not appear to be very prospective.

MINERAL PLANNING PERMISSION (as at 31/12/03)

Source: Oxfordshire County Council

- Surface planning permission (valid and expired)

SAND AND GRAVEL

Sand and gravel are defined on the basis of particle size rather than composition. In current commercial practice, following the introduction of new European standards from 1st January 2004, the term 'gravel' is more commonly used for general and concrete aggregate applications. The term 'sand' is used for material that is finer than 0.425 mm but coarser than 2.0 mm. For use in concrete, sand is now the break point between coarse and fine aggregate. Most sand and gravel is composed of particles that are not in silica, quartzite and flint, but other rock types occur.

The principal uses of sand are in fine aggregate in concrete, mortar and asphalt. The main use of gravel is as coarse aggregate in concrete. Substantial quantities of sand and gravel may also be used for construction fill.

Oxfordshire produced 1.5 million tonnes of sand and gravel in 2002, and has estimated permitted reserves of 12 million tonnes. Recent production figures may be seen on the final map.

Sand and gravel resources occur in a variety of geological environments. In Oxfordshire, these resources fall into three categories:

- river terrace deposits
- glaciofluvial deposits
- bedrock sand and gravel

The first two categories are considered as superficial deposits and were assessed in parts of Oxfordshire by BGS in the 1970s and 1980s. Resources identified in these areas are identified separately on the map, and the possible extent of sand and gravel (concealed beneath outcrops) is shown. The third category is defined by outcrops of gravel and sandstone at least 0.5 m thick. Outside these areas, available data are more limited. Generally, only exposed sand and gravel are defined, although sub-alluvial resources of sand and gravel occurring beneath modern river floodplains may be extensive in some areas, and are marked on the map. However, some 200 m wide spreads of sub-alluvial deposits are partly excluded from the map. They are limited with likely to preclude economic working of any sand and gravel present.

River terrace deposits

These deposits occur in both raised river terrace sequences and as flood plain terraces associated with and underlying present day alluvium. They are all late Anglian to Brunstadean. River terrace occur at several sites in the east of the county including the county boundary in the present location, particularly associated with the River Thames and the major tributaries thereof. The river terraces are higher above the present course of the river and are more likely to be derived from the Thames and the major tributaries. The terrace deposits are composed of sand and gravel with weak to medium clay content, and thicknesses of up to a few metres. The younger deposits are more laterally continuous since they have been cut down less by subsequent river erosion.

Bedrock sand and gravel

In Oxfordshire, there are several areas where late-working, high-volume rivers derived from glacial meltwater laid down gravelly deposits in pre-Anglian times. These earlier deposits are here named 'glaciofluvial deposits', and are described below. Previously some of these deposits may have been covered by gravel.

Through the central part of Oxfordshire to the south of Woodstock, further upstream on the banks of the River Fyne, and to the south of Oxford there are some patches of high-level glaciofluvial sand and gravel. These are thought to be of pre-Anglian age. The largest outcrop of these deposits occurs around Fiddlers where BGS boreholes revealed the presence of up to 2.5 m thickness of clay-rich sand and gravel beds. However, the outcrops are generally small and are thus not currently worked anywhere, they are unlikely to be a major resource, but are included on this map for completeness.

In the south-east of Oxfordshire to the east of Wallingford, a series of gravel known as the Wallingford Flat Gravels crops out. The age of the deposits is uncertain, but they are thought to be mostly of early Anglian age. These are largely suffusion deposits, although they may have also accumulated east of a local fault, and are found on the steep edge of the chalk escarpment. They are rich in flint, and up to 4.5 m thick. Around the Ewelme area they have been extensively quarried, although there are no current workings.




Bedrock sand and gravel

Several formations in Oxfordshire consist of locally consolidated sandstones that are being worked for building stone. In particular, the Horseshay Sand Formation of the Great Oolite Group (Middle Jurassic), the Highworth Grit in the Cornwall Formation in the Cornwall Group (Upper Jurassic) and the Forest Marble Formation of the Tertiary Forest Marble Group are currently being quarried.

The Horseshay Sand Formation crops out in a limited area in the north of the county. It consists of a medium to fine grained quartzose sand, locally cemented into calcareous or weakly limonitic sandstones with thin dark grey mudstone and siltstone beds in places. It is up to 7 m thick, and is quarried near Great Tew.

The Highworth Grit formation crops out on the southern part of Oxfordshire and runs approximately north-south west to east from Farnham to the north-east of Oxford. The whole formation is up to 30 m thick, although the principle resource, the Highworth Grit, is only a part of the formation and is present in a limited area in the north of the county. The Highworth Grit consists mainly of medium-grained quartzose sands. It is currently being quarried near Farnham and near Farnham Wood.

The Farnham Stone Formation crops out in a small area near Farnham and is composed of red and yellow siltstone and sandstone in the lower part of the formation, and is capped by a thin layer of claystone and siltstone. The latter is a coarse grained, sandy, silty and shaly sandstone with many small, smooth quartz pebbles. It is thought to be up to 10 m thick but borehole information and a quarry to the south of Farnham. Other smaller areas of Lower Greensand are also shown as bedrock resources.



CRUSHED ROCK AGGREGATE

A variety of hard rocks, when crushed, are suitable for use as aggregate. The technical suitability for different applications depends on their physical characteristics. The aggregate used for road surfacing and in concrete requires a high crushing strength and resistance to abrasion. Higher quality aggregate are required for coastal and heavy-duty applications, such as concrete for bridge piers and abutments, where they are subjected to sea water and salt. In some applications, such as concrete for bridge piers and abutments, they are subjected to sea water and salt.

Oxfordshire produced 620,000 tonnes of crushed rock aggregate in 2002, and has estimated permitted reserves of 10 million tonnes. In Oxfordshire, two main rock types are quarried for use as crushed rock aggregate: limestone and ironstone.

Limestone

Limestones are found in several formations in Oxfordshire. In particular, the Great Oolite Group of the Middle Jurassic has two limestone formations that are currently being worked for crushed rock aggregate: the Chipping Norton Limestone Formation and the White Limestone Formation. The Great Oolite Group runs from north-west to south-west across the northern part of Oxfordshire.

The Chipping Norton Limestone Formation is up to 10.7 m thick near Chipping Norton, where it forms an extensive plateau, but thins towards the north-west and east. It is a fine to medium-grained fossiliferous limestone with some minor interstratified mudstone beds. It is currently quarried at Pook, Little Highbury and Fairgreen Farm, to the north-west and south of Chipping Norton respectively.

The White Limestone Formation crops out extensively across Oxfordshire westward, a thickness of about 20 m near Woodstock, but locally having thicknesses of up to 30 m. It is mostly composed of a cream-colored, fine-grained fossiliferous limestone with the mudstones. It is currently quarried at Acton, Barton and Woodstock.

In addition, a bed of limestone from the Cornwall Group runs across southern Oxfordshire from Brighthelm to Oxford. These are generally low quality limestones, formerly known as Great Rail. Currently they are worked in association with sandstones of the Forest Marble Formation near Farnham. These limestones include the Highworth Limestone of the Cornwall Formation and the Wheatley Limestone Member of the Cornwall Formation. Other limestones in the Great Oolite Group and Portland Group of the Upper Jurassic have been worked in the past, but are not currently being quarried as a result of high mudstone content or limited thickness of beds. These include the Taynton Limestone of the Forest Marble Formation and Cornhill Formation from the Great Oolite Group, and the Portland Formation from the Portland Group. These are not shown on the map.

Ironstone


The Marlstone Rock Formation in the Lower Jurassic is currently quarried for crushed rock aggregate at three locations in north Oxfordshire. This is a limonitic massive limestone up to 7.8 m thick, and is also quarried as a building stone. The ironstone, which had an average Fe content of greater than 25% with 10% SiO₂, 10% CaO and 0.25% P₂O₅ was worked between 1880 and 1980 as a source of iron. Minor outcrops of ironstone in the east and west of the county are not shown on the map.

BRICK CLAY

The term 'brick clay' is used to describe clay used predominantly in the manufacture of bricks and, to a lesser extent, roof and clay pipes. These clays may sometimes be used in cement manufacture as a source of construction fill, and for firing and sealing landfill sites. The suitability of a clay for the manufacture of bricks depends principally on its behavior during drying, firing and firing. The wet behavior of the fired brick such as strength and frost resistance and, importantly, its architectural appearance.

Most firing bricks, engineering bricks and related clay-based building products are manufactured in large automated factories. These represent a high capital investment and are increasingly dependent therefore on mineral with predictable and consistent properties in order to achieve high yields of salable products. Blending different clays to achieve improved durability and to provide a range of colors and textures is an increasingly common feature of the industry. Continuity of supply of consistent raw materials is of paramount importance.

Currently there are no brick and tile manufacturing sites in Oxfordshire, and this brick clay is not shown as a resource on the map. However, clays that are or have been employed in neighboring counties such as Buckinghamshire Clay with flint, Whitehall Oxford Clay Formation, Kennet Valley Clay Formation, Hampshire Clay Formation, Reading Formation, and Berkshire (Glaxo) Formation, Peterborough Member of the Oxford Clay Formation are also present in Oxfordshire, and may constitute a resource in the future.



HYDROCARBONS

Conventional Oil and Gas

The county of Oxfordshire lies on the southern flanks of the ancient London-Bulford Massif where Variscan Basement is at relatively shallow depths. Across this area thin Mesozoic sequences were deposited, including the quality and extent of the potential reservoir and source rock facies found in the near Western Basin dependent on the depth. Any source rocks present will be rather thin although near Chipping Norton there has been local deep erosion for the generation of commercial hydrocarbons in these parts.

The early 20th Century sea exploration for conventional Coal Measures in the neighbouring county of Buckinghamshire, when the Chiltern Quarry 195 and Colver East boreholes were drilled in 1931 and 1932 respectively, both failed to encounter any Coal Measures strata, although significant gas shows and pressures were recorded in the latter.

Exploration for conventional hydrocarbons in Oxfordshire dates back to 1954 when Dr Roy Gifford drilled the Flitton No.1 well in the SW of the county. Other boreholes were subsequently drilled at New Hill (2046) (Abolition 1966), Sheeps Ash (1972) and Bletchley (1978). Sheeps Ash generated Westphalian Coal Measures, thereby proving part of the concealed Oxfordshire-Berkshire Coalfield that is now known to underlie much of the two counties. It is these coals that are the source of the gas in this area. The number of potential wells has increased and a number of seismic reflection surveys, illustrates that between the mid 1950s and mid 1970s much of Oxfordshire has been explored for hydrocarbons. However, little oil or gas wells have been discovered in the county, although gas has been encountered in minor amounts in a few of the holes drilled, mainly from the Lower Lias and deeper strata.

The county thus appears to have limited hydrocarbon prospectivity, as evidenced by the fact there were of early 2003, no active hydrocarbon exploration or production licenses. However, the possibility for the discovery of buried gas accumulations may well exist.

Coal Bed Methane (CBM) potential

The basement rocks in the Mesozoic-Palaeogene rocks of Oxfordshire comprise strata of Cambrian, Devonian and Carboniferous age. The Carboniferous rocks include strata of Westphalian age, forming the Oxfordshire-Berkshire Coalfield and lying at depths between 300 and 1500 metres. Containing an area of 500 km², high volatile bituminous coals are present in synclines with an intervening anticline. They have an average total thickness of 10 metres and average gas seam content of 0.2m³/t. No coal has been mined and the Coal Measures succession is considered to be an appropriate for Coal Bed Methane (CBM) gas accumulation.

Coal

Concealed Carboniferous coal-bearing rocks forming the Oxfordshire-Berkshire Coalfield occur in Oxfordshire as a broadly north-south trending syncline between Oxford in the east and Farnham in the west. The strata are mainly Upper Coal Measures and lying at depths between 300 and 1500 metres. Containing an area of 500 km², high volatile bituminous coals are present in synclines with an intervening anticline. They have an average total thickness of 10 metres and average gas seam content of 0.2m³/t. No coal has been mined and the Coal Measures succession is considered to be an appropriate for Coal Bed Methane (CBM) gas accumulation.