



DATA NOTE

# The genome sequence of the Brindled Beauty, *Lycia hirtaria* (Clerck, 1759) [version 1; peer review: awaiting peer review]

Douglas Boyes<sup>1+</sup>, Peter W.H. Holland<sup>2</sup>,  
University of Oxford and Wytham Woods Genome Acquisition Lab,  
Darwin Tree of Life Barcoding collective,  
Wellcome Sanger Institute Tree of Life programme,  
Wellcome Sanger Institute Scientific Operations: DNA Pipelines collective,  
Tree of Life Core Informatics collective, Darwin Tree of Life Consortium

<sup>1</sup>UK Centre for Ecology & Hydrology, Wallingford, England, UK

<sup>2</sup>University of Oxford, Oxford, England, UK

+ Deceased author

---

**V1** First published: 12 Jul 2023, 8:303  
<https://doi.org/10.12688/wellcomeopenres.19650.1>  
Latest published: 12 Jul 2023, 8:303  
<https://doi.org/10.12688/wellcomeopenres.19650.1>

---

## Abstract

We present a genome assembly from an individual male *Lycia hirtaria* (the Brindled Beauty; Arthropoda; Insecta; Lepidoptera; Geometridae). The genome sequence is 552.0 megabases in span. Most of the assembly is scaffolded into 14 chromosomal pseudomolecules, including the Z sex chromosome. The mitochondrial genome has also been assembled and is 15.58 kilobases in length.

## Keywords

*Lycia hirtaria*, brindled beauty, genome sequence, chromosomal, Lepidoptera



This article is included in the [Tree of Life](#) gateway.

## Open Peer Review

**Approval Status** AWAITING PEER REVIEW

Any reports and responses or comments on the article can be found at the end of the article.

**Corresponding author:** Darwin Tree of Life Consortium ([mark.blaxter@sanger.ac.uk](mailto:mark.blaxter@sanger.ac.uk))

**Author roles:** **Boyes D:** Investigation, Resources; **Holland PWH:** Writing – Original Draft Preparation, Writing – Review & Editing;

**Competing interests:** No competing interests were disclosed.

**Grant information:** This work was supported by Wellcome through core funding to the Wellcome Sanger Institute (206194, <https://doi.org/10.35802/206194>) and the Darwin Tree of Life Discretionary Award (218328, <https://doi.org/10.35802/218328>). *The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.*

**Copyright:** © 2023 Boyes D *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

**How to cite this article:** Boyes D, Holland PWH, University of Oxford and Wytham Woods Genome Acquisition Lab *et al.* **The genome sequence of the Brindled Beauty, *Lycia hirtaria* (Clerck, 1759) [version 1; peer review: awaiting peer review]** Wellcome Open Research 2023, 8:303 <https://doi.org/10.12688/wellcomeopenres.19650.1>

**First published:** 12 Jul 2023, 8:303 <https://doi.org/10.12688/wellcomeopenres.19650.1>

## Species taxonomy

Eukaryota; Metazoa; Eumetazoa; Bilateria; Protostomia; Ecdysozoa; Panarthropoda; Arthropoda; Mandibulata; Pancrustacea; Hexapoda; Insecta; Dicondylia; Pterygota; Neoptera; Endopterygota; Amphiesmenoptera; Lepidoptera; Glossata; Neolepidoptera; Heteroneura; Ditrysia; Obtectomera; Geometroidea; Geometridae; Ennominae; *Lycia*; *Lycia hirtaria* (Clerck, 1759) (NCBI:txid326963).

## Background

Several moths in the subfamily Ennominae, family Geometridae, have winged males and flightless females, with wing reduction probably having evolved multiple times (Wahlberg *et al.*, 2010). To understand the genetic basis and the selective pressures underpinning this trait, species with flightless females must be compared to close relatives with winged females. The Brindled Beauty *Lycia hirtaria* is an example of this group in which both males and females are fully winged: in each sex the forewings are smoky-grey with black cross-lines. Interspecific crosses have been made between *L. hirtaria* and several related species, although the offspring are usually infertile (Ford, 1967; Harrison, 1916). In crosses between *L. hirtaria* and species with wingless females, partially winged hybrids are sometimes obtained. When the male is *L. hirtaria*, some of these crosses also give sex ratio distortion with a predominance of phenotypic males (Harrison, 1916; Harrison, 1919; Harrison & Doncaster, 1914). In the case of infertile offspring produced by crossing *L. hirtaria* with *L. zonaria*, examination of hybrids suggests that major karyotype differences cause disruption of chromosome pairing during meiosis in the F<sub>1</sub> generation (Harrison & Doncaster, 1914).

*L. hirtaria* is found in woodland and suburban areas across northern Europe and further east through Russia to Japan (GBIF Secretariat, 2022; Wagner, 2023). In Britain and Ireland, the moth is widespread but not usually common, and is recorded most frequently in the southeast of England (Randle *et al.*, 2019). The adult moth is on the wing in early spring, peaking in April in southern England, with larvae feeding in summer on the leaves of deciduous trees including *Prunus*, *Crataegus* and *Salix*; the pupal stage overwinters. Abundance of the species in Britain has declined by over 70% since 1970 (Randle *et al.*, 2019); in the 19th century it was sufficiently abundant in London to cause widespread defoliation of trees (Newman, 1869).

The complete genome of *Lycia hirtaria* was determined as part of the Darwin Tree of Life project. The assembled genome will contribute to the growing set of resources for studying insect ecology and evolution.

## Genome sequence report

The genome was sequenced from one male *Lycia hirtaria* (Figure 1) collected from Wytham Woods, Oxfordshire, UK (51.77, -1.34). A total of 40-fold coverage in Pacific Biosciences single-molecule HiFi long reads was generated. Primary assembly contigs were scaffolded with chromosome



**Figure 1.** Photograph of the *Lycia hirtaria* (iLycHirt1) specimen used for genome sequencing.

conformation Hi-C data. Manual assembly curation corrected three missing joins or mis-joins and removed one haplotypic duplication, reducing the scaffold number by 4.76%.

The final assembly has a total length of 552.0 Mb in 19 sequence scaffolds with a scaffold N50 of 45.4 Mb (Table 1). Most (99.92%) of the assembly sequence was assigned to 14 chromosomal-level scaffolds, representing 13 autosomes and the Z sex chromosome. Chromosome-scale scaffolds confirmed by the Hi-C data are named in order of size (Figure 2–Figure 5; Table 2). While not fully phased, the assembly deposited is of one haplotype. Contigs corresponding to the second haplotype have also been deposited. The mitochondrial genome was also assembled and can be found as a contig within the multifasta file of the genome submission.

The estimated Quality Value (QV) of the final assembly is 67.1 with *k*-mer completeness of 100%, and the assembly has a BUSCO v5.3.2 completeness of 98.4% (single = 97.7%, duplicated = 0.7%), using the lepidoptera\_odb10 reference set (*n* = 5,286).

Metadata for specimens, spectral estimates, sequencing runs, contaminants and pre-curation assembly statistics can be found at <https://links.tol.sanger.ac.uk/species/326963>.

## Methods

### Sample acquisition and nucleic acid extraction

A male *Lycia hirtaria* (specimen ID Ox001108, individual iLycHirt1) was collected using a light trap in Wytham Woods, Oxfordshire (biological vice-county Berkshire), UK (latitude 51.77, longitude -1.34) on 2021-03-31. Douglas Boyes (University of Oxford) collected and identified the specimen. The specimen was snap-frozen on dry ice.


**Table 1. Genome data for *Lycia hirtaria*, ilLycHirt1.1.**

<b>Project accession data</b>		
Assembly identifier	ilLycHirt1.1	
Species	<i>Lycia hirtaria</i>	
Specimen	ilLycHirt1	
NCBI taxonomy ID	326963	
BioProject	PRJEB56733	
BioSample ID	SAMEA10107033	
Isolate information	ilLycHirt1, male: thorax (DNA sequencing), head (Hi-C scaffolding)	
<b>Assembly metrics*</b>		<b>Benchmark</b>
Consensus quality (QV)	67.1	≥ 50
<i>k</i> -mer completeness	100%	≥ 95%
BUSCO**	C:98.4%[S:97.7%,D:0.7%], F:0.4%,M:1.2%,n:5,286	C ≥ 95%
Percentage of assembly mapped to chromosomes	99.92%	≥ 95%
Sex chromosomes	Z chromosome	<i>localised homologous pairs</i>
Organelles	Mitochondrial genome assembled	<i>complete single alleles</i>
<b>Raw data accessions</b>		
PacificBiosciences SEQUEL II	ERR10395969	
Hi-C Illumina	ERR10378040	
<b>Genome assembly</b>		
Assembly accession	GCA_947563715.1	
<i>Accession of alternate haplotype</i>	GCA_947563705.1	
Span (Mb)	552.0	
Number of contigs	83	
Contig N50 length (Mb)	11.8	
Number of scaffolds	19	
Scaffold N50 length (Mb)	45.4	
Longest scaffold (Mb)	56.1	

\* Assembly metric benchmarks are adapted from column VGP-2020 of "Table 1: Proposed standards and metrics for defining genome assembly quality" from (Rhie *et al.*, 2021).

\*\* BUSCO scores based on the lepidoptera\_odb10 BUSCO set using v5.3.2.

C = complete [S = single copy, D = duplicated], F = fragmented, M = missing, n = number of orthologues in comparison. A full set of BUSCO scores is available at <https://blobtoolkit.genomehubs.org/view/Lycia%20hirtaria/dataset/CANOBA01/busco>.



**Figure 2. Genome assembly of *Lycia hirtaria*, iLycHirt1.1: metrics.** The BlobToolKit Snailplot shows N50 metrics and BUSCO gene completeness. The main plot is divided into 1,000 size-ordered bins around the circumference with each bin representing 0.1% of the 551,971,097 bp assembly. The distribution of scaffold lengths is shown in dark grey with the plot radius scaled to the longest scaffold present in the assembly (56,061,216 bp, shown in red). Orange and pale-orange arcs show the N50 and N90 scaffold lengths (45,432,280 and 37,909,223 bp), respectively. The pale grey spiral shows the cumulative scaffold count on a log scale with white scale lines showing successive orders of magnitude. The blue and pale-blue area around the outside of the plot shows the distribution of GC, AT and N percentages in the same bins as the inner plot. A summary of complete, fragmented, duplicated and missing BUSCO genes in the lepidoptera\_odb10 set is shown in the top right. An interactive version of this figure is available at <https://blobtoolkit.genomehubs.org/view/Lycia%20hirtaria/dataset/CANOBA01/snail>.

DNA was extracted at the Tree of Life laboratory, Wellcome Sanger Institute (WSI). The iLycHirt1 sample was weighed and dissected on dry ice with tissue set aside for Hi-C sequencing. Thorax tissue was cryogenically disrupted to a fine powder using a Covaris cryoPREP Automated Dry Pulveriser, receiving multiple impacts. High molecular weight (HMW) DNA was extracted using the Qiagen MagAttract HMW DNA extraction kit. HMW DNA was sheared into an average fragment size of 12–20 kb in a Megaruptor 3 system with speed setting 30. Sheared DNA was purified by solid-phase reversible immobilisation using AMPure PB beads with a 1.8X ratio of beads to sample to remove the shorter fragments and concentrate the DNA sample. The concentration of the sheared and purified DNA was assessed using a Nanodrop spectrophotometer and Qubit Fluorometer and Qubit dsDNA High Sensitivity Assay kit. Fragment size


distribution was evaluated by running the sample on the FemtoPulse system.

### Sequencing

Pacific Biosciences HiFi circular consensus DNA sequencing libraries were constructed according to the manufacturers' instructions. DNA sequencing was performed by the Scientific Operations core at the WSI on a Pacific Biosciences SEQUEL II (HiFi) instrument. Hi-C data were also generated from head tissue of iLycHirt1 using the Arima2 kit and sequenced on the Illumina NovaSeq 6000 instrument.

### Genome assembly, curation and evaluation

Assembly was carried out with Hifiasm (Cheng *et al.*, 2021) and haplotypic duplication was identified and removed with purge\_dups (Guan *et al.*, 2020). The assembly was then



**Figure 3. Genome assembly of *Lycia hirtaria*, ilLycHirt1.1: BlobToolKit GC-coverage plot.** Scaffolds are coloured by phylum. Circles are sized in proportion to scaffold length. Histograms show the distribution of scaffold length sum along each axis. An interactive version of this figure is available at <https://blobtoolkit.genomehubs.org/view/Lycia%20hirtaria/dataset/CANOBA01/blob>.

scaffolded with Hi-C data (Rao *et al.*, 2014) using YaHS (Zhou *et al.*, 2023). The assembly was checked for contamination and corrected as described previously (Howe *et al.*, 2021). Manual curation was performed using HiGlass (Kerpedjiev *et al.*, 2018) and Pretext (Harry, 2022). The mitochondrial genome was assembled using MitoHiFi (Uliano-Silva *et al.*, 2022), which runs MitoFinder (Allio *et al.*, 2020) or MITOS (Bernt *et al.*, 2013) and uses these annotations to select the final mitochondrial contig and to ensure the general quality of the sequence.


A Hi-C map for the final assembly was produced using bwa-mem2 (Vasimuddin *et al.*, 2019) in the Cooler file format (Abdennur & Mirny, 2020). To assess the assembly metrics,

the *k*-mer completeness and QV consensus quality values were calculated in Merqury (Rhie *et al.*, 2020). This work was done using Nextflow (Di Tommaso *et al.*, 2017) DSL2 pipelines “sanger-tol/readmapping” (Surana *et al.*, 2023a) and “sanger-tol/genomenote” (Surana *et al.*, 2023b). The genome was analysed within the BlobToolKit environment (Challis *et al.*, 2020) and BUSCO scores (Manni *et al.*, 2021; Simão *et al.*, 2015) were calculated.

Table 3 contains a list of relevant software tool versions and sources.

Wellcome Sanger Institute – Legal and Governance

The materials that have contributed to this genome note have been supplied by a Darwin Tree of Life Partner. The submission




**Figure 4. Genome assembly of *Lycia hirtaria*, ilLychHirt1.1: BlobToolKit cumulative sequence plot.** The grey line shows cumulative length for all scaffolds. Coloured lines show cumulative lengths of scaffolds assigned to each phylum using the buscogenes taxrule. An interactive version of this figure is available at <https://blobtoolkit.genomehubs.org/view/Lycia%20hirtaria/dataset/CANOBA01/cumulative>.

of materials by a Darwin Tree of Life Partner is subject to the ‘**Darwin Tree of Life Project Sampling Code of Practice**’, which can be found in full on the Darwin Tree of Life website [here](#). By agreeing with and signing up to the Sampling Code of Practice, the Darwin Tree of Life Partner agrees they will meet the legal and ethical requirements and standards set out within this document in respect of all samples acquired for, and supplied to, the Darwin Tree of Life Project.

Further, the Wellcome Sanger Institute employs a process whereby due diligence is carried out proportionate to the nature

of the materials themselves, and the circumstances under which they have been/are to be collected and provided for use. The purpose of this is to address and mitigate any potential legal and/or ethical implications of receipt and use of the materials as part of the research project, and to ensure that in doing so we align with best practice wherever possible. The overarching areas of consideration are:

- Ethical review of provenance and sourcing of the material
- Legality of collection, transfer and use (national and international)



**Figure 5. Genome assembly of *Lycia hirtaria*, iLycHirt1.1: Hi-C contact map of the iLycHirt1.1 assembly, visualised using HiGlass.** Chromosomes are shown in order of size from left to right and top to bottom. An interactive version of this figure may be viewed at <https://genome-note-higlass.tol.sanger.ac.uk/l/?d=eaCUaTdDS5yUWAKuI0970A>.

**Table 2. Chromosomal pseudomolecules in the genome assembly of *Lycia hirtaria*, iLycHirt1.**

INSDC accession	Chromosome	Length (Mb)	GC%
OX387375.1	1	56.06	36.0
OX387376.1	2	52.63	36.0
OX387377.1	3	50.05	36.0
OX387378.1	4	46.87	36.0
OX387379.1	5	46.41	36.0
OX387380.1	6	45.43	36.0
OX387381.1	7	43.42	36.0
OX387382.1	8	42.67	36.0
OX387383.1	9	38.71	36.5
OX387384.1	10	38.67	36.0
OX387385.1	11	37.91	36.0
OX387387.1	12	19.38	36.5
OX387388.1	13	10.32	36.5
OX387386.1	Z	23.03	36.0
OX387389.1	MT	0.02	18.5

**Table 3. Software tools: versions and sources.**

Software tool	Version	Source
BlobToolKit	4.1.5	<a href="https://github.com/blobtoolkit/blobtoolkit">https://github.com/blobtoolkit/blobtoolkit</a>
BUSCO	5.3.2	<a href="https://gitlab.com/ezlab/busco">https://gitlab.com/ezlab/busco</a>
Hifiasm	0.16.1-r375	<a href="https://github.com/chhylp123/hifiasm">https://github.com/chhylp123/hifiasm</a>
HiGlass	1.11.6	<a href="https://github.com/higlass/higlass">https://github.com/higlass/higlass</a>
Merqury	MerquryFK	<a href="https://github.com/thegenemyers/MERQURY.FK">https://github.com/thegenemyers/MERQURY.FK</a>
MitoHiFi	2	<a href="https://github.com/marcelauliano/MitoHiFi">https://github.com/marcelauliano/MitoHiFi</a>
PretextView	0.2	<a href="https://github.com/wtsi-hpag/PretextView">https://github.com/wtsi-hpag/PretextView</a>
purge_dups	1.2.3	<a href="https://github.com/dfguan/purge_dups">https://github.com/dfguan/purge_dups</a>
sanger-tol/genomenote	v1.0	<a href="https://github.com/sanger-tol/genomenote">https://github.com/sanger-tol/genomenote</a>
sanger-tol/readmapping	1.1.0	<a href="https://github.com/sanger-tol/readmapping/tree/1.1.0">https://github.com/sanger-tol/readmapping/tree/1.1.0</a>
YaHS	yahs-1.1.91eebc2	<a href="https://github.com/c-zhou/yahs">https://github.com/c-zhou/yahs</a>

Each transfer of samples is further undertaken according to a Research Collaboration Agreement or Material Transfer Agreement entered into by the Darwin Tree of Life Partner, Genome Research Limited (operating as the Wellcome Sanger Institute), and in some circumstances other Darwin Tree of Life collaborators.

## Data availability

European Nucleotide Archive: *Lycia hirtaria* (brindled beauty). Accession number PRJEB56733; <https://identifiers.org/ena.embl/PRJEB56733>. (Wellcome Sanger Institute, 2022)

The genome sequence is released openly for reuse. The *Lycia hirtaria* genome sequencing initiative is part of the Darwin Tree of Life (DToL) project. All raw sequence data and the assembly have been deposited in INSDC databases. The genome will be annotated using available RNA-Seq data and presented through the [Ensembl](#) pipeline at the European Bioinformatics Institute. Raw data and assembly accession identifiers are reported in [Table 1](#).

## Author information

Members of the University of Oxford and Wytham Woods Genome Acquisition Lab are listed here: <https://doi.org/10.5281/zenodo.4789928>.

Members of the Darwin Tree of Life Barcoding collective are listed here: <https://doi.org/10.5281/zenodo.4893703>.

Members of the Wellcome Sanger Institute Tree of Life programme are listed here: <https://doi.org/10.5281/zenodo.4783585>.

Members of Wellcome Sanger Institute Scientific Operations: DNA Pipelines collective are listed here: <https://doi.org/10.5281/zenodo.4790455>.

Members of the Tree of Life Core Informatics collective are listed here: <https://doi.org/10.5281/zenodo.5013541>.

Members of the Darwin Tree of Life Consortium are listed here: <https://doi.org/10.5281/zenodo.4783558>.

## References

- Abdennur N, Mirny LA: **Cooler: Scalable storage for Hi-C data and other genomically labeled arrays.** *Bioinformatics*. 2020; **36**(1): 311–316. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Allio R, Schomaker-Bastos A, Romiguier J, et al.: **MitoFinder: Efficient automated large-scale extraction of mitogenomic data in target enrichment phylogenomics.** *Mol Ecol Resour*. 2020; **20**(4): 892–905. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Bernt M, Donath A, Jühling F, et al.: **MITOS: Improved de novo metazoan mitochondrial genome annotation.** *Mol Phylogenet Evol*. 2013; **69**(2): 313–319. [PubMed Abstract](#) | [Publisher Full Text](#)
- Challis R, Richards E, Rajan J, et al.: **BlobToolKit - interactive quality assessment of genome assemblies.** *G3 (Bethesda)*. 2020; **10**(4): 1361–1374. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Cheng H, Concepcion GT, Feng X, et al.: **Haplotype-resolved de novo assembly using phased assembly graphs with hifiasm.** *Nat Methods*. 2021; **18**(2): 170–175. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Di Tommaso P, Chatzou M, Floden EW, et al.: **Nextflow enables reproducible computational workflows.** *Nat Biotechnol*. 2017; **35**(4): 316–319. [PubMed Abstract](#) | [Publisher Full Text](#)
- Ford EB: **Moths.** London: New Naturalist, Collins.1967.
- GBIF Secretariat: ***Lycia hirtaria* (Clerck, 1759), GBIF Backbone Taxonomy.** 2022. [Accessed 5 March 2023]. [Reference Source](#)
- Guan D, McCarthy SA, Wood J, et al.: **Identifying and removing haplotypic duplication in primary genome assemblies.** *Bioinformatics*. 2020; **36**(9): 2896–2898. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Harrison JWH: **Studies in the hybrid Bistoninae.** *J Genet*. 1916; **6**(2): 95–161. [Accessed 5 March 2023]. [Reference Source](#)
- Harrison JWH: **Studies in the hybrid Bistoninae: IV. Concerning the sex and related problems.** *J Genet*. 1919; **9**(1): 1–38. [Accessed 5 March 2023]. [Reference Source](#)
- Harrison JWH, Doncaster L: **On hybrids between moths of the geometrid sub-family Bistoninae, with an account of the behaviour of the chromosomes in gametogenesis in *Lycia* (Biston) *hirtaria*, *Ithysia* (Nyssia) *zonaria* and in their hybrids.** *J Genet*. 1914; **3**(4): 229–248. [Reference Source](#)
- Harry E: **PretextView (Paired REad TEXTure Viewer): A desktop application for viewing pretext contact maps.** 2022. [Accessed 19 October 2022]. [Reference Source](#)
- Howe K, Chow W, Collins J, et al.: **Significantly improving the quality of genome assemblies through curation.** *GigaScience*. Oxford University Press. 2021; **10**(1): g1aa153. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Kerpedjiev P, Abdennur N, Lekschas F, et al.: **HiGlass: web-based visual exploration and analysis of genome interaction maps.** *Genome Biol*. 2018; **19**(1): 125. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Manni M, Berkeley MR, Seppely M, et al.: **BUSCO update: Novel and streamlined workflows along with broader and deeper phylogenetic coverage for scoring of eukaryotic, prokaryotic, and viral genomes.** *Mol Biol Evol*. 2021; **38**(10): 4647–4654. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Newman E: **An Illustrated Natural History of British Moths.** London: Hardwicke and Bogue. 1869.
- Randle Z, Evans-Hill LJ, Parsons MS, et al.: **Atlas of Britain & Ireland's Larger Moths.** Newbury: NatureBureau. 2019.
- Rao SSP, Huntley MH, Durand NC, et al.: **A 3D map of the human genome at kilobase resolution reveals principles of chromatin looping.** *Cell*. 2014; **159**(7): 1665–1680. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Rhie A, McCarthy SA, Fedrigo O, et al.: **Towards complete and error-free genome assemblies of all vertebrate species.** *Nature*. 2021; **592**(7856): 737–746. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Rhie A, Walenz BP, Koren S, et al.: **Merquy: Reference-free quality, completeness, and phasing assessment for genome assemblies.** *Genome Biol*. 2020; **21**(1): 245. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Simão FA, Waterhouse RM, Ioannidis P, et al.: **BUSCO: assessing genome assembly and annotation completeness with single-copy orthologs.**

*Bioinformatics*. 2015; **31**(19): 3210–3212.

[PubMed Abstract](#) | [Publisher Full Text](#)

Surana P, Muffato M, Qi G: **sanger-tol/readmapping: sanger-tol/readmapping v1.1.0 - Hebridean Black (1.1.0)**. *Zenodo*. 2023a.

[Publisher Full Text](#)

Surana P, Muffato M, Sadasivan Baby C: **sanger-tol/genomenote (v1.0.dev)**. *Zenodo*. 2023b.

[Publisher Full Text](#)

Uliano-Silva M, Ferreira GJR, Krashennikova K, et al.: **MitoHiFi: a python pipeline for mitochondrial genome assembly from PacBio High Fidelity reads**. *BioRxiv*. 2022.

[Publisher Full Text](#)

Vasimuddin Md, Misra S, Li H: **Efficient Architecture-Aware Acceleration of BWA-MEM for Multicore Systems**. In: *2019 IEEE International Parallel and*

*Distributed Processing Symposium (IPDPS)*. IEEE, 2019; 314–324.

[Publisher Full Text](#)

Wagner W: ***Lycia hirtaria* (Clerck, 1759), *Lepidoptera and their ecology***. 2023. [Accessed 5 March 2023].

[Reference Source](#)

Wahlberg N, Snäll N, Viidalepp J, et al.: **The evolution of female flightlessness among Ennominae of the Holarctic forest zone (Lepidoptera, Geometridae)**. *Mol Phylogenet Evol*. 2010; **55**(3): 929–938.

[PubMed Abstract](#) | [Publisher Full Text](#)

Wellcome Sanger Institute: **The genome sequence of the Brindled Beauty, *Lycia hirtaria* (Clerck, 1759)**. European Nucleotide Archive, [dataset], accession number PRJEB56733. 2022.

Zhou C, McCarthy SA, Durbin R: **YaHS: yet another Hi-C scaffolding tool**. *Bioinformatics*. 2023; **39**(1): btac808.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)