

# ECOGRAPHY

## Research article

### Multi-colony tracking of two pelagic seabirds with contrasting flight capability illustrates how windscapes shape migratory movements at an ocean-basin scale

Françoise Amélineau<sup>1</sup>✉, Arnaud Tarrow<sup>2</sup>, Simon Lacombe<sup>2,4</sup>, Vegard S. Bråthen<sup>5</sup>, Sebastien Descamps<sup>1</sup>, Morten Ekker<sup>6</sup>, Per Fauchald<sup>2</sup>, Malin K. Johansen<sup>1</sup>, Børge Moe<sup>5</sup>, Tycho Anker-Nilssen<sup>5</sup>, Maria I. Bogdanova<sup>7</sup>, Ingar S. Bringsvor<sup>8</sup>, Olivier Chastel<sup>9</sup>, Signe Christensen-Dalsgaard<sup>5</sup>, Francis Daunt<sup>7</sup>, Nina Dehnhard<sup>5</sup>, Kjell Einar Erikstad<sup>2</sup>, Aleksey Ezhov<sup>10</sup>, Maria Gavrilov<sup>11,12,13</sup>, Erpur S. Hansen<sup>14</sup>, Mike P. Harris<sup>7</sup>, Hálfór H. Helgason<sup>1,15</sup>, Magdalene Langset<sup>5</sup>, Don-Jean Léandri-Breton<sup>9,16</sup>, Svein-Håkon Lorentsen<sup>5</sup>, Benjamin Merkel<sup>1,17</sup>, Mark Newell<sup>7</sup>, Bergur Olsen<sup>18</sup>, Tone K. Reiertsen<sup>2</sup>, Geir H. R. Systad<sup>19</sup>, Thorkell L. Thorarinsson<sup>20</sup>, Jens Åström<sup>5</sup> and Hallvard Strøm<sup>1</sup>

<sup>1</sup>Norwegian Polar Institute, Fram Centre, Tromsø, Norway

<sup>2</sup>Norwegian Institute for Nature Research, Fram Centre, Tromsø, Norway

<sup>3</sup>University Rennes, CNRS, ECOBIO [(Ecosystèmes, Biodiversité, Evolution)] – UMR 6553, Rennes, France

<sup>4</sup>Ecole Normale Supérieure de Lyon, Université Claude Bernard Lyon I, Université de Lyon, Lyon, France

<sup>5</sup>Norwegian Institute for Nature Research, Trondheim, Norway

<sup>6</sup>Norwegian Environmental Agency, Trondheim, Norway

<sup>7</sup>UK Centre for Ecology and Hydrology, Edinburgh, UK

<sup>8</sup>Norsk Ornitologisk Forening Møre og Romsdal, Sandshamn, Norway

<sup>9</sup>Centre d'Etudes Biologiques de Chizé (CEBC), UMR 7372 CNRS-Univ. La Rochelle, (CEBC), UMR 7372 CNRS-Univ. La Rochelle, France

<sup>10</sup>Murmansk Marine Biological Institute, Murmansk, Russia

<sup>11</sup>National Park Russian Arctic, Archangelsk, Russia

<sup>12</sup>Association Maritime Heritage, Saint Petersburg, Russia

<sup>13</sup>Arctic and Antarctic Research Institute, Saint-Petersburg, Russia

<sup>14</sup>South Iceland Nature Research Centre, Vestmannaeyjar, Iceland

<sup>15</sup>East Iceland Nature Research Centre, Neskaupstaður, Iceland

<sup>16</sup>Department of Natural Resource Sciences, McGill University, Ste-Anne-de-Bellevue, QC, Canada

<sup>17</sup>Akvaplan-niva, Fram Centre, Tromsø, Norway

<sup>18</sup>Faroe Marine Research Institute, Tórshavn, Faroe Islands

<sup>19</sup>Norwegian Institute for Nature Research, Bergen, Norway

<sup>20</sup>Northeast Iceland Nature Research Centre, Húsavík, Iceland

## Ecography

2024: e06496

doi: 10.1111/ecog.06496

Subject Editor: Susanne Akesson

Editor-in-Chief:

Jens-Christian Svenning

Accepted 20 August 2023



Correspondence: Françoise Amélineau ([francoise.amelineau@ens-rennes.fr](mailto:francoise.amelineau@ens-rennes.fr))

Migration is a common trait among many animals allowing the exploitation of spatiotemporally variable resources. It often implies high energetic costs to cover large distances, for example between breeding and wintering grounds. For flying or swimming animals, the adequate use of winds and currents can help reduce the associated energetic costs. Migratory seabirds are good models because they dwell in habitats characterized by strong winds while undertaking very long migrations. We tested the hypothesis that seabirds migrate through areas with favourable winds. To that end, we



[www.ecography.org](http://www.ecography.org)

used the SEATRACK dataset, a multi-colony geolocator tracking dataset, for two North Atlantic seabirds with contrasting flight capabilities, the black-legged kittiwake *Rissa tridactyla* and the Atlantic puffin *Fratercula arctica*, and wind data from the ERA5 climate reanalysis model. Both species had on average positive wind support during migration. Their main migratory routes were similar and followed seasonally prevailing winds. The general migratory movement had a loop-shape at the scale of the North Atlantic, with an autumn route (southward) along the east coast of Greenland, and a spring route (northward) closer to the British Isles. While migrating, both species had higher wind support in spring than in autumn. Kittiwakes migrated farther and benefited from higher wind support than puffins on average. The variation in wind conditions encountered while migrating was linked to the geographical location of the colonies. Generally, northernmost colonies had a better wind support in autumn while the southernmost colonies had a better wind support in spring, with some exceptions. Our study helps understanding how the physical environment shapes animal migration, which is crucial to further predict how migrants will be impacted by ongoing environmental changes.

Keywords: animal movement, bird flight, *Fratercula arctica*, migration, *Rissa tridactyla*, wind

## Introduction

Wind is involved in numerous biological processes, in particular with regards to the movements and dispersal of individual organisms, and can therefore act as a strong driver of species distribution (Chapman et al. 2010), genetic diversity (Kling and Ackerly 2021), and evolution (Di Musciano et al. 2020). Wind patterns can for instance explain the biodiversity of passively dispersed species, with prevailing winds influencing community composition over vast areas (Muñoz et al. 2004, Chapman et al. 2015, Epele et al. 2021, Kling and Ackerly 2021). While micro-organisms, plants (seeds and pollen), and some wingless arthropods can only drift passively while being transported by wind, flying animals can adjust their behaviour to wind conditions and optimize their use of the airflow to reduce their movement costs (Liechti 2006, Chapman et al. 2011). Wind conditions can affect the movement behaviour of flying birds at various spatiotemporal scales, from short foraging trips (Tarrowx et al. 2016, Hedrick et al. 2018, Collins et al. 2020) to long migrations (Sjöberg et al. 2021). Seabirds are well-known to take advantage of wind for movement, especially since their marine habitats are often characterised by strong winds (Felicísimo et al. 2008, Weimerskirch et al. 2012, Hromádková et al. 2020). Seabirds are thus particularly well suited to study the effects of wind conditions on flight behaviour, and in particular the extent to which the use of wind conditions encountered along migratory routes is optimised (Alerstam 2011).

Given the high energetic costs of flying, optimizing flight behaviour according to wind conditions may be critical for long-distance migrants (Newton 2010). Models simulating migrations show that routes with favourable winds effectively reduce travel time even when they are not the shortest ones (Kranstauber et al. 2015), and a large number of tracking studies now suggest that many long-distance migrants favour longer routes, but with better wind assistance (Shaffer et al. 2006, Egevang et al. 2010, Shamoun-Baranes et al. 2017, Vansteelant et al. 2017, Hromádková et al. 2020). These routes with favourable wind conditions are shared by multiple species and represent major migratory flyways (Newton 2010, Shamoun-Baranes et al. 2017).

The wind predictability (temporal variability in wind conditions) is an important aspect because it allows addressing questions about how migratory birds adjust to wind conditions at larger temporal scales (Kranstauber et al. 2015). At global scale, wind patterns are relatively predictable and reflect the general atmospheric circulation (Ahrens 2014). In the Northern Hemisphere, prevailing winds blow north eastwards at latitudes between 30° and 60° (westerlies), and southwards at higher latitudes. For example, in the North Atlantic westerlies flow from North America to northern Europe, while around Greenland winds blow predominantly from the north (Fig. 1). In other areas such as the Barents and Norwegian Seas, winds are more variable and blow from various directions throughout the year (Fig. 1). Consequently, wind conditions encountered during migration for seabirds from various colonies in the North Atlantic are expected to be contrasting. For example, some well-known wintering grounds such as the Grand Banks, the Greenland Sea, or the North Sea, are shared by seabirds from different colonies (Frederiksen et al. 2012, 2016, Fort et al. 2013, Fayet et al. 2017, Merkel et al. 2021) which are expected to have different migratory routes and likely encounter distinct wind conditions during migration between their breeding and wintering grounds. Such differences in encountered wind conditions during the non-breeding season could have carry-over effects on subsequent breeding attempts or influence adult survival, and ultimately impact population dynamics (Bogdanova et al. 2017, Desprez et al. 2018, Reiertsen et al. 2021).

Many studies have modelled the energetic costs of flight under varying wind conditions and at fine spatiotemporal scale based on biologging data. It was found that flying with headwinds not only decreases the ground speed, but also increases the flight duration and the associated energetic expenditure (Weimerskirch et al. 2000, Amélineau et al. 2014). Selecting routes with favourable wind conditions might thus not only reduce the duration of migration, but also the energetic costs of flight (Shamoun-Baranes et al. 2017). For flapping flight, energetic costs were also higher with stronger headwinds (Elliott et al. 2014). How migrant birds respond to wind conditions will therefore also depend on their flight mode (e.g. flapping, glide-flapping or soaring),




Figure 1. Wind patterns and main migratory movements in the North Atlantic during autumn (a) and spring (b). Main migratory movements of kittiwakes (yellow arrows) and puffins (red arrows) are based on the migratory segments presented in Fig. 2. Grey arrows represent wind direction and speed. Background colour represent the SD in wind direction (rad), with a dark colour representing a low variability in wind direction and a light colour a high variability in wind direction. Dots represent colonies with kittiwakes (yellow), puffins (red) and both species (orange). Mean wind direction and speed were calculated from the monthly means of the ERA5 dataset for the study period (2014–2020); autumn includes August to November (outbound migration) and spring December to April (inbound migration).

their body mass, aspect ratio (wing length divided by wing width) and wing loading (body mass divided by the area of the wing) (Bowlin and Wikelski 2008, Horton et al. 2018). Migrants can further adjust their behaviour to wind

conditions by waiting for optimal wind conditions to engage in migration (Gill et al. 2009), selecting an altitude with favourable winds (Mateos-Rodríguez and Liechti 2012), or, when possible, landing and waiting when winds are

unfavourable (Shamoun-Baranes et al. 2017, Geisler et al. 2022). Furthermore, if the prevailing winds are stable throughout the year, wind conditions along a given route are likely to be more favourable during either the outbound or the inbound migration. Having more favourable winds during the inbound migration might ensure higher fitness, allowing earlier arrival at the colony which is generally associated with higher breeding success (Kokko 1999, Verhulst and Nilsson 2008) or reducing energy expenditure, hence leading to a better body condition at the start of the breeding season, and subsequently a higher reproductive output (Drake et al. 2014, de Zwaan et al. 2022).

Understanding how much the physical environment influences movement is one of the key questions in marine vertebrate ecology (Hays et al. 2016) and is important for our understanding of how animals migrate (Bowlin et al. 2010). The overall goal of this study was to investigate the relationships between wind conditions and seabird migratory movements at the scale of the North Atlantic Ocean. Our study utilizes a long-term, multi-colony and multi-species tracking dataset developed under the SEATRACK project (Strøm et al. 2021). We selected as study models two pelagic seabirds that are long-range migrants but with different flight capacity, the black-legged kittiwake *Rissa tridactyla* (hereafter kittiwake) and the Atlantic puffin *Fratercula arctica* (hereafter puffin). Both species use flapping (as opposed to glide-flapping or soaring) flight, but differ in morphology (wing area, wingspan, general body shape), wing loading, and flight speed (Pennycuik 1987). While kittiwakes are surface feeders, puffins are wing-propelled divers. Puffin wings are shorter and their wing loading higher, resulting in very high flight costs (Elliott et al. 2013). Despite the high flight costs, seabirds with a higher wing loading can fly with higher wind speeds (Nourani et al. 2023), and have a lower increase in airspeed when headwinds increase (Spear and Ainley 1997a). Thus, it seems that a higher wing loading confers a higher tolerance to unfavourable wind conditions and reduces the benefits of wind support. We therefore expected that wind conditions might affect the migratory flight behaviours of puffins and kittiwakes differently.

Our main hypothesis was that kittiwakes and puffins take advantage of favourable wind conditions when they migrate. We predicted that both kittiwakes and puffins would prefer routes that provide wind support during their migration bouts and have on average a positive wind support at the species level. We expected windscapes to differ between birds of different colonies, depending on where they were located relative to prevailing winds and key wintering areas, with larger differences for colonies situated further away from each other. We also expected differences between seasons, as prevailing wind patterns in the subpolar North Atlantic region remain relatively similar throughout the year, formed by the polar easterlies and the northern extreme of the westerlies, while birds migrate in opposite directions during autumn and spring. Finally, we predicted that puffins could face less favourable wind conditions than kittiwakes on average, linked to their higher wing loading.

To test our hypothesis, we compared migratory routes of kittiwakes and puffins with wind patterns in the North Atlantic extracted from global climate models produced by the European Centre for Medium-Range Weather Forecasts (ERA5; Hersbach et al. 2020). First, we mapped and described the general migratory and wind patterns (obj. 1) as well as the average wind support encountered by migrating seabirds (obj. 2) at the scale of the North Atlantic Ocean. Then, using a cluster analysis, we investigated the similarities in encountered windscapes between colonies (obj. 3). Finally, we investigated the differences in wind support among species and seasons (autumn/outbound and spring/inbound migration) (obj. 4).

## Material and methods

### Bird tracking data

We used data collected by the SEATRACK project between 2014 and 2020. In total, 794 kittiwakes and 425 puffins from 18 colonies (Supporting information) were equipped with geolocators attached on a ring, and recaptured in following breeding seasons. Models of geolocators were c65, c65\_super, f100 and w65 produced by Migrate Technology, mk18 produced by BAS, and mk4083 and mk4093 from Biotrack (Supporting information). All of these models recorded light and saltwater immersion data. Details on the logger types, number of deployments per colony and frequency of data acquisition are presented in the Supporting information. The median number of annual tracks per individual (a track being defined as data recorded between two successive breeding seasons) was two (range 1–6 tracks per individual, Supporting information).

Positions were estimated from raw light data based on an automatic processing fully described in Bråthen et al. (2021). This method estimates positions based on a threshold method (Lisovski et al. 2020) and filters inaccurate locations based on speed, angle and position. An error of 200–250 km is expected for these latitudes (Halpin et al. 2021). As latitude is inaccurate during equinoxes (Lisovski et al. 2020), periods between 8 September–20 October and 20 February–3 April were removed. Gaps were also present in the data when tracked individuals were north of the Arctic Circle in the winter or summer due to polar night or midnight sun. Missing locations were interpolated based on the informed random movement algorithm (IRMA), fully described in Fauchald et al. (2019) and based on an algorithm that was originally developed by Technitis et al. (2015). Six kittiwakes breeding in Kara Gate (Novaya Zemlya) and migrating to the Pacific (Ezhov et al. 2021) were excluded from the current study, which is focused on the North Atlantic. For each track, we selected only one location per day (midnight) to compute the daily movement vectors. As kittiwakes and puffins fly mostly during the day, we decided to keep only positions at night; we then extracted the wind conditions at the midpoint between two successive nightly positions to get an accurate estimate of winds encountered during the day.




Figure 2. Maps showing all migratory segments (black lines) in autumn (left) and spring (right) for kittiwakes (top row) and puffins (bottom row), together with winter distribution of each breeding population plotted as colony coloured 50% kernel utilisation distribution.

Only periods when birds were migrating (hereafter migratory segments) were included in the analyses (Fig. 2). First, we excluded the breeding season based on saltwater immersion data, for each colony, species and year (Fauchald et al. 2019), as well as December and January for puffins as there were no clear migratory movements on average during these months. We then followed the method of Amélineau et al. (2021) to identify migratory segments and exclude stationary periods during the remaining months for each track. Based on the excluded periods, seasons were defined as follows for each species: autumn contained data from August to November for kittiwakes, and from July to November for puffins; spring contained data from December to April for kittiwakes, and from February to May for puffins. For each migratory segment, we calculated the detour by dividing the total length of the migratory segment (cumulative sum of the distances between successive locations) by the shortest distance between the departure and arrival locations. For each individual, the overall migratory distance was calculated as the cumulative distance between the centroids of successive stationary periods (the first and last stationary periods being the breeding season at the colony) and represents the whole non-breeding season. Distances were calculated as great circle

distances using the function *distGeo* from the ‘geosphere’ package (Hijmans 2021). For each colony and species, the winter distribution (Fig. 2) was calculated as the 50% utilisation distribution kernel calculated on the December positions with the ‘adehabitatHR’ package (Calenge 2006), using a smoothing factor (*h*) of 200km, a grid with  $50 \times 50$  km cells and a Lambert azimuthal equal area coordinate system.

Wet/dry sensors of the geolocators were used to calculate the proportion of time spent wet per 24 h, which gives an approximation of the proportion of time spent on the water and in the air.

### Wind data

We used wind from the ERA5 model (Hersbach et al. 2020), a reanalysis model from the European Centre for medium-range weather forecast (ECMWF). Data were obtained from the climate data store (<https://cds.climate.copernicus.eu/>). Wind data of the ERA5 model are available at 10 m (surface winds), 100 m, and higher altitudes. We chose winds at 10 m as previous studies found a median flight altitude of 16 m (range 0–81 m) for kittiwakes (Borkenhagen et al. 2018), and < 10 m for puffins (Johnston et al. 2014). We extracted

the hourly  $u$  (east) and  $v$  (north) components of surface wind (10 m height) on a  $0.25^\circ$  grid over the study area. We then calculated the daily mean  $u$  and  $v$  wind components. To get wind conditions encountered along the track, we calculated the midpoint between two successive locations, and extracted the daily mean wind at this midpoint. We then defined the wind support as the wind vector component in the direction of the track, and the cross-wind as the wind vector component perpendicular to the track (Fig. 3) (Safi et al. 2013), assuming a complete compensation.

To calculate expected wind conditions, we used monthly average  $u$  and  $v$  components of surface wind (10 m) available from the ERA5 model for the study period (2014–2020,  $0.25^\circ$  resolution). For each month, we calculated the monthly SD of the wind direction (index of the variability of the wind direction) with the *sd.circular* function of the ‘circular’ package in R (www.r-project.org, Agostinelli and Lund 2022) and the monthly averaged wind speed (scalar mean) for the whole study period, and extracted these values at each position.

### Cluster analysis

A K-means cluster analysis (Hartigan and Wong 1979) was performed to investigate similarities between colonies in wind conditions and seabird behaviour during migration (Supporting information). For each species and season, variables included in the cluster analysis encompassed daily wind conditions encountered (wind support, cross wind, wind speed), bird behaviour (mean proportion wet per 24 h, detour)

and average wind conditions based on interannual monthly wind during the study period 2014–2020 (SD of wind direction and scalar average wind speed at each location). Variables were aggregated by calculating, for each season and species, the median value per migratory segment (except for detour where we already had one value per segment), track, year and colony. Each variable was then normalized. The function *fviz\_nbclust* of the R package ‘factoextra’ (www.r-project.org, Kassambara and Mundt 2020) was used to detect the optimal number of clusters.

### Statistical analyses

All analyses were performed in R ver. 4.1.2 (www.r-project.org).

We investigated the variation in wind support between seasons and species. We worked at the scale of an annual track, with one median value per track and per season. We fitted linear mixed-effects models with the *lmer* function of the ‘lme4’ package (Bates et al. 2015). To take the lack of independence in our data into account (repeated measurements per individual, per colony and per year), we included a random effect of year, and a random effect of individual nested in colony (Zuur et al. 2009). We considered all combinations of covariates. We then used AIC (Akaike’s information criterion) for model selection. If the difference in AIC values between two models was  $< 2$ , the models had equal statistical support and in case of nested models, the simplest was preferred (Burnham and Anderson 2002). We detected no pattern in the distribution of model residuals indicating that the normality and homoscedasticity assumptions were met. The ‘MuMin’ package was used to generate the marginal  $R^2$  (variance explained by the fixed effects) and the conditional  $R^2$  (variance explained by the fixed and random effects) (Bartoń 2022). Unless otherwise mentioned, results are presented as median and 95% confidence interval. The presented maps have a plate carrée projection.

## Results

### Seabird migratory routes and wind patterns in the North Atlantic (Obj. 1)

The migratory movements were generally directed toward the southwest in autumn, and the northeast in spring for both species (Fig. 2). Spring migration routes differed from autumn migration routes and the overall movement had a loop-shape, which appeared to correspond very well to the wind circulation patterns over the North Atlantic during these periods (Fig. 1). The wind circulation patterns are similar during both periods, following a generally circular and anti-clockwise pattern (Fig. 1). The prevailing winds follow the eastern coast of Greenland southwards, and the main autumn migratory movements of kittiwakes and puffins followed these winds between Greenland and Iceland. Some birds of both species from Norwegian colonies flew first to the Barents Sea before migrating south through the Greenland Sea. Between Newfoundland and mainland Europe, prevailing winds blow




Figure 3. Calculation of the wind support in a case of positive wind support (a) and negative wind support (b).  $\vec{g}$ : ground speed,  $\vec{a}$ : airspeed,  $\vec{w}$ : wind speed,  $\vec{ws}$ : wind support,  $\vec{cw}$ : crosswind.

towards the northeast. The main migratory movements of kittiwakes and puffins followed these prevailing winds in spring, heading further east towards the British Isles and Norway, except for puffins overwintering north of Iceland who headed east without following the prevailing winds in this area (Fig. 1, 2). Our results provide clear evidence for an anti-clockwise migration movement pattern in kittiwakes and puffins when considering both seasons and all our study colonies (Fig. 2).

### Wind supports encountered by seabirds in the North Atlantic (Obj. 2)

To get a better overview of winds encountered during migration, and whether they helped or hindered the movement,

we plotted the average direction of bird movements and average wind support encountered per species and season in the North Atlantic (Fig. 4). For both species and seasons, there was large spatial variation in encountered wind support. Some fly paths had a higher wind support, mainly birds flying southwards along the east coast of Greenland in the autumn, and birds following the westerlies in the spring. In contrast, birds flying south of Iceland in the autumn had negative average wind support, and the same applied to birds flying northward between Iceland and Greenland in the spring. In the Barents Sea, the wind support was generally low and negative. Due to the regional variation in wind support, we expected differences in the windscape experienced by birds from distinct colonies.




Figure 4. General bird movement patterns and associated wind support for kittiwakes in autumn (top left) and spring (top right), and puffins in autumn (bottom left) and spring (bottom right). The study area was divided into a grid ( $2 \times 2^\circ$  resolution). Each vector represents the mean speed and direction of bird movement at all the bird positions within a given cell. The colour of the vector represents the average wind support encountered by all individuals located in the cell. Cells with few observations ( $\leq 5$ ) were removed to avoid bias due to low sample size. We first calculated the average speed and wind support per colony, and then averaged values for all colonies. The longer an arrow, the more birds flying in the mean direction. Short arrows indicate birds at the given location moving in different directions, and the resulting direction of the average movement is only partially informative. The thickness of the arrow represents the amount of data for each cell (thin = small amount of data, thick = large amount of data).

### Intra-colony variations in windscares (Obj. 3)

In autumn, kittiwake colonies could be grouped in five clusters (Fig. 5a–b, Supporting information). Our results show clear differences among these clusters in several parameters linked to wind and migratory behaviour. The wind support was in general slightly higher for kittiwakes breeding in the Barents Sea colonies (2.12 m s<sup>-1</sup> [1.71–2.71] for Spitsbergen and Franz Josef Land (cluster E), and 2.07 m s<sup>-1</sup> [1.77–2.49] for Anda, Hornøya and Cape Krutik (cluster C)), except for Bjørnøya and Kara Gate that had a lower wind support (1.26 m s<sup>-1</sup> [-0.49 to 1.61], cluster D). The wind support was also low for birds from Isle of May, Ålesund and Sklinna (0.89 m s<sup>-1</sup> [0.41–1.47], cluster A). The wind direction was highly variable for kittiwakes from Bjørnøya and Kara Gate (median SD: 59.6° [58.4–63.0], cluster D), whereas the lowest variability was found for colonies in the southwestern parts of the study area (median SD: 32.1° [28.6–37.8] for Iceland, Faroes and Røst (cluster A), median SD: 32.7° [29.2–35.5] for Isle of May, Ålesund and Sklinna (cluster B), Supporting information).

In spring, kittiwake colonies were grouped in four clusters (Fig. 5c–d, Supporting information). The wind support was higher for kittiwakes from the Norwegian west coast (5.52 m s<sup>-1</sup> [5.00–6.18], cluster B), and the southwest (Iceland and North Sea: 3.95 m s<sup>-1</sup> [3.42–4.46], cluster A), intermediate for the Barents Sea (excluding Bjørnøya and Kara Gate: 2.27 m s<sup>-1</sup> [1.44–3.26], cluster C) and lowest

for Bjørnøya and Kara Gate (1.39 m s<sup>-1</sup> [1.17–1.84], cluster D). The wind direction was most variable for kittiwakes from Bjørnøya and Kara Gate (median SD: 51.6° [47.6 – 57.3], cluster D), and less variable for the kittiwakes from the southwest (median SD: 30.4° [28.6 – 30.9], cluster A, Supporting information).

In autumn, puffins were grouped in four clusters (Fig. 5e–f, Supporting information). The wind support was lower for birds from the Isle of May and the Faroes (0.46 m s<sup>-1</sup> [0.04–0.75], cluster B), intermediate for birds from Iceland (0.66 m s<sup>-1</sup> [-0.08 to 1.18], cluster A) and higher for birds from the Barents Sea (0.81 m s<sup>-1</sup> [0.41–1.69], cluster D) and western Norway (0.94 m s<sup>-1</sup> [0.61–1.20], cluster C). Wind direction showed strong variability for birds migrating from the Norwegian colonies (median SD: 78.5° [74.5–83.1] for Runde, Sklinna, Røst and Anda (cluster C), median SD: 80.8° [75.1–84.2] for Hjelmøya and Hornøya (cluster D)), while it was moderately variable for birds from the Isle of May and the Faroes (median SD: 53.9° [53.3–57.9], cluster B), and less variable for birds from Iceland (median SD: 38.4° [32.7–40.1], cluster A, Supporting information).

In the spring, puffins were grouped in two clusters (Fig. 5g–h, Supporting information). Puffins from Iceland experienced higher wind support (1.50 m s<sup>-1</sup> [0.88–2.77], cluster A, versus 0.78 m s<sup>-1</sup> [0.41–1.10] for other colonies, cluster B). The wind direction was more predictable for puffins from Iceland (median SD: 40.1° [36.7–43.5], cluster A, versus 60.2° [58.4–64.7] for other colonies, cluster B).




Figure 5. Result of cluster analysis to investigate similarities in windscares encountered by kittiwakes (a)–(d) and puffins (e)–(h) from different colonies during autumn and spring migrations. Cluster analysis was made with a median value per colony of wind support, cross wind and wind speed (representing encountered wind conditions), mean proportion wet and detour (representing bird behaviour), and SD of wind direction and scalar average wind speed (representing average wind conditions). (a, c, e, g) location of the colonies and associated clusters by species and season; (b, d, f, h) violin plots showing for each cluster the distribution of the annual median of wind support (m s<sup>-1</sup>) per year and colony. Grey crosses represent median for each cluster. Black dots represent median values per colony and per year.



## Differences in wind support between seasons and species (obj. 4)

The wind support varied between species and seasons, and the seasonal difference in wind support was not the same for both species (i.e. the best model included a species  $\times$  season interaction; Table 1a–b). Both kittiwakes and puffins experienced a positive wind support on average during both autumn and spring migrations, and the wind support of kittiwakes was 2.7 times higher than the wind support of puffins (kittiwakes:  $2.66 \text{ m s}^{-1}$  [2.54–2.78] ; puffins:  $0.98 \text{ m s}^{-1}$  [0.85–1.10]). In addition, we observed seasonal differences. The wind support of kittiwakes was twice as large in spring than in autumn ( $3.61 \text{ m s}^{-1}$  [3.43–3.80] versus  $1.72 \text{ m s}^{-1}$  [1.58–1.86], Fig. 6). In comparison, puffins only had a slightly higher wind support in spring than in autumn ( $1.13 \text{ m s}^{-1}$  [0.91–1.34] versus  $0.84 \text{ m s}^{-1}$  [0.69–1.00], Fig. 6).

## Discussion

Our results highlight the importance of winds in shaping migratory routes in two seabird species despite their different flight capacity, suggesting that winds affect migration patterns across a wide range of species. Indeed, kittiwakes and puffins shared the same migratory pathways, following the prevailing winds in the North Atlantic. They had on average favourable wind conditions during their migratory movements, although puffins had generally lower wind support than kittiwakes. While both species experienced higher wind support during spring than autumn, there were also differences in wind support among colonies, depending on their location.

### Migratory corridors reflect prevailing winds favouring migratory movements

Kittiwakes and puffins had on average a positive wind support during both autumn and spring migrations. Main migratory routes were shared by both species and corresponded to areas where prevailing winds blew in a favourable direction. The general movement had a loop-shape, with birds flying southwards along the east coast of Greenland in the autumn and flying northwards along the coasts of the British Isles and

Norway in the spring. Birds that took other routes, such as birds flying northwards between Greenland and Iceland in the spring, or birds flying southwards southeast of Iceland during the autumn, had a negative wind support. The migratory corridors identified in our study are shared not only by these two species, but by other seabird species such as little auks *Alle alle*, Brünnich's guillemots *Uria lomvia* and long-tailed skuas *Stercorarius longicaudus* (Gilg et al. 2013, Amélineau et al. 2021).

Wind corridors have also been identified in other ocean basins. Some examples are the eight-shaped loop corridors of terns (Sternidae), Cory's shearwaters *Calonectris borealis* and skuas (Stercorariidae) across the Atlantic (Felicísimo et al. 2008, Wong et al. 2021), and of sooty shearwaters *Ardenna grisea* in the Pacific (Shaffer et al. 2006), as well as the circumpolar migration of wandering albatrosses *Diomedea exulans* following westerlies in the Southern Ocean (Weimerskirch et al. 2015).

Terrestrial birds and insects also follow most favourable wind corridors and can have a different route between autumn and spring migrations (Stefanescu et al. 2013, Chapman et al. 2015, Kranstauber et al. 2015, Åkesson et al. 2016, Vansteelant et al. 2017). In soaring terrestrial birds, migratory corridors are also dependent on the presence of updrafts and thermals, and tend to follow land topography favouring the formation of orographic updrafts and thermals (Nourani and Yamaguchi 2017). Flying with favourable winds is an efficient way to reduce both the costs of moving and the duration of migration.

### Geographical variation in winds and migratory behaviour

Inter-colony variation in wind conditions encountered while migrating was linked to the geographical location of the colonies. Birds from colonies situated at higher latitudes had in general higher wind support during autumn and lower wind support during spring than birds from colonies situated at lower latitudes, according to where the colonies were located relative to the favourable prevailing winds. There were however some exceptions. Puffins from Isle of May, Hjelmsøya and Hornøya migrated in several directions from the colony during the non-breeding season (dispersive migration). This

Table 1. Summary of the statistical model explaining wind support as a function of species and season. (a) model selection, (b) parameter estimation. The model with the lowest AIC is indicated in bold. Npar: number of parameters; CI: confidence interval. The marginal  $R^2$  (variance explained by the fixed effects) was 0.14, and the conditional  $R^2$  (variance explained by both fixed and random effects) was 0.22

(a) Models	npar	AIC	deltaAIC
<b>season <math>\times</math> species</b>	<b>8</b>	<b>23072.04</b>	<b>0.00</b>
season + species	7	23142.57	70.53
Season	6	23370.18	298.14
species	6	23376.20	304.16
intercept	5	23610.68	538.64
(b) Parameters	Estimate	2.5% CI	97.5% CI
Intercept	0.54	−0.04	1.11
season_spring	0.26	−0.05	0.57
species_kittiwake	1.24	0.93	1.55
season_spring:species_kittiwake	1.64	1.26	2.01




Figure 6. Wind support per species and migratory season. Boxplots show the median (thick line), the first (25%) and third (75%) quartiles (box), the lowest and highest values within 1.5 interquartile range (vertical segments) and outliers (dots). Notches represent the 95% confidence interval of the median ( $1.58 \times \text{interquartile range}/\sqrt{n}$ ).

contrasted with other colonies where birds from a given colony followed similar migratory routes and travelled over longer distances (Amélineau et al. 2021). While this behaviour is likely linked to the availability of resources, it could also reflect more unpredictable winds around these colonies. This is especially the case in the Barents Sea, where wind was least predictable. For kittiwakes, two colonies in the Barents Sea had a different migratory behaviour: Kara Gate and Bjørnøya. Contrary to all other kittiwake colonies for which the majority of individuals migrated to the Grand Banks area, many kittiwakes from Bjørnøya and Kara Gate migrated to the North Sea, and some from Kara Gate stayed at high latitudes during the winter (Fig. 2) (Amélineau et al. 2021). Kittiwakes from these two colonies had a particularly low wind support, especially in the spring, as well as a very high variability of the direction of encountered wind. This migratory behaviour contrasts with kittiwakes from other colonies of the Barents Sea such as Spitsbergen or Franz Josef Land, that travel to the Grand Banks during the winter and encounter higher wind support and more predictable winds. Overall, migration destination will be a tradeoff between the relative profitability of the breeding and wintering grounds, and the costs of travelling between them. If in some cases birds seem to be making sub-optimal decisions in terms of travel costs, it could be because the increase in profitability at the destination offsets travel costs. Such differences in migratory distance and wind conditions encountered are likely resulting in different energetic balance, and may have further implications for subsequent timing of breeding, carry-over effects, or survival (Alerstam 2011, Weimerskirch et al. 2015, Fayet et al. 2017) but see (Dufour et al. 2021). Such differences in windscapes will be larger for colonies that are further away from each other, compared to colonies that are closer to each other.

## The use of windscapes varies among species

Puffins had a lower wind support than kittiwakes during both autumn and spring migrations, suggesting that they used wind conditions differently during migration. The main differences between the two species are their foraging mode and their morphology. Puffins are divers, and their wings, used for swimming and flying, are shorter compared to kittiwakes that are surface feeders and forage from the air. As a consequence, puffins have a higher wing loading (Pennycuik 1987). Puffins could be less selective towards favourable winds due to their higher wing loading, as birds with higher wing loading have a lower increase in airspeed (and thus energetic costs) when wind speed increases or when wind direction is less favourable (Spear and Ainley 1997a). Moreover, their migratory range is different, with puffins migrating shorter distances than kittiwakes (Amélineau et al. 2021). During migratory bouts, puffins spend also less time flying daily (18% on average, versus 30% in kittiwakes) and have a lower movement rate ( $6.25 \text{ km h}^{-1}$  on average, versus  $12.37 \text{ km h}^{-1}$  in kittiwakes (Amélineau et al. 2021)). The best strategy for puffins could be to use more direct routes with low wind support, which could be beneficial for shorter distance migration, while when migrating farther and faster such as the kittiwakes, it could be more advantageous to have a route with better wind support. Their different migratory strategies are linked to a different use of wind.

More generally, wing shape seems under selective pressure in migrants: migratory birds tend to have more pointed wing-tips (Lockwood et al. 1998, Sheard et al. 2020) and migratory insects tend to have higher aspect ratio than non-migrants (Le Roy et al. 2019), although this is not always the case as other selective pressures can impact wing shape, such as the need for manoeuvrability in stationary habitats (Le Roy et al. 2019), or the use of wings to swim (Elliott et al. 2013). Ultimately, wing shape will be linked to flight abilities and the use of winds (Spear and Ainley 1997b, Dehnhard et al. 2021).

## A higher wind support in the spring

For both species, wind support was higher during spring than autumn. This has also been observed in other species such as Arctic terns *Sterna paradisaea* (Hromádková et al. 2020), and nocturnal terrestrial migrant birds through Europe (Kemp et al. 2010). The fact that winds were more favourable during the inbound migration can be linked to prevailing winds at the latitudes where these species migrate (Kemp et al. 2010), and to the detour that some transequatorial migrants do to follow prevailing winds (Felicísimo et al. 2008, Hromádková et al. 2020). Having more favourable winds during the inbound migration might ensure higher fitness, allowing earlier arrival at the colony which is generally associated with higher breeding success (Kokko 1999, Verhulst and Nilsson 2008) or reducing energy expenditure, hence leading to a better body condition at the start of the breeding season, and subsequently a higher reproductive output (Drake et al. 2014, de Zwaan et al. 2022). More generally,

stronger time selection during spring migration compared to the autumn has been found for migratory birds (Nilsson et al. 2013), which supports the positive influence of early arrival on fitness. How wind conditions encountered during the non-breeding season, and especially during the spring migration, impact arrival date, body condition and reproductive performances in migratory birds remains to be tested.

### Can winds influence seabird species distribution?

Adult kittiwakes and puffins show a high individual repeatability in their migratory routes (Guilford et al. 2011, Léandri-Breton et al. 2021). Such a repeatability in routes support the hypothesis that individuals migrate along routes that optimize wind conditions and have been selected through time. Indeed, one could expect a high individual variability in routes if birds follow winds encountered along the way, especially in regions with high wind variability. Selection of optimized routes is possible when there is genetic inheritance of migratory routes (Berthold 1991, Liedvogel et al. 2011), or cultural inheritance with a transmission via one of the parents or other individuals (Jesmer et al. 2018, Byholm et al. 2022). Additionally, the selection of optimized migratory routes could arise through learning, if the individuals refine their routes over the course of their life, as found in sea turtles (Scott et al. 2014), black kites *Milvus migrans* (Sergio et al. 2014) and Cory's shearwaters (Campioni et al. 2020). Further investigations are needed on the individual variability in migratory routes and in encountered wind supports to deepen our understanding of the selection of migratory routes at the individual scale.

If wind shapes migratory routes, we can further question whether wind could be a factor influencing species distribution. For a given species, some areas might be difficult to exploit due to unfavourable atmospheric conditions (Krietsch et al. 2020). Moreover, if different populations face contrasting wind support, one could expect differences in terms of average individual fitness among them, and further local impacts on population dynamics (Newton 2006, Reiertsen et al. 2021). Indeed, wind has been identified as a factor explaining the global distribution of Procellariiforms and of some albatross species in the North and Central Pacific (Suryan et al. 2008, Davies et al. 2010). Wind conditions also explain the spatial segregation between male and female black-browed and grey-headed albatrosses (*Thalassarche melanophris* and *T. chrysostoma*), where males with a higher wing loading forage at higher latitudes where wind speeds are higher (Phillips et al. 2004). How wind would influence the distribution of other seabirds and flying migrants in general remains to be tested, keeping in mind that many other factors are involved in shaping a species' distribution, such as food availability, habitat, competitors or predators.

In the context of climate change, wind conditions are expected to change according to the Intergovernmental Panel on Climate Change (IPCC) predictions (Pinto and Ludwig 2020, Masson-Delmotte et al. 2021). It is therefore important to understand how migratory species will respond and

cope with these changes. While phylogenetic studies suggest that migratory behaviour can evolve relatively quickly in birds, based on the great range of migratory behaviours observed within taxa (Alerstam et al. 2003), the rapidity of ongoing changes means that plasticity would be much more important than the potential for evolution. More studies investigating the plasticity of birds in different wind conditions as well as temporal variations in wind conditions are thus needed, and will require long term tracking programs.

*Acknowledgements* – We thank the numerous fieldworkers for their hard work collecting the data, and to landowners for access and logistical support.

*Funding* – The SEATRACK project ([www.seapop.no/en/seatrack](http://www.seapop.no/en/seatrack)) was funded by the Norwegian Ministry for Climate and the Environment, the Norwegian Ministry of Foreign Affairs and the Norwegian Oil and Gas Association along with eight oil companies. The SEAPOPOP program ([www.seapop.no](http://www.seapop.no), grant no. 192141) supported the fieldwork in Norway (including Svalbard and Jan Mayen). Kittiwake fieldwork in the Kongsfjord was supported by the French Polar Institute (IPEV, prog. 330 to O. Chastel). The work on the Isle of May was also supported by the Natural Environment Research Council (Award NE/R016429/1 as part of the UK-SCaPE programme delivering National Capability).

*Permits* – All bird handling was done according to the national guidelines and under the necessary field work permits for each colony.

### Author contributions

**Françoise Amélineau:** Conceptualization (lead); Data curation (equal); Formal analysis (lead); Visualization (lead); Writing – original draft (lead); Writing – review and editing (equal). **Arnaud Tarroux:** Conceptualization (lead); Data curation (equal); Formal analysis (supporting); Funding acquisition (supporting); Supervision (equal); Visualization (supporting); Writing – original draft (supporting); Writing – review and editing (equal). **Simon Lacombe:** Conceptualization (supporting); Formal analysis (supporting); Writing – review and editing (equal). **Vegard S. Bråthen:** Data curation (equal); Writing – review and editing (equal). **Sebastien Descamps:** Data curation (equal); Funding acquisition (supporting); Investigation (equal); Supervision (equal); Writing – review and editing (equal). **Morten Ekker:** Data curation (equal); Funding acquisition (supporting); Writing – review and editing (equal). **Per Fauchald:** Data curation (equal); Funding acquisition (supporting); Writing – review and editing (equal). **Malin K. Johansen:** Data curation (equal); Writing – review and editing (equal). **Børge Moe:** Data curation (equal); Funding acquisition (supporting); Investigation (equal); Writing – review and editing (equal). **Tycho Anker-Nilssen:** Investigation (equal); Writing – review and editing (equal). **Maria I. Bogdanova:** Investigation (equal); Writing – review and editing (equal). **Ingar S. Bringvor:** Investigation (equal); Writing – review and editing (equal). **Olivier Chastel:** Investigation (equal); Writing – review and editing (equal). **Signe Christensen-Dalsgaard:** Investigation (equal); Writing – review and

editing (equal). **Francis Daunt**: Investigation (equal); Writing – review and editing (equal). **Nina Dehnhard**: Investigation (equal); Writing – review and editing (equal). **Kjell Einar Erikstad**: Investigation (equal); Writing – review and editing (equal). **Aleksey Ezhov**: Investigation (equal); Writing – review and editing (equal). **Maria Gavrilov**: Investigation (equal); Writing – review and editing (equal). **Erpur S. Hansen**: Investigation (equal); Writing – review and editing (equal). **Mike P. Harris**: Investigation (equal); Writing – review and editing (equal). **Hálfdán H. Helgason**: Data curation (equal); Investigation (equal); Writing – review and editing (equal). **Magdalene Langset**: Investigation (equal); Writing – review and editing (equal). **Don-Jean Léandri-Breton**: Conceptualization (supporting); Writing – review and editing (equal). **Svein-Håkon Lorentsen**: Investigation (equal); Writing – review and editing (equal). **Benjamin Merkel**: Data curation (equal); Writing – review and editing (equal). **Mark Newell**: Investigation (equal); Writing – review and editing (equal). **Bergur Olsen**: Investigation (equal); Writing – review and editing (equal). **Tone K. Reiertsen**: Investigation (equal); Writing – review and editing (equal). **Geir H. R. Systad**: Investigation (equal); Writing – review and editing (equal). **Thorkell L. Thorarinsson**: Investigation (equal); Writing – review and editing (equal). **Jens Åström**: Data curation (equal); Writing – review and editing (equal). **Hallvard Strøm**: Data curation (equal); Funding acquisition (lead); Investigation (equal); Project administration (lead); Supervision (equal); Writing – review and editing (equal).

### Transparent peer review

The peer review history for this article is available at <https://publons.com/publon/10.1111/ecog.06496>.

### Data availability statement

All the R code is publicly available on GitHub (<https://github.com/FAmelineau/SeaWinds3>). Contains modified Copernicus Climate Change Service information [2021]. Data are available from the Dryad Digital Repository: <https://doi.org/10.5061/dryad.z34tmgpm6> (Amélineau et al. 2023).

### Supporting information

The Supporting information associated with this article is available with the online version.

### References

Agostinelli, C. and Lund, U. 2022. R package circular: circular statistics, ver. 0.4-94, <https://r-forge.r-project.org/projects/circular/>.  
 Ahrens, C. D. 2014. Essentials of meteorology: an invitation to the atmosphere. – Cengage Learning.  
 Åkesson, S., Bianco, G. and Hedenström, A. 2016. Negotiating an ecological barrier: crossing the Sahara in relation to winds by common swifts. – *Phil. Trans. R. Soc. B* 371: 20150393.  
 Alerstam, T. 2011. Optimal bird migration revisited. – *J. Ornithol.* 152: 5–23.

Alerstam, T., Hedenström, A. and Åkesson, S. 2003. Long-distance migration: evolution and determinants. – *Oikos* 103: 247–260.  
 Amélineau, F., Péron, C., Lescroël, A., Authier, M., Provost, P. and Grémillet, D. 2014. Windscape and tortuosity shape the flight costs of northern gannets. – *J. Exp. Biol.* 217: 876–885.  
 Amélineau, F. et al. 2021. Six pelagic seabird species of the North Atlantic engage in a fly-and-forage strategy during their migratory movements. – *Mar. Ecol. Prog. Ser.* 676: 127–144.  
 Amélineau, F. et al. 2023. Data from: Multi-colony tracking of two pelagic seabirds with contrasting flight capability illustrates how windscapes shape migratory movements at an ocean-basin scale. – Dryad Digital Repository, <https://doi.org/10.5061/dryad.z34tmgpm6>.  
 Bartoń, K. 2022. MuMIn: multi-model inference. – R package ver. 1.47.1, <https://CRAN.R-project.org/package=MuMIn>.  
 Bates, D., Mächler, M., Bolker, B. and Walker, S. 2015. Fitting linear mixed-effects models using lme4. – *J. Stat. Softw.* 67: 1–48.  
 Berthold, P. 1991. Genetic control of migratory behaviour in birds. – *Trends Ecol. Evol.* 6: 254–257.  
 Bogdanova, M. I., Butler, A., Wanless, S., Moe, B., Anker-Nilssen, T., Frederiksen, M., Boulinier, T., Chivers, L. S., Christensen-Dalsgaard, S. and Descamps, S. 2017. Multi-colony tracking reveals spatio-temporal variation in carry-over effects between breeding success and winter movements in a pelagic seabird. – *Mar. Ecol. Prog. Ser.* 578: 167–181.  
 Borkenhagen, K., Corman, A.-M. and Garthe, S. 2018. Estimating flight heights of seabirds using optical rangefinders and GPS data loggers: a methodological comparison. – *Mar. Biol.* 165: 1–7.  
 Bowlin, M. S. and Wikelski, M. 2008. Pointed wings, low wing-loading and calm air reduce migratory flight costs in songbirds. – *PLoS One* 3: e2154.  
 Bowlin, M. S., Bisson, I.-A., Shamoun-Baranes, J., Reichard, J. D., Sapir, N., Marra, P. P., Kunz, T. H., Wilcove, D. S., Hedenström, A. and Guglielmo, C. G. 2010. Grand challenges in migration biology. – *Integr. Comp. Biol.* 50: 261–279.  
 Bråthen, V. S., Moe, B., Amélineau, F., Ekker, M., Fauchald, P., Helgason, H. H., Johansen, M. K., Merkel, B., Tarroux, A., Åström, J. and Strøm, H. 2021. An automated procedure (v2.0) to obtain positions from light-level geolocators in large-scale tracking of seabirds. A method description for the SEATRACK project. – Norwegian Institute for Nature Research (NINA).  
 Burnham, K. P. and Anderson, D. R. 2002. Model selection and multimodel inference: a practical information-theoretic approach. – Springer.  
 Byholm, P., Beal, M., Isaksson, N., Lötberg, U. and Åkesson, S. 2022. Paternal transmission of migration knowledge in a long-distance bird migrant. – *Nat. Commun.* 13: 1566.  
 Calenge, C. 2006. The package “adehabitat” for the R software: a tool for the analysis of space and habitat use by animals. – *Ecol. Model.* 197: 516–519.  
 Campioni, L., Dias, M. P., Granadeiro, J. P. and Catry, P. 2020. An ontogenetic perspective on migratory strategy of a long-lived pelagic seabird: timings and destinations change progressively during maturation. – *J. Anim. Ecol.* 89: 29–43.  
 Chapman, J. W., Nesbit, R. L., Burgin, L. E., Reynolds, D. R., Smith, A. D., Middleton, D. R. and Hill, J. K. 2010. Flight orientation behaviors promote optimal migration trajectories in high-flying insects. – *Science* 327: 682–685.  
 Chapman, J. W., Klaassen, R. H. G., Drake, V. A., Fossette, S., Hays, G. C., Metcalfe, J. D., Reynolds, A. M., Reynolds, D.

- R. and Alerstam, T. 2011. Animal orientation strategies for movement in flows. – *Curr. Biol.* 21: R861–R870.
- Chapman, J. W., Reynolds, D. R. and Wilson, K. 2015. Long-range seasonal migration in insects: mechanisms, evolutionary drivers and ecological consequences. – *Ecol. Lett.* 18: 287–302.
- Collins, P. M., Green, J. A., Elliott, K. H., Shaw, P. J. A., Chivers, L., Hatch, S. A. and Halsey, L. G. 2020. Coping with the commute: behavioural responses to wind conditions in a foraging seabird. – *J. Avian Biol.* 2020: e02057.
- Davies, R. G., Irllich, U. M., Chown, S. L. and Gaston, K. J. 2010. Ambient, productive and wind energy, and ocean extent predict global species richness of procellariiform seabirds. – *Global Ecol. Biogeogr.* 19: 98–110.
- de Zwaan, D. R., Drake, A., Camfield, A. F., MacDonald, E. C. and Martin, K. 2022. The relative influence of cross-seasonal and local weather effects on the breeding success of a migratory songbird. – *J. Anim. Ecol.* 91: 1458–1470.
- Dehnhard, N., Klekociuk, A. R. and Emmerson, L. 2021. Interactive effects of body mass changes and species-specific morphology on flight behavior of chick-rearing Antarctic fulmarine petrels under diurnal wind patterns. – *Ecol. Evol.* 11: 4972–4991.
- Desprez, M., Jenouvrier, S., Barbraud, C., Delord, K. and Weimerskirch, H. 2018. Linking oceanographic conditions, migratory schedules and foraging behaviour during the non-breeding season to reproductive performance in a long-lived seabird. – *Funct. Ecol.* 32: 2040–2053.
- Di Musciano, M., Di Cecco, V., Bartolucci, F., Conti, F., Fratrotoli, A. R. and Di Martino, L. 2020. Dispersal ability of threatened species affects future distributions. – *Plant Ecol.* 221: 265–281.
- Drake, A., Rock, C. A., Quinlan, S. P., Martin, M. and Green, D. J. 2014. Wind speed during migration influences the survival, timing of breeding, and productivity of a neotropical migrant, *Setophaga petechia*. – *PLoS One* 9: e97152.
- Dufour, P., Wojczulanis-Jakubas, K., Lavergne, S., Renaud, J., Jakubas, D. and Descamps, S. 2021. A two-fold increase in migration distance does not have breeding consequences in a long-distance migratory seabird with high flight costs. – *Mar. Ecol. Prog. Ser.* 676: 117–126.
- Egevang, C., Stenhouse, I. J., Phillips, R. A., Petersen, A., Fox, J. W. and Silk, J. R. D. 2010. Tracking of Arctic terns *Sterna paradisaea* reveals longest animal migration. – *Proc. Natl Acad. Sci. USA* 107: 2078–2081.
- Elliott, K. H., Ricklefs, R. E., Gaston, A. J., Hatch, S. A., Speakman, J. R. and Davoren, G. K. 2013. High flight costs, but low dive costs, in auks support the biomechanical hypothesis for flightlessness in penguins. – *Proc. Natl Acad. Sci. USA* 110: 9380–9384.
- Elliott, K. H., Chivers, L. S., Bessey, L., Gaston, A. J., Hatch, S. A., Kato, A., Osborne, O., Ropert-Coudert, Y., Speakman, J. R. and Hare, J. F. 2014. Windscape shape seabird instantaneous energy costs but adult behavior buffers impact on offspring. – *Movement Ecol.* 2: 17.
- Epele, L. B., Dos Santos, D. A., Sarremejane, R., Grech, M. G., Macchi, P. A., Manzo, L. M., Miserendino, M. L., Bonada, N. and Cañedo-Argüelles, M. 2021. Blowin' in the wind: wind directionality affects wetland invertebrate metacommunities in Patagonia. – *Global Ecol. Biogeogr.* 30: 1191–1203.
- Ezhov, A. V., Gavrilov, M. V., Krasnov, Y. V., Bräthen, V. S., Moe, B., Baranskaya, A. V. and Strøm, H. 2021. Transpolar and bidirectional migration strategies of black-legged kittiwakes *Rissa tridactyla* from a colony in Novaya Zemlya, Barents Sea, Russia. – *Mar. Ecol. Prog. Ser.* 676: 189–203.
- Fauchald, P., Tarrow, A., Bräthen, V. S., Descamps, S., Ekker, M., Helgason, H. H., Merkel, B., Moe, B., Åström, J. and Strøm, H. 2019. Arctic-breeding seabirds' hotspots in space and time – a methodological framework for year-round modelling of environmental niche and abundance using light-logger data. – Norsk Institutt for Naturforskning (NINA).
- Fayet, A. L., Freeman, R., Anker-Nilssen, T., Diamond, A., Erikstad, K. E., Fifield, D., Fitzsimmons, M. G., Hansen, E. S., Harris, M. P. and Jessopp, M. 2017. Ocean-wide drivers of migration strategies and their influence on population breeding performance in a declining seabird. – *Curr. Biol.* 27: 3871–3878.
- Felícísimo, Á. M., Muñoz, J. and González-Solis, J. 2008. Ocean surface winds drive dynamics of transoceanic aerial movements. – *PLoS One* 3: e2928.
- Fort, J., Moe, B., Strøm, H., Grémillet, D., Welcker, J., Schultner, J., Jerstad, K., Johansen, K. L., Phillips, R. A. and Mosbech, A. 2013. Multicolony tracking reveals potential threats to little auks wintering in the North Atlantic from marine pollution and shrinking sea ice cover. – *Divers. Distrib.* 19: 1322–1332.
- Frederiksen, M. et al. 2012. Multicolony tracking reveals the winter distribution of a pelagic seabird on an ocean basin scale. – *Divers. Distrib.* 18: 530–542.
- Frederiksen, M. et al. 2016. Migration and wintering of a declining seabird, the thick-billed murre *Uria lomvia*, on an ocean basin scale: conservation implications. – *Biol. Conserv.* 200: 26–35.
- Geisler, J., Madsen, J., Nolet, B. A. and Schreven, K. H. T. 2022. Sea crossings of migratory pink-footed geese: seasonal effects of winds on flying and stopping behaviour. – *J. Avian Biol.* 2022: e02985.
- Gilg, O., Moe, B., Hanssen, S. A., Schmidt, N. M., Sittler, B., Hansen, J., Reneerkens, J., Sabard, B., Chastel, O., Moreau, J., Phillips, R. A., Oudman, T., Biersma, E. M., Fenstad, A. A., Lang, J. and Bollache, L. 2013. Trans-equatorial migration routes, staging sites and wintering areas of a high-arctic avian predator: the long-tailed skua (*Stercorarius longicaudus*). – *PLoS One* 8: e64614.
- Gill, R. E., Tibbitts, T. L., Douglas, D. C., Handel, C. M., Mulcahy, D. M., Gottschalck, J. C., Warnock, N., McCaffery, B. J., Batley, P. F. and Piersma, T. 2009. Extreme endurance flights by landbirds crossing the Pacific Ocean: ecological corridor rather than barrier? – *Proc. R. Soc. B* 276: 447–457.
- Guilford, T., Freeman, R., Boyle, D., Dean, B., Kirk, H., Phillips, R. and Perrins, C. 2011. A dispersive migration in the Atlantic puffin and its implications for migratory navigation. – *PLoS One* 6: e21336.
- Halpin, L. R., Ross, J. D., Ramos, R., Mott, R., Carlile, N., Golding, N., Reyes-González, J. M., Militão, T., De Felipe, F., Zajková, Z., Cruz-Flores, M., Saldanha, S., Morera-Pujol, V., Navarro-Herrero, L., Zango, L., González-Solis, J. and Clarke, R. H. 2021. Double-tagging scores of seabirds reveals that light-level geolocator accuracy is limited by species idiosyncrasies and equatorial solar profiles. – *Methods Ecol. Evol.* 12: 2243–2255.
- Hartigan, J. A. and Wong, M. A. 1979. Algorithm AS 136: a k-means clustering algorithm. – *J. R. Stat. Soc., Ser. C (Appl. Stat.)* 28: 100–108.
- Hays, G. C., Ferreira, L. C., Sequeira, A. M., Meekan, M. G., Duarte, C. M., Bailey, H., Bailleul, F., Bowen, W. D., Caley, M. J. and Costa, D. P. 2016. Key questions in marine megafauna movement ecology. – *Trends Ecol. Evol.* 31: 463–475.
- Hedrick, T. L., Pichot, C. and de Margerie, E. 2018. Gliding for a free lunch: biomechanics of foraging flight in common swifts (*Apus apus*). – *J. Exp. Biol.* 221: jeb186270.

- Hersbach, H. et al. 2020. The ERA5 global reanalysis. – *Q. J. R. Meteorol. Soc.* 146: 1999–2049.
- Hijmans, R. J. 2021. geosphere: spherical trigonometry. – R package ver. 1.5-14, <https://CRAN.R-project.org/package=geosphere>.
- Horton, K. G., Van Doren, B. M., La Sorte, F. A., Fink, D., Sheldon, D., Farnsworth, A. and Kelly, J. F. 2018. Navigating north: how body mass and winds shape avian flight behaviours across a North American migratory flyway. – *Ecol. Lett.* 21: 1055–1064.
- Hromádková, T., Pavel, V., Flousek, J. and Briedis, M. 2020. Seasonally specific responses to wind patterns and ocean productivity facilitate the longest animal migration on Earth. – *Mar. Ecol. Prog. Ser.* 638: 1–12.
- Jesmer, B. R., Merkle, J. A., Goheen, J. R., Aikens, E. O., Beck, J. L., Courtemanch, A. B., Hurley, M. A., McWhirter, D. E., Miyasaki, H. M., Monteith, K. L. and Kauffman, M. J. 2018. Is ungulate migration culturally transmitted? Evidence of social learning from translocated animals. – *Science* 361: 1023–1025.
- Johnston, A., Cook, A. S. C. P., Wright, L. J., Humphreys, E. M. and Burton, N. H. K. 2014. Modelling flight heights of marine birds to more accurately assess collision risk with offshore wind turbines. – *J. Appl. Ecol.* 51: 31–41.
- Kassambara, A. and Mundt, F. 2020. factoextra: extract and visualize the results of multivariate data analyses. – R package ver. 1.0.7, <https://CRAN.R-project.org/package=factoextra>.
- Kemp, M. U., Shamoun-Baranes, J., Gasteren, H. V., Bouten, W. and Loon, E. E. V. 2010. Can wind help explain seasonal differences in avian migration speed? – *J. Avian Biol.* 41: 672–677.
- Kling, M. M. and Ackerly, D. D. 2021. Global wind patterns shape genetic differentiation, asymmetric gene flow, and genetic diversity in trees. – *Proc. Natl Acad. Sci. USA* 118: e2017317118.
- Kokko, H. 1999. Competition for early arrival in migratory birds. – *J. Anim. Ecol.* 68: 940–950.
- Kranstauber, B., Weinzierl, R., Wikelski, M. and Safi, K. 2015. Global aerial flyways allow efficient travelling. – *Ecol. Lett.* 18: 1338–1345.
- Krietsch, J., Valcu, M. and Kempenaers, B. 2020. Wind conditions influence breeding season movements in a nomadic polygynous shorebird. – *Proc. R. Soc. B* 287: 20192789.
- Le Roy, C., Debat, V. and Llaurens, V. 2019. Adaptive evolution of butterfly wing shape: from morphology to behaviour. – *Biol. Rev.* 94: 1261–1281.
- Léandri-Breton, D.-J., Tarroux, A., Elliott, K. H., Legagneux, P., Angelier, F., Blévin, P., Bräthen, V. S., Fauchald, P., Goutte, A., Jouanneau, W., Tartu, S., Moe, B. and Chastel, O. 2021. Long-term tracking of an Arctic-breeding seabird indicates high fidelity to pelagic wintering areas. – *Mar. Ecol. Prog. Ser.* 676: 205–218.
- Liechti, F. 2006. Birds: Blowin' by the wind? – *J. Ornithol.* 147: 202–211.
- Liedvogel, M., Åkesson, S. and Bensch, S. 2011. The genetics of migration on the move. – *Trends Ecol. Evol.* 26: 561–569.
- Lisovski, S., Bauer, S., Briedis, M., Davidson, S. C., Dhanjal-Adams, K. L., Hallworth, M. T., Karagicheva, J., Meier, C. M., Merkel, B. and Ouweland, J. 2020. Light-level geolocator analyses: a user's guide. – *J. Anim. Ecol.* 89: 221–236.
- Lockwood, R., Swaddle, J. P. and Rayner, J. M. 1998. Avian wingtip shape reconsidered: wingtip shape indices and morphological adaptations to migration. – *J. Avian Biol.* 29: 273–292.
- Masson-Delmotte, V., Zhai, P., Pirani, A., Connors, S. L., Péan, C., Berger, S., Caud, N., Chen, Y., Goldfarb, L., Gomis, M. I., Huang, M., Leitzell, K., Lonnoy, E., Matthews, J. B. R., Maycock, T. K., Waterfield, T., Yelekçi, Ö., Yu, R. and Zhou, B. (Eds). 2021. Climate change 2021: the physical science basis. contribution of working group I to the sixth assessment report of the intergovernmental panel on climate change. – Cambridge Univ. Press.
- Mateos-Rodríguez, M. and Liechti, F. 2012. How do diurnal long-distance migrants select flight altitude in relation to wind? – *Behav. Ecol.* 23: 403–409.
- Merkel, B., Descamps, S., Yoccoz, N. G., Grémillet, D., Fauchald, P., Danielsen, J., Daunt, F., Erikstad, K. E., Ezhov, A. V. and Harris, M. P. 2021. Strong migratory connectivity across metapopulations of sympatric North Atlantic seabirds. – *Mar. Ecol. Prog. Ser.* 676: 173–188.
- Muñoz, J., Felicísimo, Á. M., Cabezas, F., Burgaz, A. R. and Martínez, I. 2004. Wind as a long-distance dispersal vehicle in the Southern Hemisphere. – *Science* 304: 1144–1147.
- Newton, I. 2006. Can conditions experienced during migration limit the population levels of birds? – *J. Ornithol.* 147: 146–166.
- Newton, I. 2010. The migration ecology of birds. – Academic Press.
- Nilsson, C., Klaassen, R. H. and Alerstam, T. 2013. Differences in speed and duration of bird migration between spring and autumn. – *Am. Nat.* 181: 837–845.
- Nourani, E. and Yamaguchi, N. M. 2017. The effects of atmospheric currents on the migratory behavior of soaring birds: a review. – *Ornithol. Sci.* 16: 5–15.
- Nourani, E. et al. 2023. Seabird morphology determines operational wind speeds, tolerable maxima, and responses to extremes. – *Curr. Biol.* 33: 1179–1184.e3.
- Pennycuik, C. J. 1987. Flight of auks (*Alcidae*) and other northern seabirds compared with southern procellariiformes: Ornithodolite observations. – *Deep Sea Res. Part B. Oceanogr. Lit. Rev.* 128: 335–347.
- Phillips, R. A., Silk, J. R. D., Phalan, B., Catry, P. and Croxall, J. P. 2004. Seasonal sexual segregation in two *Thalassarche* albatross species: competitive exclusion, reproductive role specialization or foraging niche divergence? – *Proc. R. Soc. B* 271: 1283–1291.
- Pinto, J. G. and Ludwig, P. 2020. Extratropical cyclones over the North Atlantic and western Europe during the Last Glacial Maximum and implications for proxy interpretation. – *Climate Past* 16: 611–626.
- Reiertsen, T. K. et al. 2021. Inter-population synchrony in adult survival and effects of climate and extreme weather in non-breeding areas of Atlantic puffins. – *Mar. Ecol. Prog. Ser.* 676: 219–231.
- Safi, K., Kranstauber, B., Weinzierl, R., Griffin, L., Rees, E. C., Cabot, D., Cruz, S., Proaño, C., Takekawa, J. Y. and Newman, S. H. 2013. Flying with the wind: scale dependency of speed and direction measurements in modelling wind support in avian flight. – *Movement Ecol.* 1: 4.
- Scott, R., Marsh, R. and Hays, G. C. 2014. Ontogeny of long distance migration. – *Ecology* 95: 2840–2850.
- Sergio, F., Tanferna, A., De Stephanis, R., Jiménez, L. L., Blas, J., Tavecchia, G., Preatoni, D. and Hiraldo, F. 2014. Individual improvements and selective mortality shape lifelong migratory performance. – *Nature* 515: 410–413.
- Shaffer, S. A., Tremblay, Y., Weimerskirch, H., Scott, D., Thompson, D. R., Sagar, P. M., Moller, H., Taylor, G. A., Foley, D. G., Block, B. A. and Costa, D. P. 2006. Migratory shearwaters integrate oceanic resources across the Pacific Ocean in an endless summer. – *Proc. Natl Acad. Sci. USA* 103: 12799–12802.

- Shamoun-Baranes, J., Liechti, F. and Vansteelant, W. M. G. 2017. Atmospheric conditions create freeways, detours and tailbacks for migrating birds. – *J. Comp. Physiol. A* 203: 509–529.
- Sheard, C., Neate-Clegg, M. H. C., Alioravainen, N., Jones, S. E. I., Vincent, C., MacGregor, H. E. A., Bregman, T. P., Claramunt, S. and Tobias, J. A. 2020. Ecological drivers of global gradients in avian dispersal inferred from wing morphology. – *Nat. Commun.* 11: 2463.
- Sjöberg, S., Malmiga, G., Nord, A., Andersson, A., Bäckman, J., Tarka, M., Willemoes, M., Thorup, K., Hansson, B. and Alerstam, T. 2021. Extreme altitudes during diurnal flights in a nocturnal songbird migrant. – *Science* 372: 646–648.
- Spear, L. B. and Ainley, D. G. 1997a. Flight speed of seabirds in relation to wind speed and direction. – *Ibis* 139: 234–251.
- Spear, L. B. and Ainley, D. G. 1997b. Flight behaviour of seabirds in relation to wind direction and wing morphology. – *Ibis* 139: 221–233.
- Stefanescu, C. et al. 2013. Multi-generational long-distance migration of insects: studying the painted lady butterfly in the Western Palearctic. – *Ecography* 36: 474–486.
- Strøm, H., Descamps, S., Ekker, M., Fauchald, P. and Moe, B. 2021. Tracking the movements of North Atlantic seabirds: steps towards a better understanding of population dynamics and marine ecosystem conservation. – *Mar. Ecol. Prog. Ser.* 676: 97–116.
- Suryan, R. M., Anderson, D. J., Shaffer, S. A., Roby, D. D., Tremblay, Y., Costa, D. P., Sievert, P. R., Sato, F., Ozaki, K. and Balogh, G. R. 2008. Wind, waves, and wing loading: morphological specialization may limit range expansion of endangered albatrosses. – *PLoS One* 3: e4016.
- Tarroux, A., Weimerskirch, H., Wang, S.-H., Bromwich, D. H., Cherel, Y., Kato, A., Ropert-Coudert, Y., Varpe, Ø., Yoccoz, N. G. and Descamps, S. 2016. Flexible flight response to challenging wind conditions in a commuting Antarctic seabird: do you catch the drift? – *Anim. Behav.* 113: 99–112.
- Technitis, G., Othman, W., Safi, K. and Weibel, R. 2015. From A to B, randomly: a point-to-point random trajectory generator for animal movement. – *Int. J. Geograph. Inform. Sci.* 29: 912–934.
- Vansteelant, W. M., Shamoun-Baranes, J., van Manen, W., van Diermen, J. and Bouten, W. 2017. Seasonal detours by soaring migrants shaped by wind regimes along the East Atlantic Flyway. – *J. Anim. Ecol.* 86: 179–191.
- Verhulst, S. and Nilsson, J.-Å. 2008. The timing of birds' breeding seasons: a review of experiments that manipulated timing of breeding. – *Phil. Trans. R. Soc. B* 363: 399–410.
- Weimerskirch, H., Guionnet, T., Martin, J., Shaffer, S. A. and Costa, D. P. 2000. Fast and fuel efficient? Optimal use of wind by flying albatrosses. – *Proc. R. Soc. B* 267: 1869–1874.
- Weimerskirch, H., Louzao, M., de Grissac, S. and Delord, K. 2012. Changes in wind pattern alter albatross distribution and life-history traits. – *Science* 335: 211–214.
- Weimerskirch, H., Delord, K., Guitteaud, A., Phillips, R. A. and Pinet, P. 2015. Extreme variation in migration strategies between and within wandering albatross populations during their sabbatical year and their fitness consequences. – *Sci. Rep.* 5: 8853.
- Wong, J., Lisovski, S., Alisauskas, R., English, W., Giroux, M., Harrison, A., Kellett, D., Lecomte, N., Maffei, M., Nagy-MacArthur, A., Ronconi, R., Smith, P., Mallory, M. and Auger-Méthé, M. 2021. Arctic terns from circumpolar breeding colonies share common migratory routes. – *Mar. Ecol. Prog. Ser.* 671: 191–206.
- Zuur, A. F., Ieno, E. N., Walker, N. J., Saveliev, A. A. and Smith, G. M. 2009. Mixed effects models and extensions in ecology with R. – Springer.