

Resolving archaeological populations with Sr-isotope mixing modelsJanet Montgomery^{‡*}, Jane A. Evans⁺ and Rachel E. Cooper[‡]

[‡] *Archaeological, Geographical and Environmental Sciences, School of Life Sciences, University of Bradford, Bradford, West Yorkshire BD7 1DP, UK*

⁺ *NERC Isotope Geosciences Laboratory, British Geological Survey, Keyworth, Nottinghamshire, NG12 5GG, UK*

*Corresponding author: email: J.Montgomery@bradford.ac.uk

Abstract

Strontium isotope analysis of tooth enamel is a useful provenancing technique to investigate the childhood origins and residential mobility of ancient people. However, where different geographical target regions have similar biosphere $^{87}\text{Sr}/^{86}\text{Sr}$ it is often difficult to resolve the $^{87}\text{Sr}/^{86}\text{Sr}$ ranges of two different groups of people and establish what constitutes the local range at each site. Here a multi-period study is presented from the Outer Hebrides, Scotland and an investigation of Neolithic and Early Bronze Age populations from the Yorkshire Wolds, NE England. The aim is to demonstrate that, despite complex human dietary strategies, simple mixing systems with only two end-members do occur in archaeological human populations in certain geological provinces and, despite overlapping $^{87}\text{Sr}/^{86}\text{Sr}$ ranges, it is possible to separate two populations based on the structure within the data set.

1. Introduction

Strontium isotope analysis of tooth enamel is a useful provenancing technique to investigate the childhood origins and residential mobility of ancient people (e.g. Bentley and Knipper, 2005; Evans and Tatham, 2004; Ezzo and Price, 2002; Montgomery et al., 2000, 2003; Price et al., 2002, 2004). Enamel is a highly mineralised, acellular, biogenic apatite that is particularly resistant to post-mortem contamination and thus preserves the integrity of lifetime signatures (Budd et al., 2000; Hoppe et al., 2003; Montgomery, 2002; Trickett et al., 2003). The $^{87}\text{Sr}/^{86}\text{Sr}$ value of enamel is derived from food and water ingested when the tissue was mineralising, that is, in early childhood. Food and water have $^{87}\text{Sr}/^{86}\text{Sr}$ values that reflect their geographical origin because Sr weathers from the host rock into groundwater, river water and overlying soils and is ultimately transferred unfractionated into plants and animals (Åberg, 1995; Capo et al., 1998; Ericson, 1985;

Price et al., 2002). This provides the mechanism to establish whether a person or animal has an isotope ratio consistent with locally grown food or locally sourced water, and if this is not the case, to conclude that they either imported substantial resources from elsewhere (perhaps more likely in modern populations) or that they had moved from a different geological province at some time after the tooth had mineralised.

An ideal outcome is that the Sr isotope ratios of the population under study divide neatly into two (or more) visibly separate groups on an isotope plot. However, this rarely happens and investigators are frequently faced with a smear of data points with no discernible break and, perhaps, one or two obvious outliers (see for example Montgomery et al., 2005; Schweissing and Grupe, 2000). In such a situation it is often very difficult to ascertain where one should draw a line between local people and those from elsewhere or if only the outliers can confidently be ascribed non-local status. It is clearly more difficult to identify two populations if some degree of overlap, arising perhaps through similar biosphere values being available in their respective home regions, is present in the range of $^{87}\text{Sr}/^{86}\text{Sr}$ values (Price et al., 2002). Given this potential for overlapping $^{87}\text{Sr}/^{86}\text{Sr}$ ranges in populations originating in different geographical regions, it is perhaps not surprising that it has often proved difficult in previous UK studies to resolve the respective ranges of two supposedly different populations (Evans and Tatham, 2004; Montgomery et al., 2005). Several authors have advocated identifying local populations by the application of a statistical or numerical cut-off based on two or more standard deviations from the mean of bone or animal values (Bentley et al., 2004; Grupe et al., 1997; Horn and Müller-Sohnius, 1999). Whilst this may work well in the specific time period, region and site they are investigating, such a parametric method presupposes that the local population will produce a Gaussian distribution of $^{87}\text{Sr}/^{86}\text{Sr}$ values which, as in the case of Pb isotope fields, is more likely to be the exception rather than the rule, rendering many statistical methods inappropriate (Baxter, 1999; Scaife et al., 1996). Moreover, the number and type of end-members and degree of variation a population exhibits may change considerably as one moves from different geological, geographical and even cultural milieu. The absolute value and the magnitude of the required cut-off will be very specific to the local area and its geology, rendering each site different.

The current model for Sr isotope uptake from the geosphere in a maritime environment such as the British Isles, is that a community living on and sourcing their food from a single, homogeneous rock unit will nevertheless exhibit a range of $^{87}\text{Sr}/^{86}\text{Sr}$ values that fall between two end-members: the local rock and rainwater (Evans and Tatham, 2004; Montgomery, 2002; Montgomery et al., 2005). This may sound unrealistic and oversimplified, but it does not require that types of foods are restricted, only that they are all grown and grazed on just one type of bedrock plus rainwater. Rainwater has seawater of $^{87}\text{Sr}/^{86}\text{Sr} \approx 0.7092$ as its source and it can be ingested in variable quantities through food and drink. The $^{87}\text{Sr}/^{86}\text{Sr}$ value of rainwater in coastal regions is very similar to seawater but over landmasses, the incorporation of terrestrial dust aerosols will progressively alter the original marine signature (Capo et al., 1998; Faure, 1986). The resulting change in $^{87}\text{Sr}/^{86}\text{Sr}$ will depend on distance from the coast and the $^{87}\text{Sr}/^{86}\text{Sr}$ value and Sr concentration in the dust sources and may vary seasonally depending on weather patterns (Åberg, 1995; Andersson et al., 1990; Bacon et al., 1995; Land et al., 2000; Miller et al., 1993). For example, Negrel et al. (2001) found that rainwater varied between 0.70901 and 0.71060 in the Massif Central region of SE France but had a mean value of 0.70943 weighted by % of rainfall.

Consequently, a sedentary, self-sufficient population will not define a normally distributed range of $^{87}\text{Sr}/^{86}\text{Sr}$ values around the local rock value but a range offset above or below the rock value (Montgomery et al., 2005). The magnitude of the range will depend on how different the local rock Sr is from 0.7092: i.e. a population from a region of 0.715 rocks will, through personal choice, cultural practice and resource availability, have the opportunity to obtain an enamel $^{87}\text{Sr}/^{86}\text{Sr}$ value between 0.7092 and 0.715, that is, a weighted average of all inputs. Alternatively, a population subsisting on Cretaceous Chalk ($^{87}\text{Sr}/^{86}\text{Sr} \sim 0.707$) will only have the opportunity to fall within the range 0.707 – 0.7092 and will, therefore, define a much smaller spread of values.

The model becomes increasingly problematical in regions where several different rock types crop out locally or heterogeneous silicate rocks such as granites and sandstones occur. In silicates, Sr released through weathering may not be representative of the whole-rock value but will be dominated by unradiogenic Sr derived from the more easily weathered minerals such as calcite and feldspar (Åberg,

1995; Bau et al., 2004; Blum et al., 1993; Evans and Tatham, 2004; Jacobson and Blum, 2000). Drinking water sources (such as rivers, deep aquifers and springs) that are not derived directly from rain may also have $^{87}\text{Sr}/^{86}\text{Sr}$ values quite different from seawater if Sr has been dissolved from the host rocks (Montgomery et al., 2006). In addition, through choice or necessity, populations may vary considerably in their food procurement strategies, level of sedentism and cultural practices and this suggests one is unlikely to find populations who have a predictable Sr ratio derived from only two dominant sources.

However, two of the authors' recent case studies suggest that simple mixing systems with only two end-members do occur in archaeological populations and, despite overlapping Sr isotope ranges, it is possible to separate two populations based on the structure within the data set. The results are presented of two case studies from very different geological provinces, which one would hypothesise, produce very different Sr isotope ratios in their respective indigenous populations. The first is a study of Neolithic and Early Bronze Age populations from the Yorkshire Wolds, NE England and the second from the Outer Hebrides (or Western Isles) of Scotland from the Neolithic to the Viking colonisation of the islands in the 8th – 9th centuries AD (Montgomery, 2002; Montgomery et al., 2003) (Fig. 1). What the structural differences found in the data sets at both locations might tell us about the people, their geographical origins and food procurement strategies at each location will be discussed. Moreover, given the evidence in the literature that skeletal Sr concentrations do not reflect in any linear way the amount of Sr ingested (Burton and Wright, 1995; Elias, 1980), why and how some archaeological populations may define a clear two-way mixing line and what underlying factors may be ultimately controlling an individual's Sr uptake are discussed.

2. Methods and materials

Enamel samples were of core enamel only. Once childhood tooth mineralization is complete, core enamel is resistant to subsequent isotopic or elemental changes either during an individual's lifetime or during burial, whereas dentine equilibrates with the burial environment (Bocherens et al., 1994; Budd et al., 2000; Glimcher et al., 1990; Hoppe et al., 2003; Montgomery, 2002). To remove soil-derived particulate on the tooth surface, all enamel surfaces were abraded to a depth $>100\ \mu\text{m}$ with acid-cleaned, tungsten carbide dental burrs. All adhering dentine and enamel-dentine

junction tissue were also removed entirely. Dentine samples were removed from the tooth crown and all surfaces abraded with tungsten carbide dental burrs.

Six samples of soil and rock were obtained from: the Iron Age burial context at Galson; the Norse and Bronze Age burial contexts at Cnip; strata identified as sealed Iron Age and Norse cultivation soils, at Galson and Cnip respectively; and a sample of gneiss from the Cnip headland. The gneiss was washed in water (Millipore Alpha Q) and ground to a coarse powder in a tungsten carbide ball mill. Modern plant samples were obtained from machair (Quaternary shell-sands), blackland (peaty soils found between the coastal machair and the interior peat) and hill-land areas on South Uist. Plants were washed in water (Millipore Alpha Q, <1 ppb total heavy metal content), dried and powdered in a freezer-mill.

All enamel, dentine, plant and soil samples were transferred in sealed containers to the class 100, HEPA-filtered laboratory at the NERC Isotope Geosciences Laboratory (NIGL), Keyworth, UK. Enamel chips were washed ultrasonically in water (Millipore Alpha Q, <1 ppb total heavy metal content) to remove adhering particulates. No chemical decontamination was carried out as, due to the high resistance of enamel to post-mortem contamination, changes in the bulk $^{87}\text{Sr}/^{86}\text{Sr}$ ratio of enamel samples following the use of such procedures has been negligible (Horn et al., 1994; Trickett et al., 2003). Approximately 2 g of each soil and rock sample was leached overnight with water (Millipore Alpha Q) and a further 2 g leached overnight in dilute acid (10% vol. acetic). Samples were centrifuged and the leachate pipetted out and dried down. Circa 50 mg of plant powder was dissolved in a combination of Teflon-distilled 16M HNO_3 and Romil Super Purity H_2O_2 . The laboratory procedure used ion exchange chromatography and Teflon-distilled reagents to isolate the Sr prior to instrumental analysis. The full method of preparation and analysis is reported in Montgomery (2002). Strontium concentrations and compositions were obtained by thermal-ionisation mass spectrometry (TIMS) using a Finnigan Mat 262 multi-collector mass spectrometer. $^{87}\text{Sr}/^{86}\text{Sr}$ was normalized to a NBS 987 value of 0.710250. The Sr contribution from within-run laboratory blanks was ≤ 150 pg. External reproducibility was estimated at $\pm 0.004\%$ (2σ).

3. Results

3.1. The Yorkshire Wolds

The precise nature of the Neolithic-Bronze Age transition and when sedentary agriculturalism commenced in Britain remain major questions in British archaeology and the archaeological aim of this study was to investigate diet and mobility within and between Neolithic and Bronze Age populations in prehistoric East Yorkshire (Cooper, 2004).

Figure 2 displays enamel data obtained from 22 individuals (23 teeth) excavated in the 19th century by J.R. Mortimer (1905) from 2 Late Neolithic and 6 Early Bronze Age funerary barrows located on the Cretaceous Chalk of the Yorkshire Wolds (Fig. 1). The area has virtually no drift deposits and one, intermittent, water source: the Gypsy Race. The Chalk was deposited during Late Cretaceous times and records $^{87}\text{Sr}/^{86}\text{Sr}$ ratios between 0.7075 and 0.7078 (McArthur et al., 2001). Direct analyses of Chalk, chalk-derived soils and water from chalk aquifers in England have provided $^{87}\text{Sr}/^{86}\text{Sr}$ ratios from 0.7075 to 0.7077 (Evans et al., 2006; Montgomery, 2002; Montgomery et al., 2005, 2006). According to the simple two-component model, a community subsisting entirely on food sourced from a chalk substrate plus an input from rainwater should define a spread of ratios from ~ 0.7075 to ~ 0.7092 .

Although the two groups of data show almost total overlap in both Sr concentration and $^{87}\text{Sr}/^{86}\text{Sr}$ ratios, making it difficult to separate them based only on individual values, the structure within the two data sets differs markedly. The Bronze Age individuals split into two groups, both of which form discrete linear arrays (labelled A and B) that appear to converge on a lower end-member ~ 0.7075 . Such arrays occur when the samples contain variable mixtures of just two sources of Sr which have different ratios and concentrations (Faure, 1986). In contrast, the Neolithic individuals form a diffuse cloud of data points with $^{87}\text{Sr}/^{86}\text{Sr}$ values ranging between 0.7079 and 0.7102. The lack of a direct linear relationship between the samples implies the presence of more than two end-members.

Tooth enamel contains Sr that was ingested over a specific and restricted period of time during childhood when the tooth analysed was mineralising. In human terms, therefore, the Bronze Age data is consistent with both groups procuring resources from only two sources of Sr and a different mix of these two Sr inputs has resulted in enamel samples containing various mixtures of the two sources. Such variability could arise through necessity with seasonal availability of resources, personal food

preferences or, given the imprecision inherent in archaeological dating, gradual change over longer timescales. Amongst the Neolithic population, however, no such control over resources seems to be present and the distribution of the data is indicative of resources having been procured from more than two Sr sources. This suggests that the earlier Neolithic population were more eclectic and opportunistic in their exploitation of resources and foraged over a wider geographical area beyond the Wolds.

For the Bronze Age population Line A has a radiogenic upper end-member ≥ 0.7120 . To date, there is little evidence that such values can be obtained from the biosphere overlying the Permo-Triassic and Jurassic sedimentary deposits immediately to the north and west of the Chalk but some evidence that such values may be obtained from the Carboniferous Coal Measures which lie further to the west (Table 2; Evans and Tatham, 2004; Montgomery, 2002; Montgomery et al., 2005, 2006). For Line A and B, Chalk could provide the common lower end-member of $^{87}\text{Sr}/^{86}\text{Sr} \sim 0.7075$. However, Line A could equally have a lower end-member of ≤ 0.7092 , which is approximately the value of rain and seawater. Seawater Sr could provide the upper end-member for Line B through exploitation of marine or coastal resources. Alternatively, terrestrial $^{87}\text{Sr}/^{86}\text{Sr}$ ratios > 0.7086 have been obtained from the plants, soils and mineral waters in regions of Permo-Triassic and Jurassic sedimentary rocks to the north and west (Table 2; Evans and Tatham, 2004; Montgomery, 2002; Montgomery et al., 2005, 2006).

It is suggested, therefore, that both Bronze Age groups conform to the simple two end-member model for sedentary agricultural communities, farming on one type of geology with a secondary input of drinking water: one group inhabiting the Wolds and the second group the higher land of the Pennine foothills some distance to the west. The end-member for the lower mixing line could simply be drinking water (suggesting a sedentary lifestyle on the Wolds), or seawater (suggesting utilisation of coastal resources), or as suggested for the Neolithic population, regions immediately adjacent to the Wolds. It must, however, be remembered that quite different sources of Sr (e.g. sedimentary rock and modern seawater) may have the same $^{87}\text{Sr}/^{86}\text{Sr}$ ratio and that whilst geochemically these may be identical, archaeologically the difference may be important. The results may, for example, indicate that Bronze Age communities were following a rigidly controlled system of food production where

crops were grown in one place and livestock grazed in another or one of seasonal transhumance. Exploitation of coastal and marine resources may explain the 0.7092 end-member in both Bronze Age groups. Equally, it could be that both groups were regularly visiting the Wolds for cultural reasons, the area is rich in ceremonial sites such as henges, and thus coming together there at certain times of year. In this regard, it should be noted that for one Bronze Age male adult, two teeth were sampled. The canine tooth (0-5 years of age) plots on Line A whilst the later forming third molar tooth (7-12 years of age) plots on Line B suggesting a move in later childhood from one resource-group to the other. It is believed unlikely that the mixing lines are evidence for a single migration event to the Wolds from elsewhere as all of the individuals analysed would have had to make this move as children whilst the specific teeth analysed were mineralising in order to incorporate different mixtures of the two sources.

3.2. *The Outer Hebrides*

The Outer Hebrides are the most northwesterly islands of the British Isles (Fig. 1). The island chain is geologically homogeneous (Lewisian gneisses and granites) and has a clearly defined geographic boundary (the sea) separating it from places of different geology (and hence isotope ratios) and, it may be ventured, different subsistence strategies. Whether the sea represented a barrier of any real significance in the periods in question and whether any social or dietary difference existed between Outer Hebrideans and people inhabiting the Inner Hebrides (e.g. the Isle of Skye) or the Scottish mainland is debatable but it clearly presents more of a physical obstacle than, for example, trying to define the area of land utilised by a community and their animals living in central England (Evans and Tatham, 2004). Quaternary shell sands (machair) overlie much of the Lewisian gneiss on the west coast of the Outer Hebrides. The machair plain extends for up to 2 km inland and throughout much of the prehistory was the focus for settlement and agriculture. The island interiors are relatively tree-less and predominantly composed of bare rock, standing water and uncultivable blanket peat that supports very little terrestrial wildlife (Armit, 1996). To date, all biosphere samples analysed (e.g. burial and ancient cultivation soils, modern plants, archaeological herbivores) have produced $^{87}\text{Sr}/^{86}\text{Sr}$ ratios between 0.7092 and 0.7104 although leaches of the gneiss rock provided ~ 0.715 (Table 2; Montgomery et al., 2003). Hebridean redshanks have produced $^{87}\text{Sr}/^{86}\text{Sr}$

values from 0.7092 to 0.7100 (Bullman, 2003). This has led to the hypothesis that the biosphere Sr of this maritime island chain is dominated by $^{87}\text{Sr}/^{86}\text{Sr} = 0.7092$ of marine (*e.g.* rainwater, fish, marine mammals, sea-splash, seaweed as fertilizer, food and fodder) and machair (*e.g.* grazing and arable crops) origin with only a limited contribution from the gneiss (Montgomery et al., 2003).

Figure 3 displays archaeological enamel data from 20 Hebridean humans and 4 herbivores. The herbivores form a cluster on the left of the plot and the humans separate into two groups; one individual (labelled E) fits into neither group but falls within the field of data obtained from Neolithic Chalk burials transferred from Figure 2. For ease of reference the two groups have been termed machair dwellers and silicate dwellers. Machair dwellers and herbivores are defined by high Sr concentrations and Sr ratios within the range of biosphere values obtained from the Hebrides (*i.e.* 0.7092 – 0.7104). In contrast, silicate dwellers define a linear array, which appears to arise from mixing between two highly disparate silicate end-members with no input from machair, marine or marine carbonate sources. To date, no biosphere sources have been found on the Outer Hebrides that could provide either the high (>0.7130) or low (<0.7078) end-members. Such an array reflects the compositional range and isotope features seen in silicate rocks of the Inner Hebrides and west coast of Scotland (Fowler et al., 2004) leading to the hypothesis that the members of the silicate group are immigrants to the Outer Hebrides (Montgomery and Evans, 2006). Clearly, this conclusion would have been difficult to make for individuals A and K, which lie just outside the defined machair field, if only $^{87}\text{Sr}/^{86}\text{Sr}$ ratios were used. Two Bronze Age burials from the Inner Hebridean islands of Mull and Arran also fall on the silicate mixing line (Fig. 3) and are thus consistent with such a hypothesis.

The machair dwellers define a slightly sloping field suggesting that their Sr is predominantly of marine or machair origin (0.7092) with only a very small contribution from the gneiss (~ 0.715). Most archaeological human enamel analysed to date from England has Sr concentrations in the range 30-100 ppm (Evans et al., 2006; Evans and Tatham, 2004; Montgomery, 2002; Montgomery et al., 2003, 2005), including those excavated from inland chalk and limestone sites, which suggests the high concentrations obtained from the machair group (150-400ppm) are not simply a result of subsisting on food grown on soils overlying marine carbonate rocks. Such high concentrations coupled with marine Sr ratios as seen in this island population

may, instead, be characteristic of people who cling to the coast and combine exploitation of marine resources such as seaweed with small-scale farming of marginal lands (Kvamme et al. 2004). The herbivores, also, are dominated by marine Sr, although they have much higher enamel concentrations than the humans. This is not unexpected in vegetarians (Bocherens et al., 1994; Tuross et al., 1989). The Hebridean herbivores do not follow the same trajectory as the humans but they do not consume the same diet. Clearly the Sr concentration in the herbivore enamel is not directly relevant to human diets, because people do not eat teeth, and meat and milk contain much less Sr than skeletal tissues (Burton et al., 2003; Burton and Wright, 1995; Tsalev, 1984). Moreover, not all dietary components will contribute equally to body fluids due to complex synergisms and antagonisms with other dietary components such as Ca, fibre and protein, which ensure that there is no simple direct transfer of Sr from the diet to the skeleton (Elias, 1980; Elias et al., 1982; Burton and Wright, 1995; Burton et al., 2003). It is, therefore, possible that the herbivores are the immediate dietary upper end-member for the human population and a proxy for the gneiss, despite falling on the left-hand side of the plot in Figure 3. The data from modern plants (Table 2) shows that blackland and hill-land grazing can provide $^{87}\text{Sr}/^{86}\text{Sr}$ values at the upper end of the Outer Hebridean biosphere range (i.e. 0.7092 – 0.7104).

3.3. Diagenesis

An alternative explanation for the presence of mixing lines in these studies would be that they are produced by postmortem incorporation of Sr from the burial environment. The unexpectedly high enamel Sr concentrations found in the Hebridean machair dwellers are a particular cause for concern although they do not exceed normal modern concentrations which are known to be geographically variable but are typically 50-300ppm (Aufderheide, 1989; Brudevold and Söremark, 1967; Budd et al., 1998; Chaudhri, 1995; Elias, 1980; Elliott and Grime, 1993; Hancock et al., 1989; Lee et al., 1999; Montgomery, 2002; Montgomery et al., 1999; Underwood, 1977). Is it possible, for example that the two groups are simply contaminated and uncontaminated samples rather than immigrant and indigenous humans? However, such a conclusion would require that human and herbivore enamel were being acted upon differently in the burial environment despite all being buried in machair soil, and require the presence of biosphere sources to provide the necessary end-members for

the silicate mixing line. In the Yorkshire Wolds study, all individuals were excavated from barrows on the Chalk and it would be difficult to make a case that diagenesis was only affecting the Bronze Age burials in this manner, especially since they had spent less time in the ground than the Neolithic burials.

For some of the teeth analysed, dentine from the crown of the tooth was also sampled. There is considerable overlap in the mineralisation periods of enamel and primary crown dentine and neither tissue subsequently remodels or re-grows (Boyd, 1989; Fincham and Simmer, 1997; Sasaki et al., 1997; Veis, 1989; Wieser et al., 1996). Consequently, in modern individuals the Sr concentration and isotope ratio of these two tissues can be expected to remain very similar and this does appear to be the case even in individuals who have subsequently been exposed to different sources of Sr (Montgomery, 2002; Tsalev, 1984). Similarly, archaeological individuals that have not been buried, such as the Neolithic skeleton HAUG-1 (Table 1) which was removed from a chambered tomb, in which very limited exposure to soils and fluids is presumed, also show this pattern. However, in teeth excavated from soil burials it is usual for dentine to contain more Sr than enamel and this increase is normally coupled with $^{87}\text{Sr}/^{86}\text{Sr}$ ratios intermediate between the enamel value and that of the burial soil (Budd et al., 2000; Montgomery, 2002; Trickett et al., 2003). Figure 4 shows paired enamel and dentine values for 3 teeth excavated from 3 different barrows. Diagenetic vectors are shown representing two-component mixing between the biogenic composition of the enamel (used as a proxy for biogenic crown dentine) and the mobile Sr in the burial environment, which has produced a postmortem shift in the crown dentine composition. The vectors appear to converge on $^{87}\text{Sr}/^{86}\text{Sr} \approx 0.7082$, suggesting that the diagenetic Sr in the burial environment does not have the composition of pure Chalk. This may reflect the recorded use of “foreign” clay in the construction of the barrows (Mortimer, 1905). It appears that the contaminated dentine samples are not on the same mixing trajectory as the enamel samples whose biogenic isotopic integrity is intact. This discrepancy is also apparent when dentine samples from the Hebrides are plotted with the enamel data. Figure 5 shows that dentine samples are moving away from the enamel compositions and the diagenetic vectors are converging on $^{87}\text{Sr}/^{86}\text{Sr} \approx 0.7100$ rather than the biosphere lower end-member indicated by the enamel samples of 0.7092. The distribution of the dentine samples is also quite different to that of the enamel samples as there are no enamel

data points within the diagenetic triangle defined by the dentine samples, where some would perhaps be expected to be seen if enamel samples were being contaminated with Sr from the same source. Enamel samples equilibrating with burial Sr would not move down the silicate mixing line and then across but would move directly towards the value for diagenetic Sr. An exception to this is the sample from a neonatal infant (labelled G), from which only a partially formed deciduous tooth crown could be sampled. This tooth crown contained no fully mineralised enamel and was dark brown and relatively soft and it is possible that it did not have the same resistance to diagenesis as the fully mineralised, hard, translucent, white tissue analysed from the remaining samples in this study.

4. Discussion

A linear mixing relationship between human enamel samples requires not only close control over the isotope ratio but also over the concentration and, whilst it may be perfectly reasonable to assume that random mixing of two food types may produce an assortment of diets on such a linear array with two identifiable end-members, because there is no opportunity for Sr to be added or taken away, obtaining such a relationship from biological skeletal tissues is considerably more complex. Although Sr concentrations in skeletal tissue may be dose-dependent (Comar et al., 1956; Boivin et al., 1996), experimental studies have concluded that there is no straightforward, linear, relationship between the amount of Sr in any given diet and the amount of Sr in the resulting tissue (Elias, 1980; Burton and Wright, 1995). Biological organisms are not rocks; physiological processes can act upon the amount of Sr that is finally deposited in tooth enamel at all the stages in its journey from the stomach to its final deposition in enamel. Moreover, Sr is not evenly distributed across all tissues; in mammals, most resides within the skeleton and given bone formation and turnover rates is unlikely to be in constant equilibrium with the diet. Enamel concentrations may be increased or reduced relative to the concentration in any one food type and not all dietary components will contribute equally; indeed some may not contribute at all. Why, then, should one expect to find such a relationship between a human population and its diet?

What is remarkable, given the discrepancy one might consequently expect between the concentration in the diet of an individual and the resulting concentration in their enamel, is the extremely good fit of the human samples on the mixing lines

presented here. From a geochemical perspective, this is highly unlikely to arise from co-incidence. As Faure (1986) asserts “The goodness of fit of the data points to a straight line is a test for the validity of the mixing hypothesis and of the assumption that neither the Sr concentrations nor the $^{87}\text{Sr}/^{86}\text{Sr}$ ratios were modified after mixing had occurred”. Faure (1986) is not writing about biological organisms, but the individuals in the case studies do indeed appear to be strongly related in such a simple binary relationship that can be explained through access to food and water that derives from two, and only two, sources of Sr. However, in order for these individuals to lie on the mixing line, not only their isotope ratio but also the concentration of Sr in their enamel must be very closely controlled by the two geological sources providing the dietary Sr and not modified subsequently. Such a finding clearly has implications for studies that use Sr/Ca ratios as an indicator of trophic level and food sources because, as already suggested by Burton et al. (2003), it raises the possibility that in certain locations geological, rather than biological, processes will control the amount of Sr in skeletal tissue.

For the Yorkshire Wolds study, archaeological dating and funerary evidence accords well with the geochemical evidence; they independently separate the individuals into the same two period-based groups suggesting the difference in food procurement strategy is a change through time (Cooper, 2004). Despite the strong linear relationship between the members of the Hebridean silicate group, radiocarbon dates from these individuals span nearly the whole of the first millennium AD and there is little archaeological evidence to link them together. As previously mentioned, one possible region that might provide such mixing and migration opportunities is the Tertiary Volcanic Province (Fig. 1) where young basaltic rocks have extruded through the ancient Lewisian crust, as on the nearby Isle of Skye and Scottish mainland (Fowler et al., 2004). Unfortunately, skeletal remains of comparable date available for analysis from such areas are either non-existent or extremely rare. Such a linear relationship may simply be a feature of this unique geological setting, as linear arrays are certainly not routinely observed in archaeological studies (e.g. Schweissing and Grupe 2003). Moreover, such a large spread in isotope ratios within a single population is unusual and makes it very difficult to apply a statistical cut-off value to identify immigrants in such geological provinces. However, the majority of published studies have been unavoidably biased towards areas where bone survives, and these are far more likely to be regions of alkaline rock, such as chalk and limestone, than

silicate rocks, which frequently produce acidic soils that are not conducive to good bone preservation. Alternatively, the factor linking these individuals may be that they originate from quite different geographical locations, but that all locations are in regions of silicate rock where there is no input from either marine carbonates or coastal marine Sr, such as the Scottish mainland. This study is still underway but irrespective of where these individuals originated from the data appear to provide evidence for an enduring migration stream to the Outer Hebrides.

5. Conclusions

$^{87}\text{Sr}/^{86}\text{Sr}$ mixing models provide a useful method of separating archaeological populations that have overlapping isotope ranges based on the structure within the data set. The method is based on the identification of dietary end-members using simple graphical means rather than using inferential statistical procedures to define populations by means of their $^{87}\text{Sr}/^{86}\text{Sr}$ ranges and standard deviations (Baxter et al., 2000). Whilst such statistical methods may work well for the site where they are developed, it is clear from the data presented here that they may not be transferable to other studies in different geological settings because the underlying assumptions about dietary Sr might be incorrect and the data may not be normally distributed. Whilst it is acknowledged that the graphical method may not work in regions of complex, heterogeneous geology or for populations that have multiple sources of dietary Sr, the studies presented here show that in some regions the bedrock geology appears to exert a close control over both the Sr concentrations and the $^{87}\text{Sr}/^{86}\text{Sr}$ ratio in human tooth enamel.

Acknowledgements

We are grateful to the following people for providing site information, advice, teeth, plants and soils: Martin Foreman, Graham Myers, Craig Barclay and Bryan Sitch at the Hull and East Riding Museum; Alison Sheridan, National Museums of Scotland; Tim Neighbour, CFA Archaeology; Mike Parker Pearson and Andrew Chamberlain, University of Sheffield; Richard Langhorne and Mary McLeod at the Museum nan Eilean, Stornoway; Mike Church, Durham University; Oliver Craig, University of York; and Helen Smith, Bournemouth University. JM acknowledges the support of the NERC through the award of a Fellowship NER/1/S/2002/00691 and NIGFSC Isotope Grant IP/804/1103. NIGL Publication No. 826.

References

- Åberg, G., 1995. The use of natural strontium isotopes as tracers in environmental studies. *Water Air Soil Poll.* 79, 309-322.
- Andersson, P., Löfvendahl, R., Åberg, G., 1990. Major element chemistry, $\delta^2\text{H}$, $\delta^{18}\text{O}$ and $^{87}\text{Sr}/^{86}\text{Sr}$ in a snow profile across central Scandinavia. *Atmos. Environ.* 24A, 2601-2608.
- Armit, I., 1996. *The Archaeology of Skye and the Western Isles*. Edinburgh University Press, Edinburgh.
- Aufderheide, A.C., 1989. Chemical analysis of skeletal remains. In: Iscan, M.Y., Kennedy, K.A.R. (Eds.), *Reconstruction of life from the skeleton*. Alan R. Liss, Inc., New York, 237-260.
- Bacon, J.R., Berrow, M.L., Shand, C.A., 1995. The use of isotopic composition in field studies of lead in upland Scottish soils (UK). *Chem. Geol.* 124, 125-134.
- Bau, M., Alexander, B., Chesley, J.T., Dulski, P., Brantley, S.L., 2004. Mineral dissolution in the Cape Cod aquifer, Massachusetts, USA: I. Reaction stoichiometry and impact of accessory feldspar and glauconite on strontium isotopes, solute concentrations, and REY distribution. *Geochim. Cosmochim. Acta* 68, 1199-1216.
- Baxter, M.J., 1999. On the multivariate normality of data arising from lead isotope fields. *J. Archaeol. Sci.* 26, 117-124.
- Baxter, M.J., Beardah, C.C., Westwood, S., 2000. Sample size and related issues in the analysis of lead isotope data. *J. Archaeol. Sci.* 27, 973-980.
- Bentley, R.A., Knipper, C., 2005. Geographical patterns in biologically available strontium, carbon and oxygen isotope signatures in Prehistoric SW Germany. *Archaeometry* 47, 629-644.
- Bentley, R.A., Price, T.D., Stephan, E., 2004. Determining the 'local' Sr-87/Sr-86 range for archaeological skeletons: a case study from Neolithic Europe. *J. Archaeol. Sci.* 31, 365-375.
- Blum, J.D., Erel, Y., Brown, K., 1993. Sr-87/Sr-86 Ratios of Sierra-Nevada Stream Waters - Implications for Relative Mineral Weathering Rates. *Geochim. Cosmochim. Acta* 57, 5019-5025.

- Bocherens, H., Brinkman, D.B., Dauphin, Y., Mariotti, A., 1994. Microstructural and geochemical investigations on Late Cretaceous archosaur teeth from Alberta, Canada. *Can. J. Earth Sci.* 31, 783-792.
- Boivin, G., Deloffre, P., Perrat, B., Panczer, G., Boudeulle, M., Mauras, Y., Allain, P., Tsouderos, Y., Meunier, P.J., 1996. Strontium distribution and interactions with bone mineral in monkey iliac bone after strontium salt (S 12911) administration." *J. Bone Mineral. Res.* 11, 1302-1311.
- Boyde, A., 1989. Enamel. In: Berkovitz, B.K.B. (Ed.), *Handbook of Microscopic Anatomy: Teeth*. Springer-Verlag, Berlin, 309-473.
- Brudevold, F., Söremark, R., 1967. Chemistry of the mineral phase of enamel. In: Miles, A.E.W. (Ed.), *Structural and Chemical Organization of Teeth*. Academic Press, London, 247-277.
- Budd, P., Montgomery, J., Barreiro, B., Thomas, R.G., 2000. Differential diagenesis of strontium in archaeological human dental tissues. *Appl. Geochem.* 15, 687-694.
- Budd, P., Montgomery, J., Cox, A., Krause, P., Barreiro, B., Thomas, R.G., 1998. The distribution of lead within ancient and modern human teeth: Implications for long-term and historical exposure monitoring. *Sci. Total Environ.* 220, 121-136.
- Bullman, 2003. The spatial and temporal distribution of shorebirds overwintering on the middle Forth Estuary, Scotland. Ph.D. Thesis, Univ. Stirling, UK.
- Burton, J., Price, T.D., Cahue, L., Wright, L.E., 2003. The use of barium and strontium abundances in human skeletal tissues to determine their geographical origins. *Internat. J. Osteoarchaeol.* 13, 88-95.
- Burton, J.H., Wright, L.E., 1995. Nonlinearity in the relationship between bone Sr/Ca and diet: paleodietary implications. *Am. J. Phys. Anthropol.* 96, 273-282.
- Capo, R.C., Stewart, B.W., Chadwick, O.A., 1998. Strontium isotopes as tracers of ecosystem processes: theory and methods. *Geoderma* 82, 197-225.
- Chaudhri, A., 1995. *Nuclear Analytical Methods in Calcified Tissue Research*. Nutrition 11, 538-541.
- Comar, C.L., Wasserman, R.H., Nold, M.M., 1956. Strontium-calcium discrimination factors in the rat. *Proc. Soc. Exp. Biol.* 92, 859-863.
- Cooper, R.E., 2004. An Assessment of Dietary Strontium Isotope Variation in Neolithic and Bronze Age East Yorkshire. M.Sc. Thesis, Univ. Bradford. UK.

- Elias, M., 1980. The feasibility of dental strontium analysis for diet-assessment of human populations. *Am. J. Phys. Anthropol.* 53, 1-4.
- Elias, R.W., Hirao, Y., Patterson, C.C., 1982. The circumvention of the natural biopurification of calcium along nutrients pathways by atmospheric inputs of industrial lead. *Geochim. Cosmochim. Acta* 46, 2561-2580.
- Elliott, T.A., Grime, G.W., 1993. Examining the diagenetic alteration of human bone material from a range of archaeological burial sites using nuclear microscopy. *Nucl. Instr. Meth.* 77, 537-547.
- Ericson, J.E., 1985. Strontium isotope characterization in the study of prehistoric human ecology. *J. Hum. Evol.* 14, 503-514.
- Evans, J.A., Tatham, S., 2004. Defining "local signature" in terms of Sr isotope composition using a tenth-twelfth century Anglo-Saxon population living on a Jurassic clay-carbonate terrain, Rutland, UK. In: Pye, K., Croft, D.J. (Eds.), *Forensic Geoscience: Principles, Techniques and Applications*. Geological Society of London Special Publication, London, 237-248.
- Evans, J.A., Stoodley, N., Chenery, C.A., 2006. A strontium and oxygen isotope assessment of a possible 4th century immigrant population in a Hampshire cemetery, southern England. *J. Archaeol. Sci.* 33, 265-272.
- Ezzo, J.A., Price, T.D., 2002. Migration, regional reorganization, and spatial group composition at Grasshopper Pueblo, Arizona. *J. Archaeol. Sci.* 29, 499-520.
- Faure, G., 1986. *Principles of Isotope Geology*. John Wiley & Sons Inc., New York.
- Fincham, A.G., Simmer, J.P., 1997. Amelogenin proteins of developing dental enamel. In: Chadwick, D.J., Cardew, G. (Eds.), *Dental Enamel*. Proc. Ciba Foundation Symp. 205. John Wiley & Sons Inc, Chichester, 118-134.
- Fowler, S.J., Bohrsen, W.A., Spera, F.J., 2004. Magmatic evolution of the Skye igneous centre, western Scotland: modelling of assimilation, recharge and fractional crystallization. *J. Petrol.* 45, 2481-2505.
- Glimcher, M.J., Cohen-Solal, L., Kossiva, D., de Ricqles, A., 1990. Biochemical analyses of fossil enamel and dentin. *Paleobiol.* 16, 219-232.
- Grupe, G., Price, T.D., Schröter, P., Söllner, F., Johnson, C.M., Beard, B.L., 1997. Mobility of Bell Beaker people revealed by strontium isotope ratios of tooth and bone: a study of southern Bavarian skeletal remains. *Appl. Geochem.* 12, 517-525.

- Hancock, R.G.V., Grynopas, M.D., Pritzker, K.P.H., 1989. The abuse of bone analysis for archaeological dietary studies. *Archaeometry* 31, 169-179.
- Hoppe, K.A., Koch, P.L., Furutani, T.T., 2003. Assessing the preservation of biogenic strontium in fossil bones and tooth enamel. *Internat. J. Osteoarchaeol.* 13, 20-28.
- Horn, P., Müller-Sohnius, D., 1999. Comment on "Mobility of Bell Beaker people revealed by strontium isotope ratios of tooth and bone: a study of southern Bavarian skeletal remains" by Gisela Grupe, T. Douglas Price, Peter Schröter, Frank Söllner, Clark M. Johnson and Brian L. Beard. *Appl. Geochem.* 14, 263-269.
- Horn, P., Hölzl, S., Storzer, D., 1994. Habitat determination on a fossil stag's mandible from the site of *Homo erectus heidelbergensis* at Mauer by use of $^{87}\text{Sr}/^{86}\text{Sr}$. *Naturwissenschaften* 81, 360-362.
- Jacobson, A.D., Blum, J.D., 2000. Ca/Sr and Sr-87/Sr-86 geo-chemistry of disseminated calcite in Himalayan silicate rocks from Nanga Parbat: Influence on river-water chemistry. *Geology* 28, 672-672.
- Kvamme, M., Kaland, P.E., Brekke, N.G., 2004. Conservation and Management of North European Coastal Heathlands. The Heathland Centre, Lygra.
- Land, M., Ingri, J., Andersson, P.S., Öhlander, B., 2000. Ba/Sr, Ca/Sr and $^{87}\text{Sr}/^{86}\text{Sr}$ ratios in soil water and groundwater: implications for relative contributions to stream water discharge. *Appl. Geochem.* 15, 311-325.
- Lee, K.M., Appleton, J., Cooke, M., Keenan, F., Sawicka-Kapusta, K., 1999. Use of laser ablation inductively coupled plasma mass spectrometry to provide element versus time profiles in teeth. *Anal. Chim. Acta* 395, 179-185.
- McArthur, J.M., Howarth, R.J., Bailey, T.R., 2001. Strontium isotope stratigraphy: LOWESS version 3: best fit to the marine Sr-isotope curve for 0-509 Ma and accompanying look-up table for deriving numerical age. *J. Geol.* 109, 155-170.
- Miller, E.K., Blum, J.D., Friedland, A.J., 1993. Determination of soil exchangeable cation loss and weathering rates using Sr isotopes. *Nature* 362, 438-441.
- Montgomery, J., 2002. Lead and Strontium Isotope Compositions of Human Dental Tissues as an Indicator of Ancient Exposure and Population Dynamics. Ph.D. Thesis, Univ. Bradford, UK.
- Montgomery, J., Evans, J.A., 2006. Immigrants on the Isle of Lewis - combining traditional funerary and modern isotope evidence to investigate social differentiation, migration and dietary change in the Outer Hebrides of Scotland.

- In: Gowland, R., Knusel, C. (Eds.), *The Social Archaeology of Funerary Remains*. Oxbow Books, Oxford, 122-142.
- Montgomery, J., Budd, P., Cox, A., Krause, P., Thomas, R.G., 1999. LA-ICP-MS evidence for the distribution of lead and strontium in Romano-British, medieval and modern human teeth: implications for life history and exposure reconstruction. In: Young, S.M.M., Pollard, A.M., Budd, P., Ixer, R.A. (Eds.), *Metals in Antiquity*. BAR International Series 792. Archaeopress, Oxford, 258-261.
- Montgomery, J., Budd, P., Evans, J., 2000. Reconstructing the lifetime movements of ancient people: a Neolithic case study from southern England. *Eur. J. Archaeol.* 3, 407-422.
- Montgomery, J., Evans, J.A., Neighbour, T., 2003. Sr isotope evidence for population movement within the Hebridean Norse community of NW Scotland. *J. Geol. Soc.* 160, 649-653.
- Montgomery, J., Evans, J.A., Powlesland, D., Roberts, C.A., 2005. Continuity or colonization in Anglo-Saxon England? Isotope evidence for mobility, subsistence practice, and status at West Heslerton. *Am. J. Phys. Anthropol.* 126(2), 123-138.
- Montgomery, J., Evans, J.A., Wildman, G., 2006. $^{87}\text{Sr}/^{86}\text{Sr}$ isotope composition of bottled British mineral waters for environmental and forensic purposes. *Appl. Geochem.* 21, 1626-1634.
- Mortimer, J.R., 1905. *Forty Years Researches in British and Saxon Burial Mounds of East Yorkshire*. A. Brown & Sons, London.
- Negrel, P., Casanova, J., Aranyossy, J.F., 2001. Strontium isotope systematics used to decipher the origin of groundwaters sampled from granitoids: the Vienne Case (France). *Chem. Geol.* 177, 287-308.
- Parker Pearson, M., Chamberlain, A., Craig, O., Marshall, P., Mulville, J., Smith, H., Chenery, C., Collins, M., Cook, G., Craig, G., Evans, J., Hiller, J., Montgomery, J., Schwenninger, J.-L., Taylor, G., Wess, T., 2005. Evidence for mummification in Bronze Age Britain. *Antiquity* 79(305), 529-546.
- Price, T.D., Burton, J.H., Bentley, R.A., 2002. The characterization of biologically available strontium isotope ratios for the study of prehistoric migration. *Archaeometry* 44, 117-135.

- Price, T.D., Knipper, C., Grupe, G., Smrcka, V., 2004. Strontium isotopes and prehistoric human migration: the Bell Beaker Period in Central Europe. *Eur. J. Archaeol.* 7, 9-40.
- Sasaki, T., Takagi, M., Yanagisawa, T., 1997. Structure and function of secretory ameloblasts in enamel formation. In: Chadwick, D.J., Cardew, G. (Eds.), *Dental Enamel. Proc. Ciba Foundation Symp.* 205. John Wiley, Chichester, 32-50.
- Scaife, B., Budd, P., McDonnell, J.G., Pollard, A.M., Thomas, R.G., 1996. A reappraisal of statistical techniques used in lead isotope analysis. In: Demirci, S., Ozer, A.M., Summers, G.D. (Eds.), *Archaeometry 94: Proc. 29th Internat. Symp. Archaeometry, Ankara 9-14 May 1994.* Tubitak, Ankara, 301-307.
- Schweissing, M.M., Grupe, G., 2000. Local or nonlocal? A research of strontium isotope ratios of teeth and bones on skeletal remains with artificial deformed skulls. *Anthropol. Anz.* 58, 99-103.
- Schweissing, M. M., Grupe, G., 2003. Stable strontium isotopes in human teeth and bone: a key to migration events of the late Roman period in Bavaria. *J. Archaeol. Sci.* 30, 1373-1383.
- Trickett, M.A., Budd, P., Montgomery, J., Evans, J., 2003. An assessment of solubility profiling as a decontamination procedure for the Sr-87/Sr-86 analysis of archaeological human skeletal tissue. *Appl. Geochem.* 18, 653-658.
- Tsalev, D.L., 1984. *Atomic Absorption Spectrometry in Occupational and Environmental Health Practice, Volume II: Determination of Individual Elements.* CRC Press Inc., Boca Raton, Florida.
- Tuross, N., Behrensmeyer, A.K., Eanes, E.D., 1989. Strontium increases and crystallinity changes in taphonomic and archaeological bone. *J. Archaeol. Sci.* 16, 661-672.
- Underwood, E.J., 1977. *Trace elements in human and animal nutrition.* Academic Press, London.
- Veis, A., 1989. Biochemical studies of vertebrate tooth mineralization. In: Mann, S., Webb, J. and Williams, R.J.P. (Eds.), *Biomineralization: chemical and biochemical perspectives.* VCH Publishers, New York, 189-222.
- Wieser, A., Romanyukha, A.A., M.O., D., Kozheurov, V.P., Petzoldt, G., 1996. Tooth enamel as a natural beta dosimeter for bone seeking radionuclides. *Radiation Protection Dosimetry* 65, 413-416.

Figure captions

Figure 1. Simplified geology map of Great Britain and Ireland showing the location of the Outer Hebrides and the Yorkshire Wolds.

Figure 2. Human enamel data from Neolithic and Bronze Age barrows of the Yorkshire Wolds. The horizontal dotted lines indicate possible end-members: the upper line is seawater (~ 0.7092) and an approximation for rainwater; the lower line on which both mixing lines appear to converge is the value for English Cretaceous Chalk (Evans et al., 2006; McArthur et al., 2001; Montgomery, 2002; Montgomery et al., 2005). For the upper mixing line $r^2 = 0.9789$, for the lower mixing line $r^2 = 0.9364$. The data point labelled ?F is the only skeleton plotted identified as being of possible female sex. 2σ errors are within symbol.

Figure 3. Human and herbivore enamel data from the Outer Hebrides, Scotland. The lower horizontal line denotes the seawater (~ 0.7092) end-member. For the silicate mixing line $r^2 = 0.9851$. The oval approximately replicates the field of Neolithic data from Figure X. All individuals were excavated from Outer Hebridean islands apart from the two Inner Hebrideans (Δ) from Arran and Mull. 2σ errors are within symbol. Data source: Table 1; Montgomery, 2002; Montgomery et al., 2003; Parker Pearson et al., 2005.

Figure 4. Human enamel and dentine data from Neolithic and Bronze Age barrows on the Yorkshire Wolds. Diagenetic vectors are shown between the lower concentration enamel and the higher concentration dentine value of 3 enamel-dentine pairs from 3 different barrows. They appear to converge on $^{87}\text{Sr}/^{86}\text{Sr} \approx 0.7082$ rather than the Chalk value of ≤ 0.7075 . 2σ errors are within symbol.

Figure 5. Human enamel and dentine data from the Outer Hebrides, Scotland. Diagenetic vectors are shown between the lower concentration enamel and the higher concentration dentine value of enamel-dentine pairs and appear to converge on $^{87}\text{Sr}/^{86}\text{Sr} \approx 0.7100$ rather than the marine value of 0.7092 . 2σ errors are within symbol. Data source: Table 2; Montgomery et al., 2003.




Figure 1




Figure 2




Figure 4




Figure 5

Table captions

Table 1. Sr isotope ratio and concentration data for archaeological teeth. Teeth are identified by the following abbreviations: incisor (I); canine (C); premolar (P); molar (M); maxillary (^{1,2,3}); mandibular (_{1,2,3}); left (L); right (R); lower case letters indicate deciduous dentition.

Table 2. Sr isotope ratio and concentration data for modern plants. Sr concentrations should be regarded as estimates only as they are dependent on water content.

ACCEPTED MANUSCRIPT

Table 1

Period	Site	Skeleton No.	Age	Sex	Tooth	Tissue	⁸⁷ Sr/ ⁸⁶ Sr	Sr ppm	1/Sr*1000
<i>Yorkshire Wolds</i>									
Neolithic	Towthorpe 273 (Duggleby Howe)	DH1/2/G	Adult	Male	I ² R	enamel	0.710201	68	14.66
		DH2/73/I	Adult	Male	M ² L	enamel	0.709570	86	11.70
						dentine	0.708938	155	6.45
		DH3/69/3	Adult	Male	P ₁ L	enamel	0.709138	50	19.92
		DH4/74/K	Adult	Male	P ₁ R	enamel	0.708594	44	22.68
		DH5/76/M	Adult	Male	P ₂ R	enamel	0.709997	49	20.37
		DH6/72/H	Juvenile	n/k	m ₂ R	enamel	0.709349	72	13.83
	DH7/75/L	Adult	Male	P ¹ R	enamel	0.709849	41	24.51	
	Calais Wold 275	CW8/7	Adult	Male	P ² L	enamel	0.708896	39	25.38
						dentine	0.708600	60	16.58
		CW9/9	Adult	Male	M ² R	enamel	0.709865	49	20.45
		CW10/8?	Adult	Male	C ¹ L	enamel	0.708437	46	21.79
		CW11/3	Adult	Male	P ₁ R	enamel	0.709325	55	18.25
CW12/?		Adult	Male	P ₁ L	enamel	0.707952	92	10.91	
Bronze Age	Calais Wold 100	CW13/2	Adult	Male	P ₁ R	enamel	0.712010	44	22.68
	Calais Wold 23	CW14/1	Adult	Male	M ³ R	enamel	0.708023	50	19.84
	Towthorpe 73	TP15/1	Adult	Male	C ¹ R	enamel	0.709195	82	12.24
	Towthorpe 72	TP16/1	Adult	?Female	C ¹ L	enamel	0.709227	72	13.83

	Towthorpe 3	TP17/D	Adult	Male	P ² R	enamel	0.710458	56	17.99
	Aldro 116	AL18/1	Adult	Male	M ² R	enamel	0.710981	69	14.53
		AL19/4	Juvenile	n/k	C ¹ R	enamel	0.708315	39	25.97
						dentine	0.708149	261	3.83
		AL20/6	Adult	Male	P ₂ L	enamel	0.709818	66	15.22
		AL21/2	Adult	Male	P ¹ R	enamel	0.708512	35	28.82
		AL22/5	Juvenile 7-12yrs	n/k	M ₁ L	enamel	0.708588	31	32.79
		AL23/3	Adult	Male	P ₂ L	enamel	0.708361	40	24.94
<i>Hebrides</i>									
Neolithic	Haugabost, Harris	HAUG-1	Adult	n/k	M ₂ R	enamel	0.709340	307	3.26
Bronze Age	Cnip, Lewis	BA-1	Adult	Male	C ¹ R	dentine	0.710403	316	3.17
	Tormore, Arran	ET54	Adult	Female	M ¹ R	enamel	0.714447	68	14.68
	Ardachy, Mull	MULL-1	Juvenile 15-20yrs	n/k	M ₁ L	enamel	0.709173	124	8.04
						dentine	0.709815	555	1.80
Iron Age	Kilpheder, S. Uist	SUK-1	Adult	Female	M ₂ L	enamel	0.708876	130	7.69
		SUK-2	Adult	Female	M ₁ L	enamel	0.709206	127	7.88
	Galson, Lewis	Gals-93	Adult	?Male	P ₁ R	dentine	0.710290	269	3.72
		Gals-96	Adult	?Male	P ₁ R	dentine	0.709380	309	3.23
		Gals-II	Adult	Female	P ₁ L	dentine	0.709459	242	4.14
		Gals-IV	Adult	Female	P ¹ L	dentine	0.710940	141	7.12
Norse	Cnip, Lewis	A	Adult	Female	P ¹ L	dentine	0.710014	222	4.50

B	Juvenile ~6yrs	n/k	I ¹ L	dentine	0.709746	522	1.92
C	Adult	Male	P ₁	dentine	0.710074	865	1.16
D	Adult	Male	P ¹ R	dentine	0.708426	195	5.13
F	Infant	n/k	di ¹ L	dentine	0.709961	572	1.75

ACCEPTED MANUSCRIPT

Table 2

Location	Sample No.	Geology (bedrock or drift)	$^{87}\text{Sr}/^{86}\text{Sr}$	Sr ppm	Longitude	Latitude
South Uist	Frobost 309	Dry machair	0.709220	21	-7.4052	57.2034
	Frobost 309	Dry machair	0.709220	18	-7.4052	57.2034
	Milton 310	Machair	0.709219	56	-7.4006	57.2164
	Milton 310	Machair	0.710019	24	-7.4006	57.2164
	Milton 311	Low machair	0.709207	87	-7.4006	57.2164
	Milton 310a	Dry machair	0.709176	141	-7.4006	57.2164
	Milton 314	PreCambrian Gneiss - blackland	0.710289	62	-7.4006	57.2164
	Bornish 321	PreCambrian Gneiss - blackland	0.709449	45	-7.4015	57.2430
	Bornish 322	PreCambrian Gneiss - hill-land	0.710396	31	-7.4015	57.2430
Yorkshire	Hull-1	Cretaceous Chalk	0.708489	38	-0.4482	53.8417
	Hull-2	Permo-Triassic Sandstones	0.708997	24	-1.3128	53.7653
	Hull-3	Palaeozoic Magnesian Limestone	0.708637	55	-1.3453	53.8167
	Hull-4	Triassic Mudstones	0.710181	21	-0.8255	53.8298
	Hull-5	Permian Mudstones	0.709372	25	-1.1504	53.7696
	Hull-6	Jurassic Lower Lias	0.709691	30	-0.6593	53.2743
	Hull-7	Palaeozoic Carboniferous Coal Measures	0.714427	7	-1.8061	53.6961
	Hull-8	Palaeozoic Carboniferous Coal Measures	0.712636	8	-1.5525	53.6145
	Scar-1	Lower Jurassic clays and sands	0.709509	12	-0.6922	54.3386
	Scar-2	Lower Jurassic clays and sands	0.709693	0.4	-0.6802	54.4255
	Scar-3	Lower Jurassic clays and sands	0.708707	30	-0.8328	54.1178
	Scar-5	Lower Jurassic clays and sands	0.709366	87	-0.9063	54.0783