




Institute of
Hydrology

1988/02518



ESS

RUFFORD PARK
RAINFALL STUDY

FURTHER ANALYSIS

E.J. STEWART

JUNE

1988

RUFFORD PARK RAINFALL STUDY

FURTHER ANALYSIS

A Report to
Elliot and Brown,
Consulting Engineers

E J Stewart
Institute of Hydrology

June 1988

1. OBJECTIVE

A previous study has been carried out to assess the rarity of the 5-hour storm which led to flooding near Rufford Park on 1 June 1983. The return period of the storm was estimated to be 5 years. The rainfall data used in the analysis suggested that the effect of the storm had been exacerbated by the wet spring which preceded it. The objective of this further study was to assess the return period of the combination of the storm and the antecedent precipitation conditions.

2. ANALYSIS

2.1 Introduction

The analysis focused on the calculation of the joint probability of the 5-hour storm and the antecedent catchment rainfall conditions. From probability theory, if Q and R are independent events, their joint probability can be expressed as

$$\Pr(Q \cap R) = \Pr(Q) \cdot \Pr(R)$$

Thus, if the probability of the storm was found to be independent of that of the antecedent precipitation, then the joint probability would be equal to their product. The annual probability of the 5-hour storm was already known. The return period of the storm was estimated to be 5 years, and the reciprocal of this (i.e. 0.2) represented the probability of the event being equalled or exceeded in any one year. Therefore, it was required to find the probability of the antecedent rainfall conditions and to determine whether this was independent of the annual exceedence probability.

2.2 Analysis of antecedent precipitation

A preliminary indication of the antecedent rainfall conditions was given by constructing a time series of daily catchment rainfalls for the first five months of 1983. In addition to this, monthly rainfall totals for the only gauge with long-term data situated within the catchment (123017) were extracted from the database. It was apparent that the catchment rainfall totals for April and May 1983 were considerably higher than the mean totals for the long-term gauge (see Table 1).

Table 1. Comparison of catchment monthly rainfall with long-term mean

| | Catchment monthly total 1983 (mm) | Mean at gauge 123017 (mm) |
|-------|-----------------------------------|---------------------------|
| April | 121.3 | 48.1 |
| May | 112.3 | 53.7 |

A series of annual maximum 1-day rainfalls at gauge number 123017 over the period 1910-1983 was constructed. The 1-day rainfall of 31 May 1983 constituted the annual maximum for that year. The antecedent catchment conditions relating to each of the annual maxima were described in terms of an antecedent precipitation index (API). Various definitions of API were considered, that with a daily recession factor (k) of 0.95 ultimately being adopted. APIs for 100 days were calculated for each annual maximum rainfall in the series and are given in Appendix 1. The API value calculated for 31 May 1983 was 44.58 mm and was ranked 51 out of the 67 values. This value is not particularly high, but represents the fact that the API is very sensitive to rainfall totals occurring immediately prior to the annual maximum. In the case of the 31 May 1983, the only appreciable antecedent rainfall occurred thirty days previously.

2.3 Analysis of independence

An analysis of the correlation between the APIs and their corresponding annual maxima was carried out in order to assess the extent of their dependence. The correlation coefficient was calculated to be only 0.184. A significance test was carried out on this value (see Appendix 2) and there was found to be no significant correlation between API and the annual maximum 1-day rainfall at the 5% level. Hence the two were taken to be independent.

2.4 Calculation of the probability of the antecedent rainfall

The API values in Appendix 1 were ranked and plotted on probability paper using the Blom plotting position. From this plot, the probability that the API was less than or equal to 44.58 mm was determined to be 0.74.

2.5 Calculation of joint probability

Since the storm rainfall and antecedent precipitation had been found

to be mutually independent, their joint probability was computed by multiplying their individual probabilities together as set out below:

$$\Pr(x > X) = 0.2 \quad (\text{Storm rainfall})$$

$$\Pr(y > Y) = 1 - 0.74 = 0.26 \quad (\text{Antecedent rainfall})$$

$$\begin{aligned} \Pr(x > X \text{ and } y > Y) &= 0.2 \times 0.26 \\ &= 0.052 \end{aligned}$$

The joint probability of 0.052 corresponds to a return period of 19 years.

3. CONCLUSION

This study of rainfall frequency indicates that the joint occurrence of the 5-hour storm of 31 May 1983 and the antecedent precipitation has a return period of 19 years. This suggests that the event represented something between a 5 year and a 20 year flood. However, more confidence in this estimate would be gained by carrying out a detailed analysis using a rainfall-runoff method.

LS/vw
17.6.88

Appendix 1

API VALUES FOR GADGET 125017 (R = 7.2%)

| Date | Annual Maximum (am) | API (am) | Date | Annual Maximum (am) | API (am) |
|------------|------------------------|-------------|------------|------------------------|-------------|
| 1 12 1916 | 58.6 | 64.17 | 1 7 1958 | 53.6 | 74.51 |
| 30 5 1918 | 54.1 | 56.16 | 16 4 1959 | 74.7 | 73.76 |
| 5 10 1918 | 47.2 | 49.12 | 3 12 1960 | 42.6 | 76.26 |
| 18 7 1919 | 51.8 | 55.49 | 26 4 1961 | 40.5 | 42.4 |
| 24 7 1916 | 53.8 | 56.72 | 6 5 1961 | 70.5 | 72.25 |
| 8 5 1917 | 53.9 | 43.55 | 6 7 1961 | 26.2 | 73.21 |
| 4 9 1918 | 24.1 | 52.51 | 14 5 1961 | 40.2 | 72.21 |
| 19 5 1919 | 54.5 | 66.47 | 21 5 1965 | 71.9 | 72.44 |
| 19 2 1920 | 41.7 | 51.27 | 21 5 1966 | 25.7 | 47.85 |
| 3 10 1921 | 42.5 | 18.25 | 14 5 1967 | 41.4 | 41.4 |
| 5 5 1922 | 64.9 | 56.13 | 14 7 1968 | 45.5 | 52.97 |
| 5 5 1923 | 25.4 | 26.29 | 19 6 1971 | 22.7 | 32.29 |
| 21 12 1924 | 22.1 | 41.24 | 20 11 1971 | 35.1 | 31.73 |
| 19 9 1925 | 37.5 | 23.69 | 1 12 1971 | 11.1 | 31.25 |
| 1 11 1926 | 21.5 | 29.83 | 24 9 1971 | 53.2 | 71.27 |
| 14 7 1927 | 30.5 | 45.46 | 1 11 1977 | 24.1 | 15.77 |
| 10 10 1928 | 34.8 | 29.44 | 16 6 1978 | 47.7 | 72.42 |
| 11 11 1929 | 32.5 | 47.16 | 25 10 1979 | 44.5 | 54.21 |
| 20 7 1930 | 39.6 | 29.29 | 7 5 1950 | 27.8 | 35.53 |
| 3 9 1931 | 72.1 | 51.59 | 24 4 1981 | 29.7 | 34.06 |
| 21 5 1932 | 58.4 | 41.41 | 22 5 1982 | 48.1 | 51.42 |
| 10 10 1933 | 55.6 | 44.15 | 31 5 1983 | 23.1 | 44.55 |
| 9 11 1934 | 18.5 | 24.33 | | | |
| 15 11 1935 | 30.2 | 70.74 | | | |
| 7 7 1936 | 25.4 | 41.50 | | | |
| 15 7 1937 | 31.3 | 14.24 | | | |
| 11 8 1938 | 30.0 | 55.89 | | | |
| 17 7 1939 | 27.4 | 43.25 | | | |
| 16 10 1940 | 30.5 | 22.85 | | | |
| 9 10 1941 | 29.5 | 19.95 | | | |
| 4 3 1942 | 26.2 | 16.26 | | | |
| 29 5 1943 | 23.4 | 23.08 | | | |
| 27 2 1944 | 29.2 | 33.86 | | | |
| 19 3 1946 | 31.2 | 55.81 | | | |
| 10 5 1947 | 22.9 | 33.93 | | | |
| 30 12 1948 | 42.7 | 26.10 | | | |
| 15 7 1949 | 40.1 | 24.97 | | | |
| 15 7 1950 | 21.8 | 20.86 | | | |
| 6 8 1951 | 69.6 | 16.96 | | | |
| 5 6 1952 | 31.5 | 23.47 | | | |
| 12 10 1953 | 24.1 | 14.39 | | | |
| 7 6 1954 | 35.3 | 69.75 | | | |
| 25 3 1955 | 25.4 | 30.59 | | | |
| 8 6 1956 | 24.1 | 27.73 | | | |
| 8 5 1957 | 42.4 | 51.73 | | | |

Appendix 2

Significance test on the correlation between API and annual maximum 1-day rainfall.

t-test

$$r = 0.184$$

$$n = 67$$

$$H_0 : \rho = 0$$

(no significant correlation)

$$H_1 : \rho \neq 0$$


Two-tailed test at 5% significance level

$$\begin{aligned} t &= \frac{r \sqrt{n-2}}{\sqrt{1-r^2}} \\ &= \frac{0.184 \sqrt{65}}{\sqrt{1-0.184^2}} \\ &= 1.509 \end{aligned}$$

$$v \text{ (degrees of freedom)} = n-2 = 65$$

$$\text{From tables, } t_{v=65, \alpha=0.05} = 1.998$$

Since $-1.998 < 1.509 < 1.998$, H_0 cannot be rejected. Therefore it must be concluded that there is no significant correlation between API and annual maximum 1-day rainfall.



INSTITUTE of HYDROLOGY

The **Institute of Hydrology** is a component establishment of the UK Natural Environment Research Council, grant-aided from Government by the Department of Education and Science. For over 20 years the Institute has been at the forefront of research: exploration of hydrological systems within complete catchment areas and into the physical processes by which rain or snow is transformed into flow in rivers. Applied studies, undertaken both in the UK and overseas, ensures that research activities are closely related to practical needs and that newly developed methods and instruments are tested for a wide range of environmental conditions

The Institute, based at Wallingford, employs 140 staff, some 100 of whom are graduates. Staff structure is multidisciplinary involving physicists, geographers, geologists, computer scientists, mathematicians, chemists, environmental scientists, soil scientists and botanists. Research departments include catchment research, remote sensing, instrumentation, data processing, mathematical modelling, hydrogeology, hydrochemistry, soil hydrology, evaporation flux studies, vegetation-atmospheric interactions, flood and low-flow predictions, catchment response and engineering hydrology

The budget of the Institute comprises £4.5 million per year. About 50 percent relates to research programmes funded directly by the Natural Environment Research Council. Extensive commissioned research is also carried out on behalf of government departments (both UK and overseas), various international agencies, environmental organisations and private sector clients. The Institute is also responsible for nationally archived hydrological data and for publishing annually **HYDROLOGICAL DATA, UNITED KINGDOM.**