


Could it happen here?

Brian Baptie

Published in SECED Newsletter, 23, 1, 16-17 (2011)

Could a magnitude 9 earthquake and tsunami like the one that struck Japan affect the British Isles? The short answer is no. Huge mega-thrust earthquakes like this only happen at plate boundary subduction zones where one of the Earth's tectonic plates is being pushed down, or subducted, beneath another. Places where this happens include Japan, Sumatra and South America all of which have had earthquakes of magnitude 8.5 or greater in the last few years that have resulted in tsunami. The British Isles sits in the middle of a tectonic plate, Eurasia. Our nearest plate boundary is at the mid-Atlantic ridge, where the earthquakes are too small to generate tsunami. The nearest subduction zones to Britain lie at the Hellenic Arc, south of Greece and in the Caribbean. Tsunami have occurred in both these regions in historic times, but did not affect the UK. The largest recorded British earthquake had a magnitude of 5.8 (6.1 ML) and was over 65,000 times smaller than the Tohoku earthquake in Japan. Although it occurred under the North Sea it was too small to generate a tsunami. This event is close to the maximum credible magnitude for a British earthquake. The UK experiences a magnitude 5 earthquake roughly every 10-20 years. These events typically cause some superficial damage.




Earthquakes greater than magnitude 8.2 (red circles). These tend to cluster at plate boundary subduction zones (yellow lines). By contrast there are no great earthquakes at mid-ocean ridges (red lines) such as the Mid-Atlantic ridge.

In contrast, [the Tohoku earthquake of 11 March 2011](#) ruptured a 400 km long segment of the plate boundary that lies east of Japan, running from the northern end of Honshu roughly south almost as far as Tokyo. East of Honshu, the Pacific plate is moving west at around 8 cm/year and is being pushed down, or subducted, underneath Japan at the Japan Trench. The plates had been locked together for many years before enough strain accumulated to

allow the fault to rupture. The average amount of slip on the fault was around 8m, resulting in Japan moving several metres east during the earthquake. This motion would have resulted in over 3 metres of uplift of the seafloor along the fault, displacing huge volumes of water and causing the giant waves or tsunami that spread out from the epicentre like ripples on a pond. The first wave took around thirty minutes to reach the coast of Japan as was followed by a number of other waves, which surged several kilometres inland.


However, it may surprise many people to learn that tsunami have occurred in Britain in the past.

Over 7000 years ago, a massive submarine slide off the coast of Norway, known as the Storegga slide, resulted in a tsunami reaching the northeast coast of Britain. Evidence of this can be found in geological deposits from northeast England to north of the Arctic Circle. These show that the wave reached over 20m above sea level at Sullom Voe, Shetland. However this quickly decreases to the south with 3-4 m in northeast Scotland and 1 m in northeast England. A repeat of this event is unlikely, since geological models suggest that another Ice Age is needed to re-establish the conditions for a similar failure.



Slides on the NW European continental margin.

In 1755, Lisbon was destroyed by a magnitude 8+ earthquake and tsunami. The tsunami reached the southwest coast of England and its arrival in Mount's Bay, Cornwall was observed by the naturalist William Borlase, who described several large waves crashing against the shore over a period of two hours. A study by BGS, [HR Wallingford](#) and the [Proudman Oceanographic Laboratory](#) commissioned by [Defra](#) looked in detail at the possible effects of another earthquake like the 1755 Lisbon earthquake. Modelling results suggest that the wave would take around 5 hours to reach Britain, with maximum wave heights of 1-2m around the majority of Cornwall. Such wave heights are similar to those resulting from typical storm surges that are experienced on a far more frequent basis.



Modelled free surface elevation of the sea surface 5 hours after a magnitude 8.7 earthquake displaces the sea-bed in the area of the Gorringe Bank, west of Lisbon.