

# Nature and timing of Late Mississippian to Mid Pennsylvanian glacio-eustatic sea-level changes of the Pennine Basin, UK

Colin N. Waters<sup>1\*</sup> and Daniel J. Condon<sup>2</sup>

<sup>1</sup>British Geological Survey, Kingsley Dunham Centre, Keyworth, Nottingham, NG12 5GG

<sup>2</sup>NERC Isotope Geoscience Laboratory, British Geological Survey, Kingsley Dunham Centre, Keyworth, Nottingham, NG12 5GG

\* E-mail: cnw@bgs.ac.uk

**Abstract:** The Pennine Basin of northern England contains a comparatively complete Serpukhovian–Moscovian succession characterised by high-resolution ammonoid zonation and cyclic paralic sedimentation. Two new ID-TIMS zircon ages from a bentonite deposited during the Arnsbergian (mid-Serpukhovian) regional substage and tonstein of earliest Bolsovian (early Moscovian) regional substage, have been determined. The weighted mean  $^{206}\text{Pb}/^{238}\text{U}$  ages of  $328.34 \pm 0.55$  and  $314.37 \pm 0.53$  Ma (total uncertainty), respectively, require modification of the timescale for the Western Europe regional chronostratigraphy.

The areal extent of acme ammonoid facies are used as a proxy for the magnitude of 47 discrete flooding events. Incised valleys (major sequence boundaries) are used as a proxy for the magnitude of sea-level falls. The frequency of these events, in the light of the new radiometric dating, indicates: (1) temporal coincidence between major glaciations in Gondwana and phases of increased frequency of sequence boundaries in the Pennine Basin; (2) high amplitude flooding surfaces have an average frequency of *c.* 400 ka; (3) average cycle durations during the Pendleian–early Arnsbergian and Chokierian–Bolsovian, of *c.* 111 ka and *c.* 150 ka, respectively, reflect short-duration eccentricities, and (4) multiple flooding surfaces with the same ammonoid assemblages may equate with sub-100 ka precession/obliquity frequencies.

**Supplementary material:** U-Pb method description and data, procedure for the calculation of the areal extent of marine bands and tables showing a full listing of biostratigraphical data used in the study are available at [www.geolsoc.org.uk/SUPXXXXXX](http://www.geolsoc.org.uk/SUPXXXXXX).

Limited data exists for constraining the Carboniferous timescale (Davydov *et al.* 2004), representing a major limitation of our understanding of biological and environmental change, their linkages, and the rates at which change occurred during that interval. The current timescale for the Carboniferous (Davydov *et al.* 2004, modified by Ogg *et al.* 2008) is largely derived from dating of international stages, defined by a conodont-based stratigraphy. However many parts of the world, including the equatorial paralic basins of Western Europe and eastern USA, and regions located close to a palaeopole (e.g. Australia) cannot successfully use this scheme due to the lack of suitable fauna. The development of truly global geological ‘timescales’ requires the calibration, via radio-isotopic dating, and integration, of several different biostratigraphical schemes.

During the Carboniferous-Permian ‘icehouse’ global scale climate oscillations occurred at varying timescales. The Gondwanan glacial record suggests a series of distinct and ‘short-lived’ (1–4 Ma) glacial episodes (Fielding *et al.* 2008) whereas the equatorial ‘Laurussia’ records both ‘long’ and ‘short’ climate fluctuation. Outstanding issues for understanding Carboniferous-Permian environmental change include determination of the timing, duration and tempo of glaciation and the equatorial response to changes in polar regions.

This paper integrates bio- and litho-stratigraphic analyses of mid Carboniferous (Serpukhovian to Moscovian) strata of the Pennine Basin of central and northern England and North Wales (Fig. 1) combined with new U-Pb dating with the overarching aim of developing an integrated biostratigraphic, geochronologic and palaeoenvironmental dataset. There are two specific aims: (1) the high-precision calibration of mid-Carboniferous biostratigraphic zonation for Western Europe; (2) assessment of cyclic marine bands in relationship to known orbital forcings via glacio-eustatic sea-level fluctuation.

Previous attempts at estimating the duration of Namurian and Westphalian cyclicity have assumed a constant forcing mechanism during the entire interval. In this study we investigate that premise and present evidence for variations in the magnitude of flooding and regressive events which aid determination of a cyclostratigraphy for the succession.

## **Summary of existing radiometric dates**

Hess & Lippolt (1986) and Berger *et al.* (1997) derived  $^{40}\text{Ar}/^{39}\text{Ar}$  sanidine plateau dates from German and Czech tonsteins which suggested that the combined duration of the Namurian and Westphalian (broadly Serpukhovian to Moscovian) was about 21 Ma. The age determinations were largely for Stephanian, Bolsovian and early to mid Namurian tonsteins (Table 1) and provided no dates at the base of either the Namurian or Westphalian regional stages.

Recalibration of the MMHb-1 mineral standard, summarised in Davydov *et al.* (2004) results in revised ages which approximate to determinations using U–Pb TIMS (Table 1). Hess & Lippolt (1986) provided age uncertainties of 1.0 to 9.2 Ma ( $2\sigma$ ); revised by Claoué-Long *et al.* (1995) to uncertainties of 7.4 to 10.0 Ma ( $2\sigma$ ) based upon consideration of the uncertainties in the age of standard mineral against which the  $^{40}\text{Ar}/^{39}\text{Ar}$  dates are calibrated. The limited biostratigraphical control in the German/Czech succession also limits the precise correlation of these dates globally.

$^{206}\text{Pb}/^{238}\text{U}$  zircon dates using the SHRIMP ion microprobe have provided Asbian (late Viséan) dates of  $334 \pm 4$  Ma ( $2\sigma$ ) from Poland (Kryza *et al.* 2010), Arnsbergian (early Namurian) dates of  $314.5 \pm 4.6$  Ma ( $2\sigma$ ) for the Pennine Basin, UK (Riley *et al.* 1995), and Bolsovian (Westphalian) dates of  $311.0 \pm 3.4$  Ma ( $2\sigma$ ) from Germany (Claoué-Long *et al.* 1995). These suggest a *c.* 3.5 Ma duration for much of the Namurian and Westphalian, combined. The potential inaccuracies related to standardisation (Ireland & Williams 2003), in addition to the reported uncertainties, for the U–Pb SHRIMP dates discussed above do not permit a precise duration of the intervals required for the advancement of Carboniferous stratigraphy.

The U–Pb SHRIMP dates are at odds with recent timescales (Davydov *et al.* 2004, modified by Ogg *et al.* 2008) and high precision ID-TIMS U–Pb zircon ages for the Donetz Basin (Davydov *et al.* 2010). The latter work provides errors of about 100 ka, of sufficient resolution to be useful in determining the duration of marine band cyclicity. However, ammonoids are rare in the Donetz Basin and the main biostratigraphical correlations are based on foraminiferal zones. Although correlation at the substage level between the Pennine and Donetz basins is established, it is not possible to directly correlate the ages with specific marine bands in the UK. The work of Davydov *et al.* (2010) is significant in that it provides a  $328.14 \pm 0.11$  Ma age for the early–mid Pendleian and a  $314.40 \pm 0.06$  Ma age for the early Bolsovian, more than 3 Ma older than previous determinations using  $^{40}\text{Ar}/^{39}\text{Ar}$  and  $^{206}\text{Pb}/^{238}\text{U}$  SHRIMP techniques (Table 1), most likely reflecting errors in the standardisation of both the  $^{40}\text{Ar}/^{39}\text{Ar}$  and  $^{206}\text{Pb}/^{238}\text{U}$  SHRIMP dates. The Bolsovian would now appear to be in excess of 4 Ma in duration (Davydov *et al.* 2010). Two U–Pb ID-TIMS zircon ages from the Silesian Basin,  $328.84 \pm 0.38$  Ma and  $328.01 \pm 0.36$  Ma (total uncertainty) from early Pendleian and late Pendleian strata, respectively, have been determined (Gastaldo *et al.* 2009). These authors extrapolate cycle duration to produce an estimate of 329.7 Ma, rounded up to 330 Ma for the base of the Serpukhovian.

The U-Pb TIMS age of  $326.8 \pm 0.98$  Ma of Trapp & Kaufmann (2002) from Germany is derived from a bentonite found within the ammonoid *Goniatites crenistria* (P1) Zone of late Asbian age (Waters *et al.* 2011).

## U-Pb Geochronology

### *Sample localities*

Bentonites found within Namurian strata comprise typically mixed-layer illite–smectite with subordinate kaolinite (i.e. K-bentonites). The trace element geochemistry is indicative of a rhyodacite-dacite composition for the late Pendleian to Arnsbergian ashfall deposits (Spears *et al.* 1999). Tonsteins are kaolinite aggregates, typically present in Westphalian strata in the Pennine Basin, interpreted as kaolinised volcanic ash-falls or reworked volcanic detritus, with both acid and basic tonsteins recognised geochemically (Spears & Kanaris-Sotiriou 1979). Bentonites and tonsteins most likely to contain primary zircons and with minimal siliciclastic contamination were selected. Eleven localities/boreholes were investigated with the aim of providing a broad range of dates for Brigantian (late Visean) to Bolsovian (late Westphalian) strata. However, following heavy mineral separation and age-screening using laser ablation inductively coupled plasma ionisation mass spectrometry (LA-ICP-MS) only two samples contained sufficient primary zircons to make dating chemical abrasion isotope-dilution thermal ionisation mass spectrometry (CA-ID-TIMS, see below) worthwhile.

The bentonite sample BLL1976 from the BGS Harewood Borehole (Fig. 1), West Yorkshire [BNG 43220 44410] at a depth of 304.10 m, were interpreted by Riley *et al.* (1995) as representing bentonite B6 of Trewin (1968). BLL1976 occurs within the upper part of the *Eumorphoceras yatesae* (E<sub>2a3</sub>) Marine Band of early Arnsbergian age and is equivalent to the sample analysed by Riley *et al.* (1995) for which their SHRIMP U/Pb date of  $314.4 \pm 4.6$  Ma was acquired using the SL13 zircon standard for U/Pb calibration.

Sample EH28155 from the Holme Pierrepont Borehole (Fig. 1), Nottinghamshire [BNG 46306 33933] at a depth of 181.8 m comes from the Sub-High Main tonstein, located 14 m above the Aegiranum Marine Band, the base of which marks the base of the Bolsovian regional substage. The stratigraphically nearest dated horizon is the Z1 tonstein from the Ruhr Coalfield, located just below the Aegiranum Marine Band. Hess & Lippolt (1986) provide a  $^{40}\text{Ar}/^{39}\text{Ar}$  sanidine plateau date of  $310.7 \pm 2.6$  Ma ( $2\sigma$ ) relative to MMHb-1 mineral standard using an age of 519.5 Ma. Claoué-Long *et al.* (1995) determined 39 measurements of 37 zircons from the Z1 tonstein, with a mean U–Pb SHRIMP age of  $311.0 \pm 3.4$  Ma ( $2\sigma$ ).

## Results

Zircons were analysed using CA-ID-TIMS methodologies employed at NERC Isotope Geoscience Laboratory (NIGL), details of which are outlined in an online supplemental material along with the tabulated results of the analytical programme. However, two important points are outlined here: (1) prior to dissolution zircons were subject to a modified chemical abrasion pretreatment for the effective elimination of Pb-loss (Mattinson 2005); and (2) the accuracy of the  $^{238}\text{U}/^{206}\text{Pb}$  dates presented herein are controlled by the gravimetric calibration of the EARTHTIME U-Pb tracer employed in this study and the determination of the  $^{238}\text{U}$  decay constant (Condon *et al.* 2007; Jaffey *et al.* 1971). Zircons separated from both bentonite samples BLL1976 and EH28155 were small ( $<50\text{ }\mu\text{m}$ ) with aspect ratios of  $\sim 1.5$  to  $\sim 3$ . For sample BLL1976 seventeen fractions (single grains) were analysed, with the resulting data presented and their interpretation discussed in more detail in the online supplemental material. In brief,  $^{206}\text{Pb}/^{238}\text{U}$  dates between 311 and 334 Ma (Fig. 2) with a distinct population (defined by 11 of the 17) of analyses yielding a weighted mean  $^{206}\text{Pb}/^{238}\text{U}$  date of  $328.34 \pm 0.30(0.43)[0.55]^1$  Ma (Mean square weighted deviation, MSWD = 2.2), which is interpreted as being the best estimate for the zircons of this sample and inferentially the age of bentonite at the sampled stratigraphic level. For sample EH28155 nine fractions (single grains) were analysed, and the resulting data are presented in Figure 2. Two of the nine analyses produced U-Pb dates older than the constraint imposed by sample BLL 1976 (see above). The remaining seven analyses yielded  $^{206}\text{Pb}/^{238}\text{U}$  dates between 306 and 317 Ma (Fig. 2) with a distinct population (defined by 4 of the 7) of concordant analyses yielding a weighted mean  $^{206}\text{Pb}/^{238}\text{U}$  date of  $314.37 \pm 0.25(0.40)[0.53]^1$  Ma (MSWD = 1.07) which is interpreted as being the best estimate for the zircons of this sample. In both samples the U/Pb dates that are older than the main population are interpreted as reflecting the analyses of zircon ante-/xeno-crysts, and grains that are younger as reflecting Pb-loss (Fig. 2) and interpretation that is supported by consideration of biostratigraphic and geochronologic constraints (e.g., Davydov *et al.* 2010).

<sup>1</sup>Errors presented in format  $\pm X(Y)[Z]$ : X- internal or analytical uncertainty in absence of all systematic error (tracer calibration and decay constants); Y includes the quadratic addition of tracer calibration error (using a conservative estimate of the  $2\sigma$  standard deviation of 0.1% for the Pb/U ratio in the tracer); Z includes the quadratic addition of both tracer calibration error and additional  $^{238}\text{U}$  decay constant errors (see online supplemental information).

These two new ages for the early Arnsbergian and early Bolsovian are significantly older than existing published U-Pb SHRIMP and  $^{40}\text{Ar}/^{39}\text{Ar}$  sanidine plateau dates for equivalent strata in Western Europe. Closer comparison with the  $^{40}\text{Ar}/^{39}\text{Ar}$  ages is evident when recalibration of the mineral standard is taken into consideration, but the errors are too great to make the ages of any use in understanding basin evolution timing. The new ages align (within  $2\sigma$  errors) with recent U-Pb ID-TIMS zircon ages from the Donetz Basin (Davydov *et al.* 2010).

## Stratigraphic analyses

### *Marine band cyclicity referenced in the Pennine Basin*

The ‘cyclicity’ of marine bands in the palaeo-equatorial Carboniferous Pennine Basin is considered by many (e.g. Holdsworth & Collinson 1988; Maynard & Leeder 1992; Martinsen *et al.* 1995; Hampson *et al.* 1997; Waters & Davies 2006) to be driven by eustatic sea-level fluctuations, a far-field response to polar environmental fluctuations in ice-sheet volume in the southern hemisphere (Veevers & Powell 1987; Isbell *et al.* 2003). Many of the marine bands recognised within the Pennine Basin can also be found in separate basins in Scotland, Ireland, northern France, Belgium, Holland and Germany (Ramsbottom 1979), indicating that sea-level rise, as opposed to regional subsidence, was the primary control of marine band formation. This marine band cyclicity is evident throughout strata of Pendleian (early Namurian) to Bolsovian (late Westphalian) age.

The marine bands occur at the base of marine to non-marine upward-coarsening cycles, equating to the parasequence of the Exxon sequence-stratigraphic model (Posamentier *et al.* 1988). The marine bands are taken to represent transgressive systems tracts and maximum flooding surfaces, with the acme marine facies coinciding with the maximum rate in rise of the sea-level curve (Posamentier *et al.* 1988). Alternatively, Martinsen *et al.* (1995) argue that due to the lengthy and sinuous nature of connections between the open sea and the basin, the condensed section represented by each of the ammonoid-bearing marine bands is likely to coincide with a maximum of the sea-level curve.

The periodicity of cyclicity has been estimated at 185 ka (Holdsworth & Collinson 1988), 120 ka (Maynard & Leeder 1992), or 65 ka based on SHRIMP U-Pb (zircon) dates (Riley *et al.* 1995). These values are consistent with eccentricity-forced modulation frequencies. The amplitude of sea-level variation has been estimated at about 42 m (Maynard & Leeder 1992) or 60 m (Church & Gawthorpe 1994), during the Namurian. The above estimated periodicities were based upon assumptions of a constant forcing during the entire interval and using time intervals for the Namurian which are shown in this study to lack required precision. For example, if we consider the  $2\sigma$  c. 4 Ma uncertainties on the SHRIMP U-Pb (zircon) dates the total duration of this

interval could range from ca. 0 to 11 Ma, or greater if SHRIMP U-Pb (zircon) dates are inaccurate, which has a great effect on the average cycle duration.

The existence of a superimposed higher-frequency obliquity or precession cyclicity, of lesser amplitude, has been proposed within the Pennine Basin (e.g. Brettle 2001; Waters *et al.* 2008; Tucker *et al.* 2009). In addition, Ramsbottom (1977) identified long-duration cyclic units, which he termed mesothems. He proposed eleven ‘mesothemic’ cycles for the British Namurian (Table 2), linked to the appearance of new ammonoid genera, and controlled by longer duration eustatic sea-level fluctuations, which he estimated to have an average duration of 1.1–1.35 Ma (Ramsbottom 1979). Similarly, Ramsbottom (1979) proposed the presence of ten mesothems within the Westphalian succession, of average duration of 1.66 Ma, with the lowermost coinciding with the uppermost Namurian mesothem (Table 3). Both Namurian and Westphalian mesothems were described as comprising pulsed marine transgressions in which each successive transgression is increasingly extensive, with the top of the mesothem marked by the most widespread marine band, commonly with a widely developed ammonoid facies (Ramsbottom 1979). The boundaries of the mesothems are marked by widespread disconformities in shelf areas. Sequence-stratigraphic terminology would consider the mesothems to broadly represent sequences (Posamentier *et al.* 1988), although the mesothem boundaries were taken at the top of major fluvial sandbodies and not at the sequence boundary at the base. The upper cycles of each Namurian mesothem were considered to be typically sandstone-dominated and associated with overall base-level fall and progressive progradation of fluvial systems further into the basin (Ramsbottom 1977). Holdsworth & Collinson (1988) provided a rigorous critique of the mesothemic concept, arguing that the linkage of sand-dominated cycles with regression and the major ammonoid turnover of taxa at mesothem boundaries could not be demonstrated. They also argued against the lateral extent of ammonoid-bearing marine bands being suitable as a means to deduce the form of major eustatic curves. Holdsworth & Collinson (1988) also considered that the example used by Ramsbottom (1977) of the transition between the Askrigg Block and Craven Basin (Fig. 1) is invalidated by the potential of tectonic uplift and subsidence influencing the areal extent of marine bands.

### *New observations on a Namurian-Westphalian marine band cyclostratigraphy*

The areal extent of marine bands is primarily a function of the magnitude of eustatic sea-level rise, influenced by basin topography and subsidence/uplift rates. Hence, to use the areal extents of marine bands as a proxy of relative magnitude of sea-level rise it is important to understand the relative significance of these factors. The Southern Uplands High and Wales-Brabant Massif,

bounding, respectively, the northern and southern margins of the Pennine Basin (Fig. 1), form topographical highs throughout the Carboniferous, and flooding events do not extend across them. The Namurian to early Westphalian interval covered by this study was a time of the onset of broad thermal subsidence between the Southern Uplands High and Wales-Brabant Massif (Leeder 1982). Waters *et al.* (1994) provided evidence that pulses of basin extension continued, though on a much reduced scale, during the thermal subsidence phase until Bolsovian times, at which time Variscan compressional structures became increasingly important, culminating in the end Carboniferous inversion of the entire basin. Deformation varied from growth folding within thick successions toward the basin depocentre to development of angular unconformities at the basin margin (Waters *et al.* 1994). Although such deformation may locally influence the absence of ammonoid fauna in the peripheral parts of the basin for some flooding events, by studying the basin as a whole it is considered that tectonism has no or only minimal influence on the extent of the majority of marine bands.

The ‘Block and Basin’ topography generated during Late Devonian to Mid Mississippian rifting (Leeder 1982) was gradually infilled by fluvio-deltaic sediments during Namurian times. However, this infilling was a diachronous process, starting in the north of the basin during the Pendleian, and the inherited basinal topography was only largely infilled by fluvio-deltaic sediments in the south by late Marsdenian times (Church & Gawthorpe 1994; Jones & Chisholm 1997; Waters & Davies 2006). Hence, flooding events, particularly during the early Namurian and in the south of the Pennine Basin, were in part constrained to the relict topographical lows associated with former half grabens and grabens, and only the highest sea-level maxima resulted in marine inundation of the block areas. By Westphalian times the fluvio-lacustrine sedimentation was associated with low-profile delta plains occupying the entire basin and in which flooding events were able to extend unconfined across the delta top. Subsidence rates were greatest in the basin depocentre around south Lancashire (Calver 1968), but this was not expressed as a topographical low (Rippon 1996). Hence, during Westphalian times, the marine band extent closely relates to the magnitude of sea-level rise.

The extents of Namurian marine bands within the Pennine Basin have been poorly delineated by previous workers, with the exception of the *Cancelloceras* (*Gastrioceras*) *cumbriense* Marine Band (Wignall 1987). During the current study, only the extent of the acme ammonoid facies was determined, as it is this that displays the diagnostic fauna which allows the unique identification of each marine band. The methodology by which the extents of the marine bands were defined is described in the online supplemental material. The extent of each of the Namurian ammonoid-bearing marine bands is shown in figures 3 and 4a–d. In contrast, marine



bands within the Westphalian succession are well documented, with the extents of the various faunal phases shown for each flooding event (Calver 1968, 1969). Only eight of the 24 Westphalian marine bands include an acme ammonoid facies (Table 3) and only those marine bands are delineated in Figure 4e–f, again showing only the extent of the ammonoid facies. Westphalian marine bands with less diagnostic fauna, such as thin-shelled ammonoids (*Anthracoceras*), marine bivalves (*Dunbarella*), brachiopods (*Lingula*) and foraminifera, record a transition in salinity from marine to brackish environments, with four of the marine bands characterised by the presence of the brachiopod crustacean *Estheria*, interpreted as occurring within a transitional zone between swamp and marine environments (Calver 1968).

The early Namurian marine bands show compartmentalisation within small areas. This in part reflects the prominent basin topography at the time, with flooding events tending to be limited in extent to the unfilled Visean sub-basins. Also, the outcrop of these successions is relatively isolated, with successions of this age poorly known at depth within the central part of the basin. Furthermore, many of the early Namurian marine bands occur within mudstone-dominated successions lacking intervening deltaic intervals, making identification of the specific marine band at times difficult. As a result, fewer early Namurian exposures were included in the study compared with later Namurian marine bands. Consequently, comparisons between early Namurian and late Namurian magnitudes of sea-levels should be done with care, but within each of these intervals, comparison of individual flooding event extents was considered justifiable.

By Marsdenian times, the ammonoid facies components of marine bands appear to be more laterally extensive, centred upon the area of greatest magnitude of subsidence in south Lancashire (Fig. 4b–c). The pre-Namurian basin topography was largely infilled and accommodation space was the product of eustatic sea-level rises in combination with broad thermal subsidence of the Pennine Basin. This pattern continued into Langsettian and Duckmantian times (Fig. 4e–f). Bolsovian marine bands appear to show a slight shift of the focus of the ammonoid facies eastwards, towards the East Midlands, and are generally less extensive. The Cambriense Marine Band represents the final marine flooding event to affect the Pennine Basin (Guion *et al.* 1995; Aitkenhead *et al.* 2002; Waters *et al.* 2011).

### *Incised valley fills in the Pennine Basin and evidence for major regressive events*

Discontinuities and unconformities are present within the Pennine Basin, though it is only within recent decades that their significance has been recognised. These major Type 1 unconformities, evident as incised palaeovalleys, represent sequence boundaries (Posamentier *et al.* 1988). Although many cycles include an upper fluvial-deltaic component, there has been a tendency to

over interpret all boundaries with an overlying coarser grained succession to represent a Type 2 sequence boundary. The significance of these surfaces, whether they be the product of sea-level falls complementing sea-level rises indicated by the marine bands, or through deltaic avulsion during distinct flooding events, remains controversial (Waters *et al.* 2008) and, as a result, Type 2 sequence boundaries are not considered in this study. A number of Namurian and Westphalian thick, multi-storey fluvial complexes have been interpreted as sediments deposited in palaeovalleys generated by incision during significant sea-level fall.

Most Namurian-aged multi-storey sandstone bodies are 10-30 km wide and the preserved thickness is 25-35 m (Davies *et al.* 1999). Much of the evidence for low base levels and fluvial incision during the early Namurian comes from the comparatively condensed successions above the Alston and Askrigg Blocks (Fig. 1). Above the Alston Block, the Rogerley Channel (Fig. 3a), of north-south orientation, is up to 4 km wide and is associated with up to 30-40 m of erosional relief, including localised removal of the marine Knucton Shell Bed (Dunham 1990). The incised valley formed at approximately the same time as a major intra-E<sub>1c</sub> angular unconformity, associated with the northward tilting and subsequent erosion of the Askrigg Block (Brandon *et al.* 1995). Within the lowermost Arnsbergian E<sub>2a1</sub> cycle of the western part of the Askrigg Block (Fig. 3b), a lowstand erosion surface is recognised below the Upper Howgate Edge Grit (Martinsen 1993; Martinsen *et al.* 1995), though with insufficient erosion to remove the underlying *Cravenoceras cowlingsense* Marine Band (Table 2). A younger Arnsbergian succession on the Askrigg Block (Fig. 3b) includes a prominent intra-E<sub>2a3</sub> unconformity below the Red Scar Grit (and equivalent sandstones). An erosional relief of up to 100 m, associated with the removal of four marine bands may in part coincide with tectonic activity (Brandon *et al.* 1995). An intra-E<sub>2c2</sub> erosion surface at the base of the Lower Follifoot Grit in the southeast of the Askrigg Block (Fig. 3c) is associated with complete removal of the E<sub>2c1</sub> cycle, with the unconformity resting upon the E<sub>2b3</sub> marine band (Martinsen 1993; Martinsen *et al.* 1995). An unconformity is demonstrated in the northern part of the Craven Basin, occurring immediately beneath the H<sub>1a3</sub> marine band, with spores of E<sub>2c4</sub> present beneath the erosive surface, but no incised valley fill sandbody is recorded (Owens *et al.* 1990). In the southeastern part of the Askrigg Block (Fig. 3d), the markedly erosive base of the fluvial channel of the Upper Follifoot Grit, with pedogenically modified interfluvies, suggest base level fall within the H<sub>1b2</sub> subzone (Martinsen 1993).

Kinderscoutian to Yeadonian sandbodies have been subject to the most scrutiny within the central part of the Pennine Basin, with two distinct settings for development of incision. Some valley fills show a marked increase in thickness towards, and immediately upstream of the mouth

of the incised valley, developed in association with turbidite-fronted deltas within steep submarine slopes. These sandbodies thicken from 20–30 m to 50–80 m over a distance of 2–5 km and are filled with giant foresets (Hampson *et al.* 1999). These valley fills occur within the earliest deltaic infill of the Pennine Basin, ranging from Kinderscoutian within the central part of the basin, to Yeadonian in the south. In contrast, extensive sheet-like sandstones, up to 45 m thick and up to 70 km wide (Hampson *et al.* 1999), develop mainly within Marsdenian to Yeadonian successions and are associated with little basinal topography.

The Kinderscoutian Lower Kinderscout Grit (Hampson 1997) and associated basal turbidite channel of the Todmorden Grit (Fig. 4a), of probable  $R_{1c}3$  age, display local erosion of two  $R_{1c}$  marine bands. The uppermost Kinderscoutian cycle includes a marked erosive surface at the base of the Upper Kinderscout Grit (Fig. 4a), which locally erodes through the Butterly ( $R_{1c}5$ ) Marine Band (Hampson 1997). The extensive sheet-like Marsdenian Midgley Grit (Fig. 4b) shows incision of the underlying  $R_{2b}3$  marine band (Brettell 2001). The erosive base of the Marsdenian Roaches Grit/Ashover Grit is associated with a palaeovalley up to 80 m deep (Jones & Chisholm 1997), which removes the underlying  $R_{2b}5$  marine band within the eastern part of the Widmerpool Gulf (Fig. 4b; Church & Gawthorpe 1994). The east-west palaeochannel of Chatsworth Grit (Fig. 4c) is 25 km wide with a steep 50 m high northern flank, though the incision has not removed underlying marine bands (Waters *et al.* 2008). The late Yeadonian Rough Rock is typically a low-sinuosity broad sheet-like fluvial sandbody. However, within the Widmerpool Gulf and East Midlands Shelf a north-south incised valley in excess of 11 km width (Fig. 4d) incises up to 5 marine bands, down to and including the  $R_{2c}2$  Marine Band (Church & Gawthorpe 1994; Hampson *et al.* 1997).

The early Langsettian Crawshaw Sandstone of the East Midlands Shelf is less than 70 km wide (Fig. 4e) and has well defined margins with identifiable interfluvial deposits and removes up to three marine bands including the Subcrenatum and  $G_{1b}1$  marine bands (Hampson *et al.* 1997). This is the last representative of this style of fluvial deposition within the Pennine Basin. Subsequent fluvial systems are characterised by less laterally extensive sandstone bodies, typically up to 20 km wide, with a maximum of 30 km, and 8–20 m thick with a maximum of 100 m (Guion *et al.* 1995; Aitken *et al.* 1999). The presence of incised valley fills within this Westphalian succession remains controversial. Regionally developed well-drained palaeosols, considered to form on the interfluvial deposits, are not common, and major fluvial sandbodies show only limited basal incision of up to 5 m, exceptionally up to 8 m (Rippon 1996). This may reflect distance from the sea, with incision of river channels having insufficient time to work upstream from the coast before the next flooding event (Aitkenhead *et al.* 2002). Alternatively, it may relate to the enclosed nature

of the basin. Only the highest global sea-level rises would result in a rapid base-level rise within the basin, with the subsequent fall in sea-level leaving an isolated basin for which base levels may fall comparatively slowly as it continues to be fed by rivers (Waters & Davies 2006).

Westphalian multi-storey sandbodies commonly show a relationship with their adjacent strata. This may include high ash contents in coals adjacent to the sandbodies, coal splitting towards channels and an increase in interbedded sandstone layers in proximity to the channel bodies (Aitken *et al.* 1999; Guion *et al.* 1995). These observations imply that overbank flooding events from the channels occurred during peat accumulation and leads these authors to believe that many of the channel systems were aggradational, as opposed to having filled previously incised valley systems.

The Duckmantian and Bolsovian succession includes a number of multistorey sandstone bodies with local basal erosional relief that represent candidate incised valley fills (Fig. 4f). The basal Duckmantian Thornhill Rock cuts through the Vanderbecker Marine Band (Lake 1999) and reaches thicknesses of 37-45 m. The Woolley Edge Rock is distinctly coarser grained and pebbly compared with earlier Westphalian sandbodies. The channel fill, about 23 km wide, shows palaeocurrents to the west or WNW and has up to 60 m of erosional truncation (Aitken *et al.* 1999), including removal of the Manton Estheria Band. The Oaks Rock, up to 40 m thick, is similarly associated with the absence of the Haughton and Sutton marine bands (Lake 1999). The early Bolsovian Mexborough Rock is the fill of an east-west channel, 30-40 m thick (locally up to 80 m) and 15-30 km wide (Aitken *et al.* 1999) and may be associated with erosion of the Main Estheria Band and Edmondia Band.

## Discussion

The recognition of cyclic sedimentation offers the potential to develop high-resolution time-thickness models for sedimentary successions, in which the resolution is determined by the dominant forcing mechanism. This approach is routinely applied to Cenozoic strata and the current Neogene Timescale (Gradstein *et al.* 2004) is based entirely upon astronomical calibration. Milankovitch orbital forcing during the Carboniferous is thought to be 413 ka and 112 ka for long- and short-duration eccentricity periodicities, respectively, 34 ka for obliquity and 21 ka and 17 ka for precession frequencies (Maynard & Leeder 1992).

The late Mississippian to Pennsylvanian is a time of high frequency-high amplitude sea-level oscillations during icehouse conditions. The presence of cyclic stratigraphy in the Pennine Basin, evidenced by the presence of cyclothems and periodic development of marine bands, has lead many workers to suggest these can be used to generate high-resolution age-models for parts of the

Carboniferous (Ramsbottom 1977; Maynard & Leeder 1992; Brettle 2001; Waters *et al.* 2008; Tucker *et al.* 2009). Recent developments in high-precision U-Pb geochronology means that it is now possible to test some of these hypotheses and develop more accurate models for the evolution of a Carboniferous cyclostratigraphy.

Most attempts at estimating the duration of the Namurian and Westphalian cyclicity have assumed a constant forcing mechanism during the entire interval; simply duration divided by the number of marine bands (fossiliferous carbonaceous mudstones or impure limestones). The new radiometric ages from this study combined with the most recent and accurate estimates of 330 Ma and 322.8 Ma for the base of the Serpukhovian and Bashkirian, respectively (Davydov *et al.*, 2010), permit the detailed analysis of cyclicity duration over specific time intervals.

There are 15 marine bands in the 1.66 Ma interval between the base of the Pendleian to the dated bentonite (BLL1976) of early Arnsbergian age (Table 2; Fig. 5). The average duration of ca. 111 ka represents a possible short-duration eccentricity modulation. Four peak flooding events are recognised ( $E_{1a1}$ ,  $E_{1c1}$ ,  $E_{2a1}$  and  $E_{2a3}$ ), which are equated with the 400 ka long-duration eccentricity frequency and identified as the orbital cycles S1 to S4.

The mid to late Arnsbergian interval between the dated bentonite (BLL1976) and the base of the Bashkirian Stage, which occurs above the  $H_{1a1}$  marine band (Riley *et al.* 1995), there are 13 marine bands over a duration of 5.54 Ma (Table 2; Fig. 5). The average of ca. 426 ka may suggest that only the long-duration eccentricity frequency is observed. However, the marine bands  $E_{2b1a-c}$ ,  $E_{2b2a-c}$ ,  $E_{2c2-4}$  and  $H_{1a1-3}$  occur as triple short-duration flooding events associated with the same ammonoid fauna (see below), and are considered to represent only four 400 ka flooding events. If these four, along with the  $E_{2b3}$  and  $E_{2c1}$  are taken to represent peak flooding events, it requires that eight 400 ka cycles are not represented in the geological record by marine flooding events (Fig. 5). It is possible that these missing cycles may be distributed evenly throughout the mid to late Arnsbergian interval. However, the very marked change in ammonoid genera between the  $E_{2c4}$  and  $H_{1a1}$  flooding events is here interpreted as indicative of a long period without marine flooding of the basin. During the Pendleian and Arnsbergian the dominant genera are *Cravenoceras*, *Eumorphoceras* and *Cravenoceratoides*, but these genera are absent from Chokierian and younger strata (Table 2).

Between the base of the Bashkirian Stage and the dated early Bolsovian tonstein (EH28155) there are 56 marine bands in the Pennine Basin (Tables 2 & 3; Fig. 5) over an interval of 8.43 Ma. This results in an average marine band cycle of ca. 150 ka, a possible short-duration eccentricity modulation. However, these high-frequency events appear to be less common during

the early to mid Kinderscoutian and late Marsdenian to Yeadonian. Twenty one peak flooding events are recognised between the H<sub>1b</sub>1 to Aegiranum marine bands, which are equated with the 400 ka long-duration eccentricity frequency and identified as the orbital cycles B2 to M2 (Fig. 5). Eight out of the 24 Westphalian marine bands are associated with ammonoid assemblages (Table 3), of which seven of these marine bands, along with the brachiopod-bearing Haughton Marine Band, are recognised as representing the 400 ka peak flooding events.

A common feature of many of the Namurian marine bands is the presence of two or three distinct beds with marine fauna, separated by non-marine barren mudstones. In many cases the separation between these bands is only a few tens of centimetres to metres scale, e.g. E<sub>1a</sub>1 (a-c) or E<sub>2a</sub>1 (a-c), but in other cases they are separated by tens of metres of succession, including prograding deltaic lobes, e.g. R<sub>1c</sub>1-3 or R<sub>2b</sub>1-3. Importantly, these distinct “leaves” show the same ammonoid assemblages and cannot be readily distinguished, unless all “leaves” are evident in a single section or through correlation of intervening deltaic sandbodies. If one assumes a constancy of rate of evolution of ammonoid species through the Namurian it would suggest that these multiple marine bands occur at higher frequencies than between marine bands with distinct ammonoid taxa. This would suggest shorter frequencies than the 111 ka or 150 ka periodicities recognised above, and may indicate evidence of precession or obliquity components of too short a duration to be determined through current radiometric dating techniques.

Throughout the Namurian and Westphalian, 17 major base-level falls are recognised, at a frequency of about 1 Ma (Fig. 5). This long-duration cyclicity broadly approximates to the periodicity of the “mesothems” of Ramsbottom (1977; 1979). These major regressive events occur at greatest frequency during late Pendleian to late Arnsbergian, Chokierian, late Kinderscoutian to early Langsettian and Duckmantian to early Bolsoviaian times, but appear to be absent throughout much of the late Arnsbergian, Alportian to mid Kinderscoutian and Langsettian.

The variations in development of unconformities within the Pennine Basin may be a far-field response to the record of alternating glacial and non-glacial climatic regimes proposed for Gondwana. The onset of the main phase of glaciation began in the early Namurian and peaked in the late Westphalian and Stephanian (Gonzalez-Bonorino & Eyles 1995). Isbell *et al.* (2003) recognised an early Visean alpine glaciation event, but considered the main continental glaciation to persist across Gondwana from early Serpukhovian to Permian times, with a phase of minimal ice during latest Bashkirian to early Moscovian times. Fielding *et al.* (2008) indicate that eastern Australia was affected by four major glaciations during the Carboniferous, each separated by non-glacial periods of similar duration. Their estimates for the ages of these

glaciations are based upon U-Pb SHRIMP dates (e.g. Claoué-Long *et al.* 1995, Roberts *et al.* 1995), now considered unsuitable (see summary of existing dates above). Within the Paganzo Basin of Argentina, three glacial pulses during the mid Visean, early Bashkirian (*c.* 323–319.57 ± 0.09 Ma) and latest Bashkirian to early Moscovian (315.46 ± 0.07–312.82 ± 0.11 Ma) are well constrained by U-Pb ID-TIMS dates from zircons (Gulbranson *et al.*, 2010). The timing and duration of glaciations remain problematic and until redating occurs in Eastern Australia, the far-field response to these glaciations may provide the most suitable method for estimating their ages.

Glaciation C1 of earliest Namurian age, proposed by Fielding *et al.* (2008) to represent the initiation of the Late Palaeozoic Ice Age, is of limited extent and short-duration. In the study area, Type 1 unconformities are developed during the late Pendleian to early Arnsbergian and tend to be limited to the basin margin (Fig. 3). These unconformities follow after three of the four peak flooding events described earlier (Fig. 5), suggesting linkage to the ~400 ka eccentricity frequency. This early Serukhovian glaciation is considered to range from orbital cycles S1 to S9, approximately 330.0 to 326.5 Ma (Fig. 5). The start of Cycle S1 coincides with the sudden cessation of platform carbonate deposition and Cycles S2 to S5, between the E<sub>1c</sub>1 and E<sub>2a</sub>3 marine bands, is marked by the first phase of thick fluvio-deltaic siliciclastic sandbodies entering the basin (Aitkenhead *et al.* 2002; Waters & Davies 2006).

An absence of Type 1 unconformities and inferred absence of eight 400 ka cycle flooding events during the mid to late Arnsbergian may represent the interval between the C1 and C2 glaciations. Isbell *et al.* (2003) considered there to be evidence of glaciations in South America and Tibet in this interval. However, Stephenson *et al.* (2010) have demonstrated using  $\delta^{18}\text{O}$  and  $\delta^{13}\text{C}$  from the Pennine Basin that widespread ice-caps were absent throughout mid Serpukhovian times. This interval is considered to range from orbital cycles S10 to S17, approximately 326.5 to 323.0 Ma (Fig. 5). Few large sandbodies prograded into the Pennine Basin during this time interval (Aitkenhead *et al.* 2002; Waters & Davies 2006).

The Glaciation C2 of mid-Namurian age, estimated to range from 322.5–319.5 Ma by Fielding *et al.* (2008) is coincident with the second glacigenic phase in Argentina (Gulbranson *et al.*, 2010). It broadly aligns with the phase of Chokierian to Alportian increased frequency of unconformities and marine flooding events. This early Bashkirian glaciation is considered in this study to range from orbital cycles B1 to B4, approximately 323.0 to 321.5 Ma (Fig. 5). This was a time when the Pennine Basin was dominated by slow hemipelagic deposition with comparatively few large sandbodies prograding into the basin (Waters & Davies 2006).

The early to mid Kinderscoutian within the Pennine Basin is marked by low magnitude variations in sea-level, no major incision events and no multiple (high-frequency) marine bands. Deposition at this time was dominated by slow accumulation of hemipelagic and distal turbidite deposits (Waters & Davies 2006). It is suggested that this may equate with the interval between Glaciations C2 and C3, during which eustatic sea-level fluctuations are more subdued, but suggests not completely free of ice-sheet development. As with the C1–C2 interglacial, this interval is associated with a marked turnover of ammonoid genera, most notably marked by the appearance of *Reticuloceras* (Table 2). This interval is considered to range from orbital cycles B5 to B9, approximately 321.5 to 319.5 Ma (Fig. 5).

The latest Namurian to earliest Westphalian Glaciation C3, estimated by Fielding *et al.* (2008) to range of 317–315 Ma, was considered to have a comparable areal extent to Glaciation C2. No equivalent of this glaciation is recorded in the Paganzo Basin of Argentina (Gulbranson *et al.*, 2010). This broadly coincides with the late Kinderscoutian to early Langsettian phase in the Pennine Basin of highest magnitude sea-level flooding events and most numerous incision surfaces, almost on a ~400 ka cyclicity, with a return to common multiple (high-frequency) marine bands. In contrast to the early Namurian event, these unconformities commonly follow trends of decreasing amplitude flooding events. The onset of this glaciation is evident through a rapid and almost basinwide progradation of coarse, pebbly and commonly sheet-like fluvio-deltaic sandbodies, with this style of deposition persisting throughout this interval (Waters & Davies 2006). This glaciation is considered to range from orbital cycles B10 to B18, approximately 319.5 to 316.0 Ma (Fig. 5).

The short-lived interglacial interval between C3 and C4 equates with the mid- to late-Langsettian succession, possibly representing an interval of up to two orbital cycles (B18 to B20). The interval follows the Listeri Marine Band, the most extensive of all of the flooding events (Fig. 5), after which sea-level fluctuations are diminished and major incision surfaces are not recorded. The interval also records a marked diminution of sandbody dimension and grainsize (Guion *et al.* 1995; Waters & Davies 2006). As for the C2–C3 interglacial, this interval appears not to be completely free of ice-sheet development.

The youngest mid-Westphalian Glaciation C4, estimated by Fielding *et al.* (2008) to range from 313–308 Ma, is thought to be in part coincident with the third glacigenic phase in Argentina (Gulbranson *et al.*, 2010). It may equate with the Duckmantian to early Bolsovian phase of high-frequency major flooding and incision events (Fig. 5). The expression of this glaciation in the Pennine Basin is considered to range from orbital cycles B20 to M3, approximately 315.2 to 314.0 Ma (Fig. 5).



## Conclusions

This study provides two new high-precision U/Pb ages of  $328.34 \pm 0.55$  Ma (total uncertainty) for the Arnsbergian regional substage (mid-Serpukhovian stage) and  $314.37 \pm 0.53$  Ma (total uncertainty) for the earliest Bolsovian regional substage (early Moscovian stage). These ages are somewhat older than existing published ages for these successions and require modification of the current timescale for the Western Europe regional chronostratigraphy, but align with recent dates provided by Davydov *et al.* (2010) for the Donetsk Basin.

The extent of acme ammonoid facies within discrete marine intervals is used as a proxy of the magnitude of these marine flooding events. The recognition of candidate incised valleys, and the number of cycles locally removed by these major sequence boundaries, is used as a proxy of the magnitude of sea-level falls. The frequency of these events, when considered in the light of the new radiometric dating indicates the following relationships:

- 1) The interval between major sequence boundaries within the Namurian and Westphalian is approximately 1 Ma. This cyclicity may be a far-field response to the record of alternating glacial and non-glacial climatic regimes proposed for Gondwana. The four major glaciations proposed for Gondwana may equate with phases of increased numbers of sequence boundaries in the Pennine Basin. It is suggested that the main glaciations occurred during the late Pendleian to late Arnsbergian (approximately 330.0 to 326.5 Ma), Chokierian to Alportian (approximately 323.0 to 321.5 Ma), late Kinderscoutian to early Langsettian age (approximately 319.5 to 316.0 Ma) and Duckmantian to early Bolsovian (approximately 315.2 to 314.0 Ma).
- 2) The interglacial intervals are associated with no development of incised valleys, no or reduced frequency of flooding events and marked turnover of ammonoid genera, considered to mark long time durations between successive flooding events.
- 3) Distinct peak flooding surfaces within Namurian strata, associated with ammonoid-bearing marine bands in the Westphalian succession, have an average frequency of 400 ka, equating with the long-duration eccentricity component described in mid- to late-Pennsylvanian strata in the USA.
- 4) Average durations of marine band cycles during the Pendleian to early Arnsbergian and Chokierian to Bolsovian, of 111 ka and 150 ka, respectively, may reflect a short-duration eccentricity component. These flooding events are associated with non-ammonoid marine fauna in the Westphalian succession.

552 5) Multiple flooding surfaces associated with the same ammonoid assemblages in the  
553 Namurian may equate with sub-100 ka precession or obliquity frequencies.

554 The interaction of cyclicities associated with long-duration switching from glacial and non-  
555 glacial climatic regimes and long- and short-duration eccentricity cycles offers the opportunity of  
556 trans-continental cyclostratigraphical correlations within late Mississippian to Pennsylvanian  
557 successions.

## 558 **Acknowledgements**

559 Mike Stephenson and Steve Noble are thanked for helpful comments on an earlier draft of this  
560 paper and J S Daly and V Davydov for their helpful reviews of the manuscript. The help of Mike  
561 Howe (National Geoscience Information Service) in providing access to BGS borehole core and  
562 palaeontological databases is much appreciated. Colin Waters and Daniel Condon publish with  
563 the permission of the Executive Director, British Geological Survey, Natural Environment  
564 Research Council. U-Pb (zircon) analyses were supported by NIGFSC grant IP/949/1106.

## References

- AITKEN, J.F., QUIRK, D.G. & GUION, P.D. 1999. Regional Correlation of Westphalian sandbodies onshore UK: implications for reservoirs in the Southern North Sea. *In:* FLEET, A.J. & BOLDY, S.A.R. (eds). *Petroleum Geology of Northwest Europe. Proceedings of the 5th Conference*. Geological Society of London, 747–756.
- AITKENHEAD, N., BARCLAY, W.J., BRANDON, A., CHADWICK, R.A., CHISHOLM, J.I., COOPER, A.H. & JOHNSON, E.W. (eds). 2002. *British regional geology: The Pennines and adjacent areas*. (4th edition). HMSO, London.
- BRANDON, A., RILEY, N.J., WILSON, A.A. & ELLISON, R.A. 1995. Three new early Namurian (E1c-E2a) marine bands in central and northern England, UK, and their bearing on correlations with the Askrigg Block. *Proceedings of the Yorkshire Geological Society*, **50**, 333–355.
- BRETTLE, M.J. 2001. Sedimentology and high-resolution sequence stratigraphy of shallow water delta systems in the early Marsdenian (Namurian) Pennine basin, Northern England. *PhD thesis, University of Liverpool*.
- BURGER, K., HESS, J.C. & LIPPOLT, H.J. 1997. Tephrochronologie mit Kaolin-Kohlentonsteinen: Mittel zur Korrelation paralischer und limnischer Ablagerungen des Oberkarbons. *Geologisches Jahrbuch*, **A147**, 3–39.
- CALVER, M.A. 1968. Distribution of Westphalian marine faunas in northern England and adjoining areas. *Proceedings of the Yorkshire Geological Society*, **37**, 1–72.
- CALVER, M.A. 1969. Westphalian of Britain. *Compte Rendu 6e Congrès Internationale Stratigraphie et Géologie Carbonifère, Sheffield, 1967*, **I**, 233–254.
- CHURCH, K.D. & GAWTHORPE, R.L. 1994. High resolution sequence stratigraphy of the late Namurian in the Widmerpool Gulf (East Midlands, UK). *Marine and Petroleum Geology*, **11**, 528–544.
- CLAOUÉ-LONG, J.C., COMPSTON, W., ROBERTS, J. & FANNING, C.M. 1995. Two Carboniferous ages: a comparison of SHRIMP zircon dating with conventional zircon ages and  $^{40}\text{Ar}/^{39}\text{Ar}$  analysis. *SEPM Special Publication*, **54**, 3–21.
- CONDON, D., SCHOENE, B., BOWRING, S., PARRISH, R., MCLEAN, N., NOBLE, S. & CROWLEY, Q. 2007. EARTHTIME; isotopic tracers and optimized solutions for high-precision U-Pb ID-TIMS geochronology. *Eos, Transactions, American Geophysical Union*, **88**.

- DAVIES, S.J., HAMPSON, G.J., FLINT, S.S. & ELLIOTT, T. 1999. Continental-scale sequence stratigraphy of the Namurian, Upper Carboniferous and its applications to reservoir prediction. *In: FLEET, A.J. & BOLDY, S.A.R. (eds). Petroleum Geology of Northwest Europe: Proceedings of the 5th Conference.* Geological Society of London, 757–770.
- DAVYDOV, V.I., WARDLAW, B.R., & GRADSTEIN, F.M. 2004. The Carboniferous Period. *In: GRADSTEIN, F.M., OGG, J.G., & SMITH, A.G. (eds). A geologic time scale 2004.* Cambridge University Press, 222–248.
- DAVYDOV, V.I., CROWLEY, J.L., SCHMITZ, M.D., & POLETAEV, V.I. 2010. High-precision U-Pb zircon age calibration of the global Carboniferous time scale and Milankovitch band cyclicity in the Donets Basin, eastern Ukraine. *Geochemistry Geophysics Geosystems*, **11**, Q0AA04, doi:10.1029/2009GC002736.
- DUNHAM, K.C. 1990. *Geology of the Northern Pennine Orefield, Volume 1: Tyne to Stainmore.* Economic Memoir of the British Geological Survey, Sheets 40, 41 and 50. (London: HMSO.)
- FIELDING, C.R., FRANK, T.D., BIRGENHEIER, L.P., RYDEL, M.C., JONES, A.T., & ROBERTS, J. 2008. Stratigraphic imprint of the Late Palaeozoic Ice Age in eastern Australia: a record of alternating glacial and nonglacial climate regime. *Journal of the Geological Society*, **165**, 129–140.
- GASTALDO, R.A., PURKYŇÁ, E., ŠIMŮNEK, Z. & SCHMITZ, M.D. 2009. Ecological persistence in the Late Mississippian (Serukhovian, Namurian A) megafloral record of the Upper Silesian Basin, Czech Republic. *Palaios*, **24**, 336–350. DOI: 10.2110/palo.2008.p08-084r.
- GONZALEZ-BONORINO, G. & EYLES, N. 1995. Inverse relation between ice extent and the late Paleozoic glacial record of Gondwana. *Geology*, **23**, 1015–1018.
- GRADSTEIN, F.M., OGG, J.G., & SMITH, A.G. 2004. *A Geologic Time Scale.* Episodes, **19**, 3–4. (Cambridge University Press).
- GUION, P.D., FULTON, I.M. & JONES, N.S. 1995. Sedimentary facies of the coal-bearing Westphalian A and B north of the Wales-Brabant High. *In: WHATELEY, M.K.G. & SPEARS, D.A. (eds.). European Coal Geology.* Geological Society Special Publication, **82**, 45–78.
- GULBRANSON, E.L., MONTAÑEZ, I.P., SCHMITZ, M.D., LIMARINO, C.O., ISBELL, J.L., MARENSSI, S.A. & CROWLEY, J.L. 2010. High-precision U-Pb calibration of Carboniferous glaciation

- and climate history, Paganzo Group, NW Argentina. *Geological Society of America Bulletin*, **122**, 1480–1498. doi: 10.1130/B30025.1.
- HAMPSON, G.J. 1997. A sequence stratigraphic model for deposition of the Lower Kinderscout Delta, an Upper Carboniferous turbidite-fronted delta. *Proceedings of the Yorkshire Geological Society*, **51**, 273–296.
- HAMPSON, G.J., ELLIOTT, T. & DAVIES, S.J. 1997. The application of sequence stratigraphy to upper Carboniferous fluvio-deltaic strata of the onshore UK and Ireland: Implications for the southern North Sea. *Journal of the Geological Society, London*, **154**, 719–733.
- HAMPSON, G.J., DAVIES, S.J., ELLIOTT, T., FLINT, S.S. & STOLLHOFEN, H. 1999. Incised valley fill sandstone bodies in Upper Carboniferous fluvio-deltaic strata: recognition and reservoir characterization of Southern North Sea analogues. In: FLEET, A.J. & BOLDY, S.A.R. (eds). *Petroleum Geology of Northwest Europe: Proceedings of the 5th Conference*. Geological Society of London, 771–788.
- HESS, J.C. & LIPPOLT, H.J. 1986.  $^{40}\text{Ar}/^{39}\text{Ar}$  ages of tonstein and tuff sanidines: new calibration points for the improvement of the Upper Carboniferous time scale. *Chemical Geology*, **59**, 143–154.
- HOLDSWORTH, B.K. & COLLINSON, J.D. 1988. Millstone Grit cyclicity revisited. In: BESLY, B.M. & KELLING, G. (eds). *Sedimentation in synorogenic basin complex; the Upper Carboniferous of North West Europe*. Blackie, Glasgow and London, 132–152.
- IRELAND, T.R. & WILLIAMS, I.S., 2003, Considerations in Zircon Geochronology by SIMS. *Reviews in Mineralogy and Geochemistry*, **53**, 215–241.
- ISBELL, J.L., MILLER, M.F., WOLFE, K.L. & LENAHER, P.A., 2003, Timing of late Paleozoic glaciation in Gondwana: Was glaciation responsible for the development of Northern Hemisphere cyclothems? In: CHAN M.A. & ARCHER A.W. (eds). *Extreme Depositional Environments: Mega End Members in Geologic Time*. Geological Society of America Special Paper **370**, 5–24.
- JAFFEY, A.H., FLYNN, K.F., GLENDENIN, L.E., BENTLEY, W.C. & ESSLING, A.M. 1971. Precision Measurement of Half-Lives and Specific Activities of  $^{235}\text{U}$  and  $^{238}\text{U}$ . *Physical Review C*, **4**, 1889–1906.

- 657 JONES, C.M. & CHISHOLM, J.I. 1997. The Roaches and Ashover Grits: sequence stratigraphic  
658 interpretation of a 'turbidite fronted delta' system. *Geological Journal*, **32**, 45–68.
- 659 KRYZA, R, MUSZER, J, HAYDUKIEWICZ, J, AUGUST, C, JURASIK, M, & RODIONOV, N. 2010. A  
660 SIMS zircon age for a biostratigraphically dated Upper Viséan (Asbian) bentonite in the  
661 Central-European Variscides (Bardo Unit, Polish Sudetes). *International Journal of Earth*  
662 *Sciences*, DOI: 10.1007/s00531-010-0529-y.
- 663 LAKE, R.D. 1999. *The Wakefield district - a concise account of the geology*. Memoir of the  
664 British Geological Survey. Sheet 78 (England and Wales).
- 665 LEEDER, M.R. 1982. Upper Palaeozoic basins of the British Isles: Caledonide inheritance versus  
666 Hercynian plate margin processes. *Journal of the Geological Society London*, **139**, 481–  
667 494.
- 668 MARTINSEN, O.J. 1993. Namurian (late Carboniferous) depositional systems of the Craven-  
669 Askrigg area, northern England: implications for sequence stratigraphic models. In:  
670 POSAMENTIER, H.W., SUMMERHAYES, C.P., HAQ, B.U. & ALLEN, G.P. (eds) *Stratigraphy*  
671 *and Facies associations in a Sequence Stratigraphic Framework*, 247–281.
- 672 MARTINSEN, O.J., COLLINSON, J.D. & HOLDSWORTH, B.K. 1995. Millstone Grit cyclicity revisited,  
673 II: sequence stratigraphy and sedimentary responses to changes of relative sea-level. In:  
674 PLINT, A.G. (ed.) *Sedimentary facies analysis. International Association of Sedimentologists*  
675 *Special Publication*, **22**, 305–327. Blackwell Scientific, Oxford.
- 676 MATTINSON, J.M. 2005. Zircon U-Pb chemical abrasion ("CA-TIMS") method: Combined annealing  
677 and multi-step partial dissolution analysis for improved precision and accuracy of zircon  
678 ages. *Chemical Geology*, **220**, 47–66.
- 679 MAYNARD, J.R. & LEEDER, M.R. 1992. On the periodicity and magnitude of Late Carboniferous  
680 glacio-eustatic sea-level changes. *Journal of the Geological Society London*, **149**, 303–  
681 311.
- 682 OGG, J.G., OGG, G., and GRADSTEIN, F.M. 2008. *The Concise Geologic Time Scale*. Cambridge  
683 University Press.
- 684 OWENS, B., VARKER, W.J., & HUGHES, R.A. 1990. Lateral biostratigraphical consistency across  
685 the Mid-Carboniferous boundary in northern England. *Courier Forschungsinstitut*  
686 *Senckenberg*, **130**, 237–244.
- 687 POSAMENTIER, H.W., JERVEY, M.T. & VAIL, P.R. 1988. Eustatic controls on clastic deposition  
688 1—conceptual framework. In: WILGNUS, C.K., HASTINGS, C.G. ST C., KENDALL, H.W.,

- 689 POSAMENTIER, H.W., ROSS, C.A. & VAN WAGONER, J.C. (eds). *Sea-level changes: an*  
690 *integrated approach*. Society of Economic Paleontologists and Mineralogists Special  
691 Publication, **42**, 39–45.
- 692 RAMSBOTTOM, W.H.C. 1977. Major cycles of transgression and regression (mesothems) in the  
693 Namurian. *Proceedings of the Yorkshire Geological Society*, **41**, 261–291.
- 694 RAMSBOTTOM, W.H.C. 1979. Rates of transgression and regression in the Carboniferous of NW  
695 Europe. *Quarterly Journal of the Geological Society of London*, **136**, 147–154.
- 696 RILEY, N.J., CLAOUÉ-LONG, J.C., HIGGINS, A.C., OWENS, B., SPEARS, A., TAYLOR, L., &  
697 VARKER, W.J. 1995. Geochronometry and geochemistry of the European mid-  
698 Carboniferous boundary global stratotype proposal, Stonehead Beck, North Yorkshire,  
699 UK. *Annales de la Société géologique de Belgique*, **116**, 275–289.
- 700 RIPPON, J.H. 1996. Sand body orientation, palaeoslope analysis and basin-fill implications in the  
701 Westphalian A-C of Great Britain. *Journal of the Geological Society, London*, **153**, 881–  
702 900.
- 703 ROBERTS, J., CLAOUÉ-LONG, J.C., JONES, P.J. & FOSTER, C.B. 1995. SHRIMP zircon age control  
704 of Gondwanan sequences in Late Carboniferous and Early Permian Australia. *In*: DUNAY,  
705 R.E. & HAILWOOD, E.A. (eds), *Non-biostratigraphical Methods of Dating and Correlation*.  
706 Geological Society Special Publication, **89**, 145–174.
- 707 SPEARS, D.A. & KANARIS-SOTIRIOU, R. 1979. A geochemical and mineralogical investigation of  
708 some British and other European tonsteins. *Sedimentology*, **26**, 407–425.
- 709 SPEARS, D.A., KANARIS-SOTIRIOU, R., RILEY, N.J. & KRAUSE, P. 1999. Namurian bentonites in  
710 the Pennine Basin, UK - origin and magmatic affinities. *Sedimentology*, **46**, 385–401.
- 711 STEPHENSON, M.H., ANGIOLINI, L., CÓZAR, P., JADOUL, F., LENG, M.J., MILLWARD, D. &  
712 CHENERY, S. 2010. Northern England Serpukhovian (early Namurian) farfield responses to  
713 southern hemisphere glaciation. *Journal of the Geological Society*, **167**, 1171–1184.
- 714 TRAPP, E. & KAUFMANN, B. 2002. Hochpräzise U–Pb Datierungen von Paroklastika im  
715 Jungpaläozoikum. *Neustadt: Schwerpunktprogram der deutschen Forschungsgemeinschaft*  
716 DFG (SPP 1054), 18–19.
- 717 TREWIN, N.H. 1968. Potassium bentonites in the Namurian of Staffordshire and Derbyshire.  
718 *Proceedings of the Yorkshire Geological Society*, **37**, 73–91.

- TUCKER, M.E., GALLAGHER, J., & LENG, M.J. 2009. Are beds in shelf carbonates millennial-scale cycles? An example from the mid-Carboniferous of northern England. *Sedimentary Geology*, **214**, 19–34.
- VEEVERS, J.J. & POWELL, C.M. 1987. Late Paleozoic glacial episodes in Gondwanaland reflected in Transgressive-Regressive depositional sequences in Euramerica. *Geological Society of America Bulletin*, **98**, 475–487.
- WATERS, C.N., SOMERVILLE, I.D., JONES, N.S. & 16 others 2011. *A revised Correlation of Carboniferous Rocks in the British Isles*. Geological Society Special Report No. 26, London.
- WATERS, C.N., CHISHOLM, J.I., BENFIELD, A.C., & O’BEIRNE, A.M. 2008. Regional evolution of a fluviodeltaic cyclic succession in the Marsdenian (Late Namurian Stage, Pennsylvanian) of the Central Pennine Basin, UK. . *Proceedings of the Yorkshire Geological Society*, **57**, 1–28.
- WATERS, C.N. & DAVIES, S.J. 2006. Carboniferous extensional basins, advancing deltas and coal swamps. In: BRENCHLEY, P.J. & RAWSON, P.F. (eds) *The Geology of England and Wales*. The Geological Society London (Second edition), 173–223.
- WATERS, C.N., GLOVER, B.W. & POWELL, J.H. 1994. Structural synthesis of S Staffordshire, UK: implications for the Variscan evolution of the Pennine Basin. *Journal of the Geological Society London*, **151**, 697–713.
- WIGNALL, P.B. 1987. A biofacies analysis of the *Gastrioceras cumbriense* Marine Band (Namurian) of the central Pennines. *Proceedings of the Yorkshire Geological Society*, **46**, 111–121.



## Figure captions

**Fig. 1.** Map showing the approximate extent of Namurian and Westphalian strata at crop, the subcrop of Westphalian strata and the main pre-Namurian structural features of the Pennine Basin, derived from Waters *et al.* (2011). The location of the Harewood and Holme Pierrepont boreholes, from which new dates have been acquired during this study, are shown. DH- Derbyshire High, WG- Widmerpool Gulf.

**Fig. 2.** U-Pb data for samples BLL1976 and EH28155. A, conventional U-Pb concordia plot of zircons analysed from samples BLL1976 and EH28155. The grey band reflects the uncertainty in the  $^{238}\text{U}$  and  $^{235}\text{U}$  decay constants (Jaffey *et al.* 1971). B, plot of  $^{238}\text{U}/^{206}\text{Pb}$  dates for single zircon crystals analyses (same data as in Figure 2a). Dashed ellipses/bars represent analyses of zircon that are considered to be xenocrysts and/or inherited crystals that are disregarded in calculation of final date, whereas as undashed ellipses/bars represent the analyses used for calculation of the weighted mean final date (see text for discussion). Data point error ellipses/bars are  $2\sigma$ .

**Fig. 3.** Distribution of ammonoid acme facies in early Namurian marine bands: a) Pendleian; b) early Arnsbergian; c) mid to late Arnsbergian; d) Chokierian; e) Alportian; f) early Kinderscoutian. Grey tone denotes marine band with maximum areal extent for each interval. Key for Incised Valleys, as for Figure 5.

**Fig. 4.** Distribution of ammonoid acme facies in late Namurian–Westphalian marine bands: a) mid to late Kinderscoutian; b) early Marsdenian; c) late Marsdenian; d) Yeadonian-  $G_{1b1}$  modified from Wignall (1987); e) Langsettian and f) Duckmantian–Bolsovian, in part based upon Calver (1968, 1969). Grey tone denotes marine band with maximum areal extent for each interval. Key for Incised Valleys, as for Figure 5.

**Fig. 5.** Magnitude and duration of sea-level oscillations. Sea-level maxima are estimated through the determination of maximum areal extent of acme ammonoid facies. Abbreviations for Westphalian marine bands: SMB Subcrenatum Marine Band; LMB Listeri Marine Band; AmMB Amaliae Marine Band; VMB Vanderbecke Marine Band; HMB Haughton Marine Band; AMB Aegiranum Marine Band; CMB Cambriense Marine Band. For marine bands lacking ammonoid fauna the magnitude of sea-level is determined through the acme marine fauna, ranging from *Estheria* to brachiopod-bivalve facies. Sea-level minima are determined through the presence of incised valleys, with the magnitude recorded by the number of underlying marine bands removed beneath the sequence boundary. Sources for incised valleys are as follows: a) Rogerley Channel (Dunham 1990); b) Upper Howgate Edge channel (Martinsen *et al.* 1995); c) Red Scar Grit

775 (Brandon *et al.* 1995); d) Lower Follifoot Grit (Martinsen 1993); e) Intra-H<sub>1a</sub> unconformity  
776 (Owens *et al.* 1990); f) Upper Follifoot Grit (Martinsen 1993); g) Todmorden Grit/Kinderscout  
777 Grit (Hampson 1997); h) Upper Kinderscout Grit (Hampson 1997); i) Midgley Grit (Brettle  
778 2001); j) Ashover Grit/Roaches Grit (Jones & Chisholm 1997; Church & Gawthorpe 1994); k)  
779 Chatsworth Grit (Waters *et al.* 2008); l) Rough Rock (Church & Gawthorpe 1994); m)  
780 Crawshaw Sandstone (Hampson *et al.* 1997); n) Thornhill Rock (Lake 1999); o) Woolley Edge  
781 Rock (Aitken *et al.* 1999); p) Oaks Rock (Lake 1999); q) Mexborough Rock (Aitken *et al.*  
782 1999). Radiometric dates are from this study and estimated ages of stage boundaries are from  
783 Davydov *et al.* (2010), with an imposed 400 ka long-duration eccentricity oscillation numbered  
784 sequentially for each international stage: S Serpukhovian; B Bashkirian; M Moscovian. The  
785 proposed four main glaciations are highlighted as grey bands.

**Nature and timing of Late Mississippian to Mid Pennsylvanian glacio-eustatic sea-level changes of the Pennine Basin, UK**

Colin N. Waters<sup>1\*</sup> and Daniel J. Condon<sup>2</sup>

<sup>1</sup>British Geological Survey, Kingsley Dunham Centre, Keyworth, Nottingham, NG12 5GG

<sup>2</sup>NERC Isotope Geoscience Laboratory, British Geological Survey, Kingsley Dunham Centre, Keyworth, Nottingham, NG12 5GG

\* E-mail: cnw@bgs.ac.uk

**Online Supplemental Information**

**Zircon U-Pb ID-TIMS Methods**

Zircons were isolated from ca. 100 grams of bentonite layer (sample BLL1976) from the BGS Harewood Borehole section and ca. 100 grams of bentonite layer (sample EH28155) from the Holme Pierrepont Borehole section, using conventional mineral separation techniques. Prior to isotope dilution thermal ionization mass spectrometry (ID-TIMS) analyses zircons were subject to a modified version of the chemical abrasion technique (Mattinson 2005). For details of sample pre-treatment, dissolution and anion exchange chemistry see Sláma *et al.* (2008). U-Pb ID-TIMS analyses herein utilized the EARTHTIME <sup>205</sup>Pb-<sup>233</sup>U-<sup>235</sup>U (ET535) tracer solution. Measurements at the NERC Isotope Geosciences Laboratory were performed on a Thermo Triton TIMS. Pb analyses were measured in dynamic mode on a MassCom SEM detector and corrected for 0.14 ±0.04%/u. mass fractionation. Linearity and dead-time corrections on the SEM were monitored using repeated analyses of NBS 982, NBS 981 and U500. Uranium was measured in static Faraday mode on 10<sup>11</sup> ohm resistors or for signal intensities < 15 mV, in dynamic mode on the SEM detector. Uranium was run as the oxide and corrected for isobaric interferences with an <sup>18</sup>O/<sup>16</sup>O composition of 0.00205 (IUPAC value and determined through direct measurement at NIGL). U-Pb dates and uncertainties were calculated using the algorithms of Schmitz & Schoene (2007) and a <sup>235</sup>U/<sup>205</sup>Pb ratio for ET535 of 100.18 ±0.1%. All common Pb in the analyses was attributed to the blank and subtracted based on the isotopic composition and associated uncertainties analysed over time. The <sup>206</sup>Pb/<sup>238</sup>U ratios and dates were corrected

for initial  $^{230}\text{Th}$  disequilibrium using a Th/U[magma] of  $4 \pm 1$  applying the algorithms of Schärer (1984) resulting in an increase in the  $^{206}\text{Pb}/^{238}\text{U}$  dates of  $\sim 100$  kyr. Errors for U-Pb dates are reported in the following format:  $\pm X(Y)[Z]$ , where X is the internal or analytical uncertainty in the absence of all systematic error (tracer calibration and decay constants), Y includes the quadratic addition of tracer calibration error (using a conservative estimate of the  $2\sigma$  standard deviation of 0.1% for the Pb/U ratio in the tracer), and Z includes the quadratic addition of both the tracer calibration error and additional  $^{238}\text{U}$  decay constant errors of Jaffey *et al.* (1971). All analytical uncertainties are calculated at the 95% confidence interval. These  $^{238}\text{U}/^{206}\text{Pb}$  dates are traceable back to SI units via the gravimetric calibration of the EARTHTIME U-Pb tracer and the determination of the  $^{238}\text{U}$  decay constant (Condon *et al.* 2007; Jaffey *et al.* 1971). The results for the analyses of samples BLL1976 and EH28155 are shown in Table 1.

### Zircon U-Pb ID-TIMS Results and Discussion

Zircons separated from both bentonite samples BLL1976 and EH28155 were small ( $< 50 \mu\text{m}$ ) with aspect ratios of  $\sim 1.5$  to  $\sim 3$ . For sample BLL1976 seventeen fractions (single grains) were analysed. Relatively large uncertainties on  $^{207}\text{Pb}/^{206}\text{Pb}$  dates ( $\sim 5$  to  $50$  Ma) limit the usefulness of concordance for assessing the accuracy of  $^{206}\text{Pb}/^{238}\text{U}$  and the potential for subtle Pb-loss and/or incorporation of minor amounts of older material.  $^{206}\text{Pb}/^{238}\text{U}$  dates for sample BLL1976 between 311 and 334 Ma with a distinct population (defined by 11 of the 17) of concordant analyses yielding a weighted mean  $^{206}\text{Pb}/^{238}\text{U}$  date of  $328.34 \pm 0.30(0.43)[0.55]$  Ma (Mean square weighted deviation (95% confidence level, MSWD = 2.2).  $^{206}\text{Pb}/^{238}\text{U}$  dates that are older than the main population are interpreted as reflecting the analyses of zircon ante-/xeno-crysts, and grains that are younger as reflecting Pb-loss (see below). The MSWD for this sample exceeds that expected for a single population (Wendt & Carl 1991) indicating excess scatter. There are two possible explanations for this scatter: (1) residual Pb-loss in younger zircons (e.g., z6, z15 and z12); or (2) older zircon U/Pb dates (e.g., z4, z19, z22, zA2) reflect pre-eruptive crystallisation of zircon (i.e., ante-/xeno-crysts). The alternative interpretations, based upon statistically coherent populations, result in weighted mean  $^{206}\text{Pb}/^{238}\text{U}$  dates that are  $\sim 200$  ka older/younger than the weighted mean  $^{206}\text{Pb}/^{238}\text{U}$  date of  $328.34 \pm 0.30$  Ma, in part as the 95% confidence error of the weighted average multiplies the ' $2\sigma$  internal' error by the square root of the MSWD therefore using the actual scatter of the data rather than the predicted scatter based upon the main population

(Ludwig 1991). Thus, in the absence of any independent criteria for the exclusion of data points from the main population we interpret the weighted mean  $^{206}\text{Pb}/^{238}\text{U}$  date of  $328.34 \pm 0.30(0.43)[0.55]$  Ma as being the best estimate for the zircons of this sample and inferentially the age of bentonite at the sampled stratigraphic level. For sample EH28155 nine fractions (single grains) were analysed, and the resulting data are presented in Figure 2 of the paper. Two of the nine analyses produced U-Pb dates older than the constraint imposed by sample BLL 1976 (see above). The remaining seven analyses yielded  $^{206}\text{Pb}/^{238}\text{U}$  dates between 306 and 317 Ma with a distinct population (defined by 4 of the 7) of analyses yielding a weighted mean  $^{206}\text{Pb}/^{238}\text{U}$  date of  $314.37 \pm 0.25(0.40)[0.53]$  Ma (MSWD = 1.07) which is interpreted as being the best estimate for the zircons of this sample. In both samples the U/Pb dates that are older than the main population are interpreted as reflecting the analyses of zircon antecrysts, and grains that are younger as reflecting Pb-loss. This interpretation is supported by consideration of biostratigraphical and geochronological constraints (e.g., Davydov *et al.* 2010) for this time interval and is typical for moderate-*n* high-precision U-Pb (zircon) data sets obtained on primary air-fall ash beds (e.g., Davydov *et al.*, 2010; Schoene *et al.* 2010).

### **Ammonoid biostratigraphy**

Ammonoid biozones in the Namurian succession are defined by the first appearance of ammonoid taxa, with the base of the biozones coinciding with the bases of specific marine bands. Ammonoid evolution rates reached an acme during the Namurian, such that the majority of marine bands comprise a distinct ammonoid fauna. The Namurian ammonoid biostratigraphy was developed largely from studies within the Central Pennine Basin by Bisat (1924; 1928) and Bisat & Hudson (1943), and later refined by Ramsbottom (1969; 1971). The scheme used for the Namurian in this study is derived from Riley *et al.* (1995). An ammonoid biozonation scheme has not been developed for the Westphalian, but key ammonoid-bearing marine bands have been used as marker beds to divide the cyclical Coal Measures lithofacies. The key marine bands, identified by Ramsbottom *et al.* (1978), are named after the diagnostic ammonoid species, replacing a plethora of local geographical names used historically. Correlation in the Westphalian is enhanced by the presence of an additional framework of marker marine bands that contain less diagnostic fauna, such as thin-shelled ammonoids (*Anthracoeras*), marine bivalves (*Dunbarella*), brachiopods (*Lingula*)

and crustacea (Ostracodes, *Estheria*). Few such marine marker bands, entirely devoid of ammonoid fauna across the basin, are recognized in the Namurian succession.

#### **Calculation of areal extent of marine bands**

Ammonoid-bearing marine band locations, either from surface exposures or borehole samples, were determined by a comprehensive trawl of BGS palaeontological registers, BGS memoirs and technical reports and scientific publications, too many to reference individually. Marine band locations were used in the study only where ammonoids were described, representing the acme fauna, and for which the marine band attribution was unequivocal. It was decided that the recognition of marine faunal facies within the Namurian succession, as carried out for the Westphalian by Calver (1968, 1969), was beyond the scope of this study.

Table 2 shows the location and ammonoid assemblages used to compile the distribution maps. The described location is that given in registers or publications, presented without modification to show the basis by which locations were determined. Borehole locations and logs, available from the BGS Single Onshore Borehole Index (SOBI) database, were used to check the attribution of the marine bands. The British National Grid of surface locations was determined in the majority of cases by comparing the description of the locality to BGS fieldslips and published 1:10,560- or 1:10,000-scale maps which showed the location of the fossil locality. This also provided a means of confirming or determining from which marine band the sample was collected. If marine band identification was equivocal it was not included in the study. This is in particular relevant to marine bands present in great thicknesses of mudstone, for example in the Bowland Shale Formation, where correlation of distinct marine bands can be more uncertain. As the identification of marine band nomenclature is dependent on recognition of the first incoming of certain taxa, it can be difficult to confirm the first appearance unless a sequential succession of several marine bands are studied in a single section, which is relatively unusual in surface sections. Within the Millstone Grit Group, the cyclic nature of sedimentation means that marine bands commonly rest upon sandstones, with significant thicknesses between flooding events. This makes correlation of individual marine bands over large distances comparatively easy, with the relationship of specific marine bands to named sandstones well established. As a consequence of this study it has been possible to compile typical ammonoid assemblage lists for each marine band, something not previously published systematically. In Table 2 the faunal list is that which appears in the source register or literature. A modern reinterpretation

of these faunal lists appears in Tables 2 to 3 in the publication. However, it is important to realize that none of the specimens, many of which will be stored at BGS Keyworth, were examined during this study to confirm the original identifications.

The distribution of the ammonoid facies in the Westphalian succession were previously constrained by Calver (1968, 1969) and these envelopes were used in this study, with modification where samples listed in Table 2 indicated the need to extend the extent of the envelope. In many cases this resulted from borehole data from the eastern part of the basin, acquired after Calver's studies.

For each marine band, the position of the ammonoid localities were presented using ArcGIS software, an envelope was drawn around the locations and the areal extent of the envelope calculated. To some extent, an understanding of the geology of the basin was required to determine the extent that the area between locations could be assumed to still be ammonoid-bearing. In particular, envelopes were not extrapolated across areas where no data were available, either through the marine band being too deeply buried to be proved by boreholes, or absent through removal by erosion.

## Production of the cyclostratigraphical model

The regional chronostratigraphy was initially scaled against absolute time using the two dates acquired during this study and the additional constraints for the age of the base of the Serpukhovian and Bashkirian, derived from Davydov *et al.* (2010). By dividing each of the age ranges (base Pendleian to early Arnbergian, early Arnbergian to Chokierian and Chokierian to early Bolsovian) by the number of recognized flooding surfaces within each interval it was possible to recognize the average cycle duration. However, consideration of just peak flooding events and Type 1 unconformities suggested that a ~400 kyr eccentricity frequency persisted throughout the study interval, a duration consistent with findings from other international studies. On this basis the Serpukhovian, Bashkirian and lower part of the Moscovian stages were subdivided into ~400 kyr cycles, numbered S1 to S17, B1 to B20 and M1 to M3, respectively. Peak flooding events were then aligned to each of the cycles. Where this proved not possible for the Mid to Late Arnbergian succession because of too few flooding events, it was decided to take an arbitrary position of spacing equally the E<sub>2b</sub>3 and E<sub>2c</sub>1 flooding events.

The magnitude of sea-level oscillations is presented using three distinct scales. The areal extent of ammonoid facies is presented as a linear scale of 1000's Km<sup>2</sup>. Non-ammonoid bearing flooding events were incorporated into the analysis by indicating the acme marine facies described within the Pennine Basin. The order presented of *Estheria*, Foraminifera, *Lingula* and brachiopod-bivalve facies, indicates increasing marine influence (Calver 1968). The absolute magnitude of sea-level fall associated with many of the Type 1 unconformities is unrecorded, whereas the number of cycles removed by erosion was determinable from publications describing these erosional surfaces. Again, this does not provide absolute magnitudes of sea-level falls, but is indicative of the relative significance of these events. It is important to realize that the resultant figure, using three different scales provides a means to visualize relative magnitudes of sea-level fluctuation but cannot be used to an oscillatory sea-level curve.



## References

- BISAT, W.S. 1924. The Carboniferous goniatites of the north of England and their zones. *Proceedings of the Yorkshire Geological Society*, **20**, 40-124.
- BISAT, W.S. 1928. The Carboniferous goniatite zones of England and their continental equivalents. *Congrès pour l'Avancement des Etudes de Stratigraphie Carbonifère, Compte Rendu, Heerlen 1927*, 117-133.
- BISAT, W.S. & HUDSON, R.G.S. 1943. The Lower Reticuloceras (R1) Goniatite succession in the Namurian of the north of England. *Proceedings of the Yorkshire Geological Society*, **24**, 383-440.
- CALVER, M.A. 1968. Distribution of Westphalian marine faunas in northern England and adjoining areas. *Proceedings of the Yorkshire Geological Society*, **37**, 1-72.
- CALVER, M.A. 1969. Westphalian of Britain. *Compte Rendu 6e Congrès Internationale Stratigraphie et Géologie Carbonifère, Sheffield, 1967*, **I**, 233-254.
- CONDON, D., SCHOENE, B., BOWRING, S., PARRISH, R., MCLEAN, N., NOBLE, S. & CROWLEY, Q. 2007. EARTHTIME; isotopic tracers and optimized solutions for high-precision U-Pb ID-TIMS geochronology. *Eos, Transactions, American Geophysical Union*, **88**.
- CROWLEY, J.L., SCHOENE, B. & BOWRING, S.A. 2007. U-Pb Dating of Zircon in the Bishop Tuff at the Millennial Scale. *Geology*, **35**, 1123-1126.
- DAVYDOV, V.I., CROWLEY, J.L., SCHMITZ, M.D., & POLETAEV, V.I. 2010. High-precision U-Pb zircon age calibration of the global Carboniferous time scale and Milankovitch band cyclicity in the Donets Basin, eastern Ukraine. *Geochemistry Geophysics Geosystems*, **11**, Q0AA04, doi:10.1029/2009GC002736.
- JAFFEY, A.H., FLYNN, K.F., GLENDENIN, L.E., BENTLEY, W.C. & ESSLING, A.M. 1971. Precision Measurement of Half-Lives and Specific Activities of  $^{235}\text{U}$  and  $^{238}\text{U}$ . *Physical Review C*, **4**, 1889-1906.
- LUDWIG, K.R. 1991. Isoplot - a plotting and regression program for radiogenic isotope data: *USGS Open File Report*, 91-445
- MATTINSON, J M. 2005. Zircon U-Pb chemical abrasion ("CA-TIMS") method: Combined annealing and multi-step partial dissolution analysis for improved precision and accuracy of zircon ages. *Chemical Geology*, **220**, 47-66.

- RAMSBOTTOM, W.H.C. 1969. The Namurian of Britain. *Compte Rendue 6 ème Congrès International de Stratigraphie et de Geologie du Carbonifère, Sheffield, 1967.*
- RAMSBOTTOM, W.H.C. 1971. Palaeogeography and goniatite palaeogeography in the Namurian and early Westphalian. *Compte Rendue, 6 ème Congrès International de Stratigraphie et de Geologie du Carbonifère, Sheffield, 1967*, 1396-1399.
- RAMSBOTTOM, W.H.C., CALVER, M.A., EAGAR, R.M.C., HODSON, F., HOLLIDAY, D.W., STUBBLEFIELD, C.J. & WILSON, R.B. 1978. A correlation of Silesian rocks in the British Isles. *Special Report of the Geological Society of London*, **10**.
- RILEY, N.J., CLAOUÉ-LONG, J.C., HIGGINS, A.C., OWENS, B., SPEARS, A., TAYLOR, L., & VARKER, W.J. 1995. Geochronometry and geochemistry of the European mid-Carboniferous boundary global stratotype proposal, Stonehead Beck, North Yorkshire, UK. *Annales de la Société géologique de Belgique*, **116**, 275-289.
- SCHÄRER, U., 1984, The effect of initial  $^{230}\text{Th}$  disequilibrium on young U–Pb ages: the Makalu case, Himalaya. *Earth and Planetary Science Letters*, **67** (2), 191-204.
- SCHOENE, B., GUEX, J., BARTOLINI, A., SCHALTEGGER, U., & BLACKBURN, T.J., 2010, Correlating the end-Triassic mass extinction and flood basalt volcanism at the 100 ka level. *Geology*, **38**, 387-390.
- SCHMITZ, M.D. & SCHOENE, B. 2007. Derivation of isotope ratios, errors, and error correlations for U-Pb geochronology using  $^{205}\text{Pb}$ – $^{235}\text{U}$ –( $^{233}\text{U}$ )-spiked isotope dilution thermal ionization mass spectrometric data. *Geochemistry Geophysics Geosystems*, **8**, 20pp.
- SLÁMA, J., KOSLER, J., CONDON, D.J., CROWLEY, J.L., GERDES, A., HANCHAR, J.M., HORSTWOOD, M.S. A., MORRIS, G.A., NASDALA, L., NORBERG, N., SCHALTEGGER, U., SCHOENE, B., TUBRETT, M. N., & WHITEHOUSE, M. J. 2008. Plesovice zircon; a new natural reference material for U-Pb and Hf isotopic microanalysis. *Chemical Geology*, **249**, 1-35.
- WENDT, I. & CARL, C. 1991. The statistical distribution of the mean squared weighted deviation. *Chemical Geology, Isotope Geoscience section*, **86**, 275-285

**Table 1.** U-Th-Pb isotopic data

| Sample<br>(a)   | Compositional Parameters |   |                                     |                               |                                |   | Radiogenic Isotope Ratios                     |   |              |  |              |  |              |                       | Isotopic Ages                                 |          |  |          |   |             |
|-----------------|--------------------------|---|-------------------------------------|-------------------------------|--------------------------------|---|---|---|--------------|--|--------------|--|--------------|-----------------------|---|----------|--|----------|---|-------------|
|                 | Th<br>U<br>(b)           | $^{206}\text{Pb}^*$<br>$\times 10^{-13}$ mol<br>(c) | mol %<br>$^{206}\text{Pb}^*$<br>(c) | Pb*<br>Pb <sub>c</sub><br>(c) | Pb <sub>c</sub><br>(pg)<br>(c) | $^{206}\text{Pb}$<br>$^{204}\text{Pb}$<br>(d) | $^{208}\text{Pb}$<br>$^{206}\text{Pb}$<br>(e) | $^{207}\text{Pb}$<br>$^{206}\text{Pb}$<br>(e) | % err<br>(f) | $^{207}\text{Pb}$<br>$^{235}\text{U}$<br>(e) | % err<br>(f) | $^{206}\text{Pb}$<br>$^{238}\text{U}$<br>(e) | % err<br>(f) | corr.<br>coef.<br>(f) | $^{207}\text{Pb}$<br>$^{206}\text{Pb}$<br>(g) | ±<br>(f) | $^{207}\text{Pb}$<br>$^{235}\text{U}$<br>(g) | ±<br>(f) | $^{206}\text{Pb}$<br>$^{238}\text{U}$<br>(g, h) | ±<br>(f)    |
| <b>EH-28155</b> |                          |   |                                     |                               |                                |   |   |   |              |  |              |  |              |                       |   |          |  |          |   |             |
| z2              | 0.407                    | 0.4794  | 98.68%                              | 22                            | 0.53                           | 1378  | 0.128   | 0.05276                                       | 0.27         | 0.3666                                       | 0.34         | 0.05040                                      | 0.10         | 0.740                 | 317.6   | 6.2      | 317.14                                       | 0.92     | 317.07  | 0.31        |
| z3              | 0.379                    | 0.4504  | 98.43%                              | 18                            | 0.60                           | 1159  | 0.120   | 0.05271                                       | 0.66         | 0.3634                                       | 0.70         | 0.05000                                      | 0.17         | 0.315                 | 315.7   | 15.1     | 314.73                                       | 1.88     | <b>314.60</b>                                   | <b>0.52</b> |
| z4              | 0.213                    | 0.3503  | 97.99%                              | 14                            | 0.60                           | 904   | 0.067   | 0.05254                                       | 0.38         | 0.3606                                       | 0.45         | 0.04977                                      | 0.12         | 0.650                 | 308.3   | 8.7      | 312.63                                       | 1.21     | 313.22  | 0.38        |
| z5              | 0.362                    | 0.5413  | 98.74%                              | 23                            | 0.57                           | 1446  | 0.115   | 0.05272                                       | 0.26         | 0.3629                                       | 0.33         | 0.04993                                      | 0.12         | 0.708                 | 316.0   | 6.0      | 314.39                                       | 0.90     | <b>314.17</b>                                   | <b>0.36</b> |
| z6              | 0.319                    | 0.1435  | 97.09%                              | 10                            | 0.36                           | 625   | 0.100   | 0.05287                                       | 0.62         | 0.3863                                       | 0.70         | 0.05300                                      | 0.18         | 0.545                 | 322.6   | 14.1     | 331.69                                       | 1.97     | 333.00  | 0.57        |
| zA6             | 0.357                    | 0.2455  | 95.99%                              | 7                             | 0.85                           | 454   | 0.112   | 0.05222                                       | 0.82         | 0.3601                                       | 0.91         | 0.05001                                      | 0.20         | 0.514                 | 294.5   | 18.7     | 312.30                                       | 2.43     | <b>314.69</b>                                   | <b>0.62</b> |
| zA7             | 0.470                    | 0.1401  | 90.22%                              | 3                             | 1.26                           | 186   | 0.145   | 0.05120                                       | 2.57         | 0.3525                                       | 2.72         | 0.04994                                      | 0.32         | 0.498                 | 249.2   | 59.2     | 306.63                                       | 7.20     | <b>314.23</b>                                   | <b>0.99</b> |
| zA8             | 0.345                    | 0.5371  | 91.46%                              | 3                             | 4.16                           | 213   | 0.109   | 0.05346                                       | 1.04         | 0.3958                                       | 1.14         | 0.05370                                      | 0.24         | 0.534                 | 347.7   | 23.4     | 338.59                                       | 3.29     | 337.27  | 0.78        |
| zA9             | 0.261                    | 0.1667  | 90.37%                              | 3                             | 1.47                           | 189   | 0.081   | 0.05171                                       | 1.85         | 0.3475                                       | 1.99         | 0.04874                                      | 0.20         | 0.704                 | 271.7   | 42.4     | 302.82                                       | 5.21     | 306.87  | 0.61        |
| <b>BLL 1976</b> |                          |   |                                     |                               |                                |   |   |   |              |  |              |  |              |                       |   |          |  |          |   |             |
| z2              | 1.464                    | 0.3195  | 98.10%                              | 20                            | 0.51                           | 960   | 0.463   | 0.05305                                       | 0.39         | 0.3821                                       | 0.45         | 0.05225                                      | 0.12         | 0.638                 | 330.4   | 8.8      | 328.61                                       | 1.28     | <b>328.35</b>                                   | <b>0.38</b> |
| z4              | 1.771                    | 0.1298  | 94.23%                              | 7                             | 0.66                           | 315   | 0.555   | 0.05255                                       | 1.40         | 0.3793                                       | 1.51         | 0.05235                                      | 0.21         | 0.564                 | 309.1   | 32.0     | 326.51                                       | 4.22     | <b>328.96</b>                                   | <b>0.67</b> |
| z5              | 1.482                    | 0.3720  | 94.55%                              | 7                             | 1.78                           | 334   | 0.463   | 0.05239                                       | 1.18         | 0.3759                                       | 1.28         | 0.05204                                      | 0.17         | 0.630                 | 302.0   | 26.9     | 324.01                                       | 3.55     | 327.08  | 0.54        |
| z6              | 1.457                    | 0.7789  | 98.34%                              | 22                            | 1.09                           | 1093  | 0.460   | 0.05296                                       | 0.36         | 0.3807                                       | 0.43         | 0.05213                                      | 0.13         | 0.618                 | 326.8   | 8.2      | 327.55                                       | 1.19     | <b>327.65</b>                                   | <b>0.42</b> |
| z12             | 1.542                    | 0.0831  | 94.72%                              | 7                             | 0.38                           | 345   | 0.488   | 0.05316                                       | 1.13         | 0.3826                                       | 1.24         | 0.05221                                      | 0.23         | 0.530                 | 335.3   | 25.7     | 328.98                                       | 3.48     | <b>328.10</b>                                   | <b>0.73</b> |
| z14             | 1.355                    | 0.0874  | 92.38%                              | 4                             | 0.60                           | 239   | 0.430   | 0.05326                                       | 1.67         | 0.3838                                       | 1.80         | 0.05227                                      | 0.26         | 0.552                 | 339.5   | 37.7     | 329.85                                       | 5.06     | <b>328.48</b>                                   | <b>0.84</b> |
| z15             | 1.319                    | 0.4603  | 99.04%                              | 38                            | 0.37                           | 1896  | 0.417   | 0.05300                                       | 0.24         | 0.3810                                       | 0.58         | 0.05214                                      | 0.51         | 0.915                 | 328.3   | 5.3      | 327.79                                       | 1.63     | <b>327.71</b>                                   | <b>1.63</b> |
| z16             | 0.104                    | 0.2838  | 98.46%                              | 17                            | 0.37                           | 1179  | 0.033   | 0.05302                                       | 0.33         | 0.3820                                       | 0.40         | 0.05226                                      | 0.12         | 0.651                 | 328.8   | 7.5      | 328.49                                       | 1.11     | <b>328.45</b>                                   | <b>0.39</b> |
| z17             | 1.267                    | 0.1359  | 93.68%                              | 5                             | 0.76                           | 288   | 0.398   | 0.05254                                       | 1.28         | 0.3748                                       | 1.39         | 0.05173                                      | 0.20         | 0.593                 | 308.8   | 29.1     | 323.19                                       | 3.84     | 325.19  | 0.65        |
| z18             | 1.447                    | 0.1733  | 97.58%                              | 15                            | 0.36                           | 754   | 0.454   | 0.05262                                       | 0.55         | 0.3805                                       | 0.62         | 0.05244                                      | 0.15         | 0.573                 | 311.9   | 12.5     | 327.37                                       | 1.73     | 329.56  | 0.48        |
| z19             | 1.145                    | 0.1573  | 97.23%                              | 12                            | 0.37                           | 658   | 0.359   | 0.05259                                       | 0.56         | 0.3793                                       | 0.64         | 0.05231                                      | 0.18         | 0.571                 | 310.6   | 12.8     | 326.54                                       | 1.80     | <b>328.78</b>                                   | <b>0.58</b> |
| z21             | 1.472                    | 0.1358  | 95.68%                              | 8                             | 0.51                           | 421   | 0.464   | 0.05289                                       | 0.97         | 0.3830                                       | 1.05         | 0.05253                                      | 0.18         | 0.541                 | 323.5   | 21.9     | 329.24                                       | 2.96     | 330.06  | 0.59        |
| z22             | 1.552                    | 0.2240  | 94.98%                              | 7                             | 0.98                           | 363   | 0.488   | 0.05274                                       | 0.95         | 0.3805                                       | 1.06         | 0.05233                                      | 0.26         | 0.516                 | 317.2   | 21.6     | 327.42                                       | 2.96     | <b>328.86</b>                                   | <b>0.85</b> |
| zA1             | 1.515                    | 0.1162  | 96.44%                              | 10                            | 0.36                           | 511   | 0.478   | 0.05303                                       | 0.69         | 0.3886                                       | 0.77         | 0.05315                                      | 0.15         | 0.608                 | 329.9   | 15.8     | 333.38                                       | 2.20     | 333.89  | 0.48        |
| zA2             | 1.535                    | 0.0427  | 93.65%                              | 6                             | 0.24                           | 287   | 0.486   | 0.05265                                       | 2.45         | 0.3591                                       | 2.56         | 0.04946                                      | 0.34         | 0.387                 | 313.4   | 55.7     | 311.51                                       | 6.87     | 311.25  | 1.05        |
| zA3             | 1.372                    | 0.0706  | 96.95%                              | 12                            | 0.18                           | 596   | 0.435   | 0.05325                                       | 0.73         | 0.3845                                       | 0.83         | 0.05236                                      | 0.26         | 0.524                 | 339.1   | 16.5     | 330.31                                       | 2.33     | <b>329.06</b>                                   | <b>0.82</b> |
| zA4             | 1.404                    | 0.0423  | 93.74%                              | 6                             | 0.23                           | 291   | 0.444   | 0.05313                                       | 1.35         | 0.3830                                       | 1.47         | 0.05228                                      | 0.32         | 0.480                 | 334.0   | 30.6     | 329.23                                       | 4.15     | <b>328.56</b>                                   | <b>1.04</b> |

- (a) z1, z2 etc. are labels for fractions composed of single zircon grains or fragments; all fractions annealed and chemically abraded after Mattinson (2005).
- (b) Model Th/U ratio calculated from radiogenic  $^{208}\text{Pb}/^{206}\text{Pb}$  ratio and  $^{207}\text{Pb}/^{235}\text{U}$  age.
- (c) Pb\* and Pbc represent radiogenic and common Pb, respectively; mol %  $^{206}\text{Pb}^*$  with respect to radiogenic, blank and initial common Pb.
- (d) Measured ratio corrected for spike and fractionation only.
- (e) Corrected for fractionation, spike, and common Pb; up to 2 pg of common Pb was assumed to be procedural blank:  $^{206}\text{Pb}/^{204}\text{Pb} = 18.60 \pm 0.80\%$ ;  $^{207}\text{Pb}/^{204}\text{Pb} = 15.69 \pm 0.32\%$ ;  $^{208}\text{Pb}/^{204}\text{Pb} = 38.51 \pm 0.74\%$  (all uncertainties 1-sigma). Excess over blank was assigned to initial common Pb.
- (f) Errors are 2-sigma, propagated using the algorithms of Schmitz & Schoene (2007).
- (g) Calculations are based on the decay constants of Jaffey et al. (1971).  $^{206}\text{Pb}/^{238}\text{U}$  and  $^{207}\text{Pb}/^{206}\text{Pb}$  ages corrected for initial disequilibrium in  $^{230}\text{Th}/^{238}\text{U}$  using Th/U [magma] = 3 using the algorithms of Schärer (1984).
- (h) dates in bold are those included in weighted mean calculations. See text for discussion.

Table 2 Biostratigraphical data used to compile the distribution of marine bands

| LOCALITY   | EASTING | NORTHIN<br>G | AMMONOIDS   |
|--|---------|--------------|---|
| <i>Emstites (Cravenoceras) leion</i> Marine Band (E1a1)                |         |              |   |
| Whinney Gill Reservoir, Skipton, Yorkshire                             | 399900  | 451000       | <i>Cravenoceras leion</i>   |
| River Ribble, Dinckley, Lancs  | 368640  | 436520       | <i>Cravenoceras leion</i>   |
| Harry Wall Gill - in stream bed W of Station - Bolton Abbey.           | 405830  | 452740       | <i>Cravenoceras leion</i>   |
| Bramley Farm - Bleaklow - Derbys.                                      | 424300  | 373400       | <i>Cravenoceras leion</i>   |
| Carla Beck - S of Rectory Carleton - Skipton - Yorks.                  | 397600  | 448100       | <i>Cravenoceras leion</i>   |
| Audley Beck - Pendleton - nr Clitheroe - Lancs.                        | 376100  | 438500       | <i>Cravenoceras leion</i>   |
| Croasdale Beck - NNW of Slaidburn                                      | 369680  | 455510       | <i>Cravenoceras leion</i>   |
| Whinney Hill Brickpits, Skipton  | 400100  | 451100       | <i>Cravenoceras leion</i>   |
| Longstone Station - disused Railway cutting West of Longstone Station. | 419530  | 371120       | <i>Cravenoceras leion</i>   |
| Greenleighton Quarry, 9 miles NNW of Belsay                            | 403400  | 591700       | <i>Cravenoceras leion</i>   |
| Bleaklow Mining Company - Exposure.                                    | 423000  | 373420       | <i>Cravenoceras leion</i>   |
| Alport Boring, near Alport   | 413610  | 391050       | <i>Cravenoceras leion</i>   |
| Hind Clough, Forest of Bowland   | 364400  | 453300       | <i>Cravenoceras leion</i>   |
| Southwest of The Hill  | 406400  | 357300       | <i>Cravenoceras leion</i>   |
| Railway cutting near Thornbridge Hall                                  | 419530  | 371130       | <i>Cravenoceras leion</i>   |
| Raper Mine   | 421690  | 365230       | <i>Cravenoceras leion</i>   |
| Bowers Hall BH SK26SW/46   | 423490  | 364560       | <i>Cravenoceras leion</i>   |
| Mootlaw Quarry, Matfen   | 402400  | 575000       | <i>Cravenoceras</i> aff. <i>lineolatum</i>  |
| Light Clough, Pendle Hill  | 375160  | 437640       | <i>Cravenoceras leion</i>   |
| Downs Gill, Coverhead  | 399000  | 476500       | ? <i>Cravenoceras</i> sp.   |
| Roosecote BH   | 323040  | 468660       | <i>Cravenoceras</i> sp.   |
| Darnbrook Beck   | 387420  | 471710       | <i>Cravenoceras leion</i>   |
| Cominco Borehole S3  | 386010  | 463500       | <i>Cravenoceras leion</i> , <i>Eumorphoceras tornquisti</i>   |
| Eshton Beck  | 394220  | 455730       | <i>Cravenoceras</i> cf. <i>leion</i>  |
| Cowside Beck   | 385650  | 466370       | <i>Cravenoceras</i> cf. <i>leion</i>  |
| Daw Haw Beck   | 385100  | 466460       | <i>Cravenoceras leion</i>   |
| NE of Lower House, near Scotch Green, Inglewhite                       | 353640  | 440780       | <i>Cravenoceras</i> cf. <i>leion</i>  |
| Inglewhite   | 354550  | 439030       | <i>Cravenoceras</i> sp.   |
| East of Hall Trees Farm, west of Chipping                              | 360710  | 442200       | <i>Cravenoceras leion</i>   |
| 200 m north of Higher Core, Bowland Fells                              | 359240  | 444370       | <i>Cravenoceras leion</i>   |
| White Fold, near Longridge Fold  | 362180  | 439340       | <i>Cravenoceras</i> sp.   |
| Little Mearley Clough  | 377900  | 441400       | <i>Cravenoceras leion</i>   |
| Higher Laithe Plantation   | 386000  | 445300       | <i>Cravenoceras leion</i>   |
| Edale BH   | 410780  | 384930       | <i>Cravenoceras leion</i> , <i>Eumorphoceras tornquisti</i>   |
| Castleton BH   | 414100  | 382930       | <i>Cravenoceras leion</i> , <i>Eumorphoceras</i> sp.  |
| Hope cement works BH, Salter Barn                                      | 416780  | 382280       | <i>Cravenoceras leion</i> , <i>Eumorphoceras</i> sp.  |
| Calow No 1 BH  | 440860  | 370410       | <i>Cravenoceras</i> cf. <i>leion</i> , <i>Eumorphoceras</i> sp.   |
| Disused railway cutting near Waterhouses                               | 407530  | 349560       | 8 bands   |
| Bullclough   | 406030  | 355020       | <i>Cravenoceras</i> sp.   |
| Ford, R. Hamps   | 406600  | 353700       | <i>Cravenoceras leion</i>   |
| SW of Parwich  | 417680  | 354060       | <i>Cravenoceras leion</i>   |
| Lees Farm BH   | 418180  | 350160       | <i>Cravenoceras leion</i> , <i>C. sp.</i> , <i>Eumorphoceras involutum</i>  |
| Farnah House, Duffield   | 432430  | 343280       | <i>Cravenoceras</i> cf. <i>leion</i>  |
| Duffield BH  | 434280  | 342170       | <i>Cravenoceras</i> Sp., <i>C. leion</i>  |
| Widmerpool No. 1 BH SK62NW/1   | 463660  | 329580       | ? <i>Cravenoceras</i> sp., <i>C. cf. leion</i> , <i>C. cf. malhamense</i> , <i>Eumorphoceras</i> sp., <i>E. pseudobilingue</i> , <i>Girtyoceras</i> |
| <i>Cravenoceras brandoni</i> Marine Band (E1b1)                        |         |              |   |
| Burn Fell  | 367800  | 453100       | <i>Cravenoceras brandoni</i>  |
| <i>Tumulites pseudobilinguis</i> Marine Band (E1b2)                    |         |              |   |
| Ramshaw Beck - Skipton   | 397500  | 448600       | aff <i>Eumorphoceras pseudobilingue</i>   |
| Little Mearley Clough, Pendle, Lancs                                   | 378500  | 441100       | <i>Eumorphoceras pseudobilingue</i>   |

|   |         |              |   |
|---|---------|--------------|---|
| Cow Close Sike, Malham  | 390810  | 462100       | <i>Eumorphoceras pseudobilingue</i>                           |
| Alport BH, Alport Castle's Farm, Glossop                      | 413612  | 391055       | <i>Eumorphoceras pseudobilingue</i>                           |
| LOCALITY  | EASTING | NORTHIN<br>G | AMMONOIDS   |
| <i>Tumulites pseudobilinguis</i> Marine Band (E1b2) Continued |         |              |   |
| Studforth Gill - Tosside - SW of Settle                       | 377340  | 457500       | <i>Eumorphoceras pseudobilingue</i> C                         |
| Croasdale Beck - NNW of Slaidburn - Yorks                     | 369680  | 455510       | <i>Eumorphoceras pseudobilingue</i>                           |
| Jenny Gill, 30yds from bottom fence, Skipton.                 | 400440  | 451110       | <i>Eumorphoceras pseudobilingue</i>                           |
| Whinney Gill Quarry - Skipton                                 | 399900  | 451000       | <i>Eumorphoceras pseudobilingue</i>                           |
| Swarth Beck, Kellet Park Wood                                 | 353060  | 470770       | <i>Tumulites pseudobilinguis</i>                              |
| Burn Fell   | 367800  | 453100       | <i>Tumulites pseudobilinguis</i>                              |
| Skibeden Beck, Bullion 840' OD, Parkers Hull - Skipton        | 402200  | 451500       | <i>Eumorphoceras pseudobilingue</i>                           |
| Southwest of Warslow Hall                                     | 408620  | 359250       | <i>Eumorphoceras pseudobilingue</i>                           |
| Isingdale Beck, east of Linton                                | 401000  | 463000       | <i>Eumorphoceras pseudobilingue</i>                           |
| North bank of R. Wharfe, opposite Linton church               | 400500  | 463028       | <i>Eumorphoceras pseudobilingue</i>                           |
| Grimwith Reservoir  | 406000  | 464000       | <i>Eumorphoceras pseudobilingue</i>                           |
| Roosecote BH  | 323040  | 468660       | <i>Eumorphoceras pseudobilingue</i>                           |
| Hare Clough Beck, Catlow                                      | 370470  | 457150       | <i>Eumorphoceras pseudobilingue</i>                           |
| Hollow Gill Wood, south of Rathmell                           | 380040  | 458510       | <i>Eumorphoceras pseudobilingue</i>                           |
| Tranlands Beck SW of Malham                                   | 389460  | 462360       | <i>Eumorphoceras pseudobilingue</i>                           |
| Tributary of Tranlands Beck                                   | 388460  | 462350       | <i>Eumorphoceras pseudobilingue</i>                           |
| Crimple Beck BH   | 427280  | 451860       | <i>Tumulites pseudobilinguis</i>                              |
| River Brock, Walmsley Bridge to Brock Bottom                  | 353720  | 441660       | <i>Tumulites cf. pseudobilinguis</i>                          |
| Right bank Fiendsdale Water                                   | 359630  | 449350       | <i>Tumulites pseudobilinguis</i>                              |
| White Fold, near Longridge Fold                               | 362180  | 439340       | <i>Tumulites</i> sp.  |
| R. Ribble west of Dinckley Hall                               | 368640  | 436520       | <i>Eumorphoceras pseudobilingue</i>                           |
| Butler Clough   | 372600  | 435400       | <i>Eumorphoceras pseudobilingue</i>                           |
| Light Clough  | 375108  | 437708       | <i>Eumorphoceras pseudobilingue</i>                           |
| Deep Clough   | 380900  | 440300       | <i>Eumorphoceras pseudobilingue</i>                           |
| Weets Hollow  | 385900  | 445100       | <i>Eumorphoceras pseudobilingue</i>                           |
| Castleton BH  | 414100  | 382930       | <i>Eumorphoceras pseudobilingue</i> & C                       |
| River Noe, 700 yds N38W of Manor House                        | 410630  | 384950       | <i>Eumorphoceras pseudobilingue</i>                           |
| Calow No 1 BH   | 440860  | 370410       | <i>Eumorphoceras pseudobilingue</i> ss. & C                   |
| WSW of Knockerdawn  | 422810  | 352010       | <i>Eumorphoceras pseudobilingue</i>                           |
| N of Bradley Nook Farm  | 423320  | 347570       | <i>Eumorphoceras pseudobilingue</i>                           |
| Duffield BH   | 434280  | 342170       | <i>Eumorphoceras pseudobilingue</i> , <i>E.</i> sp (2 leaves) |
| Upholland No 2 BH   | 350443  | 402870       | <i>Eumorphoceras pseudobilingue</i>                           |
| <i>Cravenoceras malhamense</i> Marine Band (E1c1)             |         |              |   |
| Moor Close Gill - near Malham                                 | 393330  | 463940       | <i>Cravenoceras malhamense</i>                                |
| Swinhope Mine, from shale above Little Limestone              | 382600  | 546600       | <i>Cravenoceras aff. malhamense</i>                           |
| Roosecote BH  | 323040  | 468660       | <i>Cravenoceras malhamense</i>                                |
| Burn Side   | 368800  | 454370       | <i>Cravenoceras malhamense</i>                                |
| Copped Hill Clough  | 371140  | 457210       | <i>Cravenoceras malhamense</i>                                |
| Hollow Gill Wood, south of Rathmell                           | 379900  | 458570       | <i>Cravenoceras cf. malhamense</i>                            |
| Out Gang  | 390760  | 461510       | <i>Cravenoceras malhamense</i>                                |
| South of Stockdale Beck                                       | 384550  | 463100       | <i>Cravenoceras malhamense</i>                                |
| Cominco Borehole S9 40.5-45.4m depth                          | 383090  | 463300       | <i>Cravenoceras malhamense</i>                                |
| Daw Haw Beck  | 385100  | 466460       | <i>Cravenoceras malhamense</i>                                |
| Crimple Beck  | 425460  | 451780       | <i>Cravenoceras malhamense</i>                                |
| Woodfold, Beacon Fell   | 356790  | 442190       | <i>Cravenoceras malhamense</i>                                |
| Duckey Leach BH From 294-324 ft depth                         | 373800  | 446800       | <i>Cravenoceras</i> sp.                                       |
| Railway cutting north of Wilpshire Tunnel                     | 368700  | 432900       | <i>Cravenoceras malhamense</i>                                |
| Butler Clough   | 372600  | 435400       | <i>Cravenoceras malhamense</i>                                |
| Light Clough  | 375200  | 437500       | <i>Cravenoceras malhamense</i>                                |
| Little Mearley Clough   | 378500  | 441100       | <i>Cravenoceras malhamense</i>                                |
| Deep Clough   | 380900  | 440300       | <i>Cravenoceras malhamense</i>                                |
| 310 yards N50E from Firber House                              | 383400  | 443000       | <i>Cravenoceras malhamense</i>                                |
| Weets Hollow  | 385900  | 445100       | <i>Cravenoceras malhamense</i>                                |
| Thornton Wood   | 392000  | 448000       | <i>Cravenoceras malhamense</i>                                |
| 300 yards S70W of Smearber                                    | 393600  | 449200       | <i>Cravenoceras malhamense</i>                                |
| Carla Beck, 300 yards upstream from The Grange                | 397800  | 449400       | <i>Cravenoceras malhamense</i>                                |
| Town Edge, south of Lothersdale                               | 395800  | 445800       | <i>Cravenoceras malhamense</i>                                |
|   | 403500  | 451600       | <i>Cravenoceras cf. malhamense</i>                            |

|  |         |              |   |
|--|---------|--------------|---|
| 570 yards N5W of Ramsgreave Hall                             | 367740  | 432010       | <i>Cravenoceras malhamense</i>  |
| Castleton BH   | 414100  | 382930       | <i>Cravenoceras malhamense</i>  |
| LOCALITY   | EASTING | NORTHIN<br>G | AMMONOIDS   |
| <i>Cravenoceras malhamense</i> Marine Band (E1c1) Continued  |         |              |   |
| Hope cement works BH, Salter Barn                            | 416780  | 382280       | <i>Cravenoceras malhamense</i>  |
| Abbey Mills BH 4   | 319490  | 377470       | <i>Cravenoceras</i> sp., <i>Eumorphoceras</i> sp.   |
| Horton, 450 yds E by N of The Rails                          | 391900  | 358550       | <i>Cravenoceras</i>   |
| Calow No 1 BH  | 440860  | 370410       | <i>Cravenoceras malhamense</i>  |
| Near Moorside  | 404140  | 354570       | <i>Cravenoceras</i> sp.   |
| Duffield BH  | 434280  | 342170       | <i>Cravenoceras malhamense</i> ,<br><i>Cravenoceras</i> sp.   |
| Upholland No 2 BH  | 350443  | 402870       | <i>Cravenoceras</i> sp.   |
| <i>Cravenoceras cowlingsense</i> Marine Band (E2a1)          |         |              |   |
| Cockhill - Bewerley  | 411200  | 464400       | <i>Cravenoceras cowlingsense</i>  |
| Crook Dyke - Upper Nidderdale                                | 402570  | 476400       | <i>Cravenoceras cowlingsense</i>  |
| Brigstons Gill - Upper Swaledale - Yorks                     | 383900  | 501900       | <i>Cravenoceras cowlingsense</i>  |
| Screes End, Tarnbrook  | 360300  | 455400       | <i>Cravenoceras cowlingsense</i>  |
| Great Ugly Clough, Quernmore                                 | 351190  | 461120       | <i>Cravenoceras cowlingsense</i> , <i>E.</i><br><i>grassingtonense</i>  |
| Croft House Borehole, Newton-le-Wilows                       | 419820  | 488830       | <i>Cravenoceras</i> sp.   |
| Black Scar, Penhill  | 404200  | 486900       | <i>Cravenoceras cowlingsense</i>  |
| Gate Up Gill   | 405700  | 467400       | <i>Cravenoceras cowlingsense</i>  |
| Burn Gill, Nidderdale  | 412800  | 468200       | <i>Cravenoceras cowlingsense</i>  |
| Mirk Fell Gill   | 391000  | 507000       | <i>Cravenoceras cowlingsense</i>  |
| Oak Beck, Oakdale temporary exposure                         | 427450  | 454640       | <i>Cravenoceras cowlingsense</i>  |
| Left bank, Grizedale Brook                                   | 351170  | 447820       | <i>Cravenoceras cowlingsense</i>  |
| North of Warley Wise Farm                                    | 394400  | 443600       | <i>Eumorphoceras bisulcatum</i>   |
| 50 yards east of Owl Cotes, 500 yards ESE of<br>Mire Close   | 396700  | 445000       | <i>Eumorphoceras bisulcatum</i>   |
| Cononley Beck  | 398590  | 446910       | <i>Eumorphoceras grassingtonense</i> ,<br><i>Cravenoceras cowlingsense</i>  |
| Eller Beck   | 400500  | 448800       | <i>Eumorphoceras bisulcatum</i>   |
| Edge   | 402500  | 450000       | <i>Eumorphoceras grassingtonense</i>  |
| Bradley Gill   | 400800  | 449300       | <i>Eumorphoceras grassingtonense</i>  |
| 350 yards NE of Kildwick Hall                                | 401400  | 446500       | <i>Cravenoceras cowlingsense</i>  |
| River Noe, 20 yds downstream of roadbridge<br>to Upper Booth | 410370  | 385120       | <i>Cravenoceras cowlingsense</i>  |
| Alport Boring, near Alport, Derbyshire                       | 413612  | 391055       | <i>Cravenoceras cowlingsense</i> ,<br><i>Eumorphoceras bisulcatum</i> sl.   |
| R. Hamps near Ironpits                                       | 406620  | 352040       | <i>Cravenoceras cowlingsense</i>  |
| Lea Brook, near Cauldon                                      | 407330  | 349690       | <i>Eumorphoceras grassingtonense</i>  |
| Moorside   | 404380  | 354100       | <i>Cravenoceras cowlingsense</i>  |
| Duffield BH  | 434280  | 342170       | <i>Cravenoceras cowlingsense</i> ,<br><i>Cravenoceras</i> sp.   |
| Upholland No 2 BH  | 350443  | 402870       | <i>Cravenoceras</i> sp.   |
| <i>Eumorphoceras ferrimontanum</i> Marine Band (E2a2)        |         |              |   |
| Tarnbrook Wyre, Abbeystead                                   | 356800  | 454400       | <i>Eumorphoceras ferrimontanum</i>  |
| East of Ward's Stone   | 359900  | 459100       | <i>Eumorphoceras ferrimontanum</i>  |
| Sapling Clough   | 362600  | 456300       | <i>Eumorphoceras ferrimontanum</i>  |
| Upper Dove Valley  | 406940  | 367250       | <i>Eumorphoceras ferrimontanum</i>  |
| Cogill Seave Bead, Lovely Seat                               | 388600  | 494800       | <i>Cravenoceras</i> aff. <i>cowlingsense</i>  |
| Croft House Borehole, Newton-le-Wilows                       | 419820  | 488830       | <i>Cravenoceras</i> ?   |
| Hookstone Beck   | 431300  | 454190       | <i>Eumorphoceras</i> cf. <i>ferrimontanum</i>   |
| Stone Rings Beck   | 430510  | 452750       | <i>Cravenoceras</i> sp., <i>Eumorphoceras</i><br><i>erinense</i>  |
| Barnacre Lodge   | 351580  | 446380       | <i>Cravenoceras</i> sp., <i>Eumorphoceras</i> cf.<br><i>ferrimontanum</i> , <i>E. erinense</i>                      |
| Holbeck  | 419800  | 447000       | <i>Eumorphoceras erinense</i> , <i>E.</i><br><i>ferrimontanum</i> , <i>Cravenoceras</i> sp.                         |
| River Washburn   | 422900  | 447000       | <i>Eumorphoceras erinense</i> , <i>E.</i><br><i>ferrimontanum</i> , <i>Cravenoceras</i> sp. nov,                    |
| 775 yards N of Leathley Hall                                 | 423700  | 447500       | AMMONOIDEA <i>Eumorphoceras</i><br><i>erinense</i> , <i>Cravenoceras</i> sp. nov, <i>E.</i><br><i>ferrimontanum</i> |
| R. Hamps near Crowtrees, Waterhouses                         | 407370  | 350250       | <i>Cravenoceras gairense</i> , <i>Eumorphoceras</i><br><i>bisulcatum</i> & <i>Kazakhoceras scaliger</i>             |
| Hulland  | 424600  | 346220       | <i>Eumorphoceras bisulcatum</i>   |
| 0.75 mile SW of Wirksworth Church                            | 428020  | 353050       | <i>Eumorphoceras bisulcatum</i>   |

|  |        |        |                                 |
|--|--------|--------|---------------------------------|
| Alport Boring, near Alport, Derbyshire | 413612 | 391055 | <i>Eumorphoceras bisulcatum</i> |
|--|--------|--------|---------------------------------|

| LOCALITY   | EASTING | NORTHIN<br>G | AMMONOIDS  |
|--|---------|--------------|--|
| <i>Eumorphoceras ferrimontanum</i> Marine Band (E2a2) Continued      |         |              |  |
| Duffield BH  | 434280  | 342170       | <i>Eumorphoceras bisulcatum</i> ,<br><i>Cravenoceras</i> sp., <i>Kazakhoceras scaliger</i>               |
| Upholland No 2 BH  | 350443  | 402870       | <i>Eumorphoceras bisulcatum</i> cf. <i>erinense</i> & <i>ferrimontanum</i>                               |
| <i>Cravenoceras gressinghamense</i> Marine Band (E2a2a)              |         |              |  |
| Gressingham Beck   | 356440  | 469960       | <i>Cravenoceras gressinghamense</i>  |
| Hunt's Gill  | 360560  | 467020       | <i>Cravenoceras gressinghamense</i>  |
| Badger Ford Beck   | 369270  | 469808       | <i>Cravenoceras gressinghamense</i>  |
| Duffield BH  | 434280  | 342170       | <i>Eumorphoceras</i> sp.   |
| <i>Eumorphoceras yatesae</i> Marine Band (E2a3)                      |         |              |  |
| Mill Dam Beck - E of New Bridge - Weston - Burley in Wharfedale      | 417900  | 447900       | <i>Eumorphoceras yatesae</i>   |
| Croker Hill - N of Dawsons - Cheshire                                | 392710  | 367380       | <i>Eumorphoceras yatesae</i>   |
| Throstle Nest - Silsden - Yorkshire                                  | 403800  | 446800       | <i>Eumorphoceras yatesae</i>   |
| Artle Beck   | 355200  | 462470       | <i>Eumorphoceras yatesae</i>   |
| Coppice Beck, Harrogate  | 430000  | 456200       | <i>Eumorphoceras yatesae</i>   |
| Sales Wheel, Samlesbury Hall   | 367560  | 435850       | <i>Eumorphoceras yatesae</i>   |
| Duffield BH  | 434280  | 342170       | <i>Eumorphoceras yatesae</i> , <i>E.</i> sp.   |
| Harewood BH SE34SW/37  | 432200  | 444100       | <i>Cravenoceras</i>  |
| <i>Cravenoceratoides edalensis</i> Marine Band (E2b1)                |         |              |  |
| River Noe, right bank, W of Edale Mill                               | 412850  | 385120       | <i>Cravenoceratoides edalense</i> (type)   |
| Bosley, stream W of Higher Minnend, E of Hug Bridge                  | 393730  | 364590       | <i>Cravenoceratoides edalense</i>  |
| Throstle Nest - Silsden - Yorkshire                                  | 403800  | 446800       | <i>Cravenoceratoides aff. edalense</i>   |
| Goodber Beck   | 363920  | 460780       | <i>Cravenoceratoides edalensis</i> , <i>C.</i> cf. <i>subplicatum</i>                                    |
| Bowers Hall BH   | 423490  | 364560       | <i>Cravenoceratoides edalensis</i>   |
| Knott Copy BH  | 376980  | 464490       | <i>Cravenoceratoides edalensis</i>   |
| Coppice Beck, Harrogate  | 430000  | 456200       | <i>Cravenoceratoides edalensis</i>   |
| Sales Wheel, Salesbury Hall  | 367460  | 435850       | <i>Cravenoceratoides edalensis</i> ; underlying <i>C. subplicatum</i> bed                                |
| Alport Boring, near Alport, Derbyshire                               | 413612  | 391055       | <i>Cravenoceratoides edalensis</i>   |
| Ladywash crosscut  | 422620  | 376930       | <i>Cravenoceratoides edalensis</i>   |
| R. Hamps near Winkhill   | 406870  | 350410       | <i>Cravenoceratoides edalensis</i>   |
| SW of Shiningford  | 424200  | 352360       | <i>Cravenoceratoides edalensis</i>   |
| Duffield BH  | 434280  | 342170       | <i>Cravenoceratoides edalensis</i> , <i>Cravenoceras</i> sp. nov., <i>C. subplicatum</i>                 |
| Harewood BH  | 432200  | 444100       | <i>Cravenoceratoides</i>   |
| <i>Cravenoceras nitidus</i> Marine Band (E2b2)                       |         |              |  |
| River Ribble - Dinckley - Lancs.                                     | 368640  | 436520       | <i>Cravenoceratoides nitidum</i>   |
| Keasden (or Keasdon) Beck - 0.25ml above Tunnerford Bridge - Clapham | 372440  | 465460       | <i>Cravenoceratoides cf. nitidum</i>   |
| Goodber Beck   | 363920  | 460780       | <i>Cravenoceratoides nitidum</i>   |
| Greenholes Beck  | 356480  | 463040       | <i>Eumorphoceras leirimense</i>  |
| Branstone Beck   | 367830  | 467860       | <i>Eumorphoceras leirimense</i>  |
| Crag Hall BH, Ellel Grange   | 348390  | 453450       | <i>Cravenoceratoides cf. nitidus</i>   |
| Wiggenstall  | 409020  | 360780       | <i>Cravenoceratoides nitidus</i>   |
| Pow Gill, 130 yards N of bridge at Powbank                           | 325360  | 542300       | <i>Anthrococeras glabrum</i>   |
| Old Quarry, Wath, Nidderdale   | 414600  | 468400       | <i>Cravenoceratoides nitidus</i>   |
| Cross Gill, Nidderdale   | 404200  | 470800       | <i>Cravenoceratoides nitidus</i>   |
| NW slope of Great Whernside  | 400200  | 476000       | <i>Cravenoceratoides nitidus</i>   |
| Stand Sike, Upper Nidderdale   | 405100  | 477900       | <i>Cravenoceratoides nitidus</i>   |
| Thorny Crane Gill, Colsterdale                                       | 411200  | 479600       | <i>Cravenoceratoides nitidus</i>   |
| Spruce Gill, Colsterdale   | 413500  | 480300       | <i>Cravenoceratoides nitidus</i>   |
| Ulfers Gill, Colsterdale   | 409300  | 482700       | <i>Cravenoceratoides nitidus</i>   |
| Knott Copy BH 3  | 376980  | 464490       | <i>Cravenoceratoides nitidus</i> (lower), <i>Glaphyrites</i> (middle), <i>Gl. kettlesingense</i> (upper) |
| Former brick pit at Stonefall  | 433100  | 454800       | <i>Cravenoceratoides nitidus</i>   |
| Crimple Beck between Pannal Bridge & Almsford Bridge                 | 430740  | 451660       | <i>Cravenoceratoides</i> sp., <i>Cravenoceras</i> sp., <i>Eumorphoceras</i> sp.                          |



|   |         |              |  |
|---|---------|--------------|--|
| Left bank, Grizedale Brook  | 350740  | 447260       | <i>Cravenoceratoides nitidus</i> , <i>C. holmesi</i>   |
| Stubbing Beck   | 396600  | 443800       | <i>Cravenoceratoides</i> sp., <i>E. bisulcatum</i>   |
| Harewood BH   | 432200  | 444100       | <i>Eumorphoceras</i>   |
| Worthington BH  | 440450  | 321040       | <i>Cravenoceras subplicatum</i>  |
| LOCALITY  | EASTING | NORTHIN<br>G | AMMONOIDS  |
| <i>Cravenoceras nitidus</i> Marine Band (E2b2) Continued  |         |              |  |
| Hope cement works BH, Salter Barn   | 416780  | 382280       | <i>Cravenoceratoides</i> cf., <i>nitidus</i> ,<br><i>Cravenoceras</i> cf. <i>holmesi</i> ,<br><i>Eumorphoceras</i> sp. |
| Duffield BH   | 434280  | 342170       | <i>Cravenoceratoides nitidus</i> ,<br><i>Eumorphoceras leitricense</i> ,<br><i>Cravenoceras</i> sp.                    |
| Alport Boring, near Alport, Derbyshire  | 413612  | 391055       | <i>Cravenoceratoides nitidus</i> ,<br><i>Eumorphoceras bisulcatum</i> var.   |
| Upholland No 2 BH   | 350443  | 402870       | <i>Eumorphoceras bisulcatum</i> cf.<br><i>leitricense</i>  |
| <i>Cravenoceras nititoides</i> Marine Band (E2b3)   |         |              |  |
| North of Endon  | 392190  | 354140       | <i>Cravenoceras</i> sp.  |
| E. bank of R. Crowden, 10-20 yds above<br>confluence with R. Noe  | 410220  | 385260       | <i>Cravenoceras</i> ?, <i>Eumorphoceras</i> cf.<br><i>rostratum</i>  |
| Alport Boring, near Alport, Derbyshire  | 413612  | 391055       | <i>Cravenoceratoides nititoides</i> ,<br><i>Eumorphoceras</i> cf. <i>rostratum</i>                                     |
| Hope cement works BH, Salter Barn   | 416780  | 382280       | <i>Cravenoceratoides nititoides</i>  |
| River Terrig  | 323380  | 356970       | <i>Cravenoceratoides nititoides</i>  |
| Combes Brook, S of Ballfields   | 400770  | 352920       | <i>Eumorphoceras rostratum</i>   |
| River Ecclesbourne  | 431370  | 345550       | <i>Cravenoceratoides nititoides</i> ,<br><i>Eumorphoceras rostratum</i>  |
| Duffield BH   | 434280  | 342170       | <i>Cravenoceratoides nititoides</i> ,<br><i>Eumorphoceras rostratum</i> ,<br><i>Cravenoceras</i> sp.                   |
| Harewood BH   | 432200  | 444100       | <i>Cravenoceras</i>  |
| <i>Nuculoceras stellarum</i> Marine Band (E2c1)   |         |              |  |
| Gill Beck - Cowling - N. Yorks  | 395800  | 443600       | <i>Cravenoceratoides stellarum</i>   |
| Westfield Farm - Gill Beck - Cowling - Yorks.   | 395800  | 443600       | <i>Cravenoceratoides stellarum</i>   |
| Black Scars Beck, Cowling, SSW of Skipton<br>Station  | 394100  | 443000       | <i>Cravenoceratoides stellarum</i>   |
| Cheddleton Paper Mills BH.  | 397680  | 352470       | <i>Nuculoceras stellarum</i>   |
| River Wharfe, right bank, 200 yards SE of<br>Netherby   | 433300  | 446700       | <i>Cravenoceratoides stellarum</i>   |
| Right bank of Gill Beck, 10yds SSE of<br>Westfield 1230yds W 30deg N of Holy Trinity<br>Church, Cowling | 395800  | 443600       | <i>Cravenoceratoides stellarum</i>   |
| LOCALITY  | EASTING | NORTHIN<br>G | AMMONOIDS  |
| Carsington Reservoir BH CR10  | 424460  | 350200       | <i>Nuculoceras stellarum</i>   |
| Holehouse Lane  | 392020  | 354830       | <i>Nuculoceras stellarum</i>   |
| Castleberg Scar, 30 ft below Nesfield Sst   | 409100  | 449600       | <i>Nuculoceras stellarum</i>   |
| Hole Brook  | 360500  | 429400       | <i>Cravenoceras stellarum</i> , <i>C. holmesi</i>  |
| E. bank of R. Crowden, 10-20 yds above<br>confluence with R. Noe  | 410220  | 385260       | <i>Nuculoceras stellarum</i>   |
| Alport Boring, near Alport, Derbyshire  | 413612  | 391055       | <i>Nuculoceras stellarum</i>   |
| Combes Brook  | 401880  | 353330       | <i>Nuculoceras stellarum</i>   |
| River Ecclesbourne  | 431370  | 345550       | <i>Nuculoceras stellarum</i>   |
| Duffield BH   | 434280  | 342170       | <i>Nuculoceras</i> cf. <i>stellarum</i>  |
| Harewood BH   | 432200  | 444100       | <i>Nuculoceras stellarum</i>   |
| <i>Nuculoceras nuculum</i> Marine Band (E2c2-4)   |         |              |  |
| Tansley Bore - Derbyshire   | 433126  | 359604       | <i>Nuculoceras nuculum</i>   |
| Sutton - 240yds E by N of Crag & SW of St<br>Thomas's Church - Sutton.                                  | 400000  | 443500       | <i>Nuculoceras nuculum</i>   |
| R. Darwen, Samesbury Bottoms, Blackburn   | 361720  | 429360       | <i>Nuculoceras nuculum</i>   |
| Gill Beck - Cowling - N Yorks   | 394730  | 443300       | <i>Nuculoceras nuculum</i> (upper 2 bands)   |
| Bentend Farm - in stream 450yds SSW of<br>farm - near Dane Bridge - Staffs                              | 396420  | 363280       | <i>Nuculoceras nuculum</i>   |
| Owl Head Wood - Kearby - Yorks  | 434500  | 446600       | <i>Nuculoceras nuculum</i> , <i>Cravenoceras</i><br><i>fragile</i> (type)  |
| Oakhill Clough - 250yds NNW of Stansfield<br>Hall Station - Todmorden                                   | 393790  | 424870       | <i>Nuculoceras nuculum</i> (2 bands)   |
| Mam Tor - 0.75ml NW of Castleton  | 413150  | 383450       | <i>Nuculoceras nuculum</i>   |

|  |         |              |  |
|--|---------|--------------|--|
| Tittesworth Reservoir - SW corner - Staffs.                          | 399220  | 358830       | <i>Nuculoceras nuculum</i>   |
| Sutton-on-Trent Well No.3  | 479900  | 364900       | <i>Nuculoceras nuculum</i>   |
| Shellag Point BH RTZ 1   | 245650  | 499650       | <i>Nuculoceras nuculum</i>   |
| Moor Hall, Bagnall   | 394910  | 351180       | <i>Nuculoceras nuculum</i>   |
| Stoop Farm BH  | 406500  | 368220       | <i>Nuculoceras nuculum</i> (upper band)  |
| Bowers Hall BH   | 423490  | 364560       | <i>Nuculoceras nuculum</i>   |
| LOCALITY   | EASTING | NORTHIN<br>G | AMMONOIDS  |
| <i>Nuculoceras nuculum</i> Marine Band (E2c2-4) Continued            |         |              |  |
| Field House  | 412360  | 358390       | <i>Nuculoceras nuculum</i>   |
| Old Park Wood  | 365220  | 434350       | <i>Nuculoceras nuculum</i>   |
| R. Ribble near Balderstone Hall                                      | 361310  | 433300       | <i>Nuculoceras nuculum</i> (upper band)  |
| Shawhead Beck SSE of Shaw Gate (2 bands)                             | 392300  | 441500       | <i>Nuculoceras nuculum</i> , <i>E. bisulcatum</i> ,<br><i>Cravenoceratoides fragilis</i> (middle),<br><i>Cravenoceras darwenense</i> (middle),<br><i>Kazakhoceras hawkinsi</i> (middle band) |
| Black Scars  | 394100  | 443000       | <i>Nuculoceras nuculum</i> (lower band)  |
| 340 yards E17N of Nesfield Church                                    | 409600  | 449700       | <i>Nuculoceras nuculum</i> , <i>E. bisulcatum</i> ,<br><i>Cravenoceratoides fragilis?</i> (lower band)   |
| 160 yards S 30E of Gildersber  | 407100  | 448800       | <i>Nuculoceras nuculum</i> , <i>E. bisulcatum</i> ,<br><i>Cravenoceratoides fragilis?</i> (lower band)   |
| Crowden Brook  | 410250  | 385400       | <i>Nuculoceras nuculum</i>   |
| Grinds Brook, 0.25 miles N15W of Edale church                        | 412210  | 386160       | <i>Nuculoceras nuculum</i>   |
| Alport Boring, near Alport, Derbyshire                               | 413612  | 391055       | <i>Nuculoceras nuculum</i> , <i>Eumorphoceras bisulcatum</i> (3 horizons)  |
| Dove Holes, railway cutting  | 407610  | 379330       | <i>Nuculoceras nuculum</i> , <i>Eumorphoceras bisulcatum</i>   |
| R. Derwent, 660 yds N84W of St Helen's Church, Churchtown            | 426070  | 363040       | <i>Nuculoceras nuculum</i>   |
| Cromford Station   | 430340  | 357440       | <i>Nuculoceras nuculum</i> , <i>Eumorphoceras bisulcatum</i>   |
| Combes Brook   | 400660  | 352920       | <i>Nuculoceras nuculum</i> (3 bands)   |
| Middle Cliff   | 400020  | 354770       | <i>Nuculoceras nuculum</i> , <i>Eumorphoceras bisulcatum</i> (highest band)  |
| Ipstones Edge BH   | 402580  | 351090       | <i>Nuculoceras nuculum</i> , (two bands)   |
| Biggin Brook   | 425760  | 347770       | <i>Nuculoceras nuculum</i> , <i>Eumorphoceras bisulcatum</i> (lower band)  |
| Franker Brook  | 430810  | 347290       | <i>Nuculoceras nuculum</i> (lower band)  |
| Duffield BH  | 434280  | 342170       | <i>Nuculoceras nuculum</i> (3 bands)   |
| <i>Isohomoceras subglobosum</i> Marine Band (H1a1-3)                 |         |              |  |
| Brunthwaite Beck - E of Brunthwaite & S of Parish Church - Silsden   | 405200  | 446200       | <i>Homoceras subglobosum</i>   |
| Rowley Wood, W of Low House & S of Station - Ben Rhydding            | 414500  | 447100       | <i>Homoceras cf subglobosum</i>  |
| Ilkley, NE of Pomona/SW of Middleton, Ilkley                         | 412100  | 449000       | <i>Homoceras cf subglobosum</i>  |
| Stone Head Beck/Gill Beck - East of Colne                            | 394730  | 443300       | <i>Homoceras subglobosum</i> (3 bands)   |
| Stream - between Ford of Meerbrook & NW of New Grange Farm - Staffs. | 399290  | 360330       | <i>Homoceras subglobosum</i>   |
| River Noe - below Edale Mill - Derbyshire                            | 413700  | 385450       | <i>Homoceras subglobosum</i>   |
| River Darwen 1500 yards W of chapel at Nab's Head                    | 360890  | 429270       | <i>Homoceras subglobosum</i> (3 bands)   |
| Alport Boring, Derbyshire  | 413612  | 391055       | <i>Homoceras subglobosum</i>   |
| Black Scars Beck - Cowling - 5.25mls SSW of Skipton Station          | 394100  | 443000       | <i>Homoceras subglobosum</i>   |
| River Noe - left bank - 270yds S & 77deg W of Harrop Farm            | 416660  | 385370       | <i>Homoceras subglobosum</i> (2 bands)   |
| Tunnel Entrance - 1100yds N & 12deg E of Doveholes Station           | 407700  | 479010       | <i>Homoceras subglobosum</i>   |
| Tittesworth Reservoir, 2mls North of Leek                            | 399210  | 358960       | <i>Isohomoceras subglobosum</i>  |
| Stream section at Cocker Clough Wood, north of Dolphinholme          | 350780  | 455900       | <i>Isohomoceras subglobosum</i>  |
| Well Beck, near Summersgill  | 363980  | 463600       | <i>Isohomoceras subglobosum</i> (upper band)   |
| Field House  | 412360  | 358390       | <i>Isohomoceras subglobosum</i>  |
| Haddon Park Farm   | 423020  | 367580       | <i>Isohomoceras subglobosum</i>  |
| R. Ribble near Balderstone Hall                                      | 361330  | 433250       | <i>Isohomoceras subglobosum</i> (lower/middle band)  |
| Lumb Clough Beck   | 400600  | 443600       | <i>Isohomoceras subglobosum</i>  |

|   |         |              |  |
|---|---------|--------------|--|
| Swartha Wood  | 405380  | 446560       | <i>Isohomoceras subglobosum</i>  |
| Grinds Brook, 0.25 miles N15W of Edale church   | 412210  | 386160       | <i>Isohomoceras subglobosum</i>  |
| Crowden Brook, Upper Booth  | 410270  | 385520       | <i>Isohomoceras subglobosum</i>  |
| R. Noe, 710 yds S15E of Clough Farm   | 414750  | 385970       | <i>Homoceras subglobosum</i> (3 bands)   |
| Dove Holes, railway cutting   | 407610  | 379330       | <i>Homoceras subglobosum</i> (lower band)  |
| LOCALITY  | EASTING | NORTHIN<br>G | AMMONOIDS  |
| <b><i>Isohomoceras subglobosum</i> Marine Band (H1a1-3) Continued</b>                 |         |              |  |
| Quarry, Coed Llwybr-y-bi  | 319280  | 375200       | <i>Isohomoceras subglobosum</i> (lower band)   |
| Rushton, Dingle Brook, 230 yds SSE of Harper's Farm                                   | 392070  | 361750       | <i>Homoceras subglobosum</i>   |
| Horton, 160 yds NNE of Porter's Farm  | 392600  | 359390       | <i>Homoceras subglobosum</i>   |
| Carsington Reservoir R11 BH   | 424700  | 349860       | <i>Isohomoceras subglobosum</i> (3 bands)  |
| Ashcombe Park   | 397710  | 351070       | <i>Isohomoceras subglobosum</i> (1 band seen)  |
| Boosemoor Brook   | 437380  | 340530       | <i>Homoceras subglobosum</i>   |
| Mill Plantation   | 437590  | 339550       | <i>Homoceras subglobosum</i>   |
| Duffield BH   | 434280  | 342170       | <i>Homoceras subglobosum</i> (3 bands)   |
| <b><i>Homoceras beyrichianum</i> Marine Band (H1b1)</b>                               |         |              |  |
| Old Wives Gill, NE of Tivoli, 600yds WNW of Myddleton Lodge, Ilkley                   | 410600  | 449500       | <i>Homoceras beyrichianum</i>  |
| Swartha Gill, Silsden   | 405300  | 471000       | <i>Homoceras beyrichianum</i>  |
| Ilkley, right bank Hebers Gill, 1275yds E & 3deg N of Netherwood House & W of Station | 410120  | 447790       | <i>Homoceras beyrichianum</i>  |
| Brunthwaite Beck, 220yds N & 27deg E of Brunthwaite, Silsden                          | 405300  | 446500       | <i>Homoceras beyrichianum</i>  |
| Lumb Beck, 350yds SE of Throstle Nest, Addingham                                      | 408100  | 448600       | <i>Homoceras beyrichianum</i>  |
| Alport Boring, near Alport, Derbyshire  | 413612  | 391055       | <i>Homoceras beyrichianum</i>  |
| Lowgill, 170 m downstream of road bridge, Crossdale Beck                              | 365620  | 465240       | <i>Homoceras beyrichianum</i> , <i>H. cf. diadema</i> , <i>Isohomoceras</i> sp.        |
| Well Beck, near Summersgill   | 365110  | 465460       | <i>Homoceras beyrichianum</i> , <i>H. cf. diadema</i> , <i>Isohomoceras</i> sp.        |
| Wiggenstall   | 408750  | 360970       | <i>Homoceras beyrichianum</i>  |
| Stoop Farm BH   | 406500  | 368220       | <i>Homoceras beyrichianum</i>  |
| Field House   | 412360  | 358390       | <i>Homoceras cf. beyrichianum</i>  |
| Haddon Park Farm  | 423020  | 367580       | <i>Homoceras beyrichianum</i>  |
| Gill Beck - Cowling - N Yorks.  | 394730  | 443300       | <i>Homoceras beyrichianum</i>  |
| Knott Copy BH – nonsequence immediately above marine band                             | 376980  | 464490       | <i>Homoceras beyrichianum</i>  |
| 400 yards S80E from Shaw Gate, Shawhead Beck  | 392500  | 441800       | <i>Homoceras beyrichianum</i>  |
| River Darwen  | 361720  | 429360       | <i>Homoceras beyrichianum</i>  |
| River Noe, left bank, 270yds S & 77deg W of Harrop Farm                               | 416590  | 385540       | <i>Homoceras beyrichianum</i>  |
| Franker Brook   | 430590  | 347610       | <i>Homoceras beyrichianum</i> , <i>H. cf. subglobosum</i>                              |
| Ing Gill - Primrose Hill - Middleton - Ilkley   | 411160  | 449710       | <i>Homoceras aff. subglobosum</i>  |
| <b><i>Isohomoceras</i> sp. nov. Marine Band (H1b2)</b>                                |         |              |  |
| Lowgill, 170 m downstream of road bridge, Crossdale Beck                              | 365620  | 465240       | <i>Isohomoceras</i> sp. nov.   |
| Franker Brook   | 430570  | 347740       | <i>Homoceras</i> sp. aff. <i>beyrichianum</i> , <i>H. sp. of the subglobosum</i> group |
| <b><i>Hudsonoceras proteum</i> Marine Band (H2a1)</b>                                 |         |              |  |
| River Noe - Edale - Derbyshire  | 409570  | 385540       | <i>Hudsonoceras proteum</i> , <i>Homoceras cf. smithi</i> (3 bands)                    |
| Congleton Edge - Staffordshire  | 387680  | 360570       | <i>Hudsonoceras proteum</i>  |
| Mam Tor - Castleton - Derbyshire  | 412900  | 383400       | <i>Hudsonoceras proteum</i>  |
| Wiggenstall   | 408990  | 360800       | <i>Hudsonoceras proteum</i>  |
| Blake Brook, Longnor  | 406250  | 361190       | <i>Hudsonoceras proteum</i>  |
| Pendle Water, east of the inn at Roughlee   | 384600  | 440400       | <i>Hudsonoceras proteum</i>  |
| R. Darwen, N of Samlesbury Bottoms  | 361810  | 429090       | <i>Hudsonoceras proteus</i> , <i>Homoceras smithi</i>                                  |
| Crowden Brook, 640 yds N3E of Highfield   | 410250  | 385630       | <i>Hudsonoceras proteus</i> , <i>Homoceras smithi</i>                                  |
| Grinds Brook, Grindsbrook Booth   | 412220  | 386260       | <i>Hudsonoceras proteus</i> (upper), <i>Homoceras smithi</i> (lower)                   |

|   |        |        |   |
|---|--------|--------|---|
| Harden Clough                               | 412240 | 384460 |   |
| Alport Boring,near Alport,Derbyshire        | 413612 | 391055 | <i>Hudsonoceras proteus</i>                           |
| 1010 yds S32E of Alport Castles Farm        | 414010 | 390290 | <i>Hudsonoceras proteus</i> , <i>Homoceras smithi</i> |
| Potbank Quarry, Newbold Astbury             | 386910 | 359220 | <i>Hudsonoceras proteus</i>                           |
| R. Derwent, 750 yds N89W of Stancliffe Hall | 426030 | 364010 | <i>Hudsonoceras proteus</i>                           |
| Scow Brook                                  | 424970 | 350930 | <i>Hudsonoceras proteus</i>                           |

| LOCALITY   | EASTING | NORTHIN<br>G | AMMONOIDS   |
|--|---------|--------------|---|
| <i>Homoceras undulatum</i> Marine Band (H2b1)                                    |         |              |   |
| Ladywash Mine - Eyam - Derbyshire  | 422500  | 376800       | <i>Homoceras</i> aff. <i>undulatum</i>                                  |
| Brunthwaite Beck - 220yds N & 27deg E of Brunthwaite - Silsden                   | 405300  | 446500       | <i>Homoceras</i> aff. <i>undulatum</i> , <i>H. beyrichianum</i>         |
| Lumb Clough Beck - 735yds S & 13deg W of Sutton Church - Yorks.                  | 400700  | 443500       | ? <i>Homoceras undulatum</i>  |
| Samlesbury - River Darwen  | 361830  | 429160       | <i>Homoceras undulatum</i>  |
| Shell Brook - Greasley Hollow  | 394700  | 366130       | <i>Homoceras</i> aff. <i>undulatum</i>                                  |
| Roughlee - Right bank - immediately downstream for Stepping Stones               | 384600  | 440400       | <i>Homoceras undulatum</i>  |
| Eskew Beck, Bentham  | 364890  | 468330       | <i>Homoceras undulatum</i> , <i>H. cf. smithi</i>                       |
| Tittesworth Reservoir Water Treatment Plant-N of Leek                            | 399500  | 358500       | <i>Homoceras undulatum</i>  |
| Right bank of stream, 350 yards SW of Lower Jack Field                           | 399400  | 443400       | <i>Homoceras</i> cf. <i>undulatum</i> , <i>H. cf. smithi</i>            |
| Alport Boring, near Alport, Derbyshire   | 413612  | 391055       | <i>Homoceratoides</i> cf. <i>undulatum</i> ,                            |
| 1010 yds S32E of Alport Castles Farm   | 414010  | 390290       | <i>Homoceras</i> cf. <i>undulatum</i>                                   |
| Stream section   | 322680  | 371130       | <i>Homoceras undulatum</i>  |
| Coed-y-cra   | 327540  | 370590       | <i>Homoceras undulatum</i>  |
| Rushton, Dingle Brook, 370 yds SW by W of Fold Farm                              | 392750  | 361280       | <i>Homoceras undulatum</i>  |
| Franker Brook  | 430550  | 347820       | <i>Homoceras</i> cf. <i>undulatum</i>                                   |
| <i>Vallites eostriolatus</i> Marine Band (H2c1)                                  |         |              |   |
| Right bank of River Darwen, 100yds NW of bridge, Samlesbury Bottoms, Blackburn   | 361830  | 429160       | <i>Vallites eostriolatus</i>  |
| Pendle Water, east of the inn at Roughlee  | 384600  | 440400       | <i>Homoceras eostriolatum</i>   |
| Hillside Section - 450yds E & 13deg S of Knot House - Eastburn - Sutton          | 402100  | 444200       | <i>Homoceras</i> aff. <i>undulatum</i> , <i>H. cf. striolatum</i>       |
| Alport Boring,near Alport, Derbyshire  | 413612  | 391055       | <i>Homoceras eostriolatum</i>   |
| <i>Homoceratoides prereticulatus</i> Marine Band (H2c2)                          |         |              |   |
| Roughlee - 10yds downstream from Stepping Stones                                 | 384600  | 440400       | <i>Homoceratoides prereticulatus</i>                                    |
| Mam Tor - Castleton - Derbyshire.  | 410440  | 384300       | <i>Homoceratoides prereticulatum</i>                                    |
| Holden Beck, Silsden (Holotype)  | 405940  | 445470       | <i>Homoceratoides prereticulatum</i>                                    |
| River Darwen NW of Samlesbury Bridge,near Blackburn Lancs                        | 361830  | 429160       | <i>Homoceratoides</i> cf. <i>prereticulatum</i>                         |
| Blake Brook, Longnor   | 406250  | 361190       | <i>Homoceratoides prereticulatum</i>                                    |
| River Noe - at junction with Grains Clough - Edale                               | 409530  | 385510       | <i>Homoceratoides prereticulatum</i>                                    |
| Harden Clough  | 412240  | 384460       | <i>Homoceratoides</i> aff. <i>prereticulatum</i> , <i>Homoceras</i> sp. |
| Alport Boring,near Alport,Derbyshire   | 413612  | 391055       | <i>Homoceratoides</i> aff. <i>prereticulatum</i>                        |
| Alport, ENE of Hayridge Farm   | 414120  | 389690       | <i>Homoceratoides prereticulatum</i>                                    |
| R. Noe, 200 yds W-WNW of Fulwood Holmes  | 416700  | 385000       | <i>Homoceratoides prereticulatum</i>                                    |
| Horton, 210 yds N by E of Bentend  | 396600  | 363890       | <i>Homoceratoides prereticulatus</i> , <i>H. sp.</i>                    |
| Biddulph, 420 yds E by N of Heath Hay  | 390900  | 359310       | <i>Homoceratoides prereticulatus</i>                                    |
| Johannesburg No 6 BH   | 430370  | 359010       | <i>Homoceratoides prereticulatus</i>                                    |
| <i>Hodsonites magistrorum</i> Marine Band (R1a1)                                 |         |              |   |
| Backstone Beck,SE of Ilkley Station Yorks  | 412500  | 447200       | <i>Homoceras</i> [sp. nov. A] <i>magistrorum</i>                        |
| Blake Brook, Longnor   | 406250  | 361190       | <i>Homoceras magistrorum</i>  |
| Right bank of River Darwen 100yds NW of Bridge,Samlesbury Bottoms,near Blackburn | 361830  | 429160       | <i>Homoceras magistrorum</i>  |
| Alport, ENE of Hayridge Farm   | 414120  | 389690       | <i>Homoceras</i> sp.  |
| Maplebeck Well   | 470520  | 360090       | <i>Homoceras</i> cf. <i>magistrorum</i>                                 |
| <i>Reticuloceras circumplicatile</i> Marine Band (R1a2)                          |         |              |   |
| Pendle Water - Rough Lee - Nelson  | 384500  | 440300       | <i>Reticuloceras circumplicatile</i>                                    |
| Blake Brook, Longnor   | 406250  | 361190       | <i>Reticuloceras circumplicatile</i>                                    |
| Right bank of River Darwen 100yds NW of  | 361830  | 429160       | <i>Reticuloceras circumplicatile</i>                                    |

|  |         |              |   |
|--|---------|--------------|---|
| Bridge, Samlesbury Bottoms, near Blackburn   |         |              |   |
| Mousegill Beck, Stainmore  | 383520  | 512490       | <i>Vallites henkei</i>  |
| Franker Brook  | 430560  | 347880       | <i>Reticuloceras circumplicatile</i> , <i>Vallites henkei</i> , <i>Homoceratoides</i> sp.   |
| Ferriby Brook  | 437870  | 339660       | <i>Reticuloceras circumplicatile</i> , <i>Vallites henkei</i>   |
| <i>Reticuloceras subreticulatum</i> Marine Band (R1a3)                                 |         |              |   |
| Grinds Brook, 25 yds E of Grindslow House  | 412140  | 386350       | <i>Homoceratoides</i> sp., <i>Reticuloceras</i> cf. <i>pulchellum</i> , <i>R.</i> cf. <i>subreticulatum</i>                                 |
| LOCALITY   | EASTING | NORTHIN<br>G | AMMONOIDS   |
| <i>Reticuloceras subreticulatum</i> Marine Band (R1a3) Continued                       |         |              |   |
| Franker Brook  | 430590  | 347960       | <i>Reticuloceras subreticulatum</i>   |
| Ferriby Brook  | 437970  | 339720       | <i>Reticuloceras</i> cf. <i>subreticulatum</i>  |
| <i>Reticuloceras todmordenense</i> Marine Band (R1a4)                                  |         |              |   |
| Lumbutts Clough, Woodhouse, Todmorden  | 395030  | 424220       | <i>Reticuloceras todmordenense</i>  |
| Roughlee - Pendle  | 384600  | 440400       | <i>Reticuloceras todmordenense</i>  |
| Brund BHs, Manifold valley   | 408920  | 361530       | <i>Reticuloceras</i> cf. <i>todmordenense</i>   |
| Knott Copy BH  | 376980  | 464490       | <i>Reticuloceras todmordenense</i> , <i>R. paucicrenulatum</i> , <i>R.</i> aff. <i>adpressum</i>  |
| Farnham BH   | 434690  | 459960       | <i>Reticuloceras paucicrenulatum</i>  |
| Grinds Brook, 25 yds E of Grindslow House  | 412140  | 386350       | <i>Homoceras</i> sp., <i>Reticuloceras</i> sp., <i>R. ?todmordenense</i>  |
| <i>Reticuloceras dubium</i> Marine Band (R1a5)   |         |              |   |
| Holden Beck - N of Holden Bridge - S of Silsden Parish Church                          | 405900  | 445500       | <i>Reticuloceras</i> cf. <i>dubium</i>  |
| Knott Copy BH  | 376980  | 464490       | <i>Reticuloceras dubium</i>   |
| 533yds SSW of Hags Rd Farm, Spofforth Hags - North Yorkshire                           | 433740  | 450700       | <i>Reticuloceras dubium</i>   |
| <i>Reticuloceras dubium</i> Marine Band (R1a5) Continued                               |         |              |   |
| Greenway Hall Golf Course  | 391860  | 351270       | <i>Reticuloceras</i> cf. <i>dubium</i> , <i>R.</i> sp.  |
| Bentham Station BH   | 366590  | 468930       | <i>Reticuloceras dubium</i>   |
| Blackwood End Farm, Quernmore  | 351270  | 457810       | <i>Reticuloceras dubium</i>   |
| Samlesbury - River Darwen  | 361830  | 429160       | <i>Reticuloceras dubium</i>   |
| <i>Reticuloceras eoreticulatum</i> Marine Band (R1b1)                                  |         |              |   |
| Mam Tor - Castleton - Derbyshire   | 412990  | 383460       | <i>Reticuloceras eoreticulatum</i>  |
| Black Bank Syke  | 376370  | 465270       | <i>Reticuloceras</i> sp.  |
| Roughlee (type specimen) large scar on right bank 65 yards upstream of stepping stones | 384300  | 440200       | <i>Reticuloceras eoreticulatum</i>  |
| <i>Reticuloceras nodosum</i> Marine Band (R1b2)  |         |              |   |
| Swint Clough - Alport Valley   | 413470  | 391060       | <i>Reticuloceras nodosum</i> , <i>Homoceras spiraloide</i> s, <i>H. striolatum</i>  |
| Spofforth Hags - North Yorkshire   | 433740  | 450700       | <i>Reticuloceras nodosum</i> group  |
| Porters Farm - 420yds East + 3deg South of Porters Farm - Horton                       | 392930  | 359140       | <i>Reticuloceras</i> cf. <i>nodosum</i>   |
| Brund BHs, Manifold valley   | 408920  | 361530       | <i>Reticuloceras</i> aff. <i>nodosum</i>  |
| Mam Tor - Castleton - Derbyshire   | 412990  | 383460       | <i>Reticuloceras</i> cf. <i>nodosum</i>   |
| <i>Reticuloceras stubblefieldi</i> Marine Band (R1b3)                                  |         |              |   |
| River Noe - Edale - Derbyshire   | 417640  | 383170       | <i>Reticuloceras stubblefieldi</i>  |
| 1000m ENE of Bull Bank   | 362880  | 472060       | <i>Reticuloceras</i> cf. <i>stubblefieldi</i>   |
| Brund BHs, Manifold valley   | 408920  | 361530       | <i>Reticuloceras</i> cf. <i>stubblefieldi</i>   |
| Acton Burn, north of Derwent Reservoir   | 398300  | 552880       | <i>Reticuloceras</i> cf. <i>stubblefieldi</i> (juv)   |
| Crag Gill, near White House  | 402680  | 523620       | <i>Reticuloceras stubblefieldi</i>  |
| Black Bank Syke  | 376370  | 465270       | <i>Reticuloceras stubblefieldi</i>  |
| Swint Clough - Alport Valley - Derbyshire  | 413470  | 391060       | <i>Reticuloceras</i> cf. <i>stubblefieldi</i> , <i>Hudsonoceras ornatum</i> , <i>R.</i> aff. <i>Moorei</i> , <i>R.</i> cf. <i>regularum</i> |
| Franker Brook  | 430600  | 348050       | <i>Reticuloceras stubblefieldi</i> , <i>H.</i> cf. <i>striolatum</i>  |
| Upper part of shale quarry, Earle's cement works, southern slope of the Folly, Hope    | 417000  | 382600       | <i>Reticuloceras stubblefieldi</i> (type specimen)  |
| <i>Reticuloceras reticulatum</i> Marine Band (R1c1-3)                                  |         |              |   |
| Eccup - 4.5mils NNW of Leeds - Yorkshire   | 428000  | 442000       | <i>Reticuloceras reticulatum</i>  |
| Stanbury, adit spoil, Sladen Bridge, Stanbury  | 401800  | 437200       | <i>Reticuloceras reticulatum</i> (type form)  |
| Woodfold Park Nab's Head   | 363720  | 428840       | <i>Reticuloceras</i> cf. <i>reticulatum</i>   |
| Clough - W of Ewood Hall - 1100yds NW of Todmorden Station                             | 392730  | 424720       | <i>Reticuloceras</i> cf. <i>reticulatum</i>   |
| Black Scout, Crimsworth Dean, Hebden Bridge  | 3988170 | 429820       | <i>Reticuloceras reticulatum</i>  |

|  |         |              |   |
|--|---------|--------------|---|
| Shewboard (or Shrewbroad) Clough, Todmorden                        | 393590  | 423700       | <i>Reticuloceras reticulatum</i>  |
| Shell Brook - Mareknowles - Staffordshire                          | 394770  | 365640       | <i>Reticuloceras reticulatum</i>  |
| Quarmby Clough Mills BH SE11NW/8                                   | 411453  | 4167370      | <i>Reticuloceras reticulatum</i>  |
| Greenway Hall Golf Course  | 391860  | 351270       | <i>Reticuloceras reticulatum</i>  |
| Grange Brickworks, Killinghall                                     | 428650  | 457700       | <i>Reticuloceras reticulatum</i> , <i>Vallites</i> sp., <i>V. striolatus</i>                        |
| Stockeld BH SE34NE/16  | 438030  | 449450       | <i>Reticuloceras reticulatum</i> , <i>Vallites. striolatus</i>                                      |
| Sabden Brook   | 374600  | 434500       | <i>Reticuloceras reticulatum</i> , <i>R. davis</i>  |
| LOCALITY   | EASTING | NORTHIN<br>G | AMMONOIDS   |
| <i>Reticuloceras reticulatum</i> Marine Band (R1c1-3) Continued    |         |              |   |
| Westfield Mills BH, Yeadon SE24SW/1                                | 420450  | 440940       | <i>Reticuloceras</i> cf. <i>reticulatum reticulatum</i>   |
| Co-operative Laundry SE04SE 6 BH 49.99-51.51m depth                | 405800  | 441200       | <i>Reticuloceras</i> cf. <i>reticulatum reticulatum</i>   |
| Samlesbury - River Darwen  | 361830  | 429160       | <i>Reticuloceras reticulatum</i>  |
| Samlesbury - River Darwen  | 362410  | 428640       | <i>Reticuloceras reticulatum</i> , <i>R. davis</i> , <i>Homoceras striolatum</i>                    |
| Samlesbury - River Darwen 300 yards W of Beardwood Hall            | 365960  | 428840       | <i>Reticuloceras</i> cf. <i>reticulatum</i> , <i>R. cf. regularum</i> , <i>Homoceras striolatum</i> |
| Stream below Wimberry Stone  | 401500  | 402500       | <i>Reticuloceras reticulatum</i> type   |
| River Noe  | 409250  | 385760       | <i>Reticuloceras reticulatum</i>  |
| Swint Clough - Alport Valley - Derbyshire                          | 413470  | 391060       | <i>Reticuloceras reticulatum</i> ss.  |
| Eyam View  | 421420  | 377190       | <i>Reticuloceras reticulatum</i> ss.  |
| South Leverton No 1 BH   | 479330  | 380400       | <i>Reticuloceras</i> cf. <i>reticulatum</i>   |
| Biddulph, 265 yds E of Heath Hay                                   | 390770  | 359280       | <i>Homoceratoides prereticulatus</i>  |
| Franker Brook  | 430510  | 348120       | <i>Reticuloceras</i> cf. <i>reticulatum</i>   |
| <i>Reticuloceras coreticulatum</i> Marine Band (R1c4)              |         |              |   |
| Pendle Water, right bank, Rough Lee, Forest of Pendle              | 384000  | 437000       | <i>Reticuloceras coreticulatum</i>  |
| Sabden Brook - 1250' N of confluence with River Calder             | 374600  | 434400       | <i>Reticuloceras</i> aff. <i>coreticulatum</i>  |
| Ponden Clough - 550yds upstream from Ponden Reservoir - Stanbury   | 398700  | 436700       | <i>Reticuloceras coreticulatum</i> , <i>R. reticulatum</i>  |
| Heysham Power Station BH SD45NW/229                                | 340450  | 459890       | <i>Reticuloceras coreticulatum</i> , <i>Homoceratoides divaricatus</i> , <i>R. reticulatum</i>      |
| Co-operative Laundry BH SE04SE/6 49.99-51.51m depth                | 405800  | 441200       | <i>Reticuloceras coreticulatum</i> , <i>R. reticulatum</i> , <i>Hudsonoceras ornatum</i> ,          |
| Bradup BH SE04SE/774 97.85-99.85m                                  | 409140  | 444170       | <i>Reticuloceras coreticulatum</i>  |
| Westfield Mills BH - Yeadon SE24SW/1                               | 420450  | 440940       | <i>Reticuloceras reticulatum</i> late form  |
| Clough Hole  | 401800  | 436800       | <i>Reticuloceras reticulatum</i>  |
| Wike Whin 1.25 miles WSW of Bardsey church                         | 434500  | 442200       | <i>Reticuloceras coreticulatum</i> , <i>R. reticulatum</i>  |
| Callow BH  | 426650  | 352820       | <i>Reticuloceras</i> spp., <i>R. coreticulatum</i>  |
| <i>Bilinguites gracilis</i> Marine Band (R2a1)                     |         |              |   |
| Bankfield Mills BH, Mold Green SE11NE/11                           | 414660  | 416270       | <i>Reticuloceras gracile</i>  |
| Foster Clough - Mytholmroyd Station                                | 401880  | 427210       | <i>Reticuloceras gracile</i>  |
| Mount Road, Pule Hill, 400yds E by N of Gilberts Farm, Marsden     | 403160  | 410120       | <i>Reticuloceras gracile</i>  |
| Birchover Borehole, Buxton SK26SW/16                               | 424130  | 362330       | <i>Bilinguites gracile</i>  |
| Yeadon Waterworks BH - SE24SW/14                                   | 422410  | 442470       | <i>Reticuloceras gracile</i> late mut. alpha  |
| Holme Woods Dike - 130yds S of Holme Woods - 1mile S of Holme      | 410460  | 404450       | <i>Reticuloceras gracile</i>  |
| Grinding Stone Hole - Rag Clough - W of Church - Oxenhope          | 401400  | 433800       | <i>Reticuloceras gracile</i>  |
| Alum Crag - NNE of Alum Scar - 1.1/8ml SE of Chapel - Nabs Head    | 363680  | 428050       | <i>Reticuloceras gracile</i> late mut.  |
| Butts Clough - 100yds NE of Rishworth Mills                        | 403750  | 418020       | <i>Reticuloceras gracile</i> , <i>R. reticulatum</i>  |
| Rake Dike - Holme - Holmfirth - Yorkshire                          | 409980  | 405210       | <i>Reticuloceras gracile</i>  |
| Sun Hill Clough - Oxenhope   | 400600  | 434000       | <i>Reticuloceras gracile</i>  |
| Long Ridge - 630yds SSE of North Grain with Howels Head Clough     | 404940  | 403940       | <i>Reticuloceras reticulatum</i> mut. <i>gracile</i>  |
| Star Wood, 1ml NE of Oakamoor                                      | 406120  | 346080       | <i>Reticuloceras gracile</i> (2 bands)  |
| Bank of Salter's Brook, 550 SSW of Salter's Brook Bridge, Woodhead | 413510  | 399660       | <i>Reticuloceras reticulatum</i> mut. alpha   |
| Greenway Hall Golf Course  | 391860  | 351270       | <i>Bilinguites gracilis</i>   |
| Heysham Power Station BH SD45NW/229                                | 340450  | 459890       | <i>Bilinguites gracilis</i>   |

|   |         |              |  |
|---|---------|--------------|--|
| Seat Hall BH SD66NE/2   | 366030  | 469820       | <i>Bilinguites gracilis</i>  |
| Tittesworth Farm  | 400040  | 358740       | <i>Bilinguites gracilis</i> (Lower and both middle bands of Marine Band)   |
| River Churnet north of Swainsmoor   | 402410  | 361900       | <i>Bilinguites gracilis</i> (Both middle bands of Marine Band)   |
| Boreholes near Brund  | 409630  | 361780       | <i>Bilinguites gracilis</i> (3 bands of Marine Band: <i>Reticuloceras</i> sp. nov, in Upper and lower Middle band) |
| NE of Pilsley   | 423260  | 371750       | <i>Bilinguites gracilis</i> late form  |
| Newton Bank BH  | 395820  | 395060       | <i>Bilinguites gracilis</i> late form  |
| Park Clough, Hey Green, Marsden   | 402990  | 412460       | <i>Bilinguites gracilis</i>  |
| Farnham BH SE35NE 27  | 434690  | 459960       | <i>Bilinguites gracilis</i>  |
| LOCALITY  | EASTING | NORTHIN<br>G | AMMONOIDS  |
| <i>Bilinguites gracilis</i> Marine Band (R2a1) Continued                  |         |              |  |
| Sabden Brook  | 374600  | 434300       | <i>Reticuloceras gracile</i> late form   |
| Rams Clough   | 391000  | 432100       | <i>Reticuloceras gracile</i> late form   |
| Bradup BH SE04SE/774 23.05-24.49m (upper band); 26.15-28.48m (lower band) | 409140  | 444170       | <i>Reticuloceras gracile</i> type & late form (lower band) and early (upper band)                                  |
| Aire Valley BH 29 SE04SE/17 35.5-37.0m                                    | 408990  | 440570       | <i>Reticuloceras gracile</i> early form  |
| Westfield Mills BH - Yeadon SE24SW/1                                      | 420450  | 440940       | <i>Reticuloceras gracile</i> late mut. alpha   |
| Kirk Lane Dye Works BH SE24SW/4a  | 420350  | 441020       | <i>Reticuloceras gracile</i> late mut. alpha   |
| Horsforth Water Works BH SE24SW/7a  | 423370  | 441160       | <i>Reticuloceras gracile</i> late mut. alpha   |
| Junction of Bent & Middle Moor cloughs 800 yards N of High Greave         | 399200  | 433600       | <i>Reticuloceras reticulatum</i> early mut. alpha  |
| Paul Clough 640 yards SE of Aberdeen                                      | 403100  | 433900       | <i>Reticuloceras reticulatum</i> late mut. alpha & mut. alpha  |
| Sough Hole 330 yards NW of Two Laws                                       | 397400  | 438300       | <i>Reticuloceras reticulatum</i> mut. alpha  |
| Victoria Hospital, Keighley 1040 yards SW of Cliffe Castle                | 405300  | 441500       | <i>Reticuloceras reticulatum</i> late mut. alpha   |
| Brickpit (Park Wood Brick co) 530yds S 6degs S of station, Keighley       | 406600  | 440700       | <i>Reticuloceras reticulatum</i> cf. late mut alpha, <i>R. reticulatum</i> (2 bands)                               |
| Snail Green BH  | 411800  | 442500       | <i>Reticuloceras gracile</i> mut. alpha  |
| Banksfield Dye Works BH   | 420700  | 441500       | <i>Reticuloceras gracile</i> aff. mut alpha  |
| Corringham No 3 BH  | 489050  | 393520       | <i>Bilinguites gracilis</i>  |
| Trumfleet No 1 BH SE51SE/1  | 460520  | 412640       | <i>Bilinguites gracilis</i>  |
| Moss Oil BH SE51SE/19   | 459980  | 413900       | <i>Bilinguites gracilis</i> , <i>B. gracilis</i> (early form)  |
| W bank of Long Clough, c 400yds W of bridge on Glossop-Hayfield road      | 403020  | 390730       | <i>Reticuloceras gracile</i>   |
| The Heys  | 404390  | 385540       | <i>Reticuloceras gracile</i>   |
| Whitehall Works BH  | 403550  | 382020       | <i>Reticuloceras gracile</i>   |
| Forge Works No. 3 BH  | 404170  | 382190       | <i>Reticuloceras gracile</i>   |
| Clough, 0.25 miles NNE of Ridge Hall                                      | 405950  | 379250       | <i>Reticuloceras</i> cf. <i>gracile</i>  |
| 300 yds NW of Cowlow Farm   | 406620  | 378780       | <i>Reticuloceras gracile</i> [evolute form]  |
| Stream near Dunge Farm  | 398900  | 377690       | <i>Reticuloceras gracile</i>   |
| S bank of stream 1 mile NW of Hallfield                                   | 423300  | 392890       | <i>Reticuloceras gracile</i>   |
| E bank of stream, W of Bole Edge Plantation                               | 422590  | 392000       | <i>Reticuloceras</i> cf. <i>gracile</i> and late form  |
| N side of Raddlepit Rushes in Strines Dike                                | 421040  | 389740       | <i>Reticuloceras gracile</i> late form   |
| Moscar Moor   | 422230  | 387360       | <i>Reticuloceras gracile</i> late form   |
| 210 yds NNE of Mitchell Field, E of Hathersage                            | 424870  | 381920       | <i>Reticuloceras gracile</i>   |
| Leeswood Old Hall BH  | 326360  | 361800       | <i>Bilinguites gracilis</i>  |
| Coed-y-felin stream section   | 322370  | 371500       | <i>Bilinguites gracilis</i>  |
| North Rode, N bank of R. Dane, 760 yds S by E of Ladderstile              | 390240  | 365310       | <i>Reticuloceras gracile</i> early and late forms  |
| Rushton, Dingle Brook, 350 yds SE by E of Harper's Farm                   | 392250  | 361760       | <i>Reticuloceras gracile</i> early and late forms  |
| Biddulph, 250 yds E by S of Heath Hay                                     | 390740  | 359220       | <i>Reticuloceras gracile</i> early form  |
| Abbey Mills BH 4  | 319490  | 377470       | <i>Reticuloceras gracile</i> late form   |
| Ashover, 1033 yds S20E of Raven House                                     | 435570  | 360270       | <i>Reticuloceras gracile</i> late form   |
| Tansley BH - Derbyshire   | 433126  | 359604       | <i>Reticuloceras gracile</i> , and early form  |
| Bothamsall No 1 BH  | 465860  | 373675       | <i>Reticuloceras gracile</i>   |
| Kelham Hills No 51 BH   | 476480  | 357500       | <i>Reticuloceras gracile</i>   |
| Kelham Hills No 1 BH  | 475940  | 357620       | <i>Reticuloceras gracile</i> late form   |
| Eakring No 1 BH   | 467600  | 361330       | <i>Reticuloceras gracile</i> late form   |
| Eakring No 3 BH   | 467710  | 361450       | <i>Reticuloceras gracile</i>   |
| Felthouse Wood  | 397900  | 350200       | <i>Reticuloceras gracile</i>   |
| Rotherwood BH   | 434580  | 315590       | <i>Bilinguites</i> cf. <i>gracilis</i>   |

|   |         |           |  |
|---|---------|-----------|--|
| Callow BH   | 426650  | 352820    | <i>Reticuloceras gracilis</i>  |
| Shottlegate   | 431370  | 347410    | <i>Reticuloceras gracilis</i>  |
| Load Clough, 700yds S by E of Luddenden Parish Church                               | 404400  | 425590    | <i>Reticuloceras reticulatum</i>   |
| Clark Bridge Mills - Halifax  | 409846  | 425166    | <i>Reticuloceras reticulatum</i>   |
| Butterley Clough - W bank of Gorge at top of Clough - Swellands Reservoir - Marsden | 404080  | 409010    | <i>Reticuloceras reticulatum</i> mut alpha   |
| Worthington BH SK24SW/204   | 440450  | 321040    | <i>Bilinguites gracilis</i>  |
| Asfordby Hydro BH SK72SW/71   | 472520  | 320610    | <i>Bilinguites</i> sp.   |
| <b><i>Bilinguites bilinguis</i> Marine Band (R2b1-3)</b>                            |         |           |  |
| Dry Clough, Warm Withens, Rishworth Moor, W of Rishworth                            | 399030  | 417750    | <i>Reticuloceras reticulatum</i> late mut alpha, <i>Bilinguites bilinguis</i> (R2b1-2) |
| Netherend's Beck - right bank - Sowerby   | 404110  | 422270    | <i>Reticuloceras bilingue</i> (R2b1-2)   |
| LOCALITY  | EASTING | NORTHIN G | AMMONOIDS  |
| <b><i>Bilinguites bilinguis</i> Marine Band (R2b1-3) Continued</b>                  |         |           |  |
| Phoenix Mills BH Huddersfield   | 414940  | 417500    | <i>Reticuloceras reticulatum</i> mut. <i>bilingue</i> (R2b1-2)                         |
| W bank of Long Clough, c 400yds W of bridge on Glossop-Hayfield road                | 403020  | 390730    | <i>Reticuloceras bilingue</i> early form (R2b1-2)                                      |
| Section in cut of bank 150yds N 31degs of High Cote, Riddlesden                     | 406800  | 443000    | <i>Reticuloceras reticulatum</i> mut. <i>bilingue</i> (R2b1-2)                         |
| High Marcroft Fold - Near Rochdale  | 384100  | 414800    | <i>Reticuloceras bilingue</i> (R2b1-2)   |
| March Hill - N of Dobcross  | 400800  | 413270    | <i>Bilinguites bilinguis</i> (R2b1-2)  |
| Brickpit (Park Wood Brick co) 530yds S 6degs S of station, Keighley                 | 406600  | 440700    | <i>Reticuloceras reticulatum</i> mut. <i>bilingue</i> (R2b1-2)                         |
| Rake Dyke @ 12000' OD 1mile SW of Holme   | 409670  | 404980    | <i>Reticuloceras bilingue</i> (R2b1-2)   |
| Kitchen Clough - Slaithwaite  | 408150  | 413470    | <i>Reticuloceras bilingue</i> (R2b1-2)   |
| Bank of R Dane W of Swythanley Hall Church  | 396320  | 364520    | <i>Reticuloceras bilingue</i> (R2b1-2)   |
| Bankfield Mills BH - Mold Green - Huddersfield                                      | 414660  | 416270    | <i>Reticuloceras bilingue</i> (R2b1-2)   |
| Shale scar above right bank of River Derwent about 0.5ml SW of Beeley               | 425760  | 367010    | <i>Reticuloceras bilingue</i> (R2b1-2)   |
| Fairweather Green, Four Lane Ends, Bradford   | 413410  | 433350    | <i>Reticuloceras bilingue</i> (R2b1-2)   |
| Saltaire BH - NW of Bradford  | 414100  | 438000    | <i>Reticuloceras bilingue</i> (R2b1-2)   |
| 350yds SW of Lench House, Blackwood Rishworth                                       | 400700  | 417600    | <i>Reticuloceras reticulatum</i> mut. Beta (R2b1-2)                                    |
| Pike Clough, 300yds ENE of Pike Farm Rishworth                                      | 403210  | 417570    | <i>Bilinguites bilinguis</i> (R2b1-2)  |
| Upper Deanhead Clough, Scammonden   | 402630  | 414540    | <i>Reticuloceras reticulatum</i> mut. Beta (R2b1-2)                                    |
| Hard Head Clough - Shot Scar - ENE of March Haigh Reservoir - Marsden               | 402440  | 413250    | <i>Reticuloceras reticulatum</i> mut. Beta (R2b1-2)                                    |
| Old quarry, E side of Valley Road Slaithwaite                                       | 408150  | 413470    | <i>Reticuloceras reticulatum</i> mut. Beta (R2b1-2)                                    |
| Slaithwaite Railway Station   | 408160  | 415140    | <i>Reticuloceras reticulatum</i> mut. Beta (R2b1-2)                                    |
| 600yds S 23degs W of Hooley Hey Farm, 3 and 1/3rd miles SW of Taxal Church Cheshire | 397300  | 374810    | <i>Reticuloceras bilingue</i> early form (R2b1-2)                                      |
| Borehole at Phoenix Mills Huddersfield  | 414940  | 417500    | <i>Reticuloceras reticulatum</i> mut. <i>bilingue</i> (R2b1-2)                         |
| Black Sike 0.5ml SW of Upperthong, 1ml W of Holmfirth                               | 412170  | 408080    | <i>Reticuloceras reticulatum</i> mut. Beta (R2b1-2)                                    |
| SW end of Pule Hill, 1150ft OD, 400yds NE of Gilberts Farm W of Marsden             | 403240  | 410160    | <i>Reticuloceras bilingue</i> ; <i>R. reticulatum</i> mut. early Beta (R2b1-2)         |
| Kirk Lane Dyeworks BH   | 420350  | 441020    | <i>Reticuloceras reticulatum</i> early mut. Beta (R2b1-2)                              |
| Wittonstall Clough, 500yds NE Cornholme Station                                     | 391480  | 426670    | <i>Reticuloceras reticulatum</i> mut. Beta (R2b1-2)                                    |
| Paul Clough - Stiperden House - NE Portsmouth Station                               | 390960  | 427920    | <i>Reticuloceras reticulatum</i> mut. Beta (R2b2)                                      |
| Bagnall   | 393740  | 359230    | <i>Bilinguites bilinguis</i> ss. (R2b2)  |
| Greenway Hall Golf Course   | 391860  | 351270    | <i>Bilinguites bilinguis</i> (R2b1)  |
| Middleton Towers  | 340940  | 458660    | <i>Bilinguites</i> sp. juv. (ex. gr. <i>bilinguis</i> ) (R2b1)                         |
| Birchover Borehole, Buxton  | 424130  | 362330    | <i>Bilinguites bilinguis</i> early form (R2b1)   |
| River Churnet north of Swainsmoor   | 402410  | 361900    | <i>Bilinguites bilinguis</i> early and type (R2b1)                                     |
| Boreholes near Brund  | 409630  | 361780    | <i>Bilinguites bilinguis</i> early form and type (R2b1&2)                              |



|                                    |        |        |  |
|------------------------------------|--------|--------|--|
| NE of Pilsley                      | 423260 | 371750 | <i>Bilinguites bilinguis</i> early form, <i>Hudsonoceras ornatum</i> , <i>Reticuloceras</i> sp. (R2b1) |
| Newton Bank BH                     | 395820 | 395060 | <i>Bilinguites bilinguis</i>   |
| Park Clough, Hey Green, Marsden    | 402990 | 412460 | <i>Bilinguites bilinguis</i> early form (R2b1)   |
| NW of Black Bank                   | 375750 | 464800 | <i>Bilinguites bilinguis</i>   |
| Sabden Brook                       | 374600 | 434300 | <i>Reticuloceras bilingue</i>  |
| Aire Valley BH 26                  | 410190 | 439500 | <i>Bilinguites bilinguis</i>   |
| Snail Green BH                     | 411800 | 442500 | <i>Reticuloceras bilingue</i>  |
| Horsforth Water Works BH SE24SW/7a | 423370 | 441160 | <i>Reticuloceras bilingue</i>  |
| Red Brook 914m WNW of Lydgate Mill | 396160 | 416650 | <i>Bilinguites bilinguis</i> , <i>B. cf. circumplicatilis</i>  |

| LOCALITY   | EASTING | NORTHIN<br>G | AMMONOIDS   |
|--|---------|--------------|---|
| <i>Bilinguites bilinguis</i> Marine Band (R2b1-3) Continued                  |         |              |   |
| Moorley Clough 183m S of Rough Stones Farm                                   | 394070  | 420490       | <i>Bilinguites bilinguis</i> , <i>B. cf. circumplicatilis</i>       |
| 557m S31E of Cowall, Cowall Manor  | 390690  | 355160       | <i>Bilinguites bilinguis</i> (R2b1-2)                               |
| Broomhead Reservoir dam trench   | 426880  | 396000       | <i>Reticuloceras reticulatum</i> mut. Beta (R2b1-2)                 |
| Trumfleet No 2 BH  | 460330  | 412460       | <i>Bilinguites bilinguis</i>  |
| Whitehall Works BH   | 403550  | 382020       | <i>Reticuloceras bilingue</i> early form (R2b1)                     |
| Forge Works No. 3 BH   | 404170  | 382190       | <i>Reticuloceras bilingue</i> early form (R2b1)                     |
| Blackedge Reservoir  | 406730  | 376550       | <i>Reticuloceras bilingue</i> (R2b1)                                |
| Hogshaw Brook  | 405990  | 374230       | <i>Reticuloceras bilingue</i> early form & <i>bilingue</i> (R2b1&2) |
| Stream 0.25 miles ESE of Longhill Farm                                       | 403790  | 374830       | <i>Reticuloceras bilingue</i> (R2b1)                                |
| Stream near Dunge Farm   | 398930  | 377680       | <i>Reticuloceras bilingue</i> early form (R2b1)                     |
| Broughton Brook waterfall  | 331970  | 361590       | <i>Bilinguites cf. bilinguis</i> (R2b2)                             |
| Heaton, 300 yds N by E Bearda  | 396320  | 364510       | <i>Reticuloceras bilingue</i> (R2b2)                                |
| Heaton, S bank of R. Deane, 350 yds W by N Hollinhal                         | 395380  | 363880       | <i>Reticuloceras bilingue</i> (R2b2)                                |
| Heaton, S bank of R. Deane, 450 yds NE by E Wormhill                         | 394120  | 363490       | <i>Reticuloceras bilingue</i> early form (R2b1)                     |
| Rushton, Dingle Brook, 500 yds W by S Fold Farm                              | 392610  | 361330       | <i>Reticuloceras bilingue</i> early form (R2b1)                     |
| Heaton, 500 yds NE by N Fairboroughs   | 396020  | 361270       | <i>Reticuloceras bilingue</i>                                       |
| Horton, 320 yds SE Endon Hays  | 393100  | 360470       | <i>Reticuloceras bilingue</i> early form (R2b1)                     |
| Biddulph, 240 yds ESE Heath Hay  | 390720  | 359180       | <i>Reticuloceras bilingue</i>                                       |
| Beeley Brook, 570 yds S66E of St Anne's Church, Beeley                       | 426980  | 367450       | <i>Reticuloceras bilingue</i> (R2b2)                                |
| Lindup Wood, 1050 yds S48W of St Anne's Church, Beeley                       | 425790  | 367000       | <i>Reticuloceras bilingue</i> (R2b2)                                |
| Ravensnest Wood, 770 yds N82W of Raven House                                 | 434530  | 361270       | <i>Reticuloceras bilingue</i> (R2b2)                                |
| Hole Wood, 530 yds S9W of Raven House  | 435160  | 360670       | <i>Reticuloceras bilingue</i> (R2b2)                                |
| Tansley BH - Derbyshire  | 433126  | 359604       | <i>Reticuloceras bilingue</i> early form (R2b1)                     |
| Uppertown BH - Derbyshire  | 432370  | 364250       | <i>Reticuloceras bilingue</i> early form (R2b1)                     |
| South of Cheddleton  | 397980  | 350230       | <i>Reticuloceras bilingue</i> early form (R2b1)                     |
| Cotton Dell  | 406120  | 346080       | <i>Reticuloceras bilingue</i> ss. (R2b2)                            |
| Lumb Grange  | 433140  | 346750       | <i>Reticuloceras bilinguis</i>                                      |
| Croxteth Park BH   | 340300  | 394300       | <i>Reticuloceras bilingue</i> (R2b2)                                |
| Head of Doe Holes Clough, 200yds above Deanhead Clough, Scammonden           | 402910  | 414930       | <i>Reticuloceras reticulatum</i> mut. Beta                          |
| Crimble Clough - Slaithwaite - Yorkshire                                     | 408160  | 415110       | <i>Reticuloceras reticulatum</i> (R2b2)                             |
| Clark Bridge Mills BH - Halifax  | 409846  | 425166       | <i>Reticuloceras reticulatum</i>                                    |
| Old quarry - Varley Road/Mansergh House - Slaithwaite                        | 408150  | 413470       | <i>Reticuloceras reticulatum</i>                                    |
| Stream - 0.5ml NE of Warders Tower - Knypersley Reservoir - Crowborough Wood | 390670  | 355180       | <i>Reticuloceras reticulatum</i>                                    |
| Worthington BH SK24SW/204  | 440450  | 321040       | <i>Bilinguites</i> sp., <i>B. bilinguis</i> (R2b1)                  |
| Asfordby Hydro BH SK72SW/71  | 472520  | 320610       | <i>Bilinguites cf. bilingue</i>                                     |
| <i>Bilinguites eometabilinguis</i> Marine Band (R2b4)                        |         |              |   |
| South of Stake Gutter  | 402430  | 362980       | <i>Bilinguites eometabilinguis</i>                                  |
| Birchover Borehole, Buxton SK26SW/16   | 424130  | 362330       | <i>Bilinguites eometabilinguis</i>                                  |
| Nan Scar Beck - Sunny Bank - Yorkshire                                       | 403400  | 433300       | <i>Reticuloceras reticulatum</i> late mut. Beta                     |

|  |         |              |   |
|--|---------|--------------|---|
| Gingerbread Clough   | 406100  | 439500       | <i>Reticuloceras reticulatum</i>  |
| Cullingworth, Hewenden Valley  | 407820  | 436120       | <i>Reticuloceras reticulatum</i> late mut. Beta                                     |
| Stream, 500 yds SE Lion's Paw Farm, Knypersley   | 390300  | 355600       | <i>Bilinguites eometabilinguis</i> , <i>B. bilinguis</i> late form                  |
| Cotton Dell  | 406120  | 345770       | <i>Reticuloceras eometabilingue</i> , <i>R. metabilingue</i>                        |
| Carsington Aquaduct BH M8 SK25SE/62  | 425390  | 350190       | <i>Reticuloceras eometabilingue</i> , <i>R. bilingue</i>                            |
| LOCALITY   | EASTING | NORTHIN<br>G | AMMONOIDS   |
| <i>Bilinguites eometabilinguis</i> Marine Band (R2b4) Continued                          |         |              |   |
| Asfordby Hydro BH SK72SW/71  | 472520  | 320610       | <i>Bilinguites</i> cf. <i>eometabilingue</i>  |
| <i>Bilinguites metabilinguis</i> Marine Band (R2b5)                                      |         |              |   |
| Phoenix Mills BH Huddersfield  | 414940  | 317500       | <i>Reticuloceras reticulatum</i> cf. mut. <i>metabilingue</i>                       |
| 300yds NE of Higher Hempshaws - 2mils W of Belmont                                       | 365000  | 416500       | <i>Reticuloceras eometabilingue</i> , <i>Reticuloceras reticulatum metabilingue</i> |
| Fairweather Green BH, Four Lane Ends, Bradford   | 413380  | 433300       | <i>Reticuloceras bilingue</i> late form   |
| Horsforth UD Waterworks BH - N of Horsforth  | 423370  | 441160       | <i>Reticuloceras reticulatum</i> late mut. Beta                                     |
| Bagnall  | 393640  | 350530       | <i>Bilinguites</i> cf. <i>metabilinguis</i> <i>Verneulites sigma</i>                |
| Birchover Borehole, Buxton SK26SW/16   | 424130  | 362330       | <i>Bilinguites metabilinguis</i>  |
| Brownsett  | 399250  | 363690       | <i>Bilinguites metabilinguis</i>  |
| Aire Valley BH P22 SE03NE/9  | 409850  | 439750       | <i>Bilinguites bilinguis</i> late form  |
| Aire Valley BH A4 SE04SE/15  | 408920  | 440560       | <i>Bilinguites bilinguis</i> late form  |
| Aire Valley BH A1 SE04SE/12  | 408820  | 440360       | <i>Bilinguites bilinguis</i> late form, cf. <i>B. metabilinguis</i>                 |
| 200 yards ESE of Mould Greave, Marsh, near Oxenhope                                      | 402700  | 435400       | <i>Reticuloceras reticulatum</i> late mut. Beta                                     |
| Lees Moor BH, 230 yards W of Lower Height Bingley  | 406100  | 438000       | <i>Reticuloceras reticulatum</i> late mut. Beta                                     |
| Kirk Lane Dyeworks BH SE24SW/4a  | 420350  | 441020       | <i>Reticuloceras reticulatum</i> late mut Beta                                      |
| Old lead mines 0.5 miles E of Leicester Mill Quarry                                      | 362830  | 416360       | <i>Reticuloceras metabilingue</i> , <i>Gastrioceras sigma</i>                       |
| Ryal Fold, W of Darwen Hill 280 yards SSE of the SE end of Higher Roddlesworth Reservoir | 366200  | 421520       | <i>Reticuloceras metabilingue</i>   |
| Wiggins Teape No2 BH   | 361460  | 423440       | <i>Reticuloceras metabilingue</i>   |
| Star Paper Mills B BH  | 364700  | 424950       | <i>Reticuloceras metabilingue</i>   |
| 400 yds N16W of Ladymon (?Lady Moor) Gate  | 390300  | 355800       | <i>Bilinguites metabilinguis</i> , <i>B. bilinguis</i> late form                    |
| South end of tunnel, Scout, Scout Mill   | 397280  | 401230       | <i>Reticuloceras reticulatum</i> late mut. B & gamma                                |
| Stream 140 yds WNW of Carr Meadow, Derbyshire  | 403320  | 389510       | <i>Reticuloceras metabilingue</i>   |
| Stream W of Marl House, Derbyshire   | 403090  | 388800       | <i>Reticuloceras</i> cf. <i>bilingue</i> late form, <i>R. metabilingue</i>          |
| Blackedge Reservoir  | 406730  | 376550       | <i>Reticuloceras metabilingue</i>   |
| Rocher End Brook, 420yds at N50W from Bradfield Church                                   | 426000  | 392000       | <i>Reticuloceras bilingue</i> late form, <i>Homoceras?</i>                          |
| Rivelin valley close to Wolf Wheel 1160 yds S83E of Rails                                | 430200  | 387500       | <i>Reticuloceras bilingue</i> late form   |
| Macclesfield Forest, S bank of reservoir, 1400 yds NE Thickwithers                       | 396570  | 371300       | <i>Reticuloceras metabilingue</i>   |
| Rushton, 350 yds NE Oulton   | 396570  | 371300       | <i>Reticuloceras metabilingue</i>   |
| Lindup Wood, 1050 yds S48W of St Anne's Church, Beeley                                   | 425790  | 367000       | <i>Reticuloceras bilingue</i> late form   |
| 370 yds S75E of Cromford station   | 430650  | 357280       | <i>Reticuloceras bilingue</i> late form   |
| South of Cheddleton  | 398130  | 350430       | <i>Reticuloceras bilingue</i> late form   |
| Combes Valley  | 400360  | 351450       | <i>Reticuloceras metabilingue</i> and <i>R. bilingue</i> early form                 |
| Duffield railway cutting   | 434290  | 343640       | <i>Reticuloceras bilinguis</i> late form  |
| Ferriby Brook  | 438080  | 339780       | <i>Reticuloceras bilinguis</i> late form  |
| Butts Clough - 220yds SW Clough Head Farm - Rishworth                                    | 404300  | 417810       | <i>Reticuloceras reticulatum</i> mut. Beta  |
| Blake Clough - Blake Clough Farm - Slaithwaite Moors                                     | 405200  | 413610       | <i>Reticuloceras reticulatum</i> mut. Beta  |
| Worthington BH SK24SW/204  | 440450  | 321040       | <i>Bilinguites</i> sp., <i>B. bilinguis</i> , <i>B. metabilinguis</i>               |

|  |         |           |   |
|--|---------|-----------|---|
| Asfordby Hydro BH SK72SW/71  | 472520  | 320610    | <i>Bilinguites</i> sp.  |
| <i>Bilinguites superbilinguis</i> Marine Band (R2c1)                   |         |           |   |
| Pears House Clough, 850yds S of Strines Public House - Derbyshire      | 422520  | 389890    | <i>Reticuloceras superbilingue</i>  |
| Eagle Stone - site 825yds from at S5W- 8mls NNW of Chesterfield        | 426000  | 373000    | <i>Reticuloceras reticulatum</i> mut. <i>superbilingue</i>  |
| Rocher End Brook, 530yds at 344degs from Bradfield Church, Yorkshire   | 426000  | 392000    | <i>Reticuloceras superbilinguis</i> , <i>Gastrioceras</i> sp. nov., <i>G. cf. lineatum</i> , <i>Homoceratoides cf. divaricatus</i> , <i>H.</i> sp. nov. |
| LOCALITY   | EASTING | NORTHIN G | AMMONOIDS   |
| <i>Bilinguites superbilinguis</i> Marine Band (R2c1) Continued         |         |           |   |
| Newton Bank BH   | 395820  | 395060    | <i>Bilinguites superbilinguis</i>   |
| Will Moor Clough, 240 yards S of Antley Gate                           | 391900  | 436300    | <i>Reticuloceras superbilingue</i>  |
| Trawden Brook, below Lumb Spout Waterfall                              | 392000  | 437200    | <i>Reticuloceras superbilingue</i>  |
| Aire Valley BH A1SE04SE/12   | 408820  | 440360    | <i>Bilinguites</i> ghosts   |
| Horsforth Water Works BH SE24SW/7a                                     | 423370  | 441160    | <i>Reticuloceras superbilingue</i>  |
| Haworth Moor at Near Fosse Intake 170 yards ESE of Withins             | 398300  | 435400    | <i>Reticuloceras reticulatum</i> mut. gamma   |
| Blue Scar Beck 550 yards NNE of Clough Hey                             | 400000  | 439900    | <i>Reticuloceras reticulatum</i> mut. gamma   |
| Baildon Holmes 1100 yards S67E of Baildon Green church                 | 415400  | 438200    | <i>Reticuloceras reticulatum</i> mut. gamma   |
| Sydney Works BH, Fairweather Green                                     | 413400  | 433300    | <i>Reticuloceras superbilingue</i>  |
| Sandoz Chemical Co. BH   | 416300  | 434300    | <i>Reticuloceras superbilingue</i>  |
| New Lane Mills BH Laisterdyke  | 419100  | 432700    | <i>Reticuloceras superbilingue</i>  |
| River Yarrow 300 yards NE of Hemphshaws                                | 365000  | 416500    | <i>Reticuloceras superbilingue</i> , <i>Gastrioceras sigma</i>  |
| Ryal Fold, west of Darwen Hill   | 366260  | 421700    | <i>Reticuloceras superbilingue</i>  |
| Wiggins Teape No2 BH Withnell Fold                                     | 361460  | 423440    | <i>Reticuloceras superbilingue</i>  |
| Mossley Sewage Works 450 yards S of Scout                              | 397200  | 400800    | <i>Reticuloceras reticulatum</i> mut. gamma   |
| Askern Oil BH SE51NE/1   | 456520  | 415020    | <i>Bilinguites superbilinguis</i>   |
| Moss Oil BH SE51SE/19  | 459980  | 413900    | <i>Bilinguites superbilinguis</i>   |
| Trumfleet No 1 BH  | 460520  | 412640    | <i>Bilinguites superbilinguis</i>   |
| Trumfleet No 2 BH  | 460330  | 412460    | <i>Bilinguites superbilinguis</i> , <i>Verneulites sigma</i>  |
| Tributary of Long Clough, Derbyshire                                   | 402710  | 390980    | <i>Reticuloceras superbilingue</i> , <i>Donetzoceras cf. sigma</i> , <i>Gastrioceras</i> , <i>Homoceratoides cf. divaricatus</i>                        |
| Heylee, Spire Hollins  | 403200  | 378300    | <i>Reticuloceras superbilingue</i> , <i>Gastrioceras</i> spp., <i>Homoceratoides divaricatus</i>  |
| Pyegreave Brook  | 404880  | 378060    | <i>Reticuloceras superbilingue</i>  |
| Castle Naze  | 404820  | 378570    | <i>Reticuloceras superbilingue</i> , <i>Gastrioceras</i> spp., <i>Homoceratoides fortelirifer</i>   |
| Lightwood Reservoir  | 405470  | 375290    | <i>Reticuloceras</i> sp.  |
| Yarncliff Wood   | 425140  | 379100    | <i>Donetzoceras sigma</i> , <i>Reticuloceras superbilinguis</i>   |
| Hallam Head BH   | 430091  | 389123    | <i>Donetzoceras sigma</i> , <i>Reticuloceras superbilinguis</i>   |
| Broughton Brook, SE of Corn Mill                                       | 331640  | 365410    | <i>Bilinguites superbilinguis</i>   |
| Sutton, 500 yds NE by N of Langley Print Works                         | 394350  | 371740    | <i>Reticuloceras superbilingue</i>  |
| Uppertown BH - Derbyshire  | 432370  | 364250    | <i>Reticuloceras bilingue</i> early form  |
| 3050 yds N5E of Chatsworth House                                       | 426510  | 373030    | <i>Reticuloceras superbilingue</i> , <i>Gastrioceras</i> spp. <i>Homoceratoides fortelirifer</i>  |
| Jumbel (or Jumble) Coppice - Baslow; 2230 yds N25E of Chatsworth House | 426830  | 372090    | <i>Reticuloceras superbilingue</i> , <i>Gastrioceras</i> spp. <i>Homoceratoides fortelirifer</i>  |
| Bassetbarn Farm BH   | 435540  | 364160    | <i>Reticuloceras superbilingue</i>  |
| Bothamsall No 1 BH   | 465860  | 373675    | <i>Reticuloceras cf. superbilingue</i>  |
| Shirley Hollow   | 403790  | 348090    | <i>Reticuloceras superbilingue</i>  |
| Rotherwood BH  | 434580  | 315590    | <i>Bilinguites superbilinguis</i>   |
| W. of Hankin Farm  | 432140  | 354480    | <i>Reticuloceras superbilingue</i>  |
| Ambergate railway cutting  | 434670  | 350800    | <i>Reticuloceras superbilingue</i>  |
| Blackfordby No. 1 BH   | 432250  | 318270    | <i>Reticuloceras reticulatum</i> mut. gamma   |
| Bottonley Clough - 140yds E of Bottoms Farm                            | 406340  | 419190    | <i>Reticuloceras reticulatum</i>  |

|   |         |           |  |
|---|---------|-----------|--|
| - Barkisland  |         |           |  |
| Junction of Streams - Heath House Wood - 150yds N of Heath House - Golcar               | 408990  | 415810    | <i>Reticuloceras reticulatum</i>   |
| Stream bank below Haslingden-Helmshore Road - 850yds NW of Helmshore Sation             | 377760  | 421680    | <i>Reticuloceras reticulatum</i>   |
| Worthington BH SK24SW/204   | 440450  | 321040    | <i>Bilinguites superbilinguis</i> , <i>Gastrioceras</i> sp.  |
| Asfordby Hydro BH SK72SW/71   | 472520  | 320610    | <i>Bilinguites superbilinguis</i> , cf. <i>Gastrioceras</i> sp.  |
| <b>Verneulites sigma Marine Band (R2c2)</b>   |         |           |  |
| Oxspring Borehole, SE of Oxspring SE20SE/6  | 427870  | 401360    | <i>Gastrioceras</i> ? <i>sigma</i> , <i>Reticuloceras reticulatum</i> mut. <i>superbilingue</i>  |
| LOCALITY  | EASTING | NORTHIN G | AMMONOIDS  |
| <b>Verneulites sigma Marine Band (R2c2) Continued</b>                                   |         |           |  |
| Stream bank, NNW of Bromiley, Belmont   | 365780  | 417920    | <i>Gastrioceras sigma</i>  |
| NE of Higher Hempshaws, W of Belmont  | 365000  | 416500    | <i>Gastrioceras</i> ? <i>sigma</i>   |
| Burbage Brook, Grindleford Station, Derbys.   | 425030  | 378690    | <i>Gastrioceras</i> ? <i>sigma</i>   |
| 100yds N of Chatsworth House  | 426210  | 371180    | <i>Gastrioceras</i> ? <i>sigma</i>   |
| Anglezarke Reservoir, Lancashire  | 362200  | 415900    | <i>Gastrioceras sigma</i> , <i>Homoceratoides</i>  |
| Ryal Fold, west of Darwen Hill  | 366300  | 421490    | <i>Pygmaeoceras sigma</i>  |
| Wiggins Teape No2 BH  | 361460  | 423440    | <i>Gastrioceras</i> sp.  |
| 200 yards SE of Canyards  | 425900  | 395100    | <i>Reticuloceras reticulatum</i> early mut. gamma  |
| Raynor Clough 800 yds E25N of White Lea   | 427500  | 395500    | <i>Reticuloceras reticulatum</i> early mut. gamma, <i>Gastrioceras</i> ? <i>sigma</i>  |
| More Hall (or Hull) Reservoir - trial hole for wing trench - 900yds NW of Brightholmlee | 428390  | 395570    | <i>Gastrioceras</i> cf. <i>cumbriense</i> , G. cf. <i>crenulatum</i> , G. spp, <i>Reticuloceras reticulatum</i> mut. gamma, <i>Gastrioceras</i> ? <i>sigma</i> |
| Stream N of Bankvale Mill   | 403110  | 387660    | <i>Donetzoceras sigma</i> , <i>Reticuloceras superbilingue</i>   |
| River between Strines & Dale Dike reservoirs  | 423360  | 390660    | <i>Donetzoceras sigma</i>  |
| Pear House Clough   | 422830  | 388960    | <i>Donetzoceras sigma</i>  |
| Callow Bank   | 425190  | 382290    | <i>Donetzoceras sigma</i>  |
| Yarnclyff Wood  | 425140  | 379100    | <i>Donetzoceras sigma</i>  |
| Winkle, 1300 yds SE Sutton End  | 396630  | 367980    | <i>Donetzoceras sigma</i>  |
| Rushton, 810 yds SE by E of The Cloud   | 391060  | 363200    | <i>Donetzoceras sigma</i>  |
| Biddulph, 180 yds S by W of Heath Hay   | 390480  | 359110    | <i>Donetzoceras sigma</i>  |
| Biddulph, 400 yds NNW of Cowall   | 390250  | 356030    | <i>Donetzoceras sigma</i>  |
| 3050 yds N5E of Chatsworth House  | 426150  | 373030    | <i>Pygmaeoceras sigma</i>  |
| 2230 yds N25E of Chatsworth House   | 426830  | 372090    | <i>Pygmaeoceras sigma</i> , <i>Gastrioceras</i> sp.  |
| Bassetbarn Farm BH  | 435540  | 364160    | <i>Pygmaeoceras sigma</i> , <i>Gastrioceras</i> sp., <i>Reticuloceras superbilingue</i>  |
| Combes Valley   | 401350  | 351010    | <i>Donetzoceras sigma</i>  |
| Shirley Hollow  | 403790  | 348090    | <i>Donetzoceras sigma</i>  |
| Blackfordby No. 1 BH  | 432350  | 318270    | <i>Gastrioceras</i> ? <i>sigma</i>   |
| <b>Cancelloceras cancellatum Marine Band (G1a1)</b>                                     |         |           |  |
| Royshaw Brick Works, Blackburn Station  | 368250  | 429500    | <i>Gastrioceras cancellatum</i> var. <i>crencellatum</i> ; <i>Reticuloceras reticulatum</i>  |
| Section in Dean brook, 150yds S of Higher House, 2 miles W of Belmont                   | 364350  | 415380    | <i>Gastrioceras cancellatum</i> , G. cf. <i>crencellatum</i> , R. <i>superbilingue</i>   |
| Crowborough Wood Warders Tower, Staffs.   | 390100  | 355520    | <i>Gastrioceras</i> cf. <i>cancellatum</i>   |
| Nant Figillt Farm - Rhosesmor - Flints  | 320910  | 368000    | <i>Cancelloceras cancellatum</i> , C. <i>crencellatum</i> , C. sp., <i>Homoceratoides divaricatum</i> , <i>Reticuloceras reticulatum</i> mut <i>alpha</i>      |
| Wall Grange Brick Pit - Staffs.   | 396440  | 353220    | <i>Gastrioceras cancellatum</i>  |
| Bowsey Wood BH SJ74NE/9   | 376950  | 346430    | <i>Gastrioceras</i> cf. <i>cancellatum</i>   |
| Heysham Power Station BH SD45NW/87  | 340260  | 459940    | <i>Cancelloceras cancellatum</i> , C. <i>crencellatum</i> , C. sp., <i>Homoceratoides divaricatum</i> , <i>Reticuloceras reticulatum</i> mut <i>alpha</i>      |
| Seat Hall BH SD66NE/2   | 366030  | 469820    | <i>Cancelloceras cancellatum</i> , C. <i>branneroides</i>  |
| River Greta   | 361700  | 472230    | <i>Gastrioceras</i> cf. <i>cancellatum</i> , C. <i>crencellatum</i>  |
| Newton Bank BH  | 395820  | 395060    | <i>Gastrioceras crencellatum</i>   |
| Harrop Brook  | 395990  | 378460    | <i>Gastrioceras crencellatum</i>   |

|   |         |              |   |
|---|---------|--------------|---|
| Bollington Print Works BH                                     | 393980  | 377970       | <i>Gastrioceras cancellatum</i>   |
| Orchard Farm  | 402260  | 369030       | <i>Cancelloceras cancellatum</i>  |
| Waters Farm BH  | 375370  | 467630       | <i>Cancelloceras cancellatum</i> ,<br><i>Gastrioceras crencellatum</i> , <i>G. branneroides</i>                                     |
| Farnham BH SE35NE/27  | 434690  | 459960       | <i>Cancelloceras</i> cf. <i>cancellatum</i> ,<br><i>C. crencellatum</i>   |
| Monkroyd Beck, NNE of Monkroyd                                | 393400  | 441400       | <i>Gastrioceras crencellatum</i>  |
| Aire Valley BH 28 SE13NW/22                                   | 410400  | 439110       | <i>Cancelloceras cancellatum</i> ,<br><i>C. crencellatum</i>  |
| Aire Valley BH B52 SE13NE/29                                  | 415010  | 437880       | <i>Cancelloceras cancellatum</i> ,<br><i>C. crencellatum</i>  |
| Aire Valley BH A2 SE04SE/13                                   | 408870  | 440450       | <i>Cancelloceras crencellatum</i>   |
| LOCALITY  | EASTING | NORTHIN<br>G | AMMONOIDS   |
| <i>Cancelloceras cancellatum</i> Marine Band (G1a1) Continued |         |              |   |
| Horsforth Water Works BH SE04SE/13                            | 423370  | 441160       | <i>Cancelloceras cancellatum</i>  |
| Middle Moor Clough 600 yards SW of Upper Ponden               | 397600  | 436000       | <i>Cancelloceras cancellatum</i> ,<br><i>Gastrioceras crencellatum</i> , <i>R. superbilinguis</i>                                   |
| Bingley Brick Pit   | 411200  | 441200       | <i>Cancelloceras cancellatum</i> , <i>R. superbilinguis</i>   |
| Saltaire Mills  | 414100  | 438000       | <i>Cancelloceras cancellatum</i> , <i>R. superbilinguis</i>   |
| Yeadon Brick & Tile Works                                     | 419400  | 440900       | <i>Cancelloceras cancellatum</i> , <i>R. superbilinguis</i>   |
| West View, 330 yards N of Apperley Bridge Station             | 419600  | 438700       | <i>Cancelloceras cancellatum</i> , <i>R. superbilinguis</i>   |
| Sydney Works BH, Fairweather Green                            | 413400  | 433300       | <i>Cancelloceras cancellatum</i> ,<br><i>Gastrioceras crencellatum</i> , ? <i>G. sigma</i>  |
| New Lane Mills Laisterdyke                                    | 419100  | 432700       | <i>Gastrioceras crencellatum</i> , <i>G. cf. cumbriense</i>   |
| Summit Brickworks   | 394850  | 418730       | <i>Gastrioceras crencellatum</i> ,<br><i>Cancelloceras cancellatum</i>  |
| Ring road cutting N of Meanwood Hall                          | 428300  | 438600       | <i>Gastrioceras cancellatum</i> type  |
| 60 yds WNW of Hollinshead Hall 3 miles N of Belmont           | 366230  | 419920       | <i>Gastrioceras crencellatum</i> (upper fauna)  |
| 400 yds N10W of Wheelton crossroads                           | 359990  | 421510       | <i>Gastrioceras branneroides</i> (Bed A), <i>G. cancellatum</i> , <i>R. superbilingue</i> (Bed B-C), <i>G. crencellatum</i> (Bed D) |
| Howe Brook, SW of Chorley 1560 yds NNW of Wrightington Church | 352030  | 414860       | <i>Gastrioceras crencellatum</i> (upper fauna)  |
| Acres Brook   | 397000  | 397800       | <i>Gastrioceras cancellatum</i> , <i>R. reticulatum</i> mut. <i>gamma</i>   |
| Oxspring BH, SE of Oxspring SE20SE/6                          | 427870  | 401360       | <i>Gastrioceras cancelloceras</i> ,<br><i>Reticuloceras reticulatum</i> mut. <i>gamma</i>   |
| Trumfleet No 1 BH SE51SE/1                                    | 460520  | 412640       | <i>Cancelloceras cancellatum</i> , <i>C. crencellatum</i>   |
| Askern Oil BH SE51NE/1  | 456520  | 415020       | <i>Cancelloceras crencellatum</i>   |
| Belton Oil BH   | 477710  | 408460       | <i>Cancelloceras crencellatum</i>   |
| Fernilee No. 1 BH   | 401240  | 378230       | <i>Cancelloceras cancellatum</i> , <i>C. crencellatum</i> , <i>R. superbilingue</i>   |
| Mather Clough   | 397700  | 382130       | <i>Cancelloceras cancellatum</i> , <i>C. cf. crencellatum</i>   |
| Mill Clough   | 400240  | 378070       | <i>Cancelloceras crencellatum</i>   |
| Shooter's Clough  | 400570  | 374670       | <i>Cancelloceras cancellatum</i> , <i>R. superbilingue</i>  |
| Damflask Reservoir  | 427400  | 391100       | <i>Gastrioceras cancellatum</i> , <i>G. cf. crencellatum</i> , <i>Reticuloceras superbilinguis</i>                                  |
| Rod Moor No 3 BH  | 426780  | 389160       | <i>Gastrioceras crencellatum</i> ,<br><i>Reticuloceras superbilinguis</i>   |
| Carr Brook, 1490 yds N79E of Bassett, W of Fulwood            | 429700  | 384800       | <i>Gastrioceras crencellatum</i> , <i>G. rurae</i>  |
| Limb Brook, 1250 yds N88E of Barberfields Farm                | 430800  | 382900       | <i>Gastrioceras</i> cf. <i>cancellatum</i> , <i>G. rurae</i> ?  |
| Smekley No 3 BH   | 429690  | 376498       | <i>Gastrioceras cancellatum</i> , <i>G. crencellatum</i>  |
| Tickhill No 1 BH  | 457730  | 392970       | <i>Gastrioceras cancellatum</i>   |
| Walkeringham No 1 BH  | 475550  | 391900       | <i>Gastrioceras crencellatum</i>  |

|  |         |              |   |
|--|---------|--------------|---|
| Morton No 1 BH   | 479320  | 392410       | <i>Gastrioceras crencellatum</i>  |
| Apleyhead No 1 BH  | 465510  | 376310       | <i>Gastrioceras cf. crencellatum</i>  |
| Apleyhead No 2 BH  | 465770  | 376630       | <i>Gastrioceras cf. crencellatum</i>  |
| Bothamsall No 2 BH   | 465566  | 373917       | <i>Gastrioceras crencellatum</i>  |
| Bothamsall No 4 BH   | 466193  | 374022       | <i>Gastrioceras crencellatum</i>  |
| Fishpond Wood, stream section  | 332600  | 364310       | <i>Cancelloceras cancellatum</i>  |
| Warren Dingle  | 331790  | 362340       | <i>Cancelloceras cancellatum</i> , <i>Ca. branneroides</i> , <i>Ca. sp.</i> , <i>Bilinguites superbilinguis</i>           |
| Congleton, 660 yds NE by E of Timbersbrook crossroads                              | 390030  | 362980       | <i>Cancelloceras cancellatum</i> , <i>Bilinguites superbilinguis</i> , <i>Homoceratoides sp.</i>                          |
| Biddulph, 300 yds WSW of Heath Hay   | 390260  | 359140       | <i>Gastrioceras crencellatum</i>  |
| Biddulph, W bank of R. Trent, 320 yds SW by W of Cowall                            | 390110  | 355510       | <i>Cancelloceras cancellatum</i> , <i>Bilinguites superbilinguis</i>  |
| LOCALITY   | EASTING | NORTHIN<br>G | AMMONOIDS   |
| <i>Cancelloceras cancellatum</i> Marine Band (G1a1) Continued                      |         |              |   |
| Harewood Grange, N side of R. Hipper   | 431280  | 368080       | <i>Gastrioceras crencellatum</i>  |
| Walnut Opencast C11 BH, N bank of Carr Brook, 1970 yds N86E of Butterley Reservoir | 436360  | 360130       | <i>Gastrioceras cf. crencellatum</i> , <i>G. cf. rurae</i> , <i>G. sp.</i>  |
| Calow No 1 BH  | 440860  | 370410       | <i>Gastrioceras crencellatum</i>  |
| Bothamsall No 5 BH   | 466595  | 373440       | <i>Gastrioceras cf. crencellatum</i>  |
| Farley's Wood No 2 BH  | 470969  | 369997       | <i>Gastrioceras crencellatum</i> , <i>G. crencellatum</i> , <i>G. cf. cumbriense</i> , <i>cf. Agastrioceras carinatum</i> |
| Eakring No 1 BH  | 467600  | 361330       | <i>Gastrioceras crencellatum</i>  |
| Eakring No 3 BH  | 467710  | 361450       | <i>Gastrioceras crencellatum</i>  |
| Ruelow Wood BH SK04NW/5  | 402050  | 347520       | <i>Gastrioceras crencellatum</i> (2 bands)  |
| Rugeley (Trent Valley) BH  | 405080  | 319020       | <i>Gastrioceras crencellatum</i> , with <i>G. cancellatum</i> & <i>Reticuloceras superbilingue</i> in base                |
| Whittington Heath BH   | 414780  | 308000       | <i>Gastrioceras cancellatum</i> , <i>Reticuloceras superbilingue</i>  |
| Blackfordby No. 1 BH   | 432350  | 318270       | ? <i>Gastrioceras cancellatum</i>   |
| Sandoz Chemical Co.  | 416300  | 434300       | <i>Gastrioceras crencellatum</i> , with <i>G. rurae</i> in lower bed & <i>G. cf. carinatum</i> in upper                   |
| Winksley BH  | 425070  | 471510       | <i>Gastrioceras cancellatum</i>   |
| <i>Cancelloceras cumbriense</i> Marine Band (G1b1)                                 |         |              |   |
| N side of Willow railway cutting - NNE of Chorley Station                          | 359520  | 419250       | <i>Gastrioceras cumbriense</i> , <i>G. crenulatum</i>   |
| Bigrigg - Cumberland   | 300100  | 513050       | <i>Gastrioceras cumbriense</i>  |
| Mousegill - 120yds W + 20deg S of Swinestone House - Westmorland                   | 383680  | 512420       | <i>Gastrioceras cf. cumbriense</i>  |
| Horsforth UD Waterworks BH - 2mls N of Horsforth SE24SW/7a                         | 423370  | 441160       | <i>Gastrioceras cumbriense</i>  |
| River Greta  | 361700  | 472230       | <i>Cancelloceras cf. cumbriense</i> , <i>Homoceratoides sp.</i>   |
| Newton Bank BH   | 395820  | 395060       | <i>Gastrioceras cumbriense</i>  |
| Harrop Brook   | 395830  | 378450       | <i>Gastrioceras cumbriense</i> , <i>G. crenulatum</i> , <i>Homoceratoides aff. divaricatus</i>                            |
| Orchard Farm   | 402260  | 369030       | <i>Cancelloceras cumbriense</i>   |
| Waters Farm BH   | 375370  | 467630       | <i>Cancelloceras cumbriense</i> , <i>Gastrioceras crenulatum</i>  |
| Goat Gap Syke  | 371630  | 469970       | <i>Cancelloceras cumbriense</i> , <i>Gastrioceras crenulatum</i>  |
| Farnham BH SE35NE/27   | 434690  | 459960       | <i>Cancelloceras cumbriense</i> , <i>Gastrioceras crenulatum</i>  |
| High Lea Farm, Lower Trap  | 377200  | 435400       | <i>Cancelloceras cumbriense</i> , <i>Ca. crenulatum</i> , <i>Homoceratoides divaricatus</i>                               |
| Monkroyd Beck, NNE of Monkroyd   | 393400  | 441400       | <i>Gastrioceras cumbriense</i>  |
| North bank of Swinden Water  | 390600  | 433000       | <i>Gastrioceras cumbriense</i>  |
| Brook east of Combe Hill Cross   | 395800  | 438500       | <i>Gastrioceras cumbriense</i>  |
| Aire Valley BH 43 SE13NW/23  | 410270  | 439040       | <i>Cancelloceras cumbriense</i> , <i>Ca. crenulatum</i>   |
| Aire Valley BH 29 SE13NE/29  | 415010  | 437880       | <i>Cancelloceras cumbriense</i>   |
| Oaks Farm, Yeadon  | 424040  | 441200       | <i>Cancelloceras cumbriense</i> , <i>Ca. crenulatum</i>   |
| Middle Moor Clough 600 yards SW of Upper   | 397500  | 435900       | <i>Cancelloceras cumbriense</i> ,   |


|  |         |              |   |
|--|---------|--------------|---|
| Ponden   |         |              | <i>Gastrioceras crenulatum</i> , <i>R. superbilinguis</i>   |
| Gill Beck 400 yards N14E of Ash House Farm                         | 414600  | 441800       | <i>Cancelloceras cumbriense</i> , <i>Gastrioceras crenulatum</i>                                      |
| Yeadon Brick & Tile Works  | 419400  | 440900       | <i>Cancelloceras cancellatum</i> , <i>Gastrioceras crenulatum</i>                                     |
| Sydney Works BH, Fairweather Green                                 | 413400  | 433300       | <i>Cancelloceras cumbriense</i> , <i>Gastrioceras crenulatum</i>                                      |
| New Lane Mills Laisterdyke   | 419100  | 432700       | <i>Cancelloceras cancellatum</i> , <i>Gastrioceras crenulatum</i>                                     |
| Great Heads Wood, Roundhay Park                                    | 433600  | 438500       | <i>Gastrioceras cumbriense</i>  |
| Shore Brook 320 yds SE of Higher House, 2 miles W of Belmont       | 364520  | 415300       | <i>Gastrioceras cumbriense</i> , <i>G. crenulatum</i>   |
|  |         |              |   |
| LOCALITY   | EASTING | NORTHIN<br>G | AMMONOIDS   |
| <i>Cancelloceras cumbriense</i> Marine Band (G1b1) Continued       |         |              |   |
| Yarrow valley 900 yds W40S of Euxton Hall                          | 354730  | 417950       | <i>Gastrioceras cumbriense</i> , <i>G. crenulatum</i>   |
| Booth's Farm BH  | 354220  | 417310       | <i>Gastrioceras cumbriense</i>  |
| Oxspring Borehole, SE of Oxspring SE20SE/6                         | 427870  | 401360       | <i>Gastrioceras</i> cf. <i>crenulatum</i> , <i>G. cf. cumbriense</i>                                  |
| Moss Oil BH  | 459980  | 413900       | <i>Cancelloceras crenulatum</i> , <i>Ca. cumbriense</i>   |
| Trumfleet No 1 BH  | 460520  | 412640       | <i>Cancelloceras crenulatum</i> , <i>Ca. cumbriense</i>   |
| Askern Oil BH  | 456520  | 415020       | <i>Cancelloceras cumbriense</i>   |
| N of Rowarth   | 401610  | 389840       | <i>Gastrioceras cumbriense</i>  |
| Fernilee No. 1 BH  | 401240  | 378230       | <i>Gastrioceras cumbriense</i> , <i>G. crenulatum</i>   |
| Mather Clough  | 397740  | 382250       | <i>Gastrioceras cumbriense</i> , <i>G. crenulatum</i>   |
| Stream, 0.5 miles W by S of Handley Fold                           | 397400  | 380540       | <i>Gastrioceras cumbriense</i> , <i>G. crenulatum</i>   |
| Mill Clough  | 400240  | 378070       | <i>Gastrioceras cumbriense</i>  |
| Shooter's Clough   | 400570  | 374670       | <i>Gastrioceras cumbriense</i> , <i>Homoceratoides</i> aff. <i>divaricatus</i>                        |
| Holes Clough   | 423830  | 390380       | <i>Gastrioceras cumbriense</i> , <i>G. crenulatum</i> , <i>Homoceratoides</i> aff. <i>divaricatus</i> |
| Ughill Brook, 160 yds N7W of Corker Walls                          | 426100  | 390100       | <i>Gastrioceras cumbriense</i> , <i>G. crenulatum</i>   |
| Rod Moor No 3 BH   | 426780  | 389160       | <i>Gastrioceras cumbriense</i> , <i>G. crenulatum</i>   |
| Stream 633 yds N33E of Norfolk Arms, Ringinglow                    | 429400  | 384200       | <i>Gastrioceras cumbriense</i> , <i>G. crenulatum</i>   |
| Stream 130 yds N of chapel at Longshaw Lodge                       | 426400  | 380000       | <i>Gastrioceras cumbriense</i> , <i>G. crenulatum</i>   |
| Barr Brook, 30 ft below Rough Rock                                 | 427900  | 375000       | <i>Gastrioceras cumbriense</i> , <i>G. crenulatum</i>   |
| Smeekey No 3 BH SK27NE/2   | 429690  | 376498       | <i>Gastrioceras cumbriense</i> , <i>G. crenulatum</i>   |
| Tickhill No 1 BH   | 457730  | 392970       | <i>Gastrioceras cumbriense</i>  |
| Warren Dingle  | 332270  | 362320       | <i>Cancelloceras cumbriense</i> , <i>Ca. crenulatum</i>   |
| Leeswood Old Hall  | 326410  | 361300       | <i>Cancelloceras cumbriense</i> , <i>Ca. crenulatum</i>   |
| Congleton, E bank of gully, 560 yds ENE of Timbersbrook crossroads | 389990  | 362860       | <i>Cancelloceras</i> cf. <i>cumbriense</i> , <i>Ca. crenulatum</i>                                    |
| Biddulph, W bank of stream, 640 yds SW by S of Bridestones         | 390310  | 361760       | <i>Cancelloceras</i> cf. <i>cumbriense</i>  |
| Biddulph, SE bank of R. Trent, 390 yds SW of Cowall                | 390110  | 355420       | <i>Cancelloceras cumbriense</i> , <i>Ca. cf. crenulatum</i>   |
| Abbey Mills BH 4   | 319490  | 377470       | <i>Gastrioceras cumbriense</i>  |
| Harewood Grange, N side of R. Hipper                               | 431250  | 368110       | <i>Gastrioceras cumbriense</i>  |
| 220 yds SE of Lea Hall   | 433530  | 357370       | <i>Gastrioceras cumbriense</i> , <i>G. crenulatum</i> , <i>Homoceratoides</i> aff. <i>divaricatus</i> |
| Calow No 1 BH  | 440860  | 370410       | <i>Gastrioceras cumbriense</i> , <i>G. crenulatum</i>   |
| Ruelow Wood BH   | 402050  | 347520       | <i>Gastrioceras cumbriense</i> (2 bands)  |
| Beelow Hill  | 406620  | 345060       | <i>Gastrioceras cumbriense</i>  |
| Rugeley (Trent Valley) BH  | 405080  | 319020       | <i>Gastrioceras cumbriense</i>  |
| Blackfordby No. 1 BH   | 432350  | 318270       | <i>Gastrioceras cumbriense</i>  |

|   |         |              |   |
|---|---------|--------------|---|
| Sandoz Chemical Co. BH  | 416300  | 434300       | <i>Gastrioceras cumbriense</i> ,<br><i>Homoceratoides divaricatum</i> ; <i>G. sp.</i> , <i>G. carbonarium</i> in upper part |
| Asfordby Hydro BH SK72SW 71   | 472520  | 320610       | <i>Gastrioceras cf. crenulatum</i>  |
| <i>Gastrioceras subcrenatum</i> Marine Band (G2a1)                      |         |              |   |
| Ballavarish Bh, Shellag north Bh  | 246250  | 500700       | <i>Gastrioceras subcrenatum</i>   |
| Ridgeway Bh   | 389220  | 353810       | <i>Gastrioceras subcrenatum</i>   |
| River Greta   | 364360  | 472040       | <i>Gastrioceras subcrenatum</i>   |
| Stake Clough, NW of Goyt's Moss   | 400660  | 372910       | <i>Gastrioceras subcrenatum</i>   |
| 400 yds SW of Arnold Hill reservoir, Gee Cross                          | 395000  | 393000       | <i>Gastrioceras subcrenatum</i>   |
| LOCALITY  | EASTING | NORTHIN<br>G | AMMONOIDS   |
| <i>Gastrioceras subcrenatum</i> Marine Band (G2a1) Continued            |         |              |   |
| Bakestonedale BH  | 395940  | 379510       | <i>Gastrioceras sp.</i>   |
| Mere Burn   | 408860  | 554850       | <i>Gastrioceras subcrenatum</i>   |
| River Little Don, 1km E of Langsett (type section)                      | 422150  | 400410       | <i>Gastrioceras subcrenatum</i>   |
| Harrington No. 19 BH  | 299500  | 521000       | <i>Gastrioceras subcrenatum</i>   |
| St. Bees No. 4 BH   | 295000  | 512700       | <i>Gastrioceras subcrenatum</i>   |
| Whitehaven Laundry BH   | 297600  | 516800       | <i>Gastrioceras subcrenatum</i>   |
| Waters Farm BH  | 375370  | 467630       | <i>Gastrioceras subcrenatum</i>   |
| South of Blackwood Head, Wheatley Lane                                  | 383200  | 438200       | <i>Gastrioceras subcrenatum</i> ,<br><i>Reticuloceras superbilingue</i>   |
| South bank of Colne Water, 450 yards east of Carry Bridge               | 390000  | 439800       | <i>Gastrioceras cf. subcrenatum</i>   |
| Horsforth Water Works BH SE24SW/7a                                      | 423370  | 441160       | <i>Gastrioceras subcrenatum</i>   |
| Thornton Moor   | 405300  | 432600       | <i>Gastrioceras subcrenatum</i>   |
| Cottingley Moor Bridge  | 411200  | 436200       | <i>Gastrioceras subcrenatum</i>   |
| Thackley Tunnel, NW of Apperley Bridge                                  | 418400  | 438700       | <i>Gastrioceras subcrenatum</i>   |
| Newlay cutting  | 423200  | 436800       | <i>Gastrioceras subcrenatum</i>   |
| Top Mill BH, 400 yards NW of Alerton                                    | 411800  | 434400       | <i>Gastrioceras subcrenatum</i> , <i>G. listeri</i>   |
| Sydney Works BH, Fairweather Green                                      | 413400  | 433300       | <i>Gastrioceras carbonarium</i>   |
| Horsforth UD Waterworks BH - N of Horsforth                             | 423370  | 441160       | <i>Gastrioceras subcrenatum</i>   |
| Union Mills   | 418600  | 435600       | <i>Gastrioceras cf. subcrenatum</i>   |
| Alston Works BH   | 414600  | 433400       | <i>Gastrioceras subcrenatum</i>   |
| Britannia Mills BH  | 416400  | 432500       | <i>Gastrioceras subcrenatum</i>   |
| Sandoz Chemical Co. BH  | 416300  | 434300       | <i>Gastrioceras carbonarium</i>   |
| Globe Mills BH, Leeds, 400 yards S of City station SE23SE/7             | 4297990 | 4327870      | <i>Gastrioceras spp.</i> , <i>G. retrorsum</i> , <i>R. reticuloceras</i>  |
| North bank R. Darwen, 120 yds SE of Old Hall, Feniscowles               | 363780  | 425720       | <i>Gastrioceras subcrenatum</i>   |
| Stepback Brook, W of Darwen Hill, 1300 yds S42E of the inn at Ryal Fold | 367310  | 420650       | <i>Gastrioceras subcrenatum</i>   |
| 560 yds NNW of St Stephen's Church, Chapel                              | 365760  | 424000       | <i>Gastrioceras subcrenatum</i>   |
| Howe Brook, S of Brook House  | 352280  | 413980       | <i>Gastrioceras subcrenatum</i> , <i>G. sp.</i>   |
| Heskin BH   | 353890  | 414500       | <i>Gastrioceras subcrenatum</i> , <i>G. sp.</i>   |
| Oughtibridge, 200 yds W of station                                      | 431000  | 393500       | <i>Gastrioceras subcrenatum</i>   |
| Moss Oil BH SE51SE/19   | 459980  | 413900       | <i>Gastrioceras subcrenatum</i>   |
| Fernilee No. 2 BH   | 401190  | 378630       | <i>Gastrioceras subcrenatum</i>   |
| E of Fernilee   | 401910  | 378510       | <i>Gastrioceras subcrenatum</i>   |
| Shaw Farm BH, 0.5 miles NW of Eaves Knoll                               | 399050  | 386610       | <i>Gastrioceras subcrenatum</i> ,<br><i>Homoceratoides sp. G. sp.</i>   |
| Knowle Wood   | 398160  | 388660       | <i>Gastrioceras subcrenatum</i> , <i>G. sp.</i>   |
| S of Sugworth Road, near Moscar   | 424000  | 389500       | <i>Gastrioceras subcrenatum</i> , <i>G. sp.</i> , <i>G. sp. nov.</i>  |
| Smekley No 3 BH SK27NE/2  | 429690  | 376498       | <i>Gastrioceras subcrenatum</i>   |
| Tickhill No 1 BH  | 457730  | 392970       | <i>Gastrioceras subcrenatum</i> ,<br><i>Homoceratoides aff. divaricatus</i>   |
| Ranskill No 1 BH  | 464230  | 388140       | <i>Gastrioceras subcrenatum</i>   |
| Oakenholt Paper Mill BH   | 326280  | 375120       | <i>Gastrioceras subcrenatum</i> , <i>G. cf. retrorsum</i>   |
| Alders Farm BH  | 389540  | 362080       | <i>Gastrioceras subcrenatum</i>   |
| R. Hipper, 360 yds up from Harewood Grange bridge                       | 431110  | 368350       | <i>Gastrioceras subcrenatum</i> , <i>G. sp. nov.</i> ,<br><i>G. sp.</i> , <i>Homoceratoides sp.</i>                         |
| 760 yds N33W of Stonehay Farm   | 432800  | 368080       | <i>Gastrioceras subcrenatum</i>   |
| Opencast workings SW of Alton   | 436070  | 364200       | <i>Gastrioceras subcrenatum</i>   |
| Clattercotes Wood, 400 yds N15E of Whitecarr                            | 436120  | 360140       | <i>Gastrioceras subcrenatum</i> , <i>G. sp. nov.</i> ,<br><i>Homoceratoides sp.</i>   |




|   |         |              |   |
|---|---------|--------------|---|
| Egmanton No 62 BH   | 474440  | 367770       | <i>Gastrioceras subcrenatum</i>   |
| Ruelow Wood BH  | 402050  | 347520       | <i>Gastrioceras subcrenatum</i>   |
| Wetley Rocks  | 396640  | 349440       | <i>Gastrioceras subcrenatum</i>   |
| Consall New Lock  | 400420  | 348360       | <i>Gastrioceras subcrenatum</i>   |
| Newhouse Wood, Ipstones                                       | 401770  | 348850       | <i>Gastrioceras subcrenatum</i>   |
| Crowtrees No. 8 BH SK04NW/19                                  | 404980  | 345590       | <i>Gastrioceras subcrenatum</i>   |
| Rugeley (Trent Valley) BH 100 yds ESE of Railway Inn, Rugeley | 405080  | 319020       | <i>Gastrioceras subcrenatum</i>   |
| Osbaston Hollow BH  | 441660  | 306350       | <i>Gastrioceras</i>   |
| Park Brook BH, Horsley  | 438300  | 343850       | <i>Gastrioceras subcrenatum</i>   |
| Nether Heage BH   | 435990  | 351190       | <i>Gastrioceras subcrenatum</i>   |
| Beechdale Rd (Robins Wood) BH                                 | 453610  | 341130       | <i>Gastrioceras subcrenatum</i>   |
| LOCALITY  | EASTING | NORTHIN<br>G | AMMONOIDS   |
| <i>Gastrioceras subcrenatum</i> Marine Band (G2a1) Continued  |         |              |   |
| Wilds Bridge BH SK63SE/30                                     | 467380  | 332480       | <i>Gastrioceras subcrenatum</i>   |
| Blackfordby No. 1 BH  | 432350  | 318270       | <i>Gastrioceras subcrenatum</i>   |
| Road cut  | 437640  | 324860       | <i>Gastrioceras subcrenatum</i> , G. sp                                     |
| Worthington BH SK24SW/204                                     | 440450  | 321040       | <i>Gastrioceras</i> sp., <i>Gastrioceras subcrenatum</i>                    |
| Honley Marine Band  |         |              |   |
| Ridgeway Bh SJ85SE/14   | 389220  | 353810       | <i>Gastrioceras</i> sp.   |
| Charnock Old Hall BH  | 354730  | 416560       | <i>Gastrioceras</i> sp.   |
| Cheshire Brook  | 389030  | 361430       | <i>Gastrioceras</i> sp.   |
| Listeri Marine Band   |         |              |   |
| Ridgeway Bh SJ85SE/14   | 389220  | 353810       | <i>Gastrioceras listeri</i>   |
| Robin's Clough near Knar                                      | 400420  | 367630       | <i>Gastrioceras listeri</i>   |
| Waters Farm BH  | 375370  | 467630       | <i>Gastrioceras listeri</i> , G. circumnodosum                              |
| Valley Mills BH   | 386700  | 437200       | <i>Gastrioceras listeri</i>   |
| Cockden Bridge BH   | 387600  | 434400       | <i>Gastrioceras listeri</i>   |
| Globe Mills BH, Leeds, 400 yards S of City station SE23SE/7   | 429799  | 432787       | <i>Gastrioceras listeri</i>   |
| Stepback Brook, 900 yds S35W of Darwen Tower                  | 367410  | 420920       | <i>Gastrioceras listeri</i>   |
| Tan House Farm BH   | 355420  | 416920       | <i>Gastrioceras listeri</i>   |
| Heskin BH   | 353010  | 401360       | <i>Gastrioceras listeri</i>   |
| Oxspring Borehole, SE of Oxspring Yorks SE20SE/6              | 427870  | 401360       | <i>Gastrioceras listeri</i>   |
| Moss Oil BH SE51SE/19   | 459980  | 413900       | <i>Gastrioceras listeri</i>   |
| Ringstone Clough  | 400080  | 382210       | <i>Gastrioceras listeri</i>   |
| Furness Vale Colliery   | 400380  | 383390       | <i>Gastrioceras listeri</i> , <i>Homoceratoides</i> aff. <i>divaricatus</i> |
| Pingot Clough   | 401640  | 385320       | <i>Gastrioceras listeri</i>   |
| Stirrup, west of Chisworth                                    | 398350  | 391870       | <i>Gastrioceras listeri</i>   |
| Chew  | 399400  | 392030       | <i>Gastrioceras listeri</i>   |
| Smekley No 3 BH SK27NE/2                                      | 429690  | 376498       | <i>Gastrioceras listeri</i>   |
| Connah's Quay, trial pit                                      | 328170  | 369160       | <i>Gastrioceras listeri</i>   |
| Biddulph Grange   | 389210  | 359610       | <i>Gastrioceras circumnodosum</i>   |
| Cheshire Brook  | 388970  | 361330       | <i>Gastrioceras circumnodosum</i> , G. <i>listeri</i>                       |
| Clattercotes Wood, 480 yds N53E of Whitecarr                  | 436380  | 360050       | <i>Gastrioceras listeri</i>   |
| Key Wood BH SK04NW/1  | 403910  | 345360       | <i>Gastrioceras listeri</i>   |
| Out Wood, Consall valley                                      | 398050  | 347810       | <i>Gastrioceras listeri</i>   |
| SE of Ipstones  | 402980  | 349000       | <i>Gastrioceras circumnodosum</i>   |
| Whittington Heath BH  | 414780  | 308000       | <i>Gastrioceras</i> cf. <i>listeri</i>                                      |
| Ellistown Colliery BH   | 443900  | 310560       | <i>Gastrioceras listeri</i>   |
| Wilds Bridge BH SK63SE/30                                     | 467380  | 332480       | <i>Gastrioceras listeri</i>   |
| Marriott Wood Brickpit  | 429800  | 380300       | <i>Gastrioceras listeri</i>   |
| Little Stubbin Opencast, Stubbededge                          | 436300  | 361900       | <i>Gastrioceras listeri</i> , G. <i>coronatum</i> , G. <i>retorsum</i>      |
| Sandoz Chemical Co  | 416300  | 434300       | <i>Gastrioceras listeri</i>   |
| Amaliae Marine Band   |         |              |   |
| Disused quarry, 720 yds E14S of Charnock Green                | 355980  | 416640       | <i>Gastrioceras</i> cf. <i>amaliae</i>                                      |
| Crook Fold BH   | 354700  | 415870       | <i>Gastrioceras</i> cf. <i>amaliae</i>                                      |
| Vanderbeckei Marine Band                                      |         |              |   |
| Wiggins Teape No 2 BH, Withnell Fold Chorley                  | 361460  | 423440       | <i>Anthracoeras</i> cf. <i>vanderbeckei</i>                                 |


|   |         |              |  |
|---|---------|--------------|--|
| Bankfield Mills BH - Mold Green - SE11NE/11   | 414660  | 416270       | <i>Anthracoceas aff. vanderbeckei</i>                        |
| Bowsey Wood BH SJ74NE/9   | 376950  | 346430       | <i>Anthracoceas vanderbeckei</i>                             |
| Railway cutting, 1615 yds N of Holy Cross Church, Morton (Clay Cross type locality) | 440730  | 361600       | <i>Anthracoceas cf. vanderbeckei</i>                         |
| Scaftworth BH   | 467610  | 391670       | <i>Anthracoceas vanderbeckei</i>                             |
| Disused railway cutting, Duckmanton   | 442370  | 370400       | <i>Anthracoceas vanderbeckei</i>                             |
| Manton Colliery No 7 UG BH  | 463786  | 376334       | <i>Anthracoceas vanderbeckei</i>                             |
| Carbank BH  | 463969  | 355793       | <i>Anthracoceas vanderbeckei</i>                             |
| Haughton Hall BH  | 468595  | 373305       | <i>Anthracoceas vanderbeckei</i>                             |
| Clipstone Colliery No. 1 BH   | 459530  | 363290       | <i>Anthracoceas vanderbeckei</i>                             |
| Kirton BH   | 469880  | 369130       | <i>Anthracoceas vanderbeckei</i>                             |
| Kneesall BH   | 471353  | 364380       | <i>Anthracoceas vanderbeckei</i>                             |
| Mansfield Colliery BH   | 457020  | 361450       | <i>Anthracoceas vanderbeckei</i>                             |
| Ompton BH   | 469000  | 366100       | <i>Anthracoceas vanderbeckei</i>                             |
| LOCALITY  | EASTING | NORTHIN<br>G | AMMONOIDS  |
| Vanderbeckei Marine Band Continued  |         |              |  |
| Foxfield No8 BH SJ94SE/8  | 398880  | 343220       | <i>Anthracoceas vanderbeckei</i>                             |
| Manchester Woods BH SK44SW/3  | 441600  | 344170       | <i>Anthracoceas vanderbeckei</i>                             |
| Digby Clay Pit  | 448600  | 345000       | <i>Anthracoceas vanderbeckei</i>                             |
| Jockey House BH SK67NE/18   | 468971  | 376839       | <i>Anthracoceas vanderbeckei</i>                             |
| Aegiranum Marine Band   |         |              |  |
| Wentbridge No. 2 BH SE41NE/18   | 447560  | 417570       | <i>Donetzoceras aegiranum</i>                                |
| Bowsey Wood BH SJ74NE/9   | 376950  | 346430       | <i>Donetzoceras aegiranum</i>                                |
| Nettlebank Colliery   | 388500  | 350300       | <i>Donetzoceras aegiranum</i>                                |
| Pow Gill, 130 yards N of bridge at Powbank  | 325090  | 544320       | <i>Donetzoceras aegiranum</i> ,<br><i>Anthracoceas hindi</i> |
| Stairfoot Brickworks, 3km ESE of Barnsley   | 438030  | 404980       | <i>Donetzoceras aegiranum</i>                                |
| Manvers Main Brickworks, 1650 yds E6S of Bolton upon Dearne church                  | 445300  | 400980       | <i>Gastrioceras</i> sp.                                      |
| Doles Lane BH   | 453594  | 3774910      | <i>Homoceratoides politus</i>                                |
| Whitwell Rectory  | 452480  | 376670       | <i>Homoceratoides?</i>                                       |
| Elmton Green BH   | 450560  | 373170       | <i>Gastrioceras cf. depressum</i>                            |
| Red Hill BH, 2140 yds W of St James' church   |         |              |  |
| Longdon SK01SE/9  | 406230  | 314110       | <i>Homoceratoides politus</i>                                |
| Albion Clay Pit, 300 yds SSE Dordon church  | 426300  | 300100       | <i>Gastrioceras depressum</i>                                |
| Robinson & Dowler's Pit, 300 yds SSE of Dordon church                               | 429600  | 316100       | <i>Anthracoceas cf aegiranum-hindi</i>                       |
| Donington Pit, 250 yds 106 of Swainspark  | 429900  | 317100       | <i>Anthracoceas cf aegiranum-hindi</i>                       |
| Caldwell No. 1 BH, 250 yds 106 of Swainspark  | 425990  | 317340       | <i>Gastrioceras</i> sp.                                      |
| Eymore Farm railway cutting, Upper Arley  | 376900  | 279000       | <i>Donetzoceras aegiranum</i>                                |
| Shafton Marine Band   |         |              |  |
| Maltby Main Colliery No 2 shaft   | 455120  | 392460       | <i>Anthracoceas hindi</i>                                    |
| Cambriense Marine Band  |         |              |  |
| Wentbridge No. 2 BH SE41NE/18   | 447560  | 417570       | <i>Donetzoceras cambriense</i>                               |



**Fig. 1.** Map showing the approximate extent of Namurian and Westphalian strata at crop, the subcrop of Westphalian strata and the main pre-Namurian structural features of the Pennine Basin, derived from Waters *et al.* (2011). The location of the Harewood and Holme Pierrepont boreholes, from which new dates have been acquired during this study, are shown. DH- Derbyshire High, WG- Widmerpool Gulf.







**Fig. 2.** U-Pb data for samples BLL1976 and EH28155. A, conventional U-Pb concordia plot of zircons analysed from samples BLL1976 and EH28155. The grey band reflects the uncertainty in the  $^{238}\text{U}$  and  $^{235}\text{U}$  decay constants (Jaffey *et al.* 1971). B, plot of  $^{238}\text{U}/^{206}\text{Pb}$  dates for single zircon crystals analyses (same data as in Figure 2a). Dashed ellipses/bars represent analyses of zircon that are considered to be xenocrysts and/or inherited crystals that are disregarded in calculation of final date, whereas as undashed ellipses/bars represent the analyses used for calculation of the weighted mean final date (see text for discussion). Data point error ellipses/bars are  $2\sigma$ .






**Fig. 3.** Distribution of ammonoid acme facies in early Namurian marine bands: a) Pendleian; b) early Arnsbergian; c) mid to late Arnsbergian; d) Chokierian; e) Alportian; f) early Kinderscoutian. Grey tone denotes marine band with maximum areal extent for each interval. Key for Incised Valleys, as for Figure 5.





**Fig. 4.** Distribution of ammonoid acme facies in late Namurian–Westphalian marine bands: a) mid to late Kinderscoutian; b) early Marsdenian; c) late Marsdenian; d) Yeadonian-  $G_{1b1}$  modified from Wignall (1987); e) Langsettian and f) Duckmantian–Bolsovian, in part based upon Calver (1968, 1969). Grey tone denotes marine band with maximum areal extent for each interval. Key for Incised Valleys, as for Figure 5.





**Fig. 5.** Magnitude and duration of sea-level oscillations. Sea-level maxima are estimated through the determination of maximum areal extent of acme ammonoid facies. Abbreviations for Westphalian marine bands: SMB Subcrenatum Marine Band; LMB Listeri Marine Band; AmMB Amaliae Marine Band; VMB Vanderbeckeri Marine Band; HMB Houghton Marine Band; AMB Aegiranum Marine Band; CMB Cambriense Marine Band. For marine bands lacking ammonoid fauna the magnitude of sea-level is determined through the acme marine fauna, ranging from *Estheria* to brachiopod-bivalve facies. Sea-level minima are determined through the presence of incised valleys, with the magnitude recorded by the number of underlying marine bands removed beneath the sequence boundary. Sources for incised valleys are as follows: a) Rogerley Channel (Dunham 1990); b) Upper Howgate Edge channel (Martinsen *et al.* 1995); c) Red Scar Grit (Brandon *et al.* 1995); d) Lower Follifoot Grit (Martinsen 1993); e) Intra-H<sub>1a</sub> unconformity (Owens *et al.* 1990); f) Upper Follifoot Grit (Martinsen 1993); g) Todmorden Grit/Kinderscout Grit (Hampson 1997); h) Upper Kinderscout Grit (Hampson 1997); i) Midgley Grit (Brettle 2001); j) Ashover Grit/Roaches Grit (Jones & Chisholm 1997; Church & Gawthorpe 1994); k) Chatsworth Grit (Waters *et al.* 2008); l) Rough Rock (Church & Gawthorpe 1994); m) Crawshaw Sandstone (Hampson *et al.* 1997); n) Thornhill Rock (Lake 1999); o) Woolley Edge Rock (Aitken *et al.* 1999); p) Oaks Rock (Lake 1999); q) Mexborough Rock (Aitken *et al.* 1999). Radiometric dates are from this study and estimated ages of stage boundaries are from Davydov *et al.* (2010), with an imposed 400 ka long-duration eccentricity oscillation numbered sequentially for each international stage: S Serpukhovian; B Bashkirian; M Moscovian. The proposed four main glaciations are highlighted as grey bands.

**Table 1.** Comparison of Carboniferous chronological ages from selected published literature. Picks are at base of Regional Stage or Substage unless stated otherwise. Hess & Lippolt (1986)<sup>1</sup> and Berger *et al.* (1997)<sup>2</sup> provide <sup>40</sup>Ar/<sup>39</sup>Ar plateau ages (Ma ± 1σ), with the exception of the early Arnsbergian age which is Ma ± 2σ; recalibrated ages appear in italics, as quoted in Davydov *et al.* (2004); Kryza *et al.* (2010)<sup>3</sup>, Riley *et al.* (1995)<sup>4</sup> and Claoué-Long *et al.* (1995)<sup>5</sup> provide SHRIMP (Ma ± 2σ) ages; Trapp & Kaufmann (2002)<sup>6</sup>, Gastaldo *et al.* (2009)<sup>7</sup> and Davydov *et al.* (2010)<sup>8</sup> provide U-Pb TIMS (Ma ± 2σ) ages. Stippled lines denote correlation of regional substage boundaries.

| <sup>40</sup> Ar/ <sup>39</sup> Ar plateau ages  | U-Pb SHRIMP                        | Davydov <i>et al.</i> (2004) | Ogg <i>et al.</i> (2008) | U-Pb TIMS   | This study           | International Stage | Regional Substage | Regional Stage   |
|--|------------------------------------|------------------------------|--------------------------|---|----------------------|---------------------|-------------------|------------------|
| 302.7 ± 0.5 (late) <sup>(1)</sup>  |                                    | 306.5 ± 1.0                  | 307.2 ± 1.0              | 307.26 ± 0.11 (early) <sup>(8)</sup>  |                      | Kasimovian          | Cantabrian        | Early Stephanian |
| 308.0 ± 1.8 (mid) <sup>(2)</sup><br>(310.26 ± 1.8)                                     |                                    |                              |                          |   |                      | Moscovian           | Asturian          | Westphalian      |
| 309.0 ± 3.7 (late) <sup>(1)</sup><br>309.7 ± 2.0 (mid) <sup>(2)</sup><br>(312.0 ± 2.0) |                                    |                              |                          | 310.55 ± 0.10 (late) <sup>(8)</sup>   |                      |                     | Bolsovian         |                  |
| 310.0 ± 1.0 <sup>(2)</sup><br>(313.0 ± 1.0)  | 311.0 ± 3.4 <sup>(5)</sup>         | 311.7 ± 1.1                  | 311.7 ± 1.1              | 314.40 ± 0.06 (early) <sup>(8)</sup>  | <b>314.37 ± 0.53</b> |                     |                   |                  |
| 310.7 ± 1.3 (late) <sup>(1)</sup>  |                                    |                              |                          |   |                      |                     |                   |                  |
|  |                                    |                              |                          |   |                      | Bashkirian          | Duckmantian       | Namurian         |
|  |                                    |                              |                          |   |                      |                     | Langsettian       |                  |
|  |                                    |                              |                          |   |                      |                     | Yeadonian         |                  |
|  |                                    |                              |                          |   |                      |                     | Marsdenian        |                  |
|  |                                    |                              |                          |   |                      |                     | Kinderscoutian    |                  |
|  |                                    |                              |                          |   |                      |                     | Alportian         |                  |
|  |                                    | 318.1 ± 1.3                  | 318.1 ± 1.3              |   |                      |                     | Chokierian        |                  |
| 319.9 ± 1.6 (mid) <sup>(1)</sup><br>(322.3 ± 1.7)                                      | 314.4 ± 4.6 (mid) <sup>(4)</sup>   |                              |                          |   | <b>328.34 ± 0.55</b> | Serpukhovian        | Arnsbergian       | Namurian         |
| 324.8 ± 1.2 (early) <sup>(1)</sup><br>(327.0 ± 2.2)                                    | 314.5 ± 4.6 (early) <sup>(4)</sup> |                              |                          |   |                      |                     |                   |                  |
|  |                                    |                              |                          | 328.01 ± 0.36 (late) <sup>(7)</sup><br>328.14 ± 0.11 (early-mid) <sup>(8)</sup><br>328.84 ± 0.38 (early) <sup>(7)</sup> |                      |                     | Pendleian         |                  |
|  |                                    | 326.4 ± 1.6                  | 328.3 ± 1.6              |   |                      |                     |                   |                  |
|  |                                    |                              |                          |   |                      | Late Viséan         | Brigantian        | Late Viséan      |
|  | 334 ± 4 <sup>(3)</sup>             |                              |                          | 326.8 ± 0.98 (late) <sup>(6)</sup>  |                      |                     | Asbian            |                  |

| Regional Substages                         | ZONES |                                    |            | WESTERN EUROPEAN MARINE BANDS        |   |  | ‘Meso-thems’ |
|--|-------|------------------------------------|------------|--------------------------------------|---|--|--------------|
|  | Index | Ammonoid                           | Index      | Diagnostic ammonoid                  | Former name   | Associated ammonoids   |              |
| BASHKIRIAN STAGE (NAMURIAN REGIONAL STAGE) |       |                                    |            |                                      |   |  | N11          |
| YEADONIAN                                  | G1b   |                                    |            | <i>Anthracoceras</i>                 |   | None, typically <i>Lingula</i> facies  |              |
|  |       | <i>Cancelloceras cumbriense</i>    | G1b1       | <i>Ca. cumbriense a &amp; b</i>      | <i>Gastrioceras cumbriense</i>                                  | <i>B. superbilinguis</i> , <i>Cancelloceras</i> sp., <i>Ca. crenulatum</i> , <i>Gastrioceras carbonarium</i> , <i>G. listeri</i> , <i>Homoceratoides</i> sp., <i>Hm..divaricatus</i> ,   |              |
|  | G1a   |                                    |            |                                      |   | Owd Betts Marine Band- anoxic event lacking marine fauna   |              |
|  |       | <i>Cancelloceras cancellatum</i>   | G1a1       | <i>Ca. cancellatum a, b &amp; c</i>  | <i>Gastrioceras cancellatum</i>                                 | <i>Agastrioceras carinatum</i> , <i>B. superbilinguis</i> , <i>Cancelloceras</i> sp., <i>Ca. branneroides</i> , <i>Ca. crenulatum</i> , <i>Ca. crenulatum</i> , <i>Ca. cumbriense</i> <i>Ca. Rurae</i> , <i>Homoceratoides</i> sp., <i>Hm..divaricatus</i> |              |
| MARSDENIAN                                 | R2c   | <i>Bilinguites superbilinguis</i>  | R2c2       | <i>Verneulites sigma</i>             | <i>Donetzooceras (Gastrioceras) sigma</i>                       | <i>B. superbilinguis</i> , <i>Ca. cumbriense</i> , <i>Gastrioceras</i> sp., <i>Ca. crenulatum</i> , <i>Homoceratoides</i> sp.,   | N10          |
|  |       |                                    | R2c1       | <i>B. superbilinguis</i>             | <i>R. superbilingue</i> , <i>R. reticulatum</i> mut. $\gamma$ , | <i>Gastrioceras</i> sp., <i>G. lineatum</i> , <i>Homoceratoides</i> sp. <i>Hm..divaricatus</i> , <i>Hm. fortelirifer</i> , <i>Verneulites sigma</i>  |              |
|  | R2b   | <i>Bilinguites bilinguis</i>       | R2b5       | <i>B. metabilinguis</i>              | <i>R. bilingue</i> late mut. $\beta$                            | <i>Bilinguites</i> sp., <i>B. bilinguis</i> , <i>B. eometabilinguis</i> , <i>Verneulites sigma</i>   | N9           |
|  |       |                                    | R2b4       | <i>B. eometabilinguis</i>            | <i>R. bilingue</i> late mut. $\beta$                            | <i>B. bilinguis</i>  |              |
|  |       |                                    | R2b3       | <i>B. bilinguis</i>                  | <i>R. bilingue</i> mut. $\beta$                                 | <i>Bilinguites</i> sp., <i>R. circumplicatile</i>  |              |
|  |       |                                    | R2b2       | <i>B. bilinguis</i>                  | <i>R. bilingue</i> mut. $\beta$                                 |  |              |
|  |       |                                    | R2b1       | <i>B. bilinguis</i>                  | <i>R. bilingue</i> early mut. $\beta$                           |  |              |
|  | R2a   | <i>Bilinguites gracilis</i>        | R2a1       | <i>B. gracilis a, b &amp; c</i>      | <i>R. gracile</i> , <i>R. reticulatum</i> mut. <i>a</i>         | <i>Bilinguites</i> sp., <i>R. reticulatum</i> , <i>R. gracilingue</i> , <i>R. graciloides</i>  | N8           |
|  | R1c   | <i>Reticuloceras reticulatum</i>   | R1c5       |                                      |   | Butterfly MB- <i>Lingula</i>   |              |
|  |       |                                    | R1c4       | <i>R. coreticulatum</i>              |   | <i>Anthracoceratites</i> sp., <i>Homoceratoides divaricatus</i> , <i>Hudsonoceras ornatum</i> , <i>R. reticulatum</i> (late form), <i>V. striolatus</i>  |              |
|  |       |                                    | R1c3       | <i>R. reticulatum</i>                |   | <i>Homoceratoides prereticulatus</i> ,   |              |
|  |       |                                    | R1c2       | <i>R. reticulatum</i>                |   | <i>R. davisii</i> , <i>R. regularum</i> , <i>Vallites</i> sp., <i>V. striolatus</i>  |              |
|  |       |                                    | R1c1       | <i>R. reticulatum</i>                |   |  |              |
|  | R1b   | <i>Reticuloceras eoreticulatum</i> | R1b3       | <i>R. stubblefieldi</i>              | <i>Eumorphoceras stubblefieldi</i>                              | <i>Vallites striolatus</i> , <i>Hudsonoceras ornatum</i> , <i>R. Moorei</i> , <i>R. regularum</i>  | N7           |
|  |       |                                    | R1b2       | <i>R. nodosum</i>                    |   | <i>Homoceras spiralooides</i> , <i>Vallites striolatus</i>   |              |
|  |       |                                    | R1b1       | <i>R. eoreticulatum</i>              |   | <i>Reticuloceras</i> sp.   |              |
| KINDERSCOUTIAN                             | R1a   | <i>Hodsonites magistrorum</i>      | R1a5       | <i>R. dubium</i>                     |   | <i>Reticuloceras</i> sp., <i>R. adpressum</i>  | N6           |
|  |       |                                    | R1a4       | <i>R. todmordenense</i>              |   | <i>Homoceras</i> sp., <i>Reticuloceras</i> sp., <i>R. paucicrenulatum</i> , <i>R. adpressum</i>  |              |
|  |       |                                    | R1a3       | <i>R. subreticulatum</i>             |   | <i>Homoceratoides</i> sp., <i>Reticulatum pulchellum</i>   |              |
|  |       |                                    | R1a2       | <i>R. circumplicatile</i>            |   | <i>Homoceratoides</i> sp. <i>Vallites (Homoceras) henkei</i> ,   |              |
|  |       |                                    | R1a1       | <i>Ho. magistrorum</i>               |   | <i>R. compressum</i>   |              |
|  |       |                                    |            |                                      |   |  |              |
|  |       |                                    |            |                                      |   |  |              |
|  |       |                                    |            |                                      |   |  |              |
| ALPORTIAN                                  | H2c   | <i>Vallites eostriolatus</i>       | H2c2       | <i>Homoceratoides prereticulatus</i> | <i>Hm. prereticulatum</i>                                       | <i>Homoceras</i> sp.   | N5           |
|  |       |                                    | H2c1       | <i>V. eostriolatus</i>               | <i>Homoceras eostriolatum</i>                                   | <i>Ho. undulatum</i>   |              |
|  | H2b   | <i>Homoceras undulatum</i>         | H2b1       | <i>Ho. undulatum</i>                 |   | <i>Ho. beyrichianum</i> , <i>Ho. smithii</i>   |              |
|  | H2a   | <i>Hudsonoceras proteum</i>        | H2a1       | <i>Hd. proteum a, b &amp; c</i>      | <i>Hd. proteus</i>  | <i>Homoceras smithii</i>   |              |
| CHOKIERIAN                                 | H1b   | <i>Homoceras beyrichianum</i>      | H1b2       | <i>Isohomoceras</i> sp. nov.         |   | <i>Homoceras</i> sp.   | N4           |
|  |       |                                    | H1b1 a & b | <i>H. beyrichianum</i>               |   | <i>Ho. diadema</i> , <i>Ho. subglobosum</i> , <i>Isohomoceras</i> sp.  |              |
|  | H1a   | <i>Isohomoceras subglobosum</i>    | H1a3       | <i>I. subglobosum</i>                | <i>Homoceras subglobosum</i>                                    |  |              |
|  |       |                                    | H1a2       | <i>I. subglobosum</i>                |   |  |              |
|  |       |                                    | H1a1       | <i>I. subglobosum</i>                |   |  |              |

| Regional | ZONES |  |  | WESTERN EUROPEAN MARINE BANDS |  |  |  |
|----------|-------|--|--|-------------------------------|--|--|--|
|----------|-------|--|--|-------------------------------|--|--|--|

| Substages                                    | Index | Ammonoid                            | Index      | Diagnostic ammonoid                | Former name                         | Associated ammonoids   | 'Meso-thems' |
|--|-------|-------------------------------------|------------|------------------------------------|-------------------------------------|--|--------------|
| SERPUKHOVIAN STAGE (NAMURIAN REGIONAL STAGE) |       |                                     |            |                                    |                                     |  | N3           |
| ARNSBERGIAN                                  | E2c   | <i>Nuculoceras stellarum</i>        | E2c4       | <i>N. nuculum</i>                  |                                     | <i>C. darwenense</i> , <i>Ct. fragile</i> , <i>E. bisulcatum</i> ,   |              |
|  |       |                                     | E2c3       | <i>N. nuculum</i>                  |                                     | <i>Kazakhoceras hawkinsi</i>   |              |
|  |       |                                     | E2c2       | <i>N. nuculum</i>                  |                                     |  |              |
|  |       |                                     | E2c1       | <i>N. stellarum</i>                | <i>Cravenoceratoides stellarum</i>  | <i>Fayettevillea holmesi</i> ,   |              |
|  | E2b   | <i>Cravenocera-toides edalensis</i> | E2b3       | <i>Ct. nititoides</i>              |                                     | <i>Cravenoceras</i> sp., <i>Eumorphoceras rostratum</i>  | N 2          |
|  |       |                                     | E2b2 a-c   | <i>Ct. nitidus</i>                 | <i>Ct. nitidum</i>                  | <i>Cravenoceras</i> sp., <i>C. subplicatum</i> , <i>Cravenoceratoides</i> sp., <i>Eumorphoceras</i> sp., <i>E. bisulcatum</i> , <i>E. leitrinense</i> , <i>Fayettevillea holmesi</i> , <i>Glaphyrites</i> sp., <i>Gl. Kettlesingense</i> , |              |
|  |       |                                     | E2b1 a-c   | <i>Ct. edalensis</i>               |                                     | <i>Cravenoceras</i> sp., <i>C. subplicatum</i> , <i>Fayettevillea</i> cf. <i>holmesi</i>   |              |
|  | E2a   | <i>Cravenoceras cowlingsense</i>    | E2a3       | <i>Eumorphoceras yatesae</i>       |                                     | <i>Cravenoceras</i> sp., <i>C. gairense</i> , <i>Eumorphoceras</i> sp.   | N 1          |
|  |       |                                     | E2a2β      | <i>Anthracoceras</i>               |                                     | Saleswheel Marine Band   |              |
|  |       |                                     | E2a2α      | <i>C. gressinghamense</i>          |                                     | <i>C. cf. gairense</i> , <i>Eumorphoceras</i> sp.,   |              |
|  |       |                                     | E2a2       | <i>Eumorphoceras ferrimontanum</i> | <i>Eumorphoceras bisulcatum</i>     | <i>Cravenoceras</i> sp., <i>C. gairense</i> , <i>E. erinense</i> , <i>Kazakhoceras scaliger</i>  |              |
|  |       |                                     | E2a1 a-c   | <i>C. cowlingsense</i>             | <i>Eumorphoceras bisulcatum</i>     | <i>Cravenoceras</i> sp., <i>E. grassingtonense</i>   |              |
| PENDLEIAN                                    | E1c   | <i>Cravenoceras malhamense</i>      | E1c2       |                                    |                                     | Blacko Marine Band- <i>Sanguinolites</i>   | N 1          |
|  |       |                                     | E1c1       | <i>C. malhamense</i>               |                                     | <i>Cravenoceras</i> sp., <i>Eumorphoceras</i> sp.  |              |
|  | E1b   | <i>Cravenoceras brandoni</i>        | E1b2 a & b | <i>Tumulites pseudobilinguis</i>   | <i>Eumorphoceras pseudobilingue</i> | <i>Edmooroceras angustum</i> , <i>Ed. hudsoni</i> , <i>Ed. stubblefieldi</i> , <i>Eumorphoceras</i> sp., <i>Tumulites</i> sp.  |              |
|  |       |                                     | E1b1       | <i>C. brandoni</i>                 |                                     | <i>Edmooroceras stubblefieldi</i>  |              |
|  | E1a   | <i>Emstites leion</i>               | E1a1 a-c   | <i>E. leion</i>                    | <i>Cravenoceras leion</i>           | <i>Cravenoceras</i> sp., <i>Cousteauceras rota</i> , <i>Eumorphoceras</i> sp., <i>E. involutum</i> , <i>Edmooroceras bisati</i> , <i>Ed. medusa</i> , <i>Ed. pseudocoronula</i> , <i>Ed. tornquisti</i>                                    |              |

**Table 2.** Ammonoid zones and subzones of the Namurian regional stage with diagnostic ammonoids and indices mainly from Riley *et al.* (1995). 'Mesothem' nomenclature is that of Ramsbottom (1977). Former names and a compilation of ammonoid assemblages for each marine band were used when studying sample collections and literature reviews to determine the extent of ammonoid facies within specific marine bands. a) Serpukhovian international stage; b) Bashkirian international stage (part).

**Table 3.** Marine bands of the Westphalian regional stage with diagnostic ammonoids or acme facies, mainly from Waters *et al.* (2011). ‘Mesothem’ nomenclature is that of Ramsbottom (1979). Former names and a compilation of ammonoid assemblages for each marine band were used when studying sample collections and literature reviews to determine the extent of ammonoid facies within specific marine bands.

| STAGE             | REGIONAL SUB-STAGES | WESTERN EUROPEAN MARINE BANDS |                                      |                  |   | ‘MESO-THEM’ |
|-------------------|---------------------|-------------------------------|--------------------------------------|------------------|---|-------------|
|                   |                     | MARINE BAND NAME              | Diagnostic ammonoid or acme facies   | Former name      | Associated ammonoids  |             |
| MOSCOVIAN (PART)  | BOLSOVIAN           | Cambriense                    | <i>Donetzoceras cambriense</i>       | Top              |   | W10         |
|                   |                     | Shafton                       | <i>Anthracoceras hindi</i>           |                  |   |             |
|                   |                     | Main Estheria                 | <i>Estheria</i>                      |                  |   |             |
|                   |                     | Edmondia                      | foraminifera                         |                  |   | W9          |
|                   |                     | Carway Fawr                   | foraminifera                         |                  | Proved in S. Wales only   |             |
|                   | DUCK-MANTIAN        | Aegiranum                     | <i>Donetzoceras aegiranum</i>        | Mansfield        | <i>Anthracoceras hindi</i> ,<br><i>Gastrioceras</i> sp., <i>G. depressum</i> ,<br><i>Homoceratoides politus</i>   | W8          |
|                   |                     | Sutton                        | <i>Lingula</i>                       |                  |   |             |
|                   |                     | Haughton                      | <i>Levipustula</i>                   |                  |   | W7          |
|                   |                     | Clown                         | <i>Lingula</i>                       |                  |   |             |
|                   |                     | Manton Estheria               | <i>Estheria</i>                      |                  |   |             |
|                   |                     | Maltby                        | <i>Myalina</i>                       | Two Foot         |   | W6          |
|                   |                     | Lowton Estheria               | <i>Estheria</i>                      |                  |   |             |
|                   |                     | Vanderbeckei                  | <i>Anthracoceras vanderbeckei</i>    | Clay Cross       |   | W5          |
|                   | LANGSETTIAN         | Low Estheria                  | <i>Estheria</i>                      |                  |   |             |
|                   |                     | Burton Joyce                  | <i>Caneyella</i> , <i>Posidonia</i>  |                  |   | W4          |
|                   |                     | Langley                       | <i>Lingula</i>                       | Upper Band       |   |             |
|                   |                     | Amaliae                       | <i>Gastrioceras amaliae</i>          | Norton (Tonge’s) |   | W3          |
|                   |                     | Meadow Farm                   | <i>Dunbarella</i> , <i>Posidonia</i> | Forty Yard       |   |             |
|                   |                     | Parkhouse                     | <i>Lingula</i> , <i>Caneyella</i>    |                  |   |             |
|                   |                     | Listeri                       | <i>Gastrioceras listeri</i>          | Alton            | <i>G. circumnodosum</i> , <i>G. coronatum</i> , <i>G. retrorsum</i> ,<br><i>Homoceratoides divaricatus</i>  | W2          |
|                   |                     | Honley                        | <i>Gastrioceras</i> sp.              | First Smalley    |   |             |
|                   |                     | Springwood                    | <i>Lingula</i>                       | Second Smalley   |   |             |
|                   |                     | Holbrook                      | <i>Lingula</i>                       | Lower Bassey     |   |             |
|                   |                     | Subcrenatum                   | <i>Gastrioceras subcrenatum</i>      | Pot Clay         | <i>Gastrioceras</i> sp., <i>G. carbonarium</i> , <i>G. listeri</i> , <i>G. retrorsum</i> ,<br><i>Homoceratoides</i> sp., <i>H. divaricatus</i> , <i>Reticuloceras superbilingue</i> | W1/<br>N11  |
| BASHKIRIAN (PART) | BOLSOVIAN           | Cambriense                    | <i>Donetzoceras cambriense</i>       | Top              |   | W10         |
|                   |                     | Shafton                       | <i>Anthracoceras hindi</i>           |                  |   |             |
|                   |                     | Main Estheria                 | <i>Estheria</i>                      |                  |   |             |
|                   |                     | Edmondia                      | foraminifera                         |                  |   | W9          |
|                   |                     | Carway Fawr                   | foraminifera                         |                  | Proved in S. Wales only   |             |
|                   | DUCK-MANTIAN        | Aegiranum                     | <i>Donetzoceras aegiranum</i>        | Mansfield        | <i>Anthracoceras hindi</i> ,<br><i>Gastrioceras</i> sp., <i>G. depressum</i> ,<br><i>Homoceratoides politus</i>   | W8          |
|                   |                     | Sutton                        | <i>Lingula</i>                       |                  |   |             |
|                   |                     | Haughton                      | <i>Levipustula</i>                   |                  |   | W7          |
|                   |                     | Clown                         | <i>Lingula</i>                       |                  |   |             |
|                   |                     | Manton Estheria               | <i>Estheria</i>                      |                  |   |             |
|                   |                     | Maltby                        | <i>Myalina</i>                       | Two Foot         |   | W6          |
|                   |                     | Lowton Estheria               | <i>Estheria</i>                      |                  |   |             |
|                   |                     | Vanderbeckei                  | <i>Anthracoceras vanderbeckei</i>    | Clay Cross       |   | W5          |
|                   | LANGSETTIAN         | Low Estheria                  | <i>Estheria</i>                      |                  |   |             |
|                   |                     | Burton Joyce                  | <i>Caneyella</i> , <i>Posidonia</i>  |                  |   | W4          |
|                   |                     | Langley                       | <i>Lingula</i>                       | Upper Band       |   |             |
|                   |                     | Amaliae                       | <i>Gastrioceras amaliae</i>          | Norton (Tonge’s) |   | W3          |
|                   |                     | Meadow Farm                   | <i>Dunbarella</i> , <i>Posidonia</i> | Forty Yard       |   |             |
|                   |                     | Parkhouse                     | <i>Lingula</i> , <i>Caneyella</i>    |                  |   |             |
|                   |                     | Listeri                       | <i>Gastrioceras listeri</i>          | Alton            | <i>G. circumnodosum</i> , <i>G. coronatum</i> , <i>G. retrorsum</i> ,<br><i>Homoceratoides divaricatus</i>  | W2          |
|                   |                     | Honley                        | <i>Gastrioceras</i> sp.              | First Smalley    |   |             |
|                   |                     | Springwood                    | <i>Lingula</i>                       | Second Smalley   |   |             |
|                   |                     | Holbrook                      | <i>Lingula</i>                       | Lower Bassey     |   |             |
|                   |                     | Subcrenatum                   | <i>Gastrioceras subcrenatum</i>      | Pot Clay         | <i>Gastrioceras</i> sp., <i>G. carbonarium</i> , <i>G. listeri</i> , <i>G. retrorsum</i> ,<br><i>Homoceratoides</i> sp., <i>H. divaricatus</i> , <i>Reticuloceras superbilingue</i> | W1/<br>N11  |