





INSTITUTE OF FRESHWATER ECOLOGY River Laboratory, East Stoke, Wareham, Dorset BH20 6BB

The macroinvertebrate fauna of the River Blythe at Cheswick Green

J.F. Wright PhD M.T. Furse BSc

Project leader: J.F. Wright Report date: October 1990 Report to: IFE report no: TFS project no:

Nature Conservancy Council RL/T04053g1/4 T04053g1

This is an unpublished report and should not be cited without permission, which should be sought through the Director of the Institute of Freshwater Ecology in the first instance.

The Institute of Freshwater Ecology is part of the Terrestrial and Freshwater Sciences Directorate of the Natural Environment Research Council.



Sources of information

The River Blythe, whose catchment includes areas to the south east and east of Birmingham, flows north to join the R. Cole and then the R. Tame, which in its turn discharges into the R. Trent.

In 1982 five sites were sampled on the R. Blythe as part of the River Communities Project (RCP). Field sampling was undertaken by biologists from the Severn-Trent Water Authority, using standard RCP procedures. The sampling protocol involved 3-minute pond-net samples encompassing all available habitats in each of spring, summer and autumn. Sample processing and the identification of the macroinvertebrate fauna was then carried out by staff of the River Laboratory, Wareham, Dorset.

This brief report provides information on the fauna recorded at Cheswick Green (Nat. Grid Ref. SP 127753), the most upstream location included in the survey.

In July 1989, further sampling was undertaken at the same site in a separate project on macrophyte-invertebrate relationships, sponsored by the Nature Conservancy Council. On this occasion, the pond-net samples were of 15 sec. duration and each one was taken on a separate habitat type. Samples were examined from six habitat types (algae, *Glyceria*, *Elodea*, gravel, *Ranunculus*, tree roots) and the fauna was examined as far as available keys allowed, except for the Sphaeriidae, Oligochaeta and Chironomidae. Note that these three taxonomic groups were identified in the 1982 survey.

The site

The R. Blythe at Cheswick Green is a small lowland stream, just 3.8 km from the source and -2.5 m in width. Some sections of the stream are heavily shaded by trees whilst others are more open in character. In July 1989 a 100 m section of the stream was found to have 47% cover of submerged macrophyte, 46% cover of non-macrophyte substratum and 7% emergent vegetation. Of the submerged vegetation, algae (*Cladophora*) was dominant, but *Ranunculus aquatilis* and *Elodea* nuttallii were also present in significant amounts. Pebbles/gravel was the dominant non-macrophyte substratum. The emergent vegetation included *Glyceria* x pedicillata and Apium nodiflorum but heavy grazing and trampling by cattle affected the stream margins. The six locations sampled for invertebrates varied in depth between 5 and 50 cm.

The fauna

A total of 80 taxa have been recorded at Cheswick Green as a result of the 3 season survey in 1982 and the summer-only survey in 1989. As shown in the following list, 69 taxa were recorded in 1982 and the 1989 survey contributed a further 11 taxa.

Macroinvertebrate fauna of Cheswick Green (SP 127753)

Major taxonomic group	No. taxa found in 1982 (3 seasons)	Additional taxa found summer 1989	Total taxa recorded
Gastropoda	3	3	6
Bivalvia	5		5
Oligochaeta	7		7
Hirudinea	3	1	4
Hydracarina	1		1
Crustacea	2		2
Ephemeroptera	5		5
Plecoptera	2		2
Hemiptera	2	1	3
Coleoptera	7	5	12
Megaloptera	1		1
Trichoptera	9		9
Diptera	22	_1	23
	69	11	80

The individual taxa recorded in each of spring, summer and autumn 1982 are listed at the end of this report. Taxa present in summer 1989 are also given and those previously recorded in 1982 are distinguished from those first reported in 1989.

Cheswick Green supports a reasonably diverse range of macroinvertebrates. The fauna is characteristic of a small headwater stream in the Midlands and East Anglia. Most of the species found at the site are common freshwater invertebrates and there are no nationally rare taxa.

When the faunal list is examined in relation to the frequency of occurrence of taxa in the IFE 438 site data-base, there are just 7 taxa at Cheswick Green which occurred at 20 or fewer sites within the 438 site data-base. None of these can be regarded as rare and their low frequency of occurrence in the data-base is partly because it still includes a relatively limited number of small stream sites. In addition, one hemipteran and four species of Coleoptera found at Cheswick Green are more typical of ditches and standing water than streams and rivers. The 7 taxa are listed below, together with brief comments on their status and typical habitats.

(At 16 sites in the IFE data-base)
Widespread, but not common in England
(At 12 sites in the IFE data-base)
Common in the Midlands, but more typical of standing
water
(1 site in the IFE data-base)
Ubiquitous; typical of grassy pools and ditches
(9 sites in the IFE data-base)
Ubiquitous; typical of grassy pools and ditches
(5 sites in the IFE data-base)
Ubiquitous; typical of grassy ponds

Hydrobius fuscipes	(8 sites in the IFE data-base) Ubiguitous; typical of pools with detritus	
Trichoptera		
Limnephilis extricatus	(20 sites in the IFE data-base) Common in silty and well vegetated regions of	
	streams, small rivers and canals	

The site at Cheswick Green falls within Group 8B of the running-water site classification scheme based on macroinvertebrates which has been developed by the IFE. Group 8B includes 10 sites, 8 of which are the uppermost sites examined on lowland streams in the Midlands and East Anglia. These 10 sites have a mean distance from source of 4.8 km and a mean width of 1.9 m.

MACROINVERTEBRATE FAUNA OF THE RIVER BLYTHE AT CHESWICK GREEN

a) 1982 Survey (3 seasons)

Taxa recorded in 1982 are listed in taxonomic sequence.

Positive records for individual seasons are denoted with a + in the appropriate column.

Left-hand-column - 26 April 1982 Middle column - 23 July 1982 Right-hand-column - 18 October 1982

b) 1989 Survey (18 July)

* denotes taxa in the 1982 list which were also recorded in 1989.

Taxa recorded in 1989 but not 1982 are listed separately on the right hand side of the page.

1982

1989

* Potamopyrgus jenkinsi	+++	Valvata piscinalis
* Lymnaea peregra	+++	Anisus vortex
Ancylus fluviatilis	+++	Gyraulus albus
Sphaerium corneum	· +++	
Pisidium casertanum	++0	
Pisidium personatum	++0	
Pisidium subtruncatum	+++	
Pisidium nitidum	+0+	
Nais elinguis	+++	
Nais pardalis	0+0	
Stylaria lacustris	0++	
Limnodrilus claparedeianus	00+	
Limnodrilus hoffmeisteri	+++	
Aulodrilus pluriseta	+++	
Lumbriculidae	0+0	·
* Glossiphonia complanata	+++	Helobdella stagnalis
Erpobdella octoculata	+00	
Trocheta subviridis	0+0	1
* Hydracarina	+++	
* Asellus aquaticus	0++	
* Gammarus pulex	+++	
* Baetis scambus group	0++	
* Baetis vernus	0++	
* Baetis rhodani	+++	
Habrophlebia fusca	+00	
* Ephemerella ignita	0+0	
Amphinemura standfussi	+00	
Nemoura cinerea	+00	
* Velia caprai (* = Velia sp.)	0+0	Sigara dorsalis
Nepa cinerea	0+0	
* Brychius elevatus	+00	
🔹 🕈 Haliplus lineatocollis	0+0	Haliplus fluviatilis
* Platambus maculatus	0++	Potamonectes sp.
llybius fuliginosus	0+0	Agabus sp.
* Helophorus brevipalpis	0+0	Helophorus aequalis
* Helophorus minutus	0+0	Hydrobius fuscipes
* Helophorus grandis	0+0	
* Sialia lutaria	0+0	

MACROINVERTEBRATE FAUNA OF THE RIVER BLYTHE AT CHESWICK GREEN

1982

Hydropsyche angustipennis+Hydropsyche siltalaiOHydroptila sp.O* Limnephilus lunatus groupHLimnephilus extricatusOAnabolia nervosaHPotamophylax groupOHalesus sp.HTipula montium groupO* Dicranota sp.OLimnophila (Eloeophila) sp.HApsectrotanypus trifascipennisOProcladius sp.OThienemannimyia groupHPotthastia longimana groupOBrillia modestaHEukiefferiella groupHProdiamesa olivaceaOChironomus sp.OMicrotendipes sp.OParatendipes sp.OPolypedilum sp.HMicropsectra groupHSimulium (Nevermannia) vernum group	+00
Hydropsyche siltalaiOHydroptila sp.O* Limnephilus lunatus groupHLimnephilus extricatusOAnabolia nervosaHPotamophylax groupOHalesus sp.HTipula montium groupO* Dicranota sp.OLimnophila (Eloeophila) sp.HApsectrotanypus trifascipennisOProcladius sp.OThienemannimyia groupHPotthastia longimana groupOBrillia modestaHEukiefferiella groupHProdiamesa olivaceaOChironomus sp.OMicrotendipes sp.OParatendipes sp.OPolypedilum sp.HMicropsectra groupHMicropsectra groupHSimulium (Nevermannia) vernum group	+++
Hydroptila sp.O* Limnephilus lunatus group+Limnephilus extricatusOAnabolia nervosa+Potamophylax groupOHalesus sp.+Tipula montium groupO* Dicranota sp.OLimnophila (Eloeophila) sp.+Apsectrotanypus trifascipennisOProcladius sp.OThienemannimyia group+Potthastia longimana groupOBrillia modesta+Eukiefferiella group+Prodiamesa olivaceaOChironomus sp.OMicrotendipes sp.OParatendipes sp.OPolypedilum sp.+Phaenopsectra groupOSimulium (Nevermannia) vernum group+	00+
 Limnephilus lunatus group Limnephilus extricatus Anabolia nervosa Potamophylax group Potamophylax group Halesus sp. Tipula montium group Dicranota sp. Limnophila (Eloeophila) sp. Apsectrotanypus trifascipennis Procladius sp. Thienemannimyia group Potthastia longimana group Brillia modesta Eukiefferiella group Prodiamesa olivacea Chironomus sp. Microtendipes sp. Paratendipes sp. Polypedilum sp. Phaenopsectra group Simulium (Nevermannia) vernum group 	0+0
Limnephilus extricatus 0 Anabolia nervosa + Potamophylax group 0 Halesus sp. 9 Tipula montium group 0 * Dicranota sp. 0 Limnophila (Eloeophila) sp. 4 Apsectrotanypus trifascipennis 0 Procladius sp. 0 Thienemannimyia group + Potthastia longimana group 0 Brillia modesta + Eukiefferiella group + Synorthocladius semivirens 0 Cricotopus group + Prodiamesa olivacea 0 Chironomus sp. 0 Microtendipes sp. 0 Paratendipes sp. 4 Polypedilum sp. 4 Micropsectra group 0 Simulium (Nevermannia) vernum group +	+00
Anabolia nervosa+Potamophylax group0Halesus sp.+Tipula montium group0* Dicranota sp.0Limnophila (Eloeophila) sp.+Apsectrotanypus trifascipennis0Procladius sp.0Thienemannimyia group+Potthastia longimana group0Brillia modesta+Eukiefferiella group+Prodiamesa olivacea0Chironomus sp.0Microtendipes sp.0Polypedilum sp.+Phaenopsectra sp.0Simulium (Nevermannia) vernum group+	00+
Potamophylax groupOHalesus sp.+Tipula montium groupO* Dicranota sp.OLimnophila (Eloeophila) sp.+Apsectrotanypus trifascipennisOProcladius sp.OThienemannimyia group+Potthastia longimana groupOBrillia modesta+Eukiefferiella group+Synorthocladius semivirensOCricotopus group+Prodiamesa olivaceaOChironomus sp.OParatendipes sp.OPolypedilum sp.+Phaenopsectra sp.+Micropsectra group+Simulium (Nevermannia) vernum group+	+00
Halesus sp.+Tipula montium groupO* Dicranota sp.OLimnophila (Eloeophila) sp.+Apsectrotanypus trifascipennisOProcladius sp.OThienemannimyia group+Potthastia longimana groupOBrillia modesta+Eukiefferiella group+Synorthocladius semivirensOCricotopus group+Prodiamesa olivaceaOChironomus sp.OMicrotendipes sp.OParatendipes sp.+Phaenopsectra sp.+Micropsectra group+Simulium (Nevermannia) vernum group+	0+0
Tipula montium groupO* Dicranota sp.OLimnophila (Eloeophila) sp.+Apsectrotanypus trifascipennisOProcladius sp.OThienemannimyia group+Potthastia longimana groupOBrillia modesta+Eukiefferiella group+Synorthocladius semivirensOCricotopus group+Prodiamesa olivaceaOChironomus sp.OMicrotendipes sp.OParatendipes sp.+Phaenopsectra sp.+Micropsectra group+Simulium (Nevermannia) vernum group +	+00
 Dicranota sp. Limnophila (Eloeophila) sp. Apsectrotanypus trifascipennis Procladius sp. Thienemannimyia group Potthastia longimana group Brillia modesta Eukiefferiella group Synorthocladius semivirens Cricotopus group Prodiamesa olivacea Chironomus sp. Microtendipes sp. Polypedilum sp. Phaenopsectra group Simulium (Nevermannia) vernum group 	00+
Limnophila (Eloeophila) sp. + Apsectrotanypus trifascipennis 0 Procladius sp. 0 Thienemannimyia group + Potthastia longimana group 0 Brillia modesta + Eukiefferiella group + Synorthocladius semivirens 0 Cricotopus group + Prodiamesa olivacea 0 Chironomus sp. 0 Microtendipes sp. 0 Paratendipes sp. 0 Polypedilum sp. + Phaenopsectra sp. 4 Micropsectra group 0 Simulium (Nevermannia) vernum group +	0+0
Apsectrotanypus trifascipennisOProcladius sp.OThienemannimyia group+Potthastia longimana groupOBrillia modesta+Eukiefferiella group+Synorthocladius semivirensOCricotopus group+Prodiamesa olivaceaOChironomus sp.OMicrotendipes sp.OParatendipes sp.OPolypedilum sp.+Phaenopsectra sp.HMicropsectra groupOSimulium (Nevermannia) vernum group+	+00
Procladius sp.OThienemannimyia group+Potthastia longimana groupOBrillia modesta+Eukiefferiella group+Synorthocladius semivirensOCricotopus group+Prodiamesa olivaceaOChironomus sp.OMicrotendipes sp.OParatendipes sp.+Phaenopsectra sp.+Micropsectra group-Simulium (Nevermannia) vernum group +	0+0
Thienemannimyia group+Potthastia longimana groupOBrillia modesta+Eukiefferiella group+Synorthocladius semivirensOCricotopus group+Prodiamesa olivaceaOChironomus sp.OMicrotendipes sp.OParatendipes sp.OPolypedilum sp.+Phaenopsectra sp.HMicropsectra groupOSimulium (Nevermannia) vernum group +	0+0
Potthastia longimana groupOBrillia modesta+Eukiefferiella group+Synorthocladius semivirensOCricotopus group+Prodiamesa olivaceaOChironomus sp.OMicrotendipes sp.OParatendipes sp.OPolypedilum sp.+Phaenopsectra sp.HMicropsectra groupOSimulium (Nevermannia) vernum group +	+++
Brillia modesta+Eukiefferiella group+Synorthocladius semivirensOCricotopus group+Prodiamesa olivaceaOChironomus sp.OMicrotendipes sp.OParatendipes sp.OPolypedilum sp.+Phaenopsectra sp.OSimulium (Nevermannia) vernum group+	0+0
Eukiefferiella group+Synorthocladius semivirensOCricotopus group+Prodiamesa olivaceaOChironomus sp.OMicrotendipes sp.OParatendipes sp.OPolypedilum sp.+Phaenopsectra sp.HMicropsectra groupOSimulium (Nevermannia) vernum group +	+++
Synorthocladius semivirensOCricotopus group+Prodiamesa olivaceaOChironomus sp.OMicrotendipes sp.OParatendipes sp.OPolypedilum sp.+Phaenopsectra sp.HMicropsectra groupOSimulium (Nevermannia) vernum group +	+++
Cricotopus group+Prodiamesa olivaceaOChironomus sp.OMicrotendipes sp.OParatendipes sp.OPolypedilum sp.+Phaenopsectra sp.+Micropsectra groupOSimulium (Nevermannia) vernum group +	0+0
Prodiamesa olivacea0Chironomus sp.0Microtendipes sp.0Paratendipes sp.0Polypedilum sp.+Phaenopsectra sp.+Micropsectra group0Simulium (Nevermannia) vernum group +	+++
Chironomus sp.OMicrotendipes sp.OParatendipes sp.OPolypedilum sp.+Phaenopsectra sp.+Micropsectra groupOSimulium (Nevermannia) vernum group +	0++
Microtendipes sp.OParatendipes sp.OPolypedilum sp.+Phaenopsectra sp.+Micropsectra groupOSimulium (Nevermannia) vernum group +	0+0
Paratendipes sp. 0 Polypedilum sp. + Phaenopsectra sp. + Micropsectra group 0 Simulium (Nevermannia) vernum group +	00+
Polypedilum sp. + Phaenopsectra sp. + Micropsectra group O Simulium (Nevermannia) vernum group +	0+0
Phaenopsectra sp. ++ Micropsectra group O Simulium (Nevermannia) vernum group ++	++0
Micropsectra group O Simulium (Nevermannia) vernum group +	+00
Simulium (Nevermannia) vernum group +	0++
	+00
* Simulium (Eusimulium) aureum group U	0++
* Simulium (Simulium) ornatum group 0	0++
* Empididae 0	00+

Ephydridae

·

·

DISTRIBUTION SHEET (To be completed by all Project Leaders completing commissioned research project reports. Please bind a copy of this distribution sheet as the final page in all internal (IFE) copies of the report.) AUTHORS: Wright J.P. & Furse M.T. TITLE: The macroinvertebrate fauna of the River Blythe at Cheswick Green REPORT REF: RL/T04053g1/4 MASTER COPY HELD BY: D.M. Morton REPORT ACCESS CODE (please assign a suitable code from list overleaf): Ĉ 2.DISTRIBUTION LIST (A-G standard distribution; H other) Nos. copies Date A)Contract Customer: Nature Conservancy Council 2 31.10.90 B)J.G. Jones (requires copy of title page and Т.Р. abstract only) E I C)A.D. Pickering (1 copy of all reports) 1 D/IFE Internal Coordinator for Commissioned Research 11 1 (one copy of all reports) E)Project Leader: J.F. Wright FXFBA Library, Windermere (1 copy of all reports) G)FBA Library, River Lab. (1 copy of all reports) N)OTHER:- please list below and indicate nos. copies M.T. Furse 2. 3. 4 8 1 d 8 • • TOTAL NOS COPIES MADE:

REPORT ACCESS CODES

CODE		
S	In strict confidence - restricted access	Access to named customer(s) - (could be named individuals), IFE Directorate, Project Leader and all authors.

- In confidence -С restricted access
- 'Normal' access N
- General access G

Access to customer, IFE Directorate, Project Leader, all authors, and IFE staff with permission of Project Leader.

Access to customer, and all IFE staff. Access to visitors and general public with permission of Project Leader.

General access to anyone, las required.

JB 10/89