

Defining biological assemblages (biotopes) of conservation interest in the submarine canyons of the South West Approaches (offshore United Kingdom) for use in marine habitat mapping.

Jaime S. Davies^{a,*}, Kerry L. Howell^a, Heather A. Stewart^b, Janine Guinan^{c,d} and Neil Golding^e

^aMarine Biology and Ecology Research Centre, University of Plymouth, Plymouth, PL4 8AA, UK

^bBritish Geological Survey, Murchison House, West Mains Road, Edinburgh EH9 3LA, UK

^cMarine Institute, Riuville Oranmore, Galway, Ireland

^dPresent address : INFOMAR Programme, Geological Survey of Ireland, Beggars Bush, Haddington Road, Dublin 4, Ireland

^eJoint Nature Conservation Committee, Monkstone House, City Road, Peterborough, PE1 1JY, UK

* jaime.davies@plymouth.ac.uk

Abstract

In 2007, the upper part of a submarine canyon system located in water depths between 138 and 1165 m in the South West (SW) Approaches (North East Atlantic Ocean) was surveyed over a 2 week period. High-resolution multibeam echosounder data covering 1106 km², and 44 ground-truthing video and image transects were acquired to characterise the biological assemblages of the canyons. The SW Approaches is an area of complex terrain, and intensive ground-truthing revealed the canyons to be dominated by soft sediment assemblages. A combination of multivariate analysis of seabed photographs (184-1059 m) and visual assessment of video ground-truthing identified 12 megabenthic assemblages (biotopes) at an appropriate scale to act as mapping units. Of these biotopes, 5 adhered to current definitions of habitats of conservation concern, 4 of which were classed as Vulnerable Marine Ecosystems. Some of the biotopes correspond to descriptions of communities from other megahabitat features (for example the continental shelf and seamounts), although it appears that the canyons host modified versions, possibly due to the inferred high rates of sedimentation in the canyons. Other biotopes described appear to be unique to canyon features, particularly the sea pen biotope consisting of *Kophobelemnion stelliferum* and cerianthids.

Keywords (Submarine canyons, Conservation, Deep-sea, Benthos, Habitat mapping, Biotopes, Cold-water corals, *Lophelia pertusa*)

33

34 1. Introduction

35 Submarine canyons are topographically complex features (Harris and Whiteway 2011) that
36 are incised into many of the world's continental shelves and margins (e.g. Hickey, 1995;
37 Brodeur, 2001). Canyons have been reported as containing diverse bottom types (Kottke et al.
38 2003), described as areas of high habitat heterogeneity (Schlacher et al. 2007), and are
39 suggested to enhance biodiversity on landscape scales (Vetter et al. 2010). The presence of
40 submarine canyons on the continental slope can significantly alter the hydrodynamic regime
41 of the region, thus canyons may be highly unstable environments subject to periodically
42 intense currents, debris transport, sediment slumps and turbidity flows (Shepard and Marshall,
43 1973; Inman et al. 1976; Gardner, 1989).

44

45 Canyons may act as conduits, transporting sediment and organic matter from the continental
46 shelf to the deep sea (Shepard, 1951; Heezen et al. 1955; Monaco et al. 1990), and can be
47 areas of enhanced production and species diversity as a result of the accumulation of organic
48 matter and/or upwelling of nutrient rich waters (Hickey 1995).

49

50 Submarine canyons have been suggested to play a role in generating areas of high
51 megabenthic biodiversity due to their complex topographies (Schlacher et al. 2007). Canyon
52 fauna flourish as a result of suspension feeding organisms benefiting from accelerated
53 currents within canyons (Rowe, 1971) as well as increased secondary production (Vetter et al.
54 2010) due to the exploitation of local increases in zooplankton during vertical migration
55 (Greene et al. 1988). In addition, detritivores benefit from enhanced sedimentation rates and
56 accumulated macrophytic detritus (Vetter, 1994; Harrold et al. 1998). However, a high
57 incidence of disturbance through sediment transport by intense tidal currents, turbidity

currents and detrital flows may be unfavourable to sessile invertebrate megafauna while favouring highly motile species (Rowe, 1971; Vetter and Dayton, 1999; Vetter et al. 2010). Topographic features such as canyons, which provide enhanced food supply, diverse habitats, and alter hydrodynamic activity have been described as ‘Keystone structures’ (Vetter et al. 2010). Keystone structures are defined as “distinct spatial structures providing resources, shelter or ‘goods and services’ crucial for other species” (Tews et al. 2004). Those canyons which act as keystone structures, and may be described as biodiversity hotspots, merit special attention in management (Smith et al. 2008). The inclusion of canyons as examples of topographical features that may potentially support Vulnerable Marine Ecosystems (VMEs) (FAO 2009) reflects this.

Establishing a representative network of deep-sea Marine Protected Areas offers one tool with which to address the conservation needs of the deep sea. The need to establish such networks is driven by a number of international and national policies. The United Nations Convention of the Law of the Sea (UNCLOS) is an international agreement that provides the legal basis for high seas Marine Protected Areas (UNCLOS 1982). The Convention on Biological Diversity (CBD) is an international legally binding treaty which includes within it a requirement for nations to establish a ‘comprehensive, effectively managed and ecologically representative network of Marine Protected Areas by 2020’ [(COP 10 Decision X/2) CBD 2010]. The Oslo-Paris Convention (OSPAR) is the current legal mechanism guiding international cooperation on the protection of the marine environments of the North-East Atlantic; the agreement is between 15 European countries and the European Commission. Annex V of the OSPAR convention (The convention for the protection of the Marine Environment of the North East Atlantic) lists a number of deep-sea habitats as ‘threatened or declining’, including: seamounts, *Lophelia pertusa* reefs, coral gardens, carbonate mounds, and sea pen and burrowing megafauna communities. It calls for nations to

84 establish, “an ecologically coherent network of well managed Marine Protected Areas by
85 2020” for the protection of these listed habitats.

86 Within Europe, the main legislative power for managing fisheries and marine nature
87 conservation is based on the Common Fisheries Policy and Habitats Directive (92/43/EEC).

88 The Habitats Directive (conservation of the natural habitats of wild fauna and flora) is the
89 first international tool to address the protection of selected habitats and species, listed under
90 the Directive’s Annex I (habitats) and II (species). The Habitats Directive requires member
91 states to designate and protect sites as Special Areas of Conservation (SACs). These
92 protected areas together create the *Natura 2000* sites, a network of protected areas throughout
93 the EC. Cold-water coral reefs, coral gardens and sponge dominated communities all come
94 under the definition of Annex I listed ‘reef’ habitat.

95
96 The challenge now is how to practically implement such networks given our limited
97 understanding of the deep sea ecosystem. While a number of deep-sea habitats have been
98 identified as vulnerable to anthropogenic activities (e.g. cold-water coral reefs and sponge
99 aggregations) (FAO 2008), poor knowledge of the distribution of these habitats hinders
100 conservation efforts and network planning. Additionally, it is difficult to use criteria (such as
101 those set out by the FAO) that have been developed for assessing habitat vulnerability (FAO
102 2008) as many deep-sea habitats have yet to be described, particularly in terms of their rarity,
103 resistance, resilience and vulnerability. For example, although some habitats, such as cold-
104 water coral reefs, are easily damaged from activities such as bottom trawling, it is not cold-
105 water coral reefs that are subject to repeated trawling action in the way that some soft bottom
106 deep-sea habitats are (Thrush et al. 2001). Additionally, to create the synergy needed for an
107 MPA network design, a better understanding is urgently needed of which species are present,
108 their distribution, and some detail about their connectivity; this may be achieved through the

use of physical oceanography proxies and/or knowledge about species reproduction/larval dispersal.

For nations to fulfil their legal requirements in terms of conservation they require maps that inform them of the spatial distribution of species and habitats. In light of the vast area covered by the deep sea, numerous approaches have been adopted to mapping, with a view to preserving deep-sea habitats (Harris and Whiteway, 2009; Howell, 2010). Mapping at a landscape scale (megahabitat scale of kilometres to tens of kilometres; *sensu* Greene et al. 1999), using large topographic features such as submarine canyons, allows large areas to be covered using lower resolution data, and is thus both cost and time effective. Whilst mapping at this scale may be appropriate for generalised, global conservation efforts, these mapping units have less ecological or biological meaning due to their lack of detail. Most ecological and biological processes occur at a finer scale. Therefore, the production of meaningful fine-scale habitat maps (< 1 km) which adequately take into account lateral and vertical variation within these megahabitat features is necessary. In recent years significant research effort has been focused on seamount features, adding much to our understanding of these systems (Clark et al. 2010; Howell et al. 2010a; Rowden et al. 2010; Shank, 2010). However, contrastingly, submarine canyons are more poorly sampled, and thus less well understood (De Leo et al. 2010).

To implement ecologically representative networks, biologically meaningful maps are required to inform managers on the distribution and diversity of habitats. To adequately protect species and habitats, particularly those that are listed as being of conservation interest, the approach taken needs to be at a scale that is relevant to the biology. Taking a bottom-up approach, through first defining benthic assemblages that can then act as fine-scale mapping units, cannot only be used to inform the distribution of assemblages, but may also allow the

inference of associations between biology and larger scale features (geomorphology), which may then enable these large scale features to be used for mapping across broad areas. To achieve an ecologically coherent network across regions, and globally, we need to be able to combine habitat maps originating from national and international programmes. To date deep-sea maps produced by different projects / countries are not able to be combined because of a lack of an agreed deep-sea classification system and recognised and agreed definitions of mapping units. To overcome this, standardisation of mapping practices is necessary, with consistent terms used.

To adequately protect vulnerable habitats, there is a need for clarity in the working definitions used. Habitats such as *Lophelia pertusa* reefs have been widely documented (Wilson, 1979; Mortensen et al. 1995; De Mol et al. 2002) and the definition of these habitats are more widely recognised. There are few descriptions of benthic assemblages from canyon systems (Schlacher et al. 2010), and none in the context of statistically defining units for use in habitat mapping, or assessing the potential conservation value of canyons. Consequently, the objective of this study is to: support international habitat mapping efforts through developing standardised descriptions of deep-sea biological assemblages, with a focus on assemblages that fit descriptions of 'listed' habitats, for use as functional and consistent mapping units (biotopes).

2. Material and methods

2.1 Study area

The SW Approaches study area is located on the Celtic Margin and is an area characterised by a number of submarine canyons (Figure. 1; Huthnance et al. 2001; Mulder et al. 2012). The upper reaches of three canyons were the target of this investigation. Two of those are located in UK waters: Dangeard Canyon (also known as Dangaard Canyon), and Explorer

Canyon (first in this special issue, see Stewart et al. (2014, this issue)). The head of Dangeard Canyon is around 12 km in width and ~1500 m at its deepest point, including its network of tributary gullies that feed into the main canyon which is itself around 7 km in width. The head of Explorer Canyons is around 11 km wide, compared to the main Explorer Canyon which is around 8 km in width and ~1500 m deep. We are constrained by the dataset as the canyons continue before merging downslope, feeding into the Whittard Canyon. The shelf break, which marks the boundary between the near horizontal sea floor of the continental shelf and the steeper continental slope, occurs between 180 and 250 m water depth. Mean slope angles along the Celtic Margin are 11° although locally very steep gradients to the vertical occur along canyon walls (Cunningham et al. 2005; Stewart et al. 2014). Two canyons are located in UK waters, the Dangeard (also known as Dangaard) and Explorer (first named here) canyons, and were the target of this study.

The Dangeard and Explorer canyons are separated by smooth interfluvies, which are areas of un-dissected relict continental shelf and slope (Figure 2). These interfluvies host two mini-mound provinces with individual mounds up to 3 m in height above the surrounding sea floor and 50-150 m in diameter (Stewart et al. 2014). In the canyon heads, the dendritic pattern of tributary gullies is clearly imaged in the study area forming drainage basins. Well developed “cauliflower” shaped amphitheatre rim features were identified in the canyon heads and flanks indicative of shelf-ward erosion. Stewart et al. (2014) present a geological interpretation of the study area.

2.2 Data acquisition

From 4th-18th June 2007 Dangeard and Explorer canyons and the flank of a third canyon (located in Irish waters) in the SW Approaches were surveyed onboard the *RV Celtic Explorer* (The Marine Institute, Ireland). High-resolution ground-truthing and multibeam

echosounder (MBES) data were acquired (Figures 2 and 3) over an area of 1106 km²; MBES was acquired using a hull mounted Kongsberg Simrad EM1002 system capable of collecting swath bathymetry to ~1000 m water depth (see Stewart and Davies (2007) and Stewart et al. (2014) for more details). A Seatronics drop-frame camera system was used to acquire video and image data. The camera system comprised a DTS 6000 digital video telemetry system with a live feed to the vessel, and a five megapixel Kongsberg Simrad digital stills camera (containing a Canon Powershot G5). The cameras were mounted opposite each other (with lights either side) at oblique angles to the seabed for optimal seabed coverage and to aid species identification. The frame was also fitted with a CDT sensor to record depth, altitude and temperature, and an ultra-short baseline (USBL) beacon to collect accurate positional data for the frame, allowing accurate environmental and positional data for still images to be extracted from data files. To enable quantitative analysis of data, the fields of view for both the stills and video cameras were calibrated (an image taken) at varying altitudes of the camera frame above the seabed (on seabed, 1 m, 2 m and 3 m) to enable area to be calculated. Calibration was achieved by attaching a gridded quadrat of known dimensions (grid cell size of 4.9 cm by 5.5 cm) to the base of the camera frame and the area of each still image was calculated using the appropriate calibration grid image for its altitude.

Transect locations were chosen using the processed multibeam bathymetry and backscatter data. ‘Sampling’ was stratified by depth, topographic feature, and seabed substratum (inferred from backscatter data); and where possible, replicate sampling was undertaken within and between canyons. Transect position and orientation was chosen dependent on the terrain, on the steep areas of the canyon flank it was decided that it was safer for the towed camera to travel down- rather than along-slope. The vessel’s DP was used to keep the camera on chosen transects.

Transects were *approx.* 500 m in length, and sampling occurred over a depth range of 184-1094 m. The drop-frame was deployed from the starboard side of the vessel and towed 1-3 m above the seabed at a vessel speed of *approx.* 0.5 knots (min 0.3 and max 0.7) with tows lasting between 0.5-1.5 hrs. Forty four transects were undertaken (see Table A1 for full details). Following the MESH¹ guidelines for data collection, a 2-5 minute camera stabilisation period was undertaken at the beginning of each transect to ensure the camera was moving at a constant speed. Video footage was recorded along the entire transect, and at approximately one minute intervals the drop-frame was landed and a stills image taken (sampling unit) which will be referred to here as a ‘sample’ image. Additional images were also taken to capture abrupt changes in substratum (i.e. from sand to bedrock) and to aid in species identification.

2.3 Biological data analysis

2.3.1 Quantitative analysis of image data

‘Sample’ images and those taken at abrupt changes in substratum were reviewed and poor quality images removed, predominantly due to silt clouds obscuring the image or the image being out of focus. The remaining images were quantitatively analysed using image area (derived from the calibration grids). An inherent problem with working in the deep sea is the lack of specimens to aid in identification, and without physical samples it is difficult, and in many cases impossible to identify organisms to species level from image data; however, observed organisms can be identified as distinct morphospecies (morphotypes).

All visible organisms >1 cm (at their widest point), as determined using the calibration grid for scale, were identified as distinct morphospecies and assigned an Operational Taxonomic

¹ The principal purpose of the Mapping European Seabed Habitats (MESH) project is to harmonise the way in which habitat mapping initiatives are undertaken in the northwest Europe (www.searchmesh.net).

Unit (OTU) number. OTUs were identified to the lowest possible taxonomic level, which can correspond to species, genus, family or higher taxonomic levels depending on the group. All individuals were enumerated except in the case of encrusting, colonial and lobose forms where area cover was used.

2.3.2 Community analysis

Count and cover data were treated independently prior to multivariate analysis, each were standardised to 1 m² (percent/1 m² for cover). To allow combined analysis of count and percent cover data, a standardisation function was employed to place each matrix on the same scale (Stevens and Connolly, 2004; Howell et al. 2010b). First the data were transformed to standardise the distribution of the data then each entre in the matrix was divided by the sum of the matrix total and multiplied by an appropriate factor to put the count and cover on relative scales (Prof. R. Clarke *pers. comm*). Count data were square root transformed, each entre divided by the sum of the matrix and multiplied by 200; cover data were 4th root transformed, divided by the sum of the matrix and multiplied by 100, to place both matrices on a scale of 0.01-1.019. Once each matrix was standardised, they were merged in PRIMER (v.6) and multivariate community analysis was undertaken as described below. Seabed substratum composition was assigned to each 'sample' image using the modified Folk diagram (Folk 1954; Long 2006).

Standard multivariate community analysis techniques were used to identify faunally distinct benthic assemblages within the study area. Highly mobile species such as fish, which use multiple habitats and can thus confound the result of the cluster analysis, were removed prior to data analysis. Cluster analysis with group-averaged linkage was performed using a Bray-Curtis similarity matrix derived from transformed (standardised), combined species count and percent cover data. The SIMPROF routine of the PRIMER software [similarity profile

(Clarke et al. 2008)] was used to identify significant clusters ($p < 0.01$) and the SIMPER [similarity percentages (Clarke, 1993)] routine used to identify those species that characterise those clusters. Characterising species were defined as those species with a high similarity/standard deviation ratio (Clarke, 1993), and contributed $> 5\%$ to that cluster similarity.

2.3.3 Characterising mapping units (biotopes)

There is a discrepancy between the faunal assemblages identified using community analysis methods and what is required from a practically applicable mapping unit used in producing necessarily generalised maps of variation in the biological composition of the seabed. Clusters identified by SIMPROF ($p < 0.01$) were assessed against the following criteria and rejected or accepted as faunally distinct clusters on that basis: 1) Outlier clusters were taken at a 1% Bray-Curtis similarity level on the dendrogram and discarded. 2) Clusters that contained small numbers of images (in this study less than 7 images) were deemed not sufficient to allow an adequate description of a coherent assemblage and were also discarded. 3) Those clusters that had an average similarity (SIMPER) of less than 15% were defined as not being coherent. 4) In line with existing habitat classification systems (e.g. EUNIS, (Davies and Moss, 1999-2002), SIMPROF clusters were split on the basis of substratum type. 5) SIMPROF clusters were combined at a lower similarity node on the dendrogram if it produce a more practical mapping unit (appropriate scale).

Following standard multivariate analysis, faunally distinct clusters were assessed against a second set of criteria to determine their use as mapping units. Only those clusters that subsequently met these criteria were further analysed in terms of their faunal composition. To function as a mapping unit assemblages must 1) occur at a scale relevant to the resolution of

the acoustic data and the scale of existing widely accepted benthic communities such as cold water coral reefs (e.g. 10 m scale), and 2) be easily identified from video data.

Mapping units, hereinafter referred to as 'biotopes', were defined in terms of their characterising species, as determined by SIMPER analysis, together with the range of environmental conditions over which they occurred in this study, and named according to the dominant species, in accordance with the EUNIS classification system. As a result of the small size of the sampling unit (field of view of the image 'samples') the larger conspicuous fauna were not always adequately sampled, thus additional descriptive elements were added from video observations. A 1-way Analysis of Similarity (ANOSIM) was performed on a normalised depth and temperature, Euclidean distance matrix to test if biotopes (factor) were different in terms of measured environmental variables.

To identify those biotopes which could be considered of conservation concern, biotopes were compared with current definitions of 'listed' habitats under the OSPAR Convention and the EC Habitats Directive. Specifically, to identify those which are VMEs, the guidelines of the FAO (FAO 2009) and current OSPAR definitions were used (OSPAR (Agreement 2008-6).

2.3.4 Distribution of biotopes

Video transects were reviewed and visually classified (guided by the sample image classification) using the newly defined biotopes, and changes of biotope type within a transect were mapped using ArcGIS 9.3 Abiotic data were extracted from the mapped data to define the environmental range of the distribution of each biotope.

3. Results

3.1 Biological data analysis

Twenty three hours of video footage and 5000 still images were collected over the survey area. Of these images, 1073 were ‘sample’ images [those taken at *approx.* 1 minute intervals (equating to ~30 m)]; upon inspection 199 were discarded due to poor quality.

3.1.1 Quantitative analysis of image data

Eight hundred and seventy four ‘samples’ were quantitatively analysed with 161 morphospecies identified and catalogued. Those samples where no fauna were recorded were removed prior to the multivariate analysis. Cluster analysis was performed on the remaining 746 samples. Three broad categories of substratum were revealed from the image analysis: hard substratum (16 %), reef habitats (4%) and soft substratum (80%).

3.1.2 Community analysis

The SIMPROF routine identified 43 clusters ($p < 0.01$) (see Table A2 for statistical results of clusters). Using the criteria described in Sect. 2.3.3, outlier clusters were removed (cluster a-q) and those that did not act as coherent units for mapping discarded. The remaining 11 clusters were accepted as practically applicable mapping units. Results from the cluster analysis of still image “samples”, including SIMPER analysis (characterising species) and a description of the environmental characteristics associated with each cluster are shown in Table A2 (see appendix A1 for SIMPER results).

3.1.3 Characterising mapping units (biotopes)

In total 11 biotopes were identified from the cluster analysis (Figure. 4) and related to available environmental data to describe distinct biotopes (see Table 1 for details). A 1-way ANOSIM test of environmental data (depth and temperature) for the 11 biotopes defined from image data revealed a significant difference in environmental conditions between

biotopes (Global $R = 0.265$, $p < 0.01$). Thirty one pairwise tests were significant and Fig. 5 illustrates an nMDS plot showing a variation of biotopes relating to environmental conditions. Two groups are apparent and appear to be related to depth zones, one on the left comprising of 5 biotopes (x, y, al, ac and aj) a deeper zone (654-894 m average depth of biotopes) and the other having 4 biotopes (am, aq, ap and ao) at shallower depths (326-477 m average depth of biotopes). Biotope r and ah are most dissimilar, although appear not to be strongly related to either of the main groups observed in Figure. 5.

Visual classification of video data according to the newly defined biotopes revealed an assemblage that did not fit with any of those defined (Lop.Cri: *L. pertusa* and crinoids on bedrock). Upon reviewing the data, it was apparent that image sample data had failed to capture this assemblage (due to limited areas of bedrock captured by the still images). Based on visual assessment of the assemblage it appears similar to assemblages described by Wienberg et al. (2008) and Howell et al. (2010b) and was therefore classified as such. In the interests of fully characterising the Canyons region, and given that this previously described biotope is of particular conservation importance due to the occurrence of listed species (*L. pertusa*), as well as being the only bedrock community observed in the canyons that may be classed as Annex I bedrock reef (under the EC Habitats Directive), its distribution within the canyon system is also considered here. Thus a total of 12 biotopes were described from the SW Approaches (Figure 6).

3.1.4 Distribution of biotopes

Qualitative assessment of biotope distribution, determined from visually classified video transect data, (Table 1, see also Fig A1-A2 for mapped distribution of biotopes) revealed that six of the 12 biotopes were observed in all 3 canyons, 4 soft sediment biotopes (Kop.Cer, Cer, Amp.Cer and Oph), a mixed substratum (shell hash) biotope (Mun.Lep) and Lop.Cri on

bedrock. Five biotopes fit with the ‘listed habitats’ definition. The sea pen biotope Kop.Cer was observed in all three canyons on the flank and incised channels over a depth of 463-1059 m. The bedrock associated biotope, Lop.Cri, was also observed in all canyons, occurring on incised channels, tributary gullies, flank and amphitheatre rims features over a depth of 253-1022 m. The *L. pertusa* reef biotope Lop.Mad was only observed once in Explorer canyon on flute features 795-940 m, while the dead framework biotope Lop.Hal was observed in both Explorer and Dangeard canyons on the flanks and flute features (697-927 m). The coral rubble biotope Oph.Mun was observed in Explorer and Dangeard canyons on incised channel and mini-mound features over a depth of 303-1017 m.

4. Discussion

Submarine canyons are considered to be potential biodiversity hotspots; however, to date there is very little data on canyon community composition of these features, particularly potential importance as features of conservation interest. Soft sediment habitats dominate the canyons of the SW Approaches, with 80% of analysed images and 60% of the described biotopes. Five of the biotopes could be considered of conservation interest. Of these five, only four come under the definition of VMEs, three could be classified as cold-water coral reefs under the EC Habitats Directive and OSPAR Convention, whilst the fourth could be classed as ‘Sea pen and burrowing megafauna communities’ or coral garden under the current OSPAR definition. The fifth could be considered bedrock reef under the EC Habitats Directive. Seven biotopes were soft sediment communities or faunally-sparse and thus, have little or no perceived conservation interest; of these, three have been previously described by a number of authors while four are new descriptions (see Appendix A2 for descriptions). Those habitats that are listed under policy (OSPAR and EC Habitats Directive) will be discussed in terms of a description of the new biotopes defined and related to other research,

those which are not ‘listed’ habitats will not be discussed; however full descriptions for each are given in Appendix A2.

4.1 Descriptions of ‘listed’ habitats for use as mapping units (biotopes)

4.1.1 Cold-water coral reef

Three biotopes were defined that could be considered as cold-water coral reef, these communities represent distinct reef zones (*sensu* Mortensen et al. 1995) or macrohabitats (*sensu* Greene et al. 1999) each with different associated fauna forming distinct communities.

Lophelia pertusa reef

This biotope (Lop.Mad, cluster ah) was characterised by dead *L. pertusa* framework and live patches of *L. pertusa* and *Madrepora oculata* which provide a structural habitat for associated species. Other characterising species (as identified by SIMPER) were small anemones (*Actiniaria* sp.13) and an unidentified species (Unknown sp.26) which were associated with *L. pertusa*. Additional species identified from qualitative video observations were *Pandalus borealis* and the echinoid *Cidaris cidaris*; halcampaoid anemones (*Halcampoididae* sp.1) inhabited the interspersed sediment patches in the reef. Other conspicuous fauna observed from the image and video data were large cerianthid anemones, the decapod *Bathynectes* sp. and the fish *Lepidion eques*. This assemblage was observed on steep flute features on the flank of Explorer canyon over a depth of 795-940 m and temperature of 9.41-9.92°C.

This assemblage corresponds to the ‘live *Lophelia* zone’ as described by Mortensen et al. (1995) which is the main reef habitat found on the summit of the reef and consists of predominantly live *L. pertusa* interspersed with areas of dead broken skeleton.

Lophelia pertusa is widely distributed in the North Atlantic, in oceanic waters at temperatures of 4-12°C (Roberts et al. 2006) and is predominantly found at depths of 200-1000 m but has been recorded shallower and deeper (Zibrowius, 1980). *L. pertusa* has been identified as occurring in areas subjected to fast currents such as carbonate mounds (De Mol et al. 2002), ridges and pinnacles (Howell et al. 2007). Pfannkuche et al. (2004) observed *L. pertusa* reef on the slopes of the Castor mound in the Belgica mound province (Porcupine Seabight) from 950-1036 m depth, and describe complete cover of live and dead coral colonies of *L. pertusa* and *Madrepora oculata* with antipatharians, actinians and hexactinellid sponges present. Howell et al. (2010b) described a similar *L. pertusa* reef from various locations within UK waters as being characterised by the reef-forming corals *L. pertusa* and *M. oculata*, hydroids, anemones, decapods, cerianthids and echinoderms (ophiuroids and echinoids); whilst a similar assemblage was observed from Anton Dohrn Seamount (Davies et al. subm.) consisting of *L. pertusa* (dead and live), *M. oculata*, *Cidaris cidaris* and anemones.

Whilst the assemblage defined from the SW Approaches canyons has some of the same associated species as described previously from reef habitat, the canyon assemblage appears to be subject to increased sedimentation which is clearly visible from the image and video data; although an analysis of sedimentation rates has not been carried out. Canyons are likely to experience increased rates of sediment transport as a result of hydrodynamic regime (Vetter and Dayton, 1998). The interpreted higher level of sedimentation in the study area may result in a lower proportion of live *L. pertusa* colonies and fewer suspension feeders (Brooke and Ross, 2014); however, a full comparative analysis would be required to test this.

Predominantly dead low-lying coral framework

The assemblage identified as Lop.Hal (cluster aj) was characterised by small live colonies of *L. pertusa* and dead *L. pertusa* framework with sediment infill, the sediment areas provided

microhabitats for soft sediment dwelling organisms such as cerianthid (Cerianthidae sp. 1) and halcampoid (Halcampoididae sp.1) anemones. Fauna associated with the dead framework were small growths of live *Madrepora oculata*, the bamboo coral *Acanella*, ascidians and crinoids. This assemblage was observed from the Explorer and Dangeard canyons on the flanks, and on a flute feature over a depth of 697-927 m and temperature of 8.97-9.77°C.

Mortensen et al. (1995) and Roberts et al. (2009) describe a 'Dead coral framework' zone that is characterised by suspension feeders including sponges, actinians, and other coral species (gorgonians) with smaller epifauna such as bryozoans, hydroids and barnacles. Similar assemblages have also been described from Rockall Bank (Wilson 1979; Howell et al. 2010b), Hatton Bank (Howell et al. 2010b) and Anton Dohrn Seamount (Davies et al. subm.). The 'Dead coral framework' zone (*sensu* Mortensen et al. 1995) is known to be the most diverse area of a reef (Jensen and Frederiksen, 1992; Mortensen et al. 1995). Whilst the assemblage described by the present study may be functionally similar to the dead framework assemblages of Wilson (1979), Mortensen et al. (1995) Roberts et al. (2009) and Howell et al. (2010b), based on their descriptions it would appear this assemblage is more sediment infilled, as there are more sediment dwelling organisms associated with this biotope. A similar assemblage has been reported on the upper slope and summit of Erik mound in the Belgica province from 818-855 m depth (Pfannkuche et al. 2004). Coral rubble with isolated live patches of *L. pertusa* and *M. oculata* and a low abundance of associated fauna (antipatharians and *Aphrocallistes* sp.) was described with muddy sand areas between the rubble inhabited by *Cerianthus* sp. (Pfannkuche et al. 2004).

Ophiuroids and *Munida sarsi* associated with coral rubble

Biotope Oph.Mun (cluster ap) was identified as a typical reef rubble habitat which was characterised by coral fragments in the form of rubble/biogenic gravel. The rubble was acting as a habitat for the squat lobster *Munida sarsi* and the ophiuroid Ophiuroidea sp.5. The assemblage was found associated with incised channels and mini-mound features on the interfluves in Explorer and Dangeard canyons over a depth range of 303-1017 m and a temperature of 7.98-11.5°C.

Oph.Mun biotope corresponds to 'the *Lophelia* rubble zone' described by Mortensen et al. (1995) which is the outer 'apron' of the reef where the framework has been (bio)eroded and accumulates at the base of the reef, the squat lobster *Munida sarsi* dominates this zone.

4.1.2 'Sea pen and burrowing megafauna' communities/coral gardens

Kophobelemnon stelliferum and cerianthids on mud/sand

The assemblage Kop.Cer (cluster y) was associated with mud and muddy sand substratum and was characterised by the sea pen *Kophobelemnon stelliferum* and cerianthid anemone. Other conspicuous fauna associated with this assemblage were the large *Bolocera*-like anemones (Sagartiidae sp. 3), sea pens *Halipteris* sp., a number of echinoderm species including the asteroid *Pseudarchaster* sp., the crinoid *Pentametrocrinus atlanticus* (sediment dwelling) and the holothurian *Benthogone* sp. Video observations revealed the bamboo coral *Acanella arbuscula* to be more abundant than suggested from the image analysis. Kop.Cer biotope was observed most frequently and was widespread throughout the canyons. The

assemblage was observed from all three canyon flanks, and from an incised channel in Explorer Canyon, over a depth range of 463-1059 m and a temperature of 8.87-10.85°C.

Kophobelemnion stelliferum is an upper bathyal species (Rice et al. 1992) and is known to be a deeper sea pen species (López-González and Williams, 2010) widely distributed at depth from 400-2500 m in the north Atlantic and Pacific oceans (Rice et al. 1992). Rowe (1971) reported the occurrence of a *K. stelliferum* from Hatteras canyon between 1440-2060 m and considered this species to be a ‘canyon indicator’ as it was not found away from the canyon. Whether this assemblage is unique to the canyon system here is unknown as no comparable data are available from the neighbouring continental slope.

The sea pen assemblage has not been described from the deep sea but is similar to the shallower EUNIS ‘Sea pen and burrowing megafauna in circalittoral mud’ biotope and that described by Kenchington et al. (2014). Kenchington et al. (2014) describe a biotope from the Gully Canyon characterised by 3 corals, the sea pens *Pennatula* spp. and *Halopteris* spp. and the small soft coral *Acanella arbuscula*. A xenophyophore biotope with an abundance of sea pens has also been described from Anton Dohrn Seamount (Davies et al. subm.), although this community is distinct from that observed on Anton Dohrn Seamount.

Sea pens are known to increase local biodiversity through increased habitat heterogeneity (Birkeland, 1974; Buhl-Mortensen et al. 2010). Sea pens are protected under the UK Biodiversity Action Plan (UKBAP) as ‘Mud habitats in deep water’ which corresponds to the OSPAR ‘Threatened and/or Declining Habitat’ ‘Sea pen and burrowing megafauna communities’ (OSPAR Agreement 2008-6). The newly described assemblage could also be considered both a VME (FAO 2009) and a ‘coral garden’ habitat (OSPAR 2010). The OSPAR definition is very broad and incorporates both hard and soft substratum assemblages;

this may lead to misinterpretation, and thus misrepresentation of this habitat within a network of MPAs. Soft-bottom coral gardens can be dominated by solitary scleractinians (caryophyllids), sea pens or certain types of bamboo corals (e.g. *Acanella* sp.), whilst hard-bottom coral gardens are often found to be dominated by gorgonians, stylasterids, and/or black corals (ICES, 2007). The '*Kophobelemnion stelliferum* and cerianthid' biotope described from the submarine canyons of the SW Approaches may also satisfy the criteria for being classed as a VME. This assemblage is 'unique or rare' in the sense that it may be unique to canyons, and sea pens are known to be vulnerable to fishing activities (Troffe et al. 2006) and provide structural complexity for associated species (Buhl-Mortensen et al. 2010). They may also be important nursery grounds for fish, for example, Redfish larvae have been associated with 5 species of sea pen in the northwest Atlantic (Baillon et al. 2012).

4.1.3 Other reef habitat under EC Habitats Directive

L. pertusa and crinoids on bedrock

As this biotope was described from the video, characterising species were assessed visually. Small growths of *Lophelia pertusa* (live & dead), the holothurian *Psolus squamatus* and Holothuroidea sp.4; the corkscrew antipatharian *Stichopathes* sp. and crinoids were identified as characterising species from video. The assemblage was associated with bedrock and was observed from the Dangeard, Explorer and Irish canyons associated with incised channels, amphitheatre rims, tributary gullies (canyons heads) and the flanks over a depth of 253-1022 m and temperature range of 7.93-11.42°C.

The assemblage appears to be a highly sedimented version of the 'Discrete coral' biotope described by Wienberg et al. (2008) and Howell et al. (2010b). The assemblage described by Wienberg et al. (2008) was associated with ridge features on the flanks of Rockall Bank between 650-675 m and dominated by a diverse range of corals (gorgonians, antipatharians, soft corals and stylasterids); whilst Howell et al. (2010b) describe a modified version of this

assemblage from Hatton Bank with a lower proportion of gorgonians and antipatharians but with the addition of *L. pertusa*.

4.2 Potential modelling use of biotope data

It is generally recognised that organisms show a particular affinity for certain types of topographical features or terrain (Džeroski and Drumm 2003) and multibeam bathymetry and derived terrain variables can potentially provide important information that can aid in the delineation and characterisation of biological communities (Wilson et al. 2007). Typically, surrogates used in habitat mapping are parameters that can be derived directly from the acoustic multibeam data, such as slope, aspect, rugosity, Bathymetric Position Index (BPI) and backscatter strength.

Once biotopes have been characterised, it is possible to use predictive modelling technique to map their distribution using such surrogates. This has been achieved for single species mapping (e.g. Davies & Guinotte, 2011) and has recently been applied to habitat mapping (Dolan et al. 2008; Guinan et al. 2009; Howell et al. 2011; Ross and Howell, 2012). However, to date, there are few examples of this approach being applied in the deep sea. Where this approach has been applied in the deep sea, it has generally been either on a basin-wide scale (Davies & Guinotte, 2011), or over small areas focused on specific habitats (using ROV acquired resolution acoustic data), such as cold water coral reefs (Dolan et al. 2008; Anderson et al. 2011), seeps (Sager et al. 1999; Baco et al. 2010) or vents (Desbruyères et al. 2001; Kelley et al. 2001), using project specific mapping units (or facies / biotopes).

Multibeam bathymetry data and its derived layers have proved significant in mapping and predicting the distribution of benthic assemblages in the deep sea (e.g. Ross and Howell, 2012; Knudby et al. 2013). However submarine canyons are complex topographic features that are often associated with increased sedimentation rates and sediment transport, and are

often hydrodynamically complex (Shepard 1951; Heezen et al. 1955) . The degree to which topographic variables are able to act as surrogates for the environmental parameters important in determining species and assemblage distributions within these complex environments is unknown. Studies which undertake predictive modelling mapping approaches validated using independent data are required to further elucidate the effectiveness of predictive modelling the distribution of habitats and species in submarine canyons.

5. Conclusion

With easily recognised, defined biological assemblage units, the identification of assemblages that could be considered VMEs becomes much simpler and more comprehensive, i.e. not restricted to those communities that have received the most research attention. Efforts to map the distribution of VMEs are more easily combined across studies and / or regions. In addition, the classification of all benthic assemblages into named ‘habitats’ allows a more effective assessment of representativeness of a network, and consideration of anthropogenic impacts on habitats other than those that are highly ‘charismatic’, such as cold water coral assemblages.

The SW Approaches submarine canyons harbour a range of biological assemblages, some of which correspond to those described from other megahabitat features, such as seamounts or the continental shelf. Other assemblages may be unique to canyons, but this is merely speculative as there is little comparable data. The SW Approaches canyons harbour assemblages of conservation concern, including three *L. pertusa* biotopes, one sea pen and burrowing megafauna biotope, and one bedrock reef and thus could be considered a keystone structure. The findings of this work have extended our knowledge of submarine canyons by providing much needed, comprehensive descriptions of biological assemblages, and suggest

that canyons may harbour modified versions of assemblages observed on other megahabitat features.

Acknowledgments

The authors wish to thank the captain, crew and scientific complement of the *R/V Celtic Explorer* for assistance in data collection during MESH Cruise 01-07-01, Joint copyright © 2007 Defra, JNCC, Marine Institute, BGS, UoP. These data were recorded during a collaborative survey involving the Joint Nature Conservation Committee, the Marine Institute Galway, the British Geological Survey and the University of Plymouth. The Department of the Environment, Fisheries and Rural Affairs (Defra) Natural Environment Group Science Division (CRO 361) made a significant financial contribution to this work. This work contributes to the MESH project (www.searchmesh.net) that receives European Regional Development Funding through the INTERREG IIIb Community Initiative (www.nweurope.org). HAS publishes with permission of the Director, British Geological Survey (Natural Environment Research Council).

References

- Allen, S.E., Vindeirinho, C., Thomson, R.E., Foreman, M.G.G., Mackas, D.L., 2001. Physical and biological processes over a submarine canyon during an upwelling event. *Can. J. Fish. Aquat. Sci.* 58, 671-684.
- Anderson, T.J., Nichol, S.L., Syms, C., Przeslawski, R. and Harris, P.T., 2011. Deep-sea biophysical variables as surrogates for biological assemblages, an example from the Lord Howe Rise. *Deep-sea Research II* 57, 979-991.

- Baco, A.R., Rowden, A.A., Levin, L.A., Smith, C.R. and Bowden, D.A., 2010. Initial characterization of cold seep faunal communities on the New Zealand Hikurangi margin. *Marine Geology* 272(1-4), 251-259.
- Baillon, S., Hamel, J.-F., Wareham, V.E., Mercier, A. 2012. Deep cold-water corals as nurseries for fish larvae. *Frontiers in Ecology and the Environment Research Communication*. 10 (7), 351-356.
- Birkeland, C., 1974, Interactions between a sea pen and seven of its predators. *Ecol. Monogr.* 44, 211-232.
- Brodeur, R.D., 2001. Habitat-specific distribution of Pacific ocean perch (*Sebastes alutus*) in Pribilof Canyon, Bering Sea. *Cont. Shelf Res.* 21(3), 207-224.
- Brooke, S., Ross, S.W., 2014, this issue. First observations of the cold-water coral *Lophelia pertusa* in Mid-Atlantic canyons of the USA. *Deep-Sea Research II*.
- Buhl-Mortensen, L., Vanreusel, A., Gooday, A.J., Levin, L.A., Priede, I.G., Buhl-Mortensen, P., Gheerardyn, H., King, N.J., Raes, M., 2010. Biological structures as a source of habitat heterogeneity and biodiversity on the deep ocean margins. *Mar. Ecol.* 31, 21-50.
- Cartes, J.E., Company, J.B., Maynou, F., 1994. Deep-water decapod crustacean communities in the Northwestern Mediterranean: influence of submarine canyons and season. *Mar. Biol.* 120, 221-229.

Clark, M.R., Rowden, A.A., Williams, A., Consalvey, M., Stocka, K.I., Rogers, A.D., O'Hara, T.D., White, M., Shank, T.M., Hall-Spencer, J.M., 2010. The ecology of seamounts: structure, function, and human impacts. *Annual Review of Marine Science* 2, 253-278.

Clarke, K.R., 1993. Non-parametric multivariate analyses of changes in community structure. *Australian J. Ecol.* 18, 117-143.

Clarke, K.R., Somerfield, P.J., Gorley, R.N., 2008. Testing the null hypotheses in exploratory community analyses: similarity profiles and biota-environment linkage. *J. Experimental Mar. Biol. Ecol.* 366, 56-69.

Davies, A.J. & Guinotte, J.M., 2011. Global habitat suitability for framework-forming cold-water corals. *Plos One*, 6(4), e18483. doi:10.1371/journal.pone.0018483.

Davies, C.E., Moss, D., 1999-2002. EUNIS Habitat Classification. European Environment Agency.

Davies, J.S., Howell, K.L., Stewart, H., Narayanaswamy, B.E., Jacobs, C., Spicer, J., Golding, N., 2014 (submitted). Benthic assemblages of the Anton Dohrn Seamount (NE Atlantic): defining deep-sea biotopes to support habitat mapping and management efforts with a focus on Vulnerable Marine Ecosystems.

De Leo, F.C., Smith, C.R., Rowden, A.A., Bowden, D.A., Clark, M. R., 2010. Submarine canyons: hotspots of benthic biomass and productivity in the deep sea. *Proc. Royal Soc. B: Biol. Sci.* 277, 2783-2792.

- De Mol, B., Van Rensbergen, P., Pillen, S., Van Herreweghe, K., Van Rooij, D., McDonnell, A., Huvenne, V., Ivanov, V., Swennen, R., Henriët, J.P., 2002. Large deep-water coral banks in the Porcupine Basin, southwest of Ireland. *Mar. Geol.* 188, 193-231.
- Desbruyères, D., Biscoito, M., Caprais, J.C., Colaço, A., Comtet, T., Crassous, P., Fouquet, Y., Khripounoff, A., Le Bris, N., Olu, K., Riso, R., Sarradin, P.M., Segonzac, M. and Vangriesheim, A., 2001. Variations in deep-sea hydrothermal vent communities on the Mid-Atlantic Ridge near the Azores plateau. *Deep Sea Research Part I: Oceanographic Research Papers* 48(5), 1325-1346.
- Dolan, M.F.J., Grehan, A.J., Guinan, J.C. and Brown, C., 2008. Modelling the local distribution of cold-water corals in relation to bathymetric variables: Adding spatial context to deep-sea video data. *Deep-sea Research I* 55, 1564-1579.
- Duineveld, G., Lavaleye, M., Berghuis, E., Wilde, P., 2001. Activity and composition of the benthic fauna in the Whittard Canyon and the adjacent continental slope (NE Atlantic). *Oceanologica Acta* 24(1), 69-83.
- Džeroski, S. and Drumm, D., 2003. Using regression trees to identify the habitat preference of the sea cucumber (*Holothuria leucospilota*) on Rarotonga, Cook Islands. *Ecological Modelling* 170, 219-226.
- Emery, K.O., Hülsemann, J., 1963. Submarine canyons of southern California. Part 1, Topography, water and sediment.
- Etter, R.J., Grassle, J.F., 1992. Patterns of species diversity in the deep-sea as a function of sediment particle size diversity. *Nature* 360, 576-578.

695

696 FAO (2008). Report of the FAO Workshop on Vulnerable Ecosystems and Destructive
697 Fishing on Deep-sea Fisheries. Rome, 26-29 June 2007. FAO Fisheries Report. No. 829.

698

699 FAO (2009). International Guidelines for the Management of Deep-sea Fisheries in the High
700 Seas. Rome.

701

702 Flach, E., Heip, C., 1996. Vertical distribution of macrozoobenthos within the sediment on
703 the continental slope of the Goban Spur area (NE Atlantic). Mar. Ecol. Prog. Ser. 141(1-3),
704 55-66.

705

706 Flach, E., Thomsen, L., 1998. Do physical and chemical factors structure the macrobenthic
707 community at a continental slope in the NE Atlantic? Hydrobiologia 376, 265-285.

708

709 Folk, R.L., 1954. The distinction between grain size and mineral composition in sedimentary
710 rock nomenclature. J. Geol. 62, 344-359.

711 Gage, J.D., 1986. The benthic fauna of the Rockall Trough: regional distribution and
712 bathymetric zonation. Proc. Royal Soc. Edinburgh 88B, 159-174.

713

714 Gage, J.D., 2001. Deep-sea benthic community and environmental impact assessment at the
715 Atlantic Frontier. Cont. Shelf Res. 21 (8-10), 957-986.

716

717 Gage, J.D., Lamont, P.A., Tyler, P.A., 1995. Deep-sea macrobenthic communities at
718 contrasting sites off Portugal, preliminary results. 1. Introduction and diversity comparisons.
719 Internationale Revue Der Gesamten Hydrobiologie 80(2), 235-250.

720

- Gardner, W.D., 1989. Baltimore Canyon as a modern conduit of sediment to the deep sea. *Deep-Sea Res.* 36, 323-358.
- Greene, C.H., Weibe, P.H., Burczynski, J., Youngbluth, M.J., 1988. Acoustical detection of high-density demersal krill layers in the submarine canyons off Georges Bank. *Science* 241, 359-361.
- Greene, H.G., Yoklavich, M.M., Starr, R.M., O'Connell, V.M., Wakefield, W.W., Sullivan, D.E., McRea Jr, J.E., Cailliet, G.M., 1999.. A classification scheme for deep seafloor habitats. *Oceanologica Acta* 22(6), 663-678.
- Guinan, J., Brown, C., Dolan, M.F.J. and Grehan, A.J., 2009. Ecological niche modelling of the distribution of cold-water coral habitat using underwater remote sensing data. *Ecological Informatics* 4(2), 83-92.
- Harris, P.T., Whiteway, T., 2009. High sea marine protected areas: Benthic environmental conservation priorities from a GIS analysis of global ocean biophysical data. *Ocean and Coastal Management* 52, 22-38.
- Harris, P.T., Whiteway, T., 2011. Global distribution of large submarine canyons: Geomorphic differences between active and passive continental margins. *Mar. Geol.* 285, 69-86.
- Harrold, C., Light, K., Lisin, S., 1998. Organic enrichment of submarine-canyon and continental-shelf benthic communities by macroalgal drift imported from nearshore kelp forests. *Limnology and Oceanography* 43(4), 669-678.

747

748 Headrich, R.L., Rowe, G.T., Polloni, P.T., 1975. Zonation and faunal composition of
749 epibenthic populations on the continental slope of New England. *J. Mar. Res.* 33, 191-212.

750

751 Hecker, B., Logan, D.T., Gandarillas, F.E., Gibson, P.R., 1988. Canyon and slope processes
752 study, III. Biological processes, Lamont-Doherty Geological Observatory, Columbia
753 University, New York. III: 1-134.

754

755 Heezen, B.C., Ewing, M., Menzies, R.J., 1955. The influence of Submarine Turbidity
756 Currents on Abyssal Productivity. *Oikos* 6, 170-182.

757

758 Hickey, B. M., 1995. Coastal submarine canyons. In: Muller, P. and D. Henderson (eds),
759 Proceedings of the University of Hawaii 'Aha Huliko'a Workshop on Flow Topography
760 Interactions, Honolulu, Hawaii, SOEST Spec. Pub.: 265p.

761 Houston, K.A., Headrich, R.L., 1984. Abundance and biomass of macrobenthos in the
762 vicinity of Carson Submarine Canyon, northwest Atlantic Ocean. *Mar. Biol.* 82, 301-305.

763

764 Howell, K.L. (2010). A benthic classification system to aid in the implementation of marine
765 protected area networks in the deep/high seas of the NE Atlantic. *Biol. Conserv.* 143, 1041-
766 1056.

767

768 Howell, K.L., Davies, J.S., Hughes, D.J., Narayanaswamy, B.E., 2007. Strategic
769 Environmental Assessment/Special Area for Conservation Photographic Analysis Report.
770 Department of Trade and Industry, Strategic Environmental Assessment Report, UK: 163p.

771

- 772 Howell, K.L., Mowles, S.L., Foggo, A., 2010a. Mounting evidence: near-slope seamounts are
773 faunally indistinct from an adjacent bank. *Mar. Ecol.* 31(Suppl. 1), 1-11.
774
- 775 Howell, K.L., Davies, J.S., Narayanaswamy, B.E., 2010b. Identifying deep-sea megafaunal
776 epibenthic assemblages for use in habitat mapping and marine protected area network design.
777 *J. Mar. Biol. Assoc. UK* 90(1), 33-68.
778
- 779 Howell, K.L., Holt, R., Pulido Endrino, I., Stewart, H., 2011 When the species is also a
780 habitat: comparing the predictively modelled distributions of *Lophelia pertusa* and the reef
781 habitat it forms. *Biological Conservation*. 144, 2656–2665.
782
- 783 Huthnance, J.M., Coelho, H., Griffiths, C.R., Knight, P.J., Rees, A.P., Sinha, B.,
784 Vangriesheim, A., White, M., Chatwin, P.G., 2001. Physical structures, advection and mixing
785 in the region of the Goban Spur. *Deep-Sea Res. II, Top. Stud. Oceanogr.* 48, 2979-3021.
786 ICES (2007). Report of the Working Group on Deep-Water Ecology (WGDEC), 10-14
787 March 2008. *ICES J. Mar. Sci.* 1-122.
788
- 789 Inman, D.L., Nordstrom, C.E., Flick, R.E., 1976. Currents in submarine canyons: an air-sea
790 land interaction. *Annual Review of Fluid Mechanics*, 8, 275-310.
791
- 792 Jensen, A., Frederiksen, R., 1992. The fauna associated with the bank-forming deepwater
793 coral *Lophelia pertusa* (Scleractinaria) on the Faroe shelf. *Sarsia* 77, 53-69.
794
- 795 Jones, J. B., 1992. Environmental impact of trawling on the seabed: A review. *New*
796 *Zealand J. Mar. Freshwater Res.* 26 (1), 59-67.

797

798 Kelley, D.S., Delaney, J.R. and Yoerger, D.R., 2001. Geology and venting characteristics of
799 the Mothra hydrothermal field, Endeavour segment, Juan de Fuca Ridge Geology 29, 959-
800 962.

801

802 Kenchington, E.L., Cogswell, A.T., MacIsaac, K.G., Beazley, L., Law, B.A., Kenchington,
803 T.J., 2014, this issue. Limited depth zonation among bathyal epibenthic megafauna of the
804 Gully submarine canyon, northwest Atlantic. Deep-Sea Research II,

805

806 Kottke, B., Schwenk, T., Breitzke, M., Wiedicke, M., Kudrass, H.R., Spiess, V., 2003.
807 Acoustic facies and depositional processes in the upper submarine canyon Swatch of No
808 Ground (Bay of Bengal). Deep-Sea Res. II, Top. Stud. Oceanogr. 50, 979-1001.

809

810 Knudby, A., Kenchington, E., Murill, F.J., 2013. Modeling the distribution of Geodia
811 sponges and sponge Grounds in the Northwest Atlantic. PLoS ONE 8(12):e82306.
812 Doi:10.1371/journal.pone.0082306.

813

814 Lavaleye, M.S.S., Duineveld, G.C.A., Berghuis, E.M., Kok, A., Witbaard, R., 2002. A
815 comparison between the megafauna communities on the N.W. Iberian and Celtic continental
816 margins - effects of coastal upwelling? Progr. Oceanogr. 52, 459-476.

817

818 Long, D., 2006. BGS detailed explanation of seabed sediment modified Folk classification.
819 http://ec.europa.eu/maritimeaffairs/emodnet/documents/standards/mesh_geology.pdf).

820

- López-González, P.J., Williams, G.C., 2010. A new deep-sea pennatulacean (Anthozoa: Octocorallia: Chunellidae) from the Porcupine Abyssal Plain (NE Atlantic). Helgol Mar. Res. Pub. online 30 October 2010, 1–10.
- Maurer, D., Robertson, G., Gerlinger, T., 1994. Comparison of Community Structure of Soft-Bottom Macrobenthos of the Newport Submarine Canyon, California and the Adjoining Shelf. Internationale Revue der gesamten Hydrobiologie und Hydrographie 79(4), 591-603.
- Monaco, A., Biscaye, P.E., Soyer, J., Pocklington, R., Heussner, S., 1990. Particle fluxes and ecosystem responses on a continental margin: the 1985-1988 Mediterranean ECOMARGE experiment. Cont. Shelf Res. 10, 809-839.
- Mortensen, P.B., Hovland, M., Brattegard, T., Farestveit, R., 1995. Deep water bioherms of the scleractinian coral *Lophelia pertusa* (L.) at 64° N on the Norwegian shelf: structure and associated megafauna. Sarsia 80, 145-158.
- Mulder, T., Zaragosi, S., Garlan, T., Mavel, J., Cremer, M., Sottolichio, A., Sénéchal, N., Schmidt, S., 2012. Present deep-submarine canyons activity in the Bay of Biscay (NE Atlantic). Mar. Geol. 295-298, 113-127.
- Narayanaswamy, B.E., Howell, K.L., Hughes, D.J., Davies, J.S., Roberts, J.M., Black, K.D., 2006. Strategic Environmental Assessment Area 7 - Photographic Analysis. Department of Trade and Industry, Strategic Environmental Assessment Report, UK: 178p.
- Okey, T.O., 2003. Macrobenthic colonist guilds and renegades in Monterey Canyon (USA) drift algae: partitioning multidimensions. Ecological monographs 73(3), 415-440.

OSPAR (2010). Background Document for Coral Gardens. Biodiversity Series: 38p.

OSPAR (Agreement 2008-6). OSPAR Agreement 2008-6 that replaced an earlier version of the list contained in OSPAR Agreement 2004-6 (see also the Summary Record of the 2008 meeting of the OSPAR Commission contained in OSPAR 08/24/1, Para. 7.12).

Pfannkuche, O., Bannert, B., Beck, T., Beuck, L., Dullo, W.C., Flögel, S., Freiwald, A., Gass, S., Gektidis, M., Heger, A., Jamieson, A., Kavanagh, F., King, N., Kuhanec, B., Linke, P., Martin, B., Neulinger, S., Noe, S., Queisser, W., Rüggeberg, A., Ruseler, S., Schiemer, I., Schmidt, S., Schönfeld, J., Taviani, M., Türk, M., Vertino, A., Wigham, B., 2004. Geobiological Investigations on Azooxanthellate Cold-Water Coral Reefs on the Carbonate Mounds along the Celtic Continental Slope. L. A. t. M. Meteor Cruise No. 61, 2004, Lisbon – Cork.

Pilskaln, C.H., Churchill, J.H., Mayer, L.M., 1998. Resuspension of sediment by bottom trawling in the Gulf of Maine and potential geochemical consequences. *Conserv. Biol.* 12 (6), 1223-1229.

Ramirez-Llodra, E., Ballesteros, M., Company, J.B., Dantart, L., Sardá, F., 2008. Spatio-temporal variations in the diversity, biomass and abundance of bathyal invertebrates in the Catalan Sea (Western Mediterranean). *Mar. Biol.* 153, 297-309.

Rice, A.L., Tyler, P.A., Paterson, G.J.L., 1992. The pennatulid *Kophobelemnion stelliferum* (Cnidaria: Octocorallia) in the Porcupine Seabight (North-East Atlantic Ocean). *J. Mar. Biol. Assoc. UK* 72, 417-434.

- Roberts, J.M., Wheeler, A.J., Freiwald, A., 2006. Reefs of the deep: The biology and geology of cold-water coral ecosystems. *Science* 312(5773), 543-547.
- Roberts, J.M., Wheeler, A.L., Freiwald, A., Cairns, A., 2009. Cold-water corals. The biology and Geology of Deep-Sea Coral Habitats. New York.
- Ross., R, Howell, K.L., 2012. Use of predictive habitat modelling to assess the distribution and extent of the current protection of 'listed' deep-sea habitats. *Diversity and Distributions*. 19, 433-445. DOI: 10.1111/ddi.12010
- Rowden, A.A., Selacher, T.A., Williams, A., Clark, M.R., Stewart, R., Althaus, F., Bowden, D.A., Consalvey, M., Robinson, W., Dowdney, J., 2010. A test of the seamount oasis hypothesis: seamounts support higher epibenthic megafaunal biomass than adjacent slopes. *Mar. Ecol.* 31(Suppl. 1), 1-12.
- Rowe, G. T., 1971. Observations on bottom currents and epibenthic populations in Hatteras Submarine Canyon. *Deep Sea Res. Oceanogr. Abs.* 18(6), 569-576.
- Rowe, G.T., Polloni, P.T., Haedrich, R.L., 1982. The deep-sea macrobenthos on the continental-margin of the Northwest Atlantic Ocean. *Deep-Sea Res.– I, Oceanogr. Res. Pap.* 29(2), 257-278.
- Sager, W.W., Lee, C.S., MacDonald, I.R. and Schroeder, W.W., 1999. High-frequency near-bottom acoustic reflection signatures of hydrocarbon seeps on the Northern Gulf of Mexico continental slope. *Geo-Marine Letters* 18(4), 267-276.

Sardà, F., Cartes, J.E. and Company, J.B., 1994. Spatio-temporal variation in megabenthos abundance in three different habitats of the Catalan deep-sea (Western Mediterranean). *Mar. Biol.* 120, 211-219.

Schlacher, T.A., Schlacher-Hoenlinger, M.A., Williams, A., Althaus, F., Hooper, J.N.A., Kloser, R., 2007. Richness and distribution of sponge megabenthos in continental margin canyons off southeastern Australia. *Mar. Ecol. Prog. Ser.* 340, 73-88.

Schlacher, T.A., Williams, A., Franziska, A., Schlacher-Hoenlinger, M.A., 2010. High resolution seabed imagery as a tool for biodiversity conservation planning on continental margins. *Mar.Ecol.* 31 (1), 200-221.

Shank, T.M., 2010. Seamounts. Deep-ocean laboratories of faunal connectivity, evolution, and endemism. *Oceanography* 23, 108-122.

Shepard, F.P., 1951. Mass movement in submarine canyon heads. *Trans. American Geophys. Union* 32, 405-419.

Shepard, F.P., Marshall, N.F., 1973. Currents along floors of submarine canyons. *American Assoc. Pet. Geol. Bull.* 57(2), 244-264.

Smith, C.R., De Leo, F.C., Bernardino, A.F., Sweetman, A.K., Arbizu, P.M., 2008. Abyssal food limitation, ecosystem structure and climate change. *Trends in Ecol. Evolution* 23, 518-528.

Stevens, T., Connolly, R.M., 2004. Testing the utility of abiotic surrogates for marine habitat mapping at scales relevant to management. *Biol. Conserv.* 119, 351-362.

Stewart, H.A., Davies, J.S., Guinan, J, Howell, K.L., 2014. The Dangeard and Explorer Canyons, South-West Approaches, UK: geology, sedimentology and newly discovered cold-water coral mini-mounds. *Deep-Sea Res. II, Top. Stud. Oceanogr.*, this issue.

Stewart, H.A., Davies, J.S., 2007. SW Approaches MESH Survey, R/V Celtic Explorer Cruise CE0705, BGS Project 07/06, Operations Report. British Geological Survey Commercial Report CR/07/123.

Tews, J., Brose, U., Grimm, V., Tielbörger, K., Wichmann, M.C., Schwager, M., Jeltsch, F., 2004. Animal species diversity driven by habitat heterogeneity/diversity: the importance of keystone structures. *J. Biogeogr.* 31, 79-92.

Thrush, S.F., Hewitt, J.E., Funnell, G.A., Cummings, V.J., Ellis, J. Schultz, D., Talley, D., Norkko, A., 2001. Fishing disturbance and marine biodiversity: the role of habitat structure in simple soft-sediment systems. *Mar. Ecol. Prog. Ser.* 223, 277-286.

Thrush, S.F., Dayton, P.K., 2002. Disturbance to Marine Benthic Habitats by Trawling and Dredging: Implications for Marine Ecosystems. *Annual Rev. Ecol. Systematics* 33, 449-473.

Troffe, P.M., Levings, C.D., Piercey, G.E., Keong, V., 2006. Fishing gear effects and ecology of the sea whip (*Halopteris willemoesi* (Cnidaria: Octocorallia: Pennatulacea)) in British

Columbia, Canada: preliminary observations. *Aquatic Conservation: Marine and Freshwater Ecosystems* 15, 523-533.

UNGA (2006). Resolution 61/105 Sustainable fisheries, including through the 1995 Agreement for the Implementation of the Provisions of the United Nations Convention on the Law of the Sea of 10 December 1982 relating to the Conservation and Management of Straddling Fish Stocks and Highly Migratory Fish Stocks, and related instruments. UNGA A/RES/61/105.

Veale, L.O., Hill, A.S., Hawkins, S.J., Brand, A.R., 2000. Effects of long-term physical disturbance by commercial scallop fishing on sub-tidal epifaunal assemblages and habitats. *Mar. Biol.* 137, 325-337.

Vetter, E.W., 1994. Hotspots of benthic production. *Nature* 372, 47.

Vetter, E.W., Dayton, P.K., 1998. Macrofaunal communities within and adjacent to a detritus-rich submarine canyon system. *Deep-Sea Res. II, Top. Stud. Oceanogr.* 45, 25-54

Vetter, E.W., Dayton, P.K., 1999. Organic enrichment by macrophyte detritus, and abundance patterns of megafaunal populations in submarine canyons. *Mar. Ecol. Prog. Ser.* 186, 137-148.

Vetter, E.W., Smith, C.R., De Leo, F. C., 2010. Hawaiian hotspots: enhanced megafaunal abundance and diversity in submarine canyons on the oceanic islands of Hawaii. *Mar. Ecol.* 31, 183-199.

Wienberg, C., Beuck, L., Heidkamp, S., Hebbeln, D., Freiwald, A., Pfannkuche, O., Monteys, X., 2008. Franken Mound: facies and biocoenoses on a newly-discovered "carbonate mound" on the western Rockall Bank, NE Atlantic. *Facies* 54, 1-24.

Wilson, J.B., 1979. Patch development of the deep-water coral *Lophelia pertusa* (L.) on Rockall Bank. *J. Mar. Biol. Assoc. UK* 59, 165-177.

Wilson, W.H., 1991. Competition and predation in marine soft sediment communities. *Annual Review of Ecol. and Systematics* 21, 221–241.

Wilson, M.F J., O'Connell, B., Brown, C., Guinan, J.C. and Grehan, A.J., 2007. Multiscale terrain analysis of multibeam bathymetry data for habitat mapping on the Continental Slope. *Marine Geodesy* 30(1), 3-35.

Zibrowius, H., 1980. Les Scléractiniaires de la Méditerranée et de l'Atlantique nord-oriental. *Mémoires de l'Institut Océanographique, Monaco* 11, 1-284.

Figure legends

Figure 1: The study area on the Celtic Margin encompassing Dangaard and Explorer canyons and the eastern flank of a third canyon in Irish waters. Bathymetric contours are provided by GEBCO, the 200 m depth contour (dashed line) marks the approximate position of the continental shelf break. The UK median line corresponds to the UK continental shelf limit.

Figure 2: Plan (a) and 3D view (b) of multibeam bathymetry acquired over the survey area, meso-scale geomorphology (*sensu* Greene et al. 1999) is labelled. Fig. 2b is visualised in Fledermaus™ software, for scale of features see Fig. 2a.

Figure 3: Multibeam bathymetry data and video transects acquired over the SW Approaches survey area. Black dots represent video transects and are labelled with transect names.

Figure 4: Dendrogram of hierarchical cluster analysis of species data, clusters identified using the SIMPROF routine ($p < 0.01$). Dendrogram (a) shows those clusters identified as outliers at a 1% Bray Curtis similarity level and (b) remaining clusters for rejection/acceptance process. SIMPROF clusters have been collapsed for illustrative purposes.

Figure 5: Example images of biotopes showing fauna characteristic of each assemblage. Codes given to biotopes correspond to SIMPROF clusters in brackets: Bat.Hyd (r), Amp.Cer (al), Kop.Cer (y), Unk.Cer (ac), Lop.Cri (not defined from cluster analysis), Lop.Hal (aj), Lop.Mad (ah), Cer (x), Oph (am), Ser.Bra (ao), Mun.Lep (aq), Oph.Mun (ap). Lop.Cri was not identified from the cluster analysis, but described from the video.

Figure 6: nMDS ordination plot of pairwise ANOSIM test for depicting difference in environmental variables between biotopes. Cluster letters correspond to biotope codes: r (Bat.Hyd), al (Amp.Cer), y (Kop.Cer), ac (Unk.Cer), aj (Lop.Hal), ah (Lop.Mad), x (Cer), am (Oph), ao (Ser.Bra), aq (Mun.Lep), ap (Oph.Mun).

Figure A1: Mapped distribution of defined biotopes in the SW Approaches. Figures a-f represent the biotope mapped along the transects: (a) Amp.Cer, (b) Bat.Hyd, (c) Cer, (d) Kop.Cer, (e) Lop.Cri, (f) Lop.Hal.

Figure A2: Mapped distribution of defined biotopes in the SW Approaches. Figures a-f represent the biotope mapped along the transects: : (a) Lop.Mad, (b) Mun.Lep, (c) Oph, (d) Oph.Mun, (e) Ser.Bra, (f) Unk.Cer.

1024

1025 **Greyscale legends**

1026 Figure 3: Multibeam bathymetry data and video transects acquired over the SW Approaches
1027 survey area. White dots represent video transects and are labelled with transect names.

1028

1029 **Table legends**

1030 Table 1: Summary of mapped biotope data, abiotic data extracted from video metadata,
1031 geomorphology and substratum extracted from ArcGIS 9.3 layers.* refers to the biotope
1032 described from the video footage.

1033

1034 Table A1: Transects undertaken in the SW Approaches canyons: transect code, site (canyon),
1035 start and end of transect, length, number of statistical images analysed per transect, average
1036 depth and temperature (standard deviation) per transect, topographical feature sampled by
1037 transect and generalised seabed substrate within transects.

1038

1039 Table A2: Clusters identified from multivariate hierarchical analysis with associated
1040 environmental parameters, and SIMPER results identifying the taxa that characterise the
1041 clusters.

1042

1043

1044

1045

1046

1047

1048

1049

Appendix A1: SIMPER results for the SW Approaches

Full lists of species present in each assemblage described in Sect. 4. Characterising species, as identified by the SIMPER routine, are indicated in bold. ##### denotes where the number is infinitive or cannot be calculated, as in the case of Sim/SD, where the SD is zero and cannot be divided.

Group a

All the similarities are zero

Group b

Less than 2 samples in group

Group c

Average similarity: 42.26

Species	Av.Abund	Av.Sim	Sim/SD	Contrib%	Cum.%
Sabellidae sp. 1	0.46 42.26		#####	100.00	100.00

Group d

Less than 2 samples in group

Group e

Less than 2 samples in group

Group f

Average similarity: 100.00

Species	Av.Abund	Av.Sim	Sim/SD	Contrib%	Cum.%
Benthogone sp.	0.16 100.00		#####	100.00	100.00

Group g

Less than 2 samples in group

Group h

Less than 2 samples in group

Group i

Less than 2 samples in group

Group j

Less than 2 samples in group

1104								
1105	Group k							
1106	Average similarity: 25.93							
1107								
1108	Species	Av.Abund	Av.Sim	Sim/SD	Contrib%	Cum.%		
1109	<i>Protoptilum</i> sp.	0.22 16.67		0.58	64.27	64.27		
1110	<i>Pseudarchaster</i> sp.	0.17 9.27		0.58		35.73 100.00		
1111								
1112								
1113	Group l							
1114	Average similarity: 68.45							
1115								
1116	Species	Av.Abund	Av.Sim	Sim/SD	Contrib%	Cum.%		
1117	<i>Edwardsiidae</i> sp. 1	0.27 68.45		4.76		100.00 100.00		
1118								
1119								
1120	Group m							
1121	Average similarity: 44.15							
1122								
1123	Species	Av.Abund	Av.Sim	Sim/SD	Contrib%	Cum.%		
1124	<i>Halcampoididae</i> sp. 3	0.32 24.63		0.58	55.78	55.78		
1125	Unknown sp. 13	0.22 19.53		0.58	44.22	100.00		
1126								
1127								
1128	Group n							
1129	Average similarity: 49.42							
1130								
1131	Species	Av.Abund	Av.Sim	Sim/SD	Contrib%	Cum.%		
1132	Unknown sp. 15	0.19 49.42		#####	100.00 100.00			
1133								
1134								
1135	Group o							
1136	Average similarity: 50.48							
1137								
1138	Species	Av.Abund	Av.Sim	Sim/SD	Contrib%	Cum.%		
1139	<i>Sagartiidae</i> sp. 3	0.29 48.48		1.78	96.05	96.05		
1140	<i>Kophobelemnnon stelliferum</i>	0.06 1.70		0.22		3.38		
1141	99.42							
1142	<i>Calveriosoma fenestratum</i>	0.02 0.29		0.09	0.58	100.00		
1143								
1144								
1145	Group p							
1146	Average similarity: 18.04							
1147								
1148	Species	Av.Abund	Av.Sim	Sim/SD	Contrib%	Cum.%		
1149	<i>Actiniaria</i> sp. 14	0.05 10.48		0.39	58.07	58.07		
1150	<i>Cerianthidae</i> sp. 3	0.10 6.14		0.44		34.04		
1151	92.11							
1152	Crinoidea sp. 1	0.07 1.42		0.26	7.89	100.00		
1153								
1154								
1155	Group q							
1156	Average similarity: 10.73							
1157								
1158	Species	Av.Abund	Av.Sim	Sim/SD	Contrib%	Cum.%		
1159	<i>Caryophyllia</i> sp. 2	0.11 4.00		0.32		37.27		
1160	37.27							
1161	<i>Porifera encrusting</i> sp. 1	0.09 3.60		0.31	33.50	70.77		
1162	<i>Hydrozoa</i> (flat branched)	0.15 2.15		0.24		19.99		
1163	90.75							
1164	<i>Bathynectes</i> sp.	0.04 0.40		0.13	3.71	94.47		
1165	<i>Bolocera tuediae</i>	0.05 0.30		0.13	2.77	97.23		

1166	Cerithioidea sp.	0.05	0.30	0.13	2.77	100.00
1167						
1168						
1169	Group r					
1170	Average similarity: 25.07					
1171						
1172	Species	Av.Abund	Av.Sim	Sim/SD	Contrib%	Cum.%
1173	cf. <i>Bathylasma</i> sp.	0.42	16.33	0.58	65.13	65.13
1174	Hydrozoa (bushy)	0.14	8.74	0.57	34.87	100.00
1175						
1176						
1177	Group s					
1178	Average similarity: 14.78					
1179						
1180	Species	Av.Abund	Av.Sim	Sim/SD	Contrib%	Cum.%
1181	Terebellidae sp. 1	0.26	14.94	0.79	60.27	
1182	60.27					
1183	Actiniaria sp. 17	0.15	8.96	0.39	36.16	96.43
1184	Serpulidae sp. 1	0.04	0.47	0.17	1.91	98.34
1185	<i>Bonellia viridis</i>	0.06	0.41	0.17	1.66	100.00
1186						
1187	Group t					
1188	Average similarity: 38.99					
1189						
1190	Species	Av.Abund	Av.Sim	Sim/SD	Contrib%	Cum.%
1191	Amphipoda sp. 1	0.25	38.99	#####	100.00	100.00
1192						
1193						
1194	Group u					
1195	Average similarity: 20.08					
1196						
1197	Species	Av.Abund	Av.Sim	Sim/SD	Contrib%	Cum.%
1198	<i>Colus</i> sp. 2	0.35	20.08	1.28	100.00	100.00
1199						
1200						
1201	Group v					
1202	Average similarity: 49.37					
1203						
1204	Species	Av.Abund	Av.Sim	Sim/SD	Contrib%	Cum.%
1205	<i>Pachycerianthus multiplicatus</i>	0.38	42.08	3.23	85.22	85.22
1206	Cerianthidae sp. 1	0.11	7.30	0.58	14.78	100.00
1207						
1208						
1209	Group w					
1210	Less than 2 samples in group					
1211						
1212						
1213	Group x					
1214	Average similarity: 54.39					
1215						
1216	Species	Av.Abund	Av.Sim	Sim/SD	Contrib%	Cum.%
1217	Cerianthidae sp. 1	0.31	54.10	2.63	99.47	
1218	99.47					
1219	Sagartiidae sp. 3	0.01	0.06	0.05	0.10	99.57
1220	<i>Echinus</i> spp.	0.01	0.05	0.05	0.10	
1221	99.67					
1222	<i>Munida sarsi</i>	0.01	0.05	0.05	0.09	
1223	99.76					
1224	Cerianthidae sp. 3	0.01	0.03	0.03	0.05	
1225	99.81					
1226	Unknown sp. 26	0.01	0.02	0.03	0.04	99.85

1227	<i>Ophiothrix fragilis</i>		0.01	0.02		0.03		0.04
1228	99.89							
1229	<i>Pseudarchaster</i> sp.		0.01	0.02		0.03		0.03
1230	99.92							
1231	<i>Caryophyllia</i> sp. 2		0.01	0.02		0.03		0.03
1232	99.95							
1233	<i>Kophobelemnion stelliferum</i>		0.00	0.02		0.03		0.03
1234	99.98							
1235	Halcampoididae sp. 1	0.01	0.01		0.03		0.02	100.00
1236								
1237								
1238	Group y							
1239	Average similarity: 49.80							
1240								
1241	Species	Av.Abund		Av.Sim	Sim/SD	Contrib%		Cum.%
1242	<i>Kophobelemnion stelliferum</i>	0.34	42.07		2.54		84.46	84.46
1243	Cerianthidae sp. 1		0.14	7.03		0.55		14.12
1244	98.59							
1245	Ophiuroidea sp.1	0.04	0.41		0.11		0.82	99.41
1246	Halcampoididae sp.3		0.02	0.13		0.06		0.27
1247	99.68							
1248	<i>Pentametrocrinus atlanticus</i>	0.02	0.06		0.06		0.13	99.81
1249	Crinoidea sp. 2	0.03	0.04		0.04		0.08	99.89
1250	<i>Ophiactis balli</i>	0.01	0.04		0.04		0.07	99.96
1251	<i>Acanella</i> sp.		0.01	0.02		0.04		0.04
1252	100.00							
1253								
1254								
1255	Group z							
1256	Average similarity: 41.11							
1257								
1258	Species	Av.Abund		Av.Sim	Sim/SD	Contrib%		Cum.%
1259	<i>Ophiactis balli</i>	0.71	38.06		2.29		92.59	92.59
1260	Cerianthidae sp. 1		0.07	1.82		0.27		4.42
1261	97.01							
1262	<i>Munida sarsi</i>		0.08	0.60		0.20		1.46
1263	98.47							
1264	Serpulidae sp. 1	0.05	0.20		0.11		0.49	98.96
1265	<i>Actinauge richardi</i>		0.02	0.13		0.06		0.32
1266	99.28							
1267	Halcampoididae sp. 1	0.05	0.12		0.11		0.29	99.57
1268	Zoanthidea sp. 1	0.03	0.06		0.06		0.16	99.72
1269	Unknown sp. 26	0.02	0.06		0.06		0.15	99.87
1270	<i>Echinus</i> spp.		0.01	0.03		0.06		0.07
1271	99.94							
1272	Hydrozoa (bushy)		0.02	0.02		0.06		0.06
1273	100.00							
1274								
1275								
1276	Group aa							
1277	Less than 2 samples in group							
1278								
1279								
1280	Group ab							
1281	Average similarity: 31.57							
1282								
1283	Species	Av.Abund		Av.Sim	Sim/SD	Contrib%		Cum.%
1284	Sabellidae sp. 2	0.76	31.57		9.59		100.00	100.00
1285								
1286								
1287	Group ac							
1288	Average similarity: 47.47							

1289									
1290	Species	Av.Abund		Av.Sim	Sim/SD	Contrib%		Cum.%	
1291	Unknown sp. 26	1.03 36.24			2.41	76.36		76.36	
1292	Cerianthidae sp. 1	0.26 6.79			0.93		14.31		
1293	90.67								
1294	<i>Ophiactis balli</i>	0.32 2.32			0.36	4.88		95.55	
1295	<i>Lophelia pertusa</i> (dead structure)	0.10 0.74			0.20	1.57		97.11	
1296	Halcampoididae sp. 1	0.09 0.58			0.23	1.21		98.33	
1297	Amphiuridae sp. 1		0.05 0.26			0.14	0.55		
1298	98.88								
1299	Ophiuroidea sp. 1	0.06 0.19			0.13	0.41		99.29	
1300	<i>Munida sarsi</i>		0.03 0.12			0.11	0.26		
1301	99.55								
1302	<i>Lophelia pertusa</i>	0.04 0.07			0.09	0.15		99.69	
1303	Terebellidae sp. 1	0.01 0.04			0.08	0.09		99.78	
1304	<i>Psolus squamatus</i>	0.02 0.03			0.08	0.06		99.84	
1305	<i>Echinus</i> spp.		0.02 0.02			0.07	0.05		
1306	99.89								
1307	Sagartiidae sp. 3	0.01 0.01			0.03	0.02		99.91	
1308	Brachiopoda sp. 1		0.01 0.01			0.03	0.02		
1309	99.92								
1310	<i>Bathynectes</i> sp.	0.01 0.01			0.03	0.01		99.94	
1311	Ascidacea sp. 2	0.01 0.01			0.03	0.01		99.95	
1312	<i>Bolocera tuediae</i>	0.01 0.01			0.03	0.01		99.96	
1313	Crinoidea sp. 1	0.00 0.01			0.03	0.01		99.97	
1314	Galatheididae sp. 1	0.01 0.01			0.03	0.01		99.98	
1315	<i>Pandalus borealis</i>		0.02 0.00			0.03	0.01		
1316	99.99								
1317	Actiniaria sp. 9	0.01 0.00			0.03	0.01		100.00	
1318									
1319									
1320	Group ad								
1321	Average similarity: 59.02								
1322									
1323	Species	Av.Abund		Av.Sim	Sim/SD	Contrib%		Cum.%	
1324	<i>Lophelia pertusa</i> (dead structure)	0.41 58.36			3.76	98.89	98.89		
1325	<i>Munida sarsi</i>		0.04 0.66			0.26	1.11	100.00	
1326									
1327									
1328	Group ae								
1329	Less than 2 samples in group								
1330									
1331									
1332									
1333	Group af								
1334	Less than 2 samples in group								
1335									
1336									
1337	Group ag								
1338	Less than 2 samples in group								
1339									
1340									
1341	Group ah								
1342	Average similarity: 66.25								
1343									
1344	Species	Av.Abund		Av.Sim	Sim/SD	Contrib%		Cum.%	
1345	<i>Lophelia pertusa</i> (dead structure)	0.78 21.36			5.39	32.24	32.24		
1346	<i>Lophelia pertusa</i>	0.55 14.17			3.49	21.39	53.63		
1347	<i>Madrepora oculata</i>		0.46 11.60			2.99	17.51		
1348	71.14								
1349	Unknown sp. 26	0.46 5.49			0.70	8.28			
1350	79.42								

1351	Actiniaria sp. 13	0.28	4.06	0.99	6.12	
1352	85.54					
1353	<i>Pandalus borealis</i>	0.15	2.79	1.20	4.21	
1354	89.75					
1355	Cerianthidae sp. 1	0.14	2.51	1.09	3.79	
1356	93.54					
1357	Halcampoididae sp. 1	0.18	2.16	0.70	3.25	
1358	96.79					
1359	<i>Cidaris cidaris</i>	0.10	1.47	0.68	2.22	
1360	99.01					
1361	<i>Bathynectes</i> sp.	0.04	0.32	0.33	0.49	
1362	99.50					
1363	Hydrozoa (bushy)	0.05	0.21	0.21	0.31	
1364	99.81					
1365	<i>Koehlermetra porrecta</i>	0.04	0.04	0.07	0.06	
1366	99.87					
1367	Hydrozoa (flat branched)	0.02	0.03	0.08	0.05	
1368	99.92					
1369	<i>Porania pulvillus</i>	0.01	0.03	0.08	0.04	
1370	99.96					
1371	Gastropoda sp. 1	0.01	0.01	0.05	0.01	
1372	99.97					
1373	<i>Munida sarsi</i>	0.01	0.01	0.05	0.01	
1374	99.98					
1375	<i>Brisingella coronata</i> /	0.03	0.01	0.05	0.01	
1376	99.99					
1377	<i>Brisinga endecacnemos</i>					
1378	<i>Henricia sanguinolenta</i>	0.01	0.01	0.05	0.01	
1379	100.00					
1380						
1381						
1382	Group ai					
1383	Average similarity: 61.28					
1384						
1385	Species	Av.Abund	Av.Sim	Sim/SD	Contrib%	Cum.%
1386	Unknown sp. 26	2.70 34.64		3.31	56.53	56.53
1387	<i>Lophelia pertusa</i> (dead structure)	0.81 11.70		8.18	19.09	75.61
1388	<i>Madrepora oculata</i>	0.45 6.09		3.62		9.93
1389	85.55					
1390	<i>Lophelia pertusa</i>	0.42 2.49		0.58	4.07	89.61
1391	Actiniaria sp. 13	0.46 2.40		0.58	3.92	93.53
1392	Edwardsiidae sp. 1	0.28 1.98		0.58		3.23
1393	96.77					
1394	Halcampoididae sp. 1	0.32 1.98		0.58	3.23	100.00
1395						
1396						
1397	Group aj					
1398	Average similarity: 54.00					
1399						
1400	Species	Av.Abund	Av.Sim	Sim/SD	Contrib%	Cum.%
1401	<i>Lophelia pertusa</i> (dead structure)	0.55 30.59		3.35	56.65	56.65
1402	Halcampoididae sp. 1	0.20 8.73		1.27	16.16	72.80
1403	<i>Lophelia pertusa</i>	0.21 6.92		0.76	12.82	85.63
1404	Cerianthidae sp. 1	0.24 6.16		0.77	11.40	97.03
1405	<i>Madrepora oculata</i>	0.17 1.60		0.26	2.97	
1406	100.00					
1407						
1408						
1409	Group ak					
1410	Average similarity: 66.33					
1411						
1412	Species	Av.Abund	Av.Sim	Sim/SD	Contrib%	Cum.%

1413	Halcampoididae sp. 5	0.30	66.33	4.38	100.00	100.00
1414						
1415	Group al					
1416	Average similarity: 53.22					
1417						
1418	Species	Av.Abund	Av.Sim	Sim/SD	Contrib%	Cum.%
1419	Amphiuridae sp. 1	0.62	40.91		2.56	57.53
1420	57.53					
1421	Cerianthidae sp. 1	0.46	20.85		1.18	41.59
1422	99.12					
1423	<i>Munida sarsi</i>	0.05	0.34		0.13	0.64
1424	99.76					
1425	Ophiuroidea sp. 5	0.02	0.05	0.05	0.08	99.84
1426	Terebellidae sp. 1	0.01	0.03	0.03	0.05	99.89
1427	<i>Kophobelemnnon stelliferum</i>	0.01	0.02	0.02	0.03	99.93
1428	Brachiopoda sp. 1	0.01	0.01		0.02	0.02
1429	99.95					
1430	<i>Pachycerianthus multiplicatus</i>	0.01	0.01	0.02	0.02	99.97
1431	Edwardsiidae sp. 1	0.01	0.01		0.03	0.02
1432	99.99					
1433	<i>Caryophyllia</i> sp. 3	0.01	0.00		0.02	0.01
1434	100.00					
1435						
1436						
1437	Group am					
1438	Average similarity: 47.39					
1439						
1440	Species	Av.Abund	Av.Sim	Sim/SD	Contrib%	Cum.%
1441	Ophiuroidea sp. 1	1.12	46.57		2.13	98.27
1442	98.27					
1443	Amphiuridae sp. 1	0.07	0.48		0.14	1.01
1444	99.28					
1445	<i>Munida sarsi</i>	0.02	0.07		0.05	0.15
1446	99.43					
1447	<i>Ophiactis balli</i>	0.03	0.07	0.06	0.14	99.56
1448	Cerianthidae sp. 1	0.02	0.05		0.06	0.10
1449	99.67					
1450	<i>Caryophyllia</i> sp. 1	0.03	0.03		0.04	0.07
1451	99.74					
1452	Serpulidae sp. 1	0.02	0.03	0.04	0.06	99.80
1453	Porifera encrusting sp. 1	0.01	0.01	0.02	0.02	99.82
1454	Ophiuroidea sp. 5	0.02	0.01	0.03	0.02	99.84
1455	<i>Kophobelemnnon stelliferum</i>	0.01	0.01		0.03	0.02
1456	99.86					
1457	<i>Actinauge richardi</i>	0.01	0.01		0.02	0.02
1458	99.88					
1459	<i>Caryophyllia smithii</i>	0.01	0.01		0.02	0.02
1460	99.90					
1461	<i>Leptometra celtica</i>	0.01	0.01		0.02	0.01
1462	99.91					
1463	Crinoidea sp. 5	0.01	0.01	0.02	0.01	99.92
1464	Polychaeta sp. 7	0.00	0.00	0.01	0.01	99.93
1465	Actiniaria sp. 17	0.01	0.00	0.01	0.01	99.94
1466	Terebellidae sp. 1	0.01	0.00	0.01	0.01	99.94
1467	Majidae sp. 1	0.00	0.00		0.01	0.01
1468	99.95					
1469	<i>Ophiothrix fragilis</i>	0.01	0.00		0.01	0.01
1470	99.96					
1471	Sagartiidae sp. 3	0.00	0.00	0.01	0.01	99.96
1472	Cerianthidae sp. 3	0.01	0.00		0.01	0.01
1473	99.97					

1474	<i>Ophiactis abyssicola</i>		0.01	0.00		0.01		0.01
1475	99.97							
1476	Polychaeta sp. 5	0.01	0.00		0.01		0.00	99.98
1477	<i>Astropecten irregularis</i>	0.00	0.00		0.01		0.00	99.98
1478	<i>Virgularia mirabilis</i>		0.00	0.00		0.01		0.00
1479	99.98							
1480	Paguridae spp.	0.00	0.00		0.01		0.00	99.99
1481	Unknown sp. 15	0.00	0.00		0.01		0.00	99.99
1482	Brachiopoda sp. 1		0.00	0.00		0.01		0.00
1483	99.99							
1484	<i>Caryophyllia</i> sp. 2		0.00	0.00		0.01		0.00
1485	99.99							
1486	<i>Pandalus borealis</i>		0.00	0.00		0.01		0.00
1487	99.99							
1488	Polychaeta sp. 1	0.01	0.00		0.01		0.00	99.99
1489	<i>Pentametrocrinus atlanticus</i>	0.00	0.00		0.01		0.00	99.99
1490	Unknown sp. 13	0.00	0.00		0.01		0.00	99.99
1491	<i>Tubularia</i> sp. 2	0.00	0.00		0.01		0.00	100.00
1492								
1493								
1494	Group an							
1495	Average similarity: 49.67							
1496								
1497	Species	Av.Abund		Av.Sim	Sim/SD	Contrib%		Cum.%
1498	Crinoidea sp. 5	0.39	45.53		2.16	91.66		91.66
1499	<i>Stichopathes</i> cf. <i>gravieri</i>	0.12	4.14		0.44	8.34		100.00
1500								
1501	Group ao							
1502	Average similarity: 27.51							
1503								
1504	Species	Av.Abund		Av.Sim	Sim/SD	Contrib%		Cum.%
1505	Serpulidae sp. 1	0.39	20.63		0.99	74.99		74.99
1506	Brachiopoda sp. 1		0.11	4.52		0.26		16.42
1507	91.40							
1508	<i>Munida sarsi</i>		0.12	1.57		0.23		5.69
1509	97.10							
1510	<i>Caryophyllia smithii</i>		0.05	0.63		0.14		2.28
1511	99.38							
1512	Ophiuroidea sp. 1	0.03	0.10		0.06	0.36		99.73
1513	<i>Actinauge richardi</i>		0.01	0.07		0.06		0.27
1514	100.00							
1515								
1516								
1517	Group ap							
1518	Average similarity: 41.38							
1519								
1520	Species	Av.Abund		Av.Sim	Sim/SD	Contrib%		Cum.%
1521	Ophiuroidea sp. 5		1.13	37.87		2.36		61.53
1522	61.53							
1523	<i>Munida sarsi</i>		0.75	12.22		1.33		35.36
1524	96.89							
1525	<i>Leptometra celtica</i>		0.06	0.35		0.17		0.86
1526	97.74							
1527	Amphiuridae sp. 1		0.07	0.25		0.12		0.62
1528	98.36							
1529	Hydrozoa (bushy)		0.05	0.25		0.12		0.61
1530	98.97							
1531	Serpulidae sp. 2	0.05	0.14		0.07	0.34		99.31
1532	Paguridae spp.	0.04	0.10		0.13	0.25		99.55
1533	Brachiopoda sp. 1		0.03	0.07		0.07		0.16
1534	99.72							

1535	<i>Echinus</i> spp.	0.03	0.06	0.07	0.15	
1536	99.87					
1537	Cerianthidae sp. 1	0.03	0.05	0.07	0.13	
1538	100.00					
1539						
1540						
1541	Group aq					
1542	Average similarity: 33.11					
1543						
1544	Species	Av.Abund	Av.Sim	Sim/SD	Contrib%	Cum.%
1545	<i>Munida sarsi</i>	0.31	28.05		1.21	84.74
1546	<i>Leptometra celtica</i>	0.16	3.18		0.30	9.60
1547	94.33					
1548	Crinoidea sp. 5	0.07	0.89	0.18	2.70	97.03
1549	Cerianthidae sp. 1	0.03	0.42	0.14	1.25	
1550	98.29					
1551	<i>Caryophyllia smithii</i>	0.03	0.20	0.08	0.61	
1552	98.90					
1553	Ophiuroidea sp. 1	0.03	0.13	0.08	0.38	99.28
1554	<i>Ophiactis balli</i>	0.03	0.11	0.07	0.33	99.61
1555	<i>Caryophyllia</i> sp. 2	0.02	0.05	0.04	0.15	
1556	99.76					
1557	<i>Echinus</i> spp.	0.01	0.03	0.03	0.08	
1558	99.84					
1559	Porifera encrusting sp. 31	0.01	0.02	0.03	0.07	99.90
1560	Porifera encrusting sp. 3	0.01	0.01	0.03	0.04	99.94
1561	<i>Actinauge richardi</i>	0.01	0.01	0.03	0.03	
1562	99.98					
1563	Ophiuroidea sp. 5	0.01	0.01	0.03	0.02	100.00

Appendix A2: Biotope descriptions for non-listed habitats defined from the SW

Approaches

cf. *Bathylasma* sp. and hydroid assemblage on bedrock

The biotope Bat.Hyd, identified as cluster r, was characterised by barnacles (cf. *Bathylasma* sp.) and Hydrozoa (bushy) associated with steep bedrock outcrop towards the base of Explorer canyon at a depth of 902-912 m and a temperature of 8.99-9°C. Bat.Hyd assemblage was only observed for a short period during a single camera-transect.

Bathylasma is a widespread bathyal species in the NE Atlantic (Gage 1986). A number of assemblages have been described from the region; Pfannkuche et al. (2004) describe a *Bathylasma* cf. *hirsutum* assemblage associated with drop stones between 636-650 m water depth on a prominent escarpment feature in the Belgica mound province; however, Gage

(1986) describes a *Bathylasma hirsutum* assemblage with the brachiopod *Dallina septigera* and *Macandrevia cranium* from rocky, high current areas on the Wyville-Thomson Ridge and the summit of the Anton Dohrn Seamount in a water depth band ranging from 200-700 m. He also noted the remains of plates of *Bathylasma hirsutum* covering the substratum of the floor of a gorge between the Wyville-Thomson Ridge and Ymir ridge and suggested this species may cover the walls of this gorge. Howell et al. (2010b) describe an assemblage characterised by large barnacles (noted as possibly *Bathylasma hirsutum*) and brachiopods (noted as possibly *Dallina septigera*) on the summit of the Anton Dohrn Seamount at approx. 600 m water depth.

Amphiuridae ophiuroids and cerianthid anemones on bioturbated mud/sand

The biotope Amp.Cer, identified as cluster al, was characterised by occasional cerianthid anemones and amphiuridae sp.1 ophiuroids on bioturbated mud and sand and was observed throughout the canyons over a wide depth range of 184-943 m and temperature of 9.59-11.69°C associated with the canyon head, flanks and was also observed on from one transect on the continental shelf. Note, this assemblage has not been previously described from the deep sea.

Annelids, hydroids and cerianthids on bedrock ledges

The biotope Unk.Cer, identified as cluster ac, was characterised by cerianthid anemones, annelid worms and hydroid species associated with bedrock ledges. Ophiuroid species and the squat lobster *Munida sarsi* were also commonly observed. The biotope was observed from both Dangaard and Explorer canyons from the canyon head and incised channels (canyon floor) associated with bedrock ledges over a depth range of 238-1070 m and a

temperature of 8.36-11.51°C. This kind of biotope has not been previously described in the deep sea.

Cerianthids on sediment draped bedrock

The biotope Cer, identified as cluster x, was characterised by cerianthid anemones associated with areas of bedrock covered with a sand veneer – thus preventing the attachment of fauna and acting as a soft sediment habitat. The assemblage was observed on wide range of geomorphological features including canyon head, flank, amphitheatre rims and incised channels. It was observed from the three canyons over a water depth and temperature range of 360-1064 m and 8.98-11.3°C, respectively. This assemblage has not been previously described from the deep sea. This assemblage has a similar distribution to the ‘Cerianthid anemones on bioturbated mud/sand’ biotope.

Burrowing (*Amphiura* sp.) and surface dwelling ophiuroids on mud/sand

The biotope Oph, identified as cluster am, was characterised by surface dwelling ophiuroids associated with soft sediment (mud-sand). Burrowing ophiuroids (*Amphiura* sp.) were also identified as being characteristic of this biotope from video observations. The assemblage was found on the flanks, incised channels and amphitheatre rims; and occurred in the three canyons at water depths of 184-1094 m and temperatures of 7.67-11.69°C. This assemblage has not been previously described from the deep sea.

Serpulids and brachiopods on mixed substratum

The biotope Ser.Bra, identified as cluster ao, was associated with cobble and pebble substratum with serpulid polychaetes (*Serpulidae* sp. 1) and brachiopods (*Brachiopoda* sp. 1) attached to the hard substratum and squat lobsters (*Munida sarsi*) associated with the

surrounding soft sediment. The assemblage was observed only on the smooth flank of Dangaard canyon between 691-764 m and over a temperature range of 10.1-10.5°C.

The Ser.Bra assemblage is similar to that described by Howell et al. (2010b) as ‘brachiopods on mixed substrate’ which was widely observed between 266-803 m water depth on a number of features in UK waters. Narayanaswamy et al. (2006) also reported a similar assemblage from Anton Dohrn Seamount, where abundant brachiopods were associated with coarse sediment on the seamount summit.

Munida sarsi and *Leptometra celtica* on mixed substratum

The biotope Mun.Lep, identified as cluster aq, was associated with mixed and biogenic gravel (shell hash) substratum on the canyon head and interflaves features from all three canyons. It occurred over a wide depth and temperature range (183-792 m; 9.79-11.79°C) and was characterised by the crinoid *Leptometra celtica*, the squat lobster *Munida sarsi*. This assemblage occurred on the interflaves between the mini-mounds features and was also associated with tributary gullies.

Leptometra celtica were more abundant at the canyon heads and on the edge of the flanks, which suggests they are positioning themselves within optimal conditions for feeding. The occurrence of *Leptometra celtica* has been reported by a number of authors; Lavaleye et al. (2002) reported abundant crinoids at 190 m from the NW Iberian Margin and 200 m from the Goban Spur, and Flach et al. (1998) found the crinoid to be the dominant fauna at a station at 208 m water depth from the continental Shelf (Goban Spur).

Assemblage code	Cluster	Assemblage name	Depth (m)	Temperature (°C)	Topographical Feature	Substratum	Canyon
Bat.Hyd	r	cf. <i>Bathylasma</i> sp. and hydroid	902-912 m	8.99-9°C	Incised channel	Bedrock	Explorer
Kop.Cer	y	<i>Kophobelemnion stelliferum</i> and	463-1059	8.87-10.85°C	Flank and incised channel	Mud and muddy sand	Explorer, Irish
Cer	x	Cerianthids on sediment draped	360-1064	8.98-11.3°C	Canyon head, amphitheatre rims,	Bedrock with sand veneer	Explorer, Irish
Unk.Cer	ac	Annelids, hydroids and cerianthids on	238-1070	8.36-11.51°C	Canyon head and incised	Bedrock and bedrock with	Explorer and
Lop.Mad	ah	<i>Lophelia pertusa</i> reef	795-940 m	9.41-9.92°C	Flute feature	Coral framework	Explorer
Lop.Hal	aj	Predominantly dead low-lying	697-927 m	8.97-9.77°C	Flank and flute feature (end of	Coral rubble, bedrock and	Explorer and
Amp.Cer	al	Amphiuridae ophiuroids and	184-943 m	9.59-11.69°C	Flank, canyon head and	Mud and sand	Explorer, Irish
Oph	am	Burrowing (<i>Amphiura</i> sp.) and	184-1094	7.67-11.69°C	Flank, tributary gullies,	Mud and sand	Explorer, Irish
Ser.Bra	ao	Serpulids and brachiopods on	691-764 m	10.1-10.5°C	Flank	Mixed	Dangaard
Oph.Mun	ap	Ophiuroids and <i>Munida sarsi</i>	303-1017	7.98-11.5°C	Incised channels and	Biogenic gravel (coral	Explorer and
Mun.Lep	aq	<i>Munida sarsi</i> and <i>Leptometra celtica</i>	183-792 m	9.79-11.79°C	Interfluves and canyon head	Mixed, biogenic	Explorer, Irish
Lop.Cri	*	<i>L. pertusa</i> and crinoids on	253-1022	7.93-11.42°C	Incised channels, tributary gullies,	Bedrock	Explorer, Irish

Table 1: Summary of mapped biotope data, abiotic data extracted from video metadata, geomorphology and substratum extracted from ArcGIS 9.3 layers.* refers to the biotope described from the video footage.

Transect	Canyon	Start position	End position	Transect length (m)	# of images analysed	Average depth (m) (SD)	Average temperature (°C) (SD)	Topographical unit	Generalised seabed substrate
C_1_1	Irish	48.496292 - 9.872838	48.48386 - 9.872061	1382.6	62	847.3 (77.4)	9.47 (0.23)	Flank	Mud-rich sediments
C_1_2	Irish	48.560068 - 9.834236	48.562948 - 9.830817	407.6	13	294.2 (4.9)	11.36 (0.002)	Flank	Sandy gravel
C_1_3	Irish	48.569407 - 9.841935	48.569343 - 9.846046	300	12	379.5 (26.35)	11.29 (0.008)	Flank	Sand-rich sediments with bedrock cropping out where

									slope angle greatest (amphitheatre rim)
C_1_4	Irish	48.5606 29 - 9.85788 1	48.5628 05 - 9.86140 9	341.7	19	520.2 (103.6)	10.54 (0.07)	Flank	Bedrock cropping out where slope steepest. Gravel-rich sediments immediately down slope of the outcrop becoming mud dominated as water depths increase
C_2_1	Dangeard	48.4200 6 - 9.57345	48.4161 2 - 9.57048	488.8	25	298.3 (26.4)	11.46 (0.12)	Flank	Sand dominated sediments with increasing proportion of gravel in vicinity of slump headwall
C_2_2	Dangeard	48.4036 6 - 9.54235	48.4019 2 - 9.54559	308.1	18	652.7 (0.69)	10.34 (0.008)	Canyon head	Sand
C_2_3	Dangeard	48.3925 7 - 9.57007	48.3893 6 -9.5751	513.6	31	776.1 (13.3)	9.85 (0.09)	Canyon head	Sand-rich sediments and gravelly sand. Bedrock cropping out where slope steepest
C_2_4	Dangeard	48.3835 8 - 9.67091	48.3809 7 - 9.66819	344.5	16	402.1 (3.63)	11.02 (0.03)	Flank	Gravelly sand
C_2_5	Dangeard	48.3724 5 - 9.68566	48.3680 2 - 9.68436	498.7	20	591.06 (26.42)	10.46 (0.2)	Flank	Mud-rich sediments
C_2_6	Dangeard	48.3587 98 - 9.72382 5	48.3545 8 - 9.72160 1	476.6	7	803.1 (21.12)	9.28 (1.89)	Flank	Mud-rich sediments
C_2_7	Explorer	48.3782 - 9.77602	48.3794 5 -	468.7	17	756.3 (22.2)	9.79 (0.09)	Flank	Sand

			9.78212						
C_2_8	Explorer	48.44012 - 9.68199	48.4446 - 9.68242	496.7	18	917.6 (6.6)	9.26 (0.13)	Flank	Sandy gravel becoming sand dominated as water depths increase. Bedrock cropping out where slope steepest
C_2_9	Explorer	48.471652 - 9.621832	48.473303 - 9.625192	288.4	9	644.8 (129.29)	10.36 (0.43)	Flank	Sand
C_2_10	Explorer	48.482903 - 9.574269	48.483972 - 9.578344	317.5	14	463.1 (149.8)	10.78 (0.45)	Flank	Sand
C_2_11	Explorer	48.498926 - 9.613132	48.494222 - 9.608227	633.5	27	895.8 (4.6)	9.05 (0.03)	Flank	Bedrock cropping out where slope steepest. Sand and gravelly sand observed on the gully bottom
C_2_12	Explorer	48.513442 - 9.504434	48.515962 - 9.499202	472.5	24	274.7 (21.2)	11.26 (0.02)	Canyon head	Sand-rich sediments with increasing proportion of gravel upslope of gully wall where bedrock observed cropping out where slope steepest
C_2_13	Explorer	48.522986 - 9.590885	48.519347 - 9.591295	405.3	19	463.4 (76.5)	10.95 (0.18)	Canyon head	Sand-rich sediments. Bedrock cropping out where slope steepest
C_2_14	Explorer	48.477797 -	48.472027 -	669.1	39	839.2 (27.69)	9.83 (0.04)	Flank	Majority of substrate obscured

		9.65665 8	9.65408						by encrusting fauna. Bedrock occasional y observed
C_2_1 5	Explore r	48.4680 16 - 9.73698 1	48.4672 49 - 9.73012 9	512.9	15	533.7 (46.5)	10.61 (0.16)	Flank	Sand-rich sediments becoming mud-rich as water depths increase
C_2_1 6	Explore r	48.4249 31 - 9.87074 1	48.4222 78 - 9.87560 7	462.6	14	827.9 (42.57)	9.4 (0.21)	Flank	Mud-rich sediments
C_2_1 7	Explore r	48.4523 97 - 9.80016	48.4490 09 - 9.80020 5	373.1	15	463.2 (203.4)	10.81 (0.82)	Flank	Mud-rich sediments
C_2_1 8	Explore r	48.4641 9 - 9.71451 5	48.4600 47 - 9.71696 3	491.7	18	751.3 (32.02)	9.75 (0.002)	Flank	Mud-rich sediments. Bedrock cropping out in slump headwall
C_2_1 9	Explore r	48.4961 12 - 9.64301 7	48.4914 18 - 9.64367 1	519.6	16	684.2 (16.7)	10.16 (0.01)	Flank	Mud-rich sediments
C_2_2 0	Explore r	48.4633 47 - 9.64709 7	48.4667 64 - 9.65027 2	439.2	12	884.8 (38.4)	9.56 (0.04)	Canyon floor	Bedrock cropping out at sea bed.
C_2_2 1	Dangea rd	48.4254 03 - 9.60910 3	48.4237 73 - 9.61222 7	286.6	14	257.9 (54.3)	11.52 (0.14)	Interfluve	Sand
C_2_2 2	Dangea rd	48.3976 24 - 9.64959 4	48.3952 15 - 9.64783 6	297.9	16	334.8 (4.27)	11.17 (0.03)	Interfluve	Gravelly sand
C_2_2 3	Dangea rd	48.3466 38 - 9.77915 6	48.3417 37 - 9.78388 2	599.7	26	769.8 (16.9)	9.71 (0.07)	Flank	Mud-rich sediments. Bedrock cropping out and increasing gravel content where slope steepest

C_2_2_4	Dangeard	48.376516 - 9.639434	48.373696 - 9.640584	321.9	11	746.2 (33.1)	9.61 (0.13)	Flank	Mud-rich sediments
C_2_2_5	Dangeard	48.377362 - 9.60101	48.374678 - 9.598054	353.7	10	750.4 (16.9)	9.8 (0.03)	Flank	Mud-rich sediments
C_2_2_6	Dangeard	48.438655 - 9.483831	48.434266 - 9.485687	471.4	24	318.8 (8.1)	11.5 (0.07)	Canyon head	Sand-rich sediments with bedrock cropping out where slope angle greatest in slump headwall
C_2_2_7	Explorer	48.575601 - 9.483186	48.573985 - 9.486711	313.1	15	187.8 (3.1)	11.68 (0.001)	Continental shelf	Sand
C_2_2_8	Explorer	48.554515 - 9.537413	48.552849 - 9.530352	551.1	27	260.7 (17.9)	11.4 (0.002)	Canyon head	Gravelly sand. Bedrock cropping out where slope steepest in gully wall
C_3_1	Dangeard	48.308271 - 9.552125	48.310245 - 9.554577	492.9	20	208.9 (2.3)	11.48 (1.91)	Interfluvial	Gravelly sand with smaller sections of sandy gravel
C_3_2_b	Dangeard	48.307334 - 9.604809	48.311251 - 9.601967	485.2	27	306.7 (0.75)	11.49 (0.001)	Interfluvial	Gravelly sand. Sandy gravel over mini-mounds
C_3_3	Dangeard	48.401267 - 9.455041	48.397704 - 9.45152	474.3	18	240.2 (26.1)	11.5 (0.02)	Canyon head	Gravelly sand with bedrock cropping out where slope angle greatest in slump headwall
C_3_4	Dangeard	48.360553 - 9.480041	48.361247 - 9.483062	236.9	9	240 (1.14)	11.37 (0.01)	Canyon head	Sandy gravel and gravelly sand with bedrock cropping out where

									slope angle greatest (amphitheatre rim)
C_3_5	Dangeard	48.3620 28 - 9.49748 1	48.3597 28 - 9.49850 5	251.4	13	389.8 (13.9)	11.01 (0.29)	Canyon head	Sand with bedrock cropping out where slope angle greatest.
C_3_6	Dangeard	48.3612 2 - 9.55597 1	48.3620 2 - 9.55843 4	178.7	23	976.3 (10.6)	7.917 (0.14)	Canyon floor	Bedrock cropping out at sea bed with veneer of sand- and gravel-rich sediments in places
C_3_7	Dangeard	48.2918 5 - 9.64142	48.2968 5 - 9.64276	564.1	27	352.8 (0.82)	11.19 (0.02)	Interfluvial	Gravelly sand. Sandy gravel over mini- mounds
C_3_8	Dangeard	48.3317 3 - 9.63122	48.3352 3 -9.6355	498.4	23	536.1 (55.6)	10.91 (0.12)	Flank	Sand
C_3_9	Dangeard	48.3119 3 - 9.70631	48.3163 - 9.70437	506.1	17	722.6 (38.1)	10.16 (0.11)	Flank	Sand
C_3_10	Dangeard	48.3014 7 - 9.73274	48.3058 3 - 9.73504	511.7	26	799.4 (52.4)	10.11 (0.15)	Flank	Sand
C_3_11	Dangeard	48.2805 4 - 9.74772	48.2812 - 9.75445	501.6	25	724 (22.12)	10.27 (0.17)	Flank	Gravelly sand
C_3_12	Dangeard	48.3471 94 - 9.53403 2	48.3520 59 - 9.53400 6	537	23	774.5 (57.9)	9.71 (0.13)	Canyon head	Bedrock cropping out at sea bed with veneer of mud-rich sediments in places


Table A1: Transects undertaken in the SW Approaches canyons: transect code, site (canyon), start and end of transect, length, number of statistical images analysed per transect, average depth and temperature (standard deviation) per transect, topographical feature sampled by transect and generalised seabed substrate within transects.



Cluster	No. images	Useful mapping unit	SIMPER similarity	Temp range (°C)	Average Temp (SD)	Depth range	Average Depth (SD)	Characterising species
a	2	N	0	9.6-11.3	10.487 (1.21)	316-840	578 (370.5)	
b	1	N		11.54	11.546	256	256	
c	2	N	42.26	10.4-11.7	11.118 (0.90)	210-695	452.5 (342.9)	Sabellidae sp. 1
d	1	N		9.252	9.252	850	850	
e	1	N		11.49	11.496	309	309	
f	3	N	100	9.0-10.4	9.951 (0.80)	508-866	694.3 (151.3)	<i>Benthogone</i> sp.
g	1	N		11.49	11.497	311	311	
h	1	N		11.54	11.542	256	256	
i	1	N		10.00	10.007	788	788	
j	1	N		9.91	9.91	762	762	
k	3	N	25.93	9.7-10.4	9.970 (0.38)	602-755	695.6 (82.1)	<i>Protophilum</i> sp., <i>Pseudarchaster</i>
l	4	N	68.45	11.22	11.388 (0.23)	212-401	342.5 (88.6)	Edwardsiidae sp. 1
m	3	N	44.15	8.8-9.1	9.02 (0.12)	885-1006	947.3 (60.5)	Halcampoididae sp. 3, Unknown
n	2	N	49.42	11.549	11.549	321-323	322 (1.4)	Unknown sp. 15


o	16	N	50.48	9.3-10.2	9.732 (0.24)	714-928	800.6 (51.7)	Sagartiidae sp. 3
p	6	N	18.04	8.9-11.1	10.499 (0.83)	331-1059	596.5 (249.7)	Actiniaria sp. 14, Cerianthidae sp. 3
q	11	N	10.73	7.7-11.6	10.465 (1.24)	185-1009	543 (305.4)	<i>Caryophyllia</i> sp. 2, Porifera encrusting sp. 1, <i>Hydazoon</i> (flat knobby)
r	7	Y	25.07	8.9-11.6	10.062 (1.32)	190-909	625.7 (341.4)	cf. <i>Bathylasma</i> sp., Hydrozoa (bushy)
s	9	N	14.78	10.3-11.4	11.11 (0.47)	238-800	407 (222.8)	Terebellidae sp. 1, Actiniaria sp. 17
t	2	N	38.99	9.2-9.7	9.745 (0.03)	729-782	755.5 (37.4)	Amphipoda sp. 1
u	4	N	20.08	8.1-10.1	9.379 (0.94)	741-1015	852.5 (122.6)	<i>Colus</i> sp. 2
v	3	N	49.37	10.5-11.3	11.026 (0.43)	378-601	452.6 (128.4)	<i>Pachycerianthus multiplicatus</i> , Cerianthidae
w	1	N		11.174	11.174	333	333	sp. 1
x	49	Y	54.39	9.0-11.5	9.922 (0.59)	308-954	738.6 (164.7)	Cerianthidae sp. 1
y	39	Y	49.80	9.1-10.3	9.544 (0.29)	609-953	836.7 (89.3)	<i>Kophobelemnnon stelliferum</i> , Cerianthidae sp. 1
z	23	N	41.11	8.0-11.5	10.31 (1.20)	295-1054	615.6 (288.2)	<i>Ophiactis balli</i>
aa	1	N		9.599	9.599	938	938	
ab	3	N	31.57	8.0-9.8	9.207 (1.01)	781-1012	869 (124.9)	Sabellidae sp. 2
ac	46	Y	47.47	7.7-10.7	9.294 (0.82)	316-1048	829.7 (166.7)	Unknown sp. 26, Cerianthidae sp. 1


ad	6	N	59.02	9.0-11.7	9.517 (1.06)	184- 942	778.8 (294.2)	<i>Lophelia pertusa</i> (dead structure)
ae	1	N		9.763	9.763	699	699	
af	1	N		9.878	9.878	798	798	
ag	1	N		9.011	9.011	874	874	
ah	30	Y	66.25	9.5-9.9	9.780 (0.09)	797- 938	860.9 (43.7)	<i>Lophelia pertusa</i> (dead structure), <i>Lophelia pertusa</i> , <i>Madrepora oculata</i> , Unknown
ai	3	N	61.28	9.5-9.7	9.646 (0.08)	914- 936	922.3 (11.9)	Unknown sp. 26, <i>Lophelia pertusa</i> (dead structure),
aj	7	Y	54.00	9.0-9.8	9.377 (0.39)	816- 942	894.6 (55.6)	<i>Madrepora oculata</i> , <i>Lophelia pertusa</i> (dead structure), Halcampoididae sp.
ak	3	N	66.33	9.7-11.3	10.523 (1.09)	417- 782	640.3 (195.7)	<i>Lophelia pertusa</i> , Cerianthidae, Halcampoididae sp. 5
al	71	Y	53.22	7.6-11.5	10.163 (0.98)	254- 1008	654.3 (218.9)	Amphiuridae sp. 1, Cerianthidae sp. 1
am	276	Y	47.39	8.9-11.8	10.803 (0.64)	205- 1021	477.3 (195.37)	Ophiuroidea sp. 1
an	6	N	49.67	10.5- 11.4	10.988 (0.38)	257- 600	433.1 (159.6)	Crinoidea sp. 5, <i>Stichopathes</i> cf. <i>gravieri</i>
ao	24	Y	27.51	9.4-11.8	10.943 (0.68)	189- 803	464.1 (214.9)	Serpulidae sp. 1, Brachiopoda sp. 1, <i>Munida sarsi</i>
ap	20	Y	41.38	9.6-11.6	10.926 (0.73)	252- 791	423.9 (212.9)	Ophiuroidea sp. 5, <i>Munida sarsi</i>
aq	51	Y	31.11	9.0-11.7	11.303 (0.40)	192- 825	326.4 (124.0)	<i>Munida sarsi</i> , <i>Leptometra celtica</i>


Table A2: Clusters identified from multivariate hierarchical analysis with associated environmental parameters, and SIMPER results identifying the taxa that characterise the clusters.















2D Stress: 0.08

ah



r

