

# Model metadata summary report for the Colchester LithoFrame 10-50 model

Geology and Landscapes Programme Open Report OR/13/001



### BRITISH GEOLOGICAL SURVEY

GEOLOGY & REGIONAL GEOPHYSICS PROGRAMME OPEN REPORT OR/13/001

# Model metadata summary report for the Colchester LithoFrame 10-50 model.

S. Mathers

Keyworth, Nottingham British Geological Survey 2013

The National Grid and other Ordnance Survey data © Crown Copyright and database rights 2012 Ordnance Survey Licence No: 100021290

#### Keywords

3D Modelling, metadata

#### Front cover & frontispiece

*Cover picture* the complete calculated block model of the Colchester area viewed from the southwest. *Frontispiece* the model from the southeast.without artificial deposits.

#### Bibliographical reference

MATHERS, S.J.2013. Model metadata summary report for the Ipswich-Sudbury LithoFrame 10-50 model. *British Geological Survey Open Report*, OR/13/001 12pp.

Copyright in materials derived from the British Geological Survey's work is owned by the Natural Environment Research Council (NERC) and/or the authority that commissioned the work. You may not copy or adapt this publication without first obtaining permission. Contact the BGS Intellectual Property Rights Section, British Geological Survey, Keyworth, e-mail ipr@bgs.ac.uk. You may quote extracts of a reasonable length without prior permission, provided a full acknowledgement is given of the source of the extract.

© NERC 2013. All rights reserved

### **BRITISH GEOLOGICAL SURVEY**

The full range of our publications is available from BGS shops at Nottingham, Edinburgh, London and Cardiff (Welsh publications only) see contact details below or shop online at www.geologyshop.com

The London Information Office also maintains a reference collection of BGS publications, including maps, for consultation.

We publish an annual catalogue of our maps and other publications; this catalogue is available online or from any of the BGS shops.

The British Geological Survey carries out the geological survey of Great Britain and Northern Ireland (the latter as an agency service for the government of Northern Ireland), and of the surrounding continental shelf, as well as basic research projects. It also undertakes programmes of technical aid in geology in developing countries.

The British Geological Survey is a component body of the Natural Environment Research Council.

### British Geological Survey offices

#### **BGS Central Enquiries Desk**

Tel	0115 936 3143
email	enquiries@bgs.ac.uk

**Kingsley Dunham Centre, Keyworth, Nottingham NG12 5GG** Tel 0115 936 3241 Fax 0115 936 3488

Fax 0115 936 3276

email sales@bgs.ac.uk

Murchison House, West Mains Road, Edinburgh EH9 3LA

Tel 0131 667 1000 Fax 0131 668 2683 email scotsales@bgs.ac.uk

Natural History Museum, Cromwell Road, London SW7 5BD

 Tel
 020 7589 4090
 Fax
 020 7584 8270

 Tel
 020 7942 5344/45
 email
 bgslondon@bgs.ac.uk

Columbus House, Greenmeadow Springs, Tongwynlais, Cardiff CF15 7NE Tal. 020 2052 1062

Гel	029 2052 1962	Fax 029 2052 1963

# Maclean Building, Crowmarsh Gifford, Wallingford OX10 8BB

Tel 01491 838800 Fax 01491 692345

### Geological Survey of Northern Ireland, Colby House, Stranmillis Court, Belfast BT9 5BF

1ei 028 9038 8402 I ax 028 9038 840	el	028 9038 8462	Fax	028 9038 846
-------------------------------------	----	---------------	-----	--------------

www.bgs.ac.uk/gsni/

#### Parent Body

Natural Environment Research Council, Polaris House, North Star Avenue, Swindon SN2 1EU

Tel 01793 411500 Fax 01793 411501 www.nerc.ac.uk

Website www.bgs.ac.uk Shop online at www.geologyshop.com

## Contents

Co	ntents	2
Sui	nmary	3
1	Modelled volume, purpose and scale	3
2	Modelled surfaces/volumes	4
3	Modelled faults	5
4	Model datasets	5
5	Dataset integration	7
6	Model development log	7
7	Model workflow	8
8	Model assumptions, geological rules used etc	8
9	Model limitations	9
10	Model images10	0
11	Model uncertainty	2
Glo	ssary12	2
Ref	erences1	3

# Summary

This report is the published product of a regional GSI3D model of the Colchester area of southern East Anglia and north Essex. The model was assembled between 2006-08 concurrently with surveying of the parts of the area within the Colchester and Brightlingsea 1:50 000 mapsheets (BGS, 2010) by S.J. Mathers, the TL92 sheet was appended in 2011.

# 1 Modelled volume, purpose and scale



Figure 1. Extent of the Colchester model in mauve, OS topography © Crown Copyright.

The model covers the 1: 25 000 sheets TL 92 and TM 01,11,21,02,12,22, (Figure 1) covering approximately 500 km<sup>2</sup>. It was built in 2006-08 for the six TM 25K sheets as a systematic model build using combinations of the individual tiles (01-11-21, 02 and 12-22) and then amalgamated in 2008. In 2011 TL92 was added to produce a single unified model. The model was constructed using 1: 10 000 digital linework with all the TM sheets produced as part of an integrated mapping-modelling resurvey. The TL 92 portion of the model will enabled revision of the approved 1: 10 000 superficial and bedrock linework, the simple linework means that the 50K maps of these areas included in the combined Colchester, and Brightlingsea mapsheet closely resembles the 10K dataset.

The model was funded as a science budget activity.

The model extends down to the base of the Cretaceous succession in the area at about -300m OD, the Gault Formation is the lowest modelled unit. The model contains 3 artificial 17 superficial, 5 bedrock units, and a single lens unit within the Anglian glacigenic sequence at Marks Tey.

The final project file is saved as Colchester\_V21.gsipr.

No faults are modelled within the area although one single small fault has been drawn into sections as a steep ramp in the base of the affected bedrock units on the TL92 mapsheet.

The model has been calculated using GSI3D 2012 utilising the superficial (sections and envelopes) calculation option. Figure 2 below shows the arrangement of cross-sections drawn to construct this model.



### Figure 2. Cross sections in the Colchester model.

The complete model takes about 10 minutes to calculate on a Dell Precision M6700

# 2 Modelled surfaces/volumes

Geological units in the model are as follows

code	name
mgr	Made ground
wgr	Worked ground
wmgr	Worked and made ground
stobco	Shoreface and beach upper layer (along coast)
alv	Main Freshwater Alluvium
itdu	Intertidal Deposits undifferentiated
stob	Shoreface and beach deposits lower layer
isc	Interglacial silt and clay
head1	Head (silt reworked loess?)
brk	Brickearth (local overbank deposits on terraces)
rtdu	River Terrace Deposits undifferentiated
lde1	Lacustrine deposits (Clacton and Marks Tey)
gstc	Glacial Silt and Clay supramain till
gsg	Glacial Sand and Gravel supramain till

loft gstcb1 gsgb1 gcd igd kes BEDROCK	Lowestoft Till (main sheet) Glacial Silt and Clay below till 1 Glacial Sand and Gravel below till 1 Glacial Channel deposits Interglacial deposits (Little Oakley) Kesgrave Sand and Gravel
rcg	Red Crag Formation
tham	Thames Group
llte	Lower London Tertiaries
ck	Chalk Group undiff
glt	Gault Formation
LENS	
gsg1t	Glacial Sand and Gravel in main till sheet (top)
gsg1	Glacial Sand and Gravel in main till sheet (base)

The sequence shares a common \*.gvs with the adjacent Ipswich-Sudbury area model.

Multiple layers of storm beach deposits were needed to cope with interdigitated sequences and Holocene evolution. The Anglian glacigenic sediments shown above in blue comprise the Lowestoft Formation and within them a single lens is identified (gsg1). Legend colours in many cases loosely correspond to the 50K published map sheets. The difficulty in separating the thin Thanet Sand Formation from the Reading Formation (Lambeth Group) has resulted in the use of the older Lower London Tertiaries term being used for this package of strata. The Harwich and London Clay Formations cannot be mapped out separately and are lumped together as the Thames Group.

The model abuts the Ipswich-Sudbury model to the north (Mathers, 2012), shared docker sections ensure a common interpretation along the join at northing 30.

# 3 Modelled faults

No faults are modelled within the area although one single small normal fault has been drawn into sections as a steep ramp in the base of the affected bedrock units on the TL92 mapsheet.

# 4 Model datasets

### GVS & GLEG files

These files were assembled in a combination of Notepad, Wordpad and Excel and iterated as the model expanded and new units were encountered. The current files are Regional\_V9.gvs and Regional\_v10.gleg. The model colour schema is shown below in Figure 3.



Figure 3. The colour schema used.

### **Boreholes**

Borehole records examined as indicated below. Many start heights were revised in SOBI and some grid references corrected by the data management team. Closely clustered sets of boreholes were not all coded but the deepest and most representative were included; any significant local variation in sequence was also recorded by coding. Entries were all made directly into the corporate Boge data entry application with sima as the interpreter in all cases.

TL92 borehole coding TL92NW 1-43 TL92NE 1-388, TL92SW 1-125 and TL92SE 1-388 were examined by S J Mathers in 2011.

TM01 borehole coding TM01NW1-7, TM01NE1-96, TM01SW 1-6 and TM01SE 1-2 were examined by S J Mathers in 2011.

TM11 borehole coding TM11NW1-8, TM11NE1-7, TM11SW 1-5, and TM11SE1-9 were examined by S J Mathers in 2011.

TM21 borehole coding TM21NW1 was examined by S J Mathers in 2011.

TM02 borehole coding TM02NW1-219, TM02NE1-139, TM02SW1-99 and TM02SE1-118 were examined by S J Mathers in 2011.

TM12 borehole coding TM12NW1-49, TM12NE1-30, TM12SW1-48 and TM12SE1-21 were examined by S J Mathers in 2011.

TM22borehole coding TM22NW1-7, TM22SW1-30 and TM22SE1-2 were examined by S J Mathers in 2011.

### Rasters

Geo referenced raster backdrops taken from Figure 14 of the Braintree memoir (Ellison & Lake 1986) have been used to align and construct portions of sections Marks Tey\_Section A,B and C.

### **Geological Linework**

Modern 1: 10 000 scale digital linework was used throughout the TM sheets. Sheets TM 01, 11, 21, 02, 12, and 22 were surveyed by S.J. Mathers in 2005-8 together with revisions of parts of TL92 NE and SE as far west as easting 98. Farther west the 1:10 560 survey data for the Braintree 1: 50 000 mapsheet (BGS 1982, Ellison & Lake, 1986) were utilised for the TL92 tile and considerably revised by the modelling. This has produced new linework for the whole of TL92 that should now be used to update the master 10K linework for both the Quaternary and bedrock themes.

### DTM

Each 25K model tile was produced individually using a CEH or Nextmap DTM mainly with a 50m grid size. The merged model was initially capped by a CEH 100m dtm, this was then replaced by a Nextmap 100m and then the final models is capped but not snapped to a trimmed BGS Bald Earth dtm (with a 50m resolution) produced by combining elements of the OS Panorama dataset and the Nextmap 5M dataset. The data was extracted using the wizard provided within the software. The current file name Bald Earth\_2 [TIN].

## 5 Dataset integration

The individual 25K model tiles were aggregated into two sets one for the TL sheet and one for the TM sheets as described above, these were subsequently merged into the complete model in 2011 which covers approximately 500 sq km of land area. Consistency between tiles was achieved by the use of docker sections iterated between the two tiles and then envelope merging using the polygon editing tools. The final workspace was assembled to include all sections, standard, helper and dockers, merged envelopes for all units, and a regional Nextmap 50m dtm supercede by the BGS Bald Earth model.

## 6 Model development log

The amalgamated model from 2011 supercedes the individual 25K sheet tiles The individual tiles in their pre amalgamation form with supporting files are stored as archived versions. The subsequent editing of the amalgamated model has not been transmitted back to the earlier individual model tiles. All entries by S J Mathers.

### 10 May 2011

Completed sections and imported a 50m grid to calculate the whole model **Southern\_merge\_v12.gsipr** 

### 19 August 2011

Added the TL 92 tile, prepared a new dtm for enhanced area (untrimmed) and a cut back one for calculation (whole), added artificial deposits wgr, mgr, wmgr from new Colchester\_Brightlingsea sheet, bedrock units are now faulted, saved as v15. Calculates well

### 8 January 2013

Added a BGS bald earth model at 50m resolution trimmed back to model area

Tidied the gsg1 lense at Marks Tey and produced envelope for the top this now works

Model is not totally snapped

Project saved as Southern merge\_16 with faults and baselines and Southern merge\_17 with the single fault stripped out and baselines converted to polygons

Calculation takes about 40 minutes

Mgr, wgr and wmgr have envelopes and are shown in sections

Several screen grabs made for this report all at VE x10

Added 2 llte helpers due to an upward bulge at Marks Tey

### 30-31 January 2013

Further revisions (extra sections, new dtm extent, unit name changes) following checking by Mark Barron saved as Colchester\_v21.

### 25 April 2013

Further minor revisions following snapping by Ian Cooke saved as Colchester\_v28

## 7 Model workflow

A standard GSI3D workflow (Kessler & Mathers, 2004; Kessler et al. 2009) for superficial geological models was followed, in the parts of the model surveyed as an integrated exercise (all TM) borehole coding was followed by the field survey, the 1: 10 000 geological lines were captured digitally by CartoGIS from field slips and/or fair drawn linework on plastic. The resulting 1:10 000 scale provisional shape files were then provided for use in modelling. Edits to the shape files were made iteratively as a result of the modelling, in particular subcrops at rockhead were revised with the benefit of a 3D visualisation of the geology.. It has been assembled and edited using the current versions of the GSI3D software since 2006 i.e. 2.0 - 2.6 - 2011 - 2012 (bedrock).

## 8 Model assumptions, geological rules used etc

Buried river terrace deposits were assumed to extend beneath and match of the distribution of overlying alluvium and intertidal deposits in river channels except in estuaries were more confined buried channels are found. The deposits die out upstream and are not found in the upper reaches of small streams and brooks.

Only the silt rich ?reworked Aeolian head deposits have been modelled the other head deposits shown on the mapsheeet are thin and have not been included in the modelling exercise. Worked, Made and Worked and made ground have been constructed in sections and mapped polygons have been inserted into the geological units however helper sections have not been constructed to allow 3D volumes of these areas within the model stack. calculation.

Helper sections usually only correlate the units they are intended to inform

The Chalk group is treated as a single unit

The Lower London Tertiaries comprise mainly Reading Formation overlying deposits of the Thanet Sand Formation as noted from surface outcrops and some boreholes.

# 9 Model limitations

- Head in most forms excluded from the calculation of the model
- Artificial deposits not modelled sufficiently to produce 3D volumes but inserted in some sections and polygons have been included in geological units.
- Chalk Group, Lower London Tertiaries and Thames Group are not subdivided in the model at this stage.
- No attempt has been made to classify levels of the River Terrace Deposits or the Kesgrave Sand and Gravel into distinct terrace levels.
- The model is suitable for use at resolutions higher than 100K but not beyond 5K or for any detailed site specific investigations.
- The model extends to the base of the Cretaceous across the whole area, in this area the Cretaceous rests with marked angular unconformity on Palaeozoic basement rocks.
- The model is not snapped between envelopes and sections

# 10Model images



**Figure 4. Bedrock geology for colours see Figure 3**. (note maroon Red Crag blue Thames Group, red Lower London Tertiaries, green Chalk royal blue Gault Clay at base.). VE x10.



**Figure 5. Pre Anglian glaciation units for colours see Figure 3.** (note Kesgrave Sand and Gravel dark pink). VE x10



**Figure 6. Including Anglian glaciation and Marks Tey and Clacton Hoxnian interglacial deposits for colours see Figure 3.** (note Lowestoft Till sky blue, Glacial sand and gravel layers all pale pink and Glacial Silt and Clay layers blue, lacustrine deposits at Marks Tey brown) VE x10.



Figure 7. The Complete model from the southwest for colours see Figure 3. VE x10



Figure 8. The model without artificial deposits from the southeast for colours see Figure 3. VE x10.

# 11 Model uncertainty.

To-date no attempt has been made to quantify the uncertainty in this model

# Glossary

DTM	Digital Terrain Model – Model of surface of the solid Earth (generally the boundary between geosphere and atmosphere or hydrosphere). This is traditionally derived from OS contours and spot heights and should therefore exclude all buildings, trees, hedges, crops, animals etc. Sometimes also referred to as 'bald earth' models
Envelope	Defined here as the extent of a geological deposit in plan view (2D): forming a distribution map of the particular unit, a presence – absence map.
GSI3D	Geological Surveying and Investigation in 3D

# References

British Geological Survey holds most of the references listed below, and copies may be obtained via the library service subject to copyright legislation (contact libuser@bgs.ac.uk for details). The library catalogue is available at: <u>http://geolib.bgs.ac.uk</u>.

BGS 1982 Braintree 1: 50 000 Mapsheet 223 Solid & Drift Geology

BGS 2010 Colchester & Brightlingsea 1:50 000 Mapsheet 224/242 Bedrock and Superficial

ELLISON, R.A. AND LAKE R.D. 1986. Geology of the country around Braintree Memoir of the British Geological Survey, Sheet 223

KESSLER, H. AND MATHERS S.J. 2004. The 3-D geological map - finally capturing the Geologists' Vision. *Geoscientist* 14/10 4-6.

KESSLER, H., MATHERS, S.J. AND H.-G. SOBISCH. 2009. The capture and dissemination of integrated 3D geospatial knowledge at the British Geological Survey using GSI3D software and methodology. *Computers & Geosciences* 35, 1311-1321.

MATHERS, S.J. 2012 Model metadata summary report for the Ipswich-Sudbury LithoFrame 10-50 model. *British Geololgical Survey Open Report OR/12/080*.