



**British
Geological Survey**

NATURAL ENVIRONMENT RESEARCH COUNCIL

Bromsgrove Aquifer Groundwater Modelling Study: Results from Task 1.1 3D Visualisation and Geological Framework of the Bromsgrove Aquifer

Geology, Geotechnics and Palaeontology Programme

Commissioned Report CR/09/023

BRITISH GEOLOGICAL SURVEY

GEOLOGY, GEOTECHNICS AND PALAEOLOGY PROGRAMME
COMMISSIONED REPORT CR/09/023

The National Grid and other Ordnance Survey data are used with the permission of the Controller of Her Majesty's Stationery Office.
Licence No: 100017897/ 2009.

Keywords

Bromsgrove, Sherwood Sandstone.

National Grid Reference

SW corner 380000,280000
Centre point 395000,270000
NE corner 405000,280000

Map

Sheet 167, 168, 182, 183, 1:99
000 scale, Map name

Front cover

Cover picture details, delete if no cover picture.

Bibliographical reference

NEWELL, A J, SMITH, N J. 2009. Bromsgrove Aquifer Groundwater Modelling Study: Results from Task 1.1 3D Visualisation and Geological Framework of the Bromsgrove Aquifer. *British Geological Survey Commissioned Report*, CR/09/023. 18pp.

Copyright in materials derived from the British Geological Survey's work is owned by the Natural Environment Research Council (NERC) and/or the authority that commissioned the work. You may not copy or adapt this publication without first obtaining permission. Contact the BGS Intellectual Property Rights Section, British Geological Survey, Keyworth, e-mail ipr@bgs.ac.uk. You may quote extracts of a reasonable length without prior permission, provided a full acknowledgement is given of the source of the extract.

Maps and diagrams in this book use topography based on Ordnance Survey mapping.

Bromsgrove Aquifer Groundwater Modelling Study: Results from Task 1.1 3D Visualisation and Geological Framework of the Bromsgrove Aquifer

A J Newell, N J Smith

Contributor/editor

E. Hough, K Ambrose

BRITISH GEOLOGICAL SURVEY

The full range of our publications is available from BGS shops at Nottingham, Edinburgh, London and Cardiff (Welsh publications only) see contact details below or shop online at www.geologyshop.com

The London Information Office also maintains a reference collection of BGS publications, including maps, for consultation.

We publish an annual catalogue of our maps and other publications; this catalogue is available online or from any of the BGS shops.

The British Geological Survey carries out the geological survey of Great Britain and Northern Ireland (the latter as an agency service for the government of Northern Ireland), and of the surrounding continental shelf, as well as basic research projects. It also undertakes programmes of technical aid in geology in developing countries.

The British Geological Survey is a component body of the Natural Environment Research Council.

British Geological Survey offices

BGS Central Enquiries Desk

Tel 0115 936 3143 Fax 0115 936 3276
email enquiries@bgs.ac.uk

Kingsley Dunham Centre, Keyworth, Nottingham NG12 5GG

Tel 0115 936 3241 Fax 0115 936 3488
email sales@bgs.ac.uk

Murchison House, West Mains Road, Edinburgh EH9 3LA

Tel 0131 667 1000 Fax 0131 668 2683
email scotsales@bgs.ac.uk

Natural History Museum, Cromwell Road, London SW7 5BD

Tel 020 7589 4090 Fax 020 7584 8270
Tel 020 7942 5344/45 email bgs_london@bgs.ac.uk

Columbus House, Greenmeadow Springs, Tongwynlais, Cardiff CF15 7NE

Tel 029 2052 1962 Fax 029 2052 1963

Forde House, Park Five Business Centre, Harrier Way, Sowton EX2 7HU

Tel 01392 445271 Fax 01392 445371

Maclean Building, Crowmarsh Gifford, Wallingford OX10 8BB

Tel 01491 838800 Fax 01491 692345

Geological Survey of Northern Ireland, Colby House, Stranmillis Court, Belfast BT9 5BF

Tel 028 9038 8462 Fax 028 9038 8461

www.bgs.ac.uk/gsni/

Parent Body

Natural Environment Research Council, Polaris House, North Star Avenue, Swindon SN2 1EU

Tel 01793 411500 Fax 01793 411501
www.nerc.ac.uk

Website www.bgs.ac.uk

Shop online at www.geologyshop.com

Contents

- Contents 2
- 1 Introduction 4
- 2 Source Data and 3D Modelling Information 5
 - 2.1 Borehole Data 5
 - 2.2 Bedrock Geology Linework 6
 - 2.3 Modelling Workflow 6
- 3 Geology of the Bedrock Units 7
 - 3.1 Stratigraphy 7
 - 3.2 Geological structure 10
- 4 Conclusions and Recommendations 12
- References 13
- Appendix 1 Borehole Stratigraphy 14
- Appendix 2 Figures as Separate PDF file 15

FIGURES (located in accompanying pdf file)

Figure 1 Maps showing the geographical features and elevation of the area of interest (blue polyline).....15

Figure 2 Distribution of boreholes used in the investigation area that are held in the BGS Single Onshore Borehole Index (SOBI).15

Figure 3 Diagram showing the workflow employed in the generation of surfaces 15

Figure 4 Perspective view of the topography and geology of the Bromsgrove area 15

Figure 5 Structural blocks of the Bromsgrove area.....15

Figure 6 Cross-sections derived from the bedrock geology model.....15

Figure 7 Geophysical and lithological logs from the Wildmoor No. 2 Borehole. 15

Figure 8 Correlation of gamma-ray logs from the Wildmoor No. 2, Wildmoor (1914) Burcot No. 4 and Brockhill No. 1 boreholes. The distinctive high gamma zone appears to be related to an interval of clayey sandstones within the formation (depths in red are downhole depth in feet).15

Figure 9 Correlation of lithological and geophysical logs in the Bromsgrove Sandstone Formation in the Bromsgrove structural block.....15

Figure 10 Correlation of lithological and geophysical logs in the Bromsgrove Sandstone Formation across the wider Bromsgrove area.15

Figure 11 Distribution of the Droitwich Halite.15

Figure 12 A-G Contour maps on the bases of the main formations and thickness maps of the Mercia Mudstone Group and Sherwood Sandstone Formation.....15

TABLES

Table 1 List of boreholes with geophysical logs.....6

Table 2 Stratigraphy of the Bromsgrove aquifer block.....8

1 Introduction

The Bromsgrove Sandstone aquifer is over-abstracted. This has resulted in a long-term fall in groundwater levels, the reduction or loss of baseflow and the derogation of surface water features. To support flows in Battlefield Brook (a BAP site and notable amenity feature in Bromsgrove), Bow Brook (BAP site) and water levels in Hewell Grange Lake (SSSI), four alleviation of low-flow (ALF) boreholes are operated (one by the Environment Agency and three by Severn Trent Water). These ALF boreholes were installed as short term measures, prior to the implementation of a long-term solution to reduce the impacts of groundwater abstraction on surface water features which is under discussion between the water company and the Environment Agency.

In 2001 an existing groundwater model of the Bromsgrove Sandstone aquifer (developed by Birmingham University in 1990) was adopted and updated as part of the Environment Agency Midlands Region Groundwater Modelling Strategy. However, monitoring data collected since 2002 has shown that this groundwater model does not accurately simulate groundwater flows and levels in critical areas. The Bromsgrove aquifer groundwater modelling project aims to develop a new groundwater model that will be used to determine a more optimal groundwater abstraction regime which benefits the surface water environment, with the minimum of overall groundwater abstraction reduction and affordability.

The British Geological Survey (BGS) was contracted to undertake Environment Agency Task 1.1 of the Bromsgrove aquifer groundwater modelling study, namely the production of a three dimensional geological model of the investigation area. The model was specified to cover the outcrop of the Bromsgrove Sandstone Formation, the outcrop of the Clent Formation to the north and the confined Sherwood Sandstone Group to the west. The geographical limits of the area are approximately Droitwich Spa and Astwood Bank in the south (Northing 261550) and Rubery in the north (Northing 279560), Elmley Lovett (Easting 387134) in the west and Redditch (Easting 405456) in the east. The outline of the project area is given in Figure 1. The 3D geological model will be used in a concurrent Task (Task 1.2) to develop the conceptual model of groundwater flow between the principal formations of the Bromsgrove Sandstone aquifer system, as well as providing the geometrical information for building the groundwater model (Task 2).

This report outlines the methods used in the BGS 3D geological visualisation work and provides a brief summary of the stratigraphy, facies relationships and structure of the bedrock geology. Much of the information in the report has not been published before, and results from an extensive reinterpretation of existing borehole lithological descriptions and geophysical logs.

The model integrates information from BGS 1:50000 geological sheets (E167 Dudley, E168 Birmingham, E182 Droitwich and E183 Redditch), borehole descriptions derived from core or cuttings, geophysical logs and NEXTMap digital terrain data. Published information on the regional geological framework was also incorporated into the model (e.g. Old et al., 1991, Old et al. 1987, Powell et al. 2000).

2 Source Data and 3D Modelling Information

2.1 BOREHOLE DATA

The BGS Single Onshore Borehole Index (SOBI) holds approximately 4700 boreholes for the investigation area (including a variable buffer several kilometres wide). This large dataset was filtered for boreholes greater than 30 m in length due to the fact that the 3D visualisation involved thick (greater than 100 m) bedrock formations. This process removed most of the shallow geotechnical and site investigation boreholes that were unlikely to provide information relevant to the project. Depth filtering reduced the total number of boreholes to 183, and each of these scanned paper records was interrogated for useful geological information (Figure 2).

In total, 60 boreholes contain information that was used to build a 3D geological model, henceforth referred to as the bedrock geology model. This represents a relatively low density of control points, however, to a large extent this was compensated for by the extremely detailed lithological descriptions provided by many of the records. Unusually, many of the deep abstraction boreholes in the Bromsgrove area were cored, and these cores were logged in detail mostly during the 1950s-1980s by Professor L J Wills (University of Birmingham), G Warrington (BGS) and others with an interest in the Permo-Triassic stratigraphy of the area. The majority of these detailed bed-by-bed lithological descriptions have been entered into Oracle database tables as part of this task. No re-examination of core was undertaken as part of this work.

BGS held no geophysical logs for the investigation area. However, a selection of logs were provided by the Environment Agency/Severn Trent Water at varying scales from 1:200, 375, 750, 800, 1000, 1250 and 2000. Logs of three observation boreholes (Sanders Park (EA Observation Borehole Reference No. 0349), New Road Deep (0577) and Wildmoor (1914)) were supplied digitally and were plotted from Recall software. Considerable effort was expended in trying to source the original digital files for the remaining boreholes, however, these proved unobtainable. A selection of the remaining paper records were therefore scanned and digitised so they could be rescaled and plotted using WellCAD and Gocad.

Borehole	Type*	EA OBH ref	EASTING	NORTHING	TD	Drill year
Hewell Grange	ALF		400920	269430	147.8	2005
Battlefield Brook SP	ALF		394800	270500	100	1991
Wildmoor No1	PWS		395490	275180	263.3	1953
Wildmoor No2	PWS		395450	275190	264.4	1954
Wildmoor	OBH	1914	395961	274641	26.9	2004
Washingstocks No1	PWS		395930	273240	192.3	1899
Washingstocks No2	PWS		395800	273190	91.4	1900
Washingstocks No3	PWS		395930	273220	191.1	1924
Webheath No1	PWS		400980	266930	390.8	1970
Webheath No2	PWS		400800	266900	222.8	1974
Webheath No3	PWS		401180	266890	396.2	1972
Burcot No1	PWS		398470	271610	194.2	1938
Burcot No2	PWS		398480	271630	122.5	1894
Sugarbrook No1	PWS		396020	268150	288.0	1963
Sugarbrook No2	PWS		396100	268210	281.9	1965
Sugarbrook No3	PWS		396190	268160	398.4	1970

Sugarbrook No4	PWS		396100	268070	375.2	1969
Brockhill No1	PWS		400100	270160	186.5	1942
Brockhill No2	PWS		400180	270140	186.5	1942
New Road Deep	OBH	0577	396290	270260	147.5	1993
Sanders Park	OBH	0349	395310	270770	150	pre1978

* PWS = public water supply, ALF = Alleviation of low flow, OBH = observation borehole

Table 1 List of boreholes with geophysical logs

2.2 BEDROCK GEOLOGY LINEWORK

The extent of each modelled unit was initially constrained by existing geological mapping, as represented by BGS DiGMapGB-50. This 1:50 000 scale dataset provides a 2D representation of the bedrock geology (members, formations, groups and faults) either at surface or beneath superficial and artificial deposits. For modelling purposes, the DiGMapGB-50 geological linework was simplified to reduce node density and attributed with an elevation value using the NEXTMap terrain model.

It is important to note that while the modelled outcrop traces provided by the surfaces generally conform to the mapped outcrop traces as shown on BGS 1:50 000 scale maps, they are not an exact match:

1. In some areas the outcrop trace has been considerably simplified to streamline the modelling, for example in the structurally complex area to the east of the Lickey Hills (Block E on Figure 5). In this area, there is insufficient borehole control to justify an attempt to convert the many small outcrop polygons into 3D surfaces.
2. Mapping of the Droitwich (182) sheet is considerably older than all other areas and may not be as accurate and detailed as other geological map data used in this study.
3. Outcrop traces represent the intersection of a geological surface with the terrain model surface (which approximates to the geological horizon at ground surface). The outcrop trace can therefore vary depending on the type and resolution of the terrain model that is used. The geological map was constructed using a different terrain model (OS contours) to that used in the modelling (NEXTMap DTM downgraded to 50 m cell size and converted to a TIN in Gocad). Geological linework projected vertically onto a different terrain model than that used in the original mapping invariably produces an extremely irregular (in the Z dimension) and unlikely copline.
4. In this study, the Gocad 3D modelling application has been given considerable freedom to generate a synthetic outcrop trace by using only widely-spaced (>1 km) outcrop control points and then clipping (eroding) the geological surface with the terrain surface. This removes some of the difficulties created by mapped outcrop polylines which do not sit realistically on the terrain model. Observing the similarities and differences between the modelled and observed outcrop traces can also provide a useful means of validating the surface (e.g. highlighting a missed structure) or indeed indicating where the original geological linework may require modification.

2.3 MODELLING WORKFLOW

The bedrock geology model was constructed using GSI3D and Gocad 3D visualisation software. These applications facilitate the construction of faulted geological surfaces based on a combination of mapped geological linework, subsurface picks from borehole descriptions/geophysics and manually-inserted phantom points that are used to constrain the

surface in a geologically reasonable way. Figure 3 shows the workflow that was used to generate the surfaces:

1. Lithological and lithostratigraphical information from paper borehole records was entered into Oracle database tables.
2. Borehole records, geological maps and a terrain model (NEXTMap 50) were loaded into GSI3D and a grid of cross-sections was generated. The building of cross-sections provides stratigraphic control on the generation of phantom points between the relatively widely-spaced borehole control points. Each cross-section polyline was attributed with a formation name.
3. The XYZ coordinates of cross-section polylines were exported from GSI3D to Gocad. In Gocad a mesh with a predefined triangle size was created and then deformed using Discrete Smooth Interpolation (DSI) to best-fit the points set for each surface. A triangle size of 1 km was considered sufficient given the relative sparseness of borehole control points and the apparent absence of folding with a wavelength of less than 1 km in the Permo-Triassic cover. The triangulated surfaces were pre-cut with simplified fault traces prior to DSI to build in the effects of faulting. Faults were modelled as simple, continuous vertical planes. In reality it is likely that the faults are inclined and composed of multiple offset segments. This is a pragmatic simplification given that the bedrock geology model will provide the geometric framework for a groundwater model.
4. Geophysical log data were loaded into Gocad and then both the surfaces and the log correlations were evaluated. Faults which did not appear necessary were removed. The dip and azimuth of modelled surfaces were checked against outcrop measurements derived from BGS maps and memoirs.
5. Insight gained from the 3D visualisation often resulted in a need to adjust borehole interpretations and the derived cross-sections. Several iterative loops of borehole interpretation, section building and surface construction were required to generate the final bedrock geology model. The latest version of the bedrock geology model may be subject to further (re)interpretation as new data become available, and the inherent uncertainties in building subsurface geological models should be taken into account when using this model.

3 Geology of the Bedrock Units

3.1 STRATIGRAPHY

Table 2 shows the five main stratigraphic intervals covered by the study. No single borehole within the area penetrates the complete stratigraphical range of these formations, and their precise thicknesses is in most cases uncertain. Full details on the historical evolution of these lithostratigraphical terms can be found elsewhere (e.g. Warrington, 1970; Warrington et al. 1980).

Triassic	Mercia Mudstone Group		Mudstones, thin sandstones and gypsum, includes the Droitwich Halite Member in the Worcester Basin
	Sherwood Sandstone	Bromsgrove Sandstone Formation	Conglomerates at base fining upwards to sandstones and mudstones
		Wildmoor Sandstone Formation	Friable sandstones and thin mudstones

	Group	Kidderminster Formation	Conglomerates with sandy matrix fining upward to sandstone
Lower Permian?	Clent Formation		Breccias with clayey matrix

Table 2 Stratigraphy of the Bromsgrove aquifer block

3.1.1 Clent Formation

The Clent Formation typically comprises poorly sorted breccia with angular and subangular fine-grained volcanic clasts set in a reddish brown mudstone matrix. These breccias range up to 150 m in thickness and commonly lie marginal to present basinal areas. Within the investigation area, they are only mapped on the western slope of the Lickey/Clent Hills (Figure 5). They appear to pre-date the development of Permo-Triassic extensional basins and have been interpreted as Late Carboniferous/Early Permian alluvial fan deposits (Powell et al. 2000).

3.1.2 Kidderminster Formation

The Kidderminster Formation comprises a succession of fluvial conglomerates and sandstones formerly known as the ‘Bunter Pebble Beds’. The base of the Kidderminster Formation is generally a clast to matrix-supported conglomerate with pebbles of grey-brown quartzite, vein quartz and rare chert, rhyolite and tuff. The conglomerate beds show a range of internal structures including poorly developed horizontal stratification, and planar and trough cross stratification. Beds of conglomerate interfinger with sandstones, pebbly sandstones and red-brown micaceous mudstones. The thickest known sequence of the formation is 166.7 m in the Kempsey Borehole (Whitaker, 1980) sited a little to the south of the study area. The characteristic high sonic velocity of the basal Kidderminster Formation probably results from extensive carbonate and/or silica cementation. Palaeozoic limestone clasts within the conglomerate could have acted as a local source of carbonate. In the upper part of the formation, the proportion of conglomerate decreases and it is dominated by red-brown, fine- to coarse-grained, slightly micaceous sandstones in which pebbles are dispersed or form discrete lenses or channel lags.

The junction with the overlying Wildmoor Sandstone Formation is transitional and is drawn conventionally at the point above which the sequence becomes pebble free (Figure 7). However, the recognition of a strong geophysical marker (high gamma-ray zone) toward the base of the Wildmoor Sandstone in the Wildmoor No. 2, Wildmoor (1914), Burcot No. 4 and Brockhill No. 1 boreholes (Figure 8) shows that the highest pebble bed can occur at different stratigraphical levels in different areas. A more reliable criterion would appear to be the first downhole appearance of conglomerates which is generally associated with an increase in resistivity (Figure 7). This may be up to 65 m below the first appearance of pebbles, as in the case of the Wildmoor No. 2 Borehole. In the model, the Kidderminster Formation/Wildmoor Sandstone contact has been taken at the first downhole appearance of pebbles in the Wildmoor No. 2 Borehole which in this borehole coincides with a gradual change toward a more uniformly high sonic velocity. A comparable level has been picked in other boreholes using (1) the known thickness between this level and the overlying high gamma marker in the Wildmoor Sandstone and (2) limited geophysical log correlation (this would be the preferred method but more work is required to acquire, digitise and interpret all of the geophysical logs).

3.1.3 Wildmoor Sandstone Formation

The Wildmoor Sandstone Formation (formerly known as the ‘Upper Mottled Sandstone’) comprises reddish brown, very fine to fine-grained, micaceous sandstones. The base of the formation is generally gradational with upper sandstones of the Kidderminster Formation. The upper boundary is a sharp erosional disconformity with the Bromsgrove Sandstone. The formation is 284.7 m thick in the Worcester Basin (Kempsey Borehole, to the south of the study

area; Whittaker, 1980). The sandstones range from friable to moderately well cemented, with a dolomitic to patchily calcareous cement. Sedimentary structures include cross-lamination and low-angle trough and planar cross-bedding. Sandstone beds are commonly separated by thin, but laterally persistent, mudstone beds. The formation is generally interpreted as fluvial in this region, but it may also contain an aeolian component.

An important outcome of this study is the recognition of the strong geophysical marker (high gamma-ray zone) approximately 100 m above the base of the formation, as described in Section 3.1.2 above (also Figure 8). Lithological records are often unclear (largely due to extensive core loss of the friable sandstones), but this 25-30 m thick interval appears to correspond to a zone of clayey sandstones.

3.1.4 Bromsgrove Sandstone FORMATION

The Bromsgrove Sandstone Formation, formerly known as the 'Lower Keuper Sandstone', is the youngest formation in the Sherwood Sandstone Group. The formation rests sharply on the Wildmoor Sandstone and passes gradationally upwards into the Mercia Mudstone Group. The Bromsgrove Sandstone is lithologically distinct from the Wildmoor Sandstone Formation, the most conspicuous feature being the reappearance of conglomerates in the lower part of the formation, the incoming of fresh K-feldspar as a detrital mineral component (Warrington, 1970), and a greater incidence of thin mudstone beds. The erosion surface at the base of the Bromsgrove Sandstone, the Hardegsden Disconformity, has long been recognised as a key unconformity in UK Triassic stratigraphy.

The formation reaches a maximum thickness of 493.8 m in the Worcester Basin (Kempsey Borehole, to the south of the study area; Whittaker, 1980) and thins to the north and east where it is typically 50-100 m thick. The formation is lithologically variable and includes conglomerates, cross-bedded sandstones, siltstones and claystones deposited in a predominantly fluvial environment. Fining up cycles are common. It has traditionally been subdivided into three units, namely the 'Basement Beds', 'Building Stones', and 'Waterstones', which have been formalised as the Burcot, Finstall and Sugarbrook members (Old et al., 1991). The Sugarbrook Member is laterally equivalent to the Tarporley Siltstone Formation of the Mercia Mudstone Group elsewhere in the UK. The interbedding of sandstones and mudstones throughout the formation results in a characteristic serrated gamma-ray profile, with a progressive increase in value as the Bromsgrove Sandstone fines upwards into the Mercia Mudstone Group.

Sugarbrook No. 3 Borehole is the key reference borehole, having spudded into the Mercia Mudstone Group and thoroughly lithofacies-logged. The gamma ray log correlates well with the adjacent Sugarbrook No. 4 Borehole (Figure 9). The recognition of internal stratigraphical markers or zones in the Bromsgrove Sandstone was an important finding of this study, as it was previously uncertain whether any boreholes at other locations in the Bromsgrove area actually prove the contact between the Bromsgrove Sandstone and the Wildmoor Sandstone. Structural markers independent from this boundary were required. Figure 9 shows how the stratigraphy has been subdivided into three zones.

The lower zone (broadly equivalent to the Burcot Member) consists of conglomerates, pebbly sandstones and mudstones with common early-diagenetic carbonate nodules (calcrete). As might be expected in a fluvial environment with switching channel belts, pebbly sandstones and conglomerates can occur at different stratigraphical levels in different locations. A consequence of this is that the downward transition from pebbly lithologies into sandstones may have been incorrectly interpreted as a Bromsgrove-Wildmoor boundary in some previous borehole interpretations.

The upper zone comprises mainly sandstones and mudstones with rare calcrete but locally common plant remains including equisetalean pteridophytes and coniferalean gymnosperms (Old et al. 1991). This part of the formation has been largely eroded from the Washingstocks-

Whitford Public Water Supply (PWS) area, and thickens southward from Sanders Park toward Sugarbrook PWS Borehole (Figure 9).

There is some evidence from gamma-ray logs and lithological descriptions that a (generally) loosely-bounded 50 m thick central zone of more abundant and thicker mudstones separates the upper plant-rich sandstones from the lower calccrete-rich pebbly sandstones. This interval occurs at a depth of approximately 200-250 m in the Sugarbrook PWS Boreholes and can be traced up-dip to Washingstocks PWS Borehole where, in this case, a sharply-defined high-gamma ray/muddy zone occurs near the top of the borehole at 20-70 m downhole depth (Figure 9).

The general two/three-fold division of the Bromsgrove Sandstone can be extended to other areas in the Bromsgrove region, such as along a section line from Ombersley Park, Westwood and Chaddlesley Corbett to the north and west of Sugarbrook PWS, through to Webheath PWS and eastwards across the Lickey Fault (Figure 10). Many faults and fractures are recorded in the Webheath cores, and parts of the upper plant-rich sandstones may be faulted out in these boreholes.

3.1.5 Mercia Mudstone Group

The Mercia Mudstone Group (MMG) is composed predominantly of red, blocky mudstones and siltstones with subordinate laminated, green and grey beds. Thin beds of greenish grey, dolomitic siltstone and sandstone (often termed 'skerries') occur at intervals throughout. The group exceeds 500 m thick in the deeper basins of the Midlands; the Stratford seismic reflectin profile suggested thicknesses in excess of 1000m adjacent to the Inkberrow Fault on the margin of the Worcester Basin in the western part of the study area. Within the Mercia Mudstone, the Arden Sandstone Formation forms a prominent marker horizon, which is up to 11 m thick. Gypsum is common throughout the group, occurring in small primary nodules and secondary veins. At least one prominent bed of gypsum occurs near the top of the group. Halite is also developed, with local thick (150 m) halite-bearing units in the northern part of the Worcester Graben at Droitwich (Figure 11). This is the lateral equivalent of the Arden Sandstone. The distribution of the Droitwich Halite Member is not shown on the geological map of this area, but is well-known from a line of brine-producing boreholes between Droitwich and Stoke Prior Salt Works (Poole and Williams, 1980), and from subsidence features at the surface. The south-eastern dip of the halite across the Shernal Green Fault has been modelled using information from the ICI Saleway Borehole. There is no control on the presence or location of the halite adjacent to the Western Boundary/Lickey Fault (Figure 11). The modelled form of the halite in this area is based only on the concept that the distribution of the Droitwich Halite is controlled by the fault-bounded basin. Evidence from mapping in the Worcester area shows the limits of the Droitwich Halite are fault controlled on its western margin and the Inkebrrow Fault is the eastern limit (Barclay et al., 1997).

3.2 GEOLOGICAL STRUCTURE

The project area covers the northern part of the Worcester Graben in the west, the southern extent of the South Staffordshire Coalfield in the north and the western part of the Knowle Basin in the east. The South Staffordshire Coalfield is bounded to the west by a series of north-north-west-trending, down-west faults forming the eastern margin of the Worcester Graben. These faults merge with the north-south trending Inkberrow Fault, where the Lower Jurassic crops out near the southern margin of the area as shown in Figure 4.

The main consequence for the hydrogeology is that there are two considerably varying geological sequences. The sequence within the Worcester Graben includes a very thick Bridgnorth Sandstone, Kidderminster, Wildmoor Sandstone, Bromsgrove Sandstone formations and Mercia Mudstone Group, as proved in the Kempsey Borehole (Whittaker, 1980) and the Stratford seismic reflection profile (Chadwick, 1985). To the east of the Inkberrow-Lickey Fault

line, a thinner sequence probably overlies a basement high (Lickey Ridge), comprising thin or absent Wildmoor Sandstone, with Kidderminster Formation unconformably overlying Carboniferous to Permian breccias (Clent Formation). These breccias may have been reached in the deepened Burcot No 3 Borehole, although there is a possibility that these sandy breccias represent a locally-derived facies of the Kidderminster Formation. It is probable that the Clent Formation does not occur in the basin and the Bridgnorth Sandstone does not occur on the high. However, there is no seismic reflection data within the project area to confirm this. The Stratford seismic reflection profile was acquired primarily to resolve this relationship between the Worcester Graben, Lickey Ridge and Knowle Basin, but was sited some distance to the south (Chadwick, 1985).

For the purposes of 3D visualisation, the stratigraphy and structure has been simplified into six main structural blocks (shown as A-F on Figure 5)

- A. To the west of the Western Boundary Fault and Lickey Fault systems, a Wildmoor Sandstone Formation to Jurassic sequence dips at around 5 degrees to the south-east. Although not shown on the BGS Droitwich 1:50000 map sheet, the Mercia Mudstone Group includes the Droitwich Halite Member which crops as a zone of 'wet rockhead' (i.e., salt is dissolved at shallow depths) between Droitwich and Stoke Prior and is intersected at depth in the south under a Jurassic and Triassic cover by the Saleway Borehole.
- B. Bromsgrove is located on a lozenge-shaped, fault-bounded block composed of Bromsgrove Sandstone dipping gently (<5 degrees) to the south-west beneath a cover of Mercia Mudstone. The western margin of the block is formed by the Western Boundary Fault which probably drops the top of the Bromsgrove Sandstone by around 200 m on its western side (Figure 6), although there is very limited borehole control on this estimate. The Lickey Fault forms the eastern margin of the block. In the northern part of Block B the Lickey Fault downthrows Bromsgrove Sandstone against Wildmoor Sandstone, and in the south, Mercia Mudstone against Bromsgrove Sandstone.
- C. The older Permo-Triassic formations (Clent Formation, Kidderminster Formation and Wildmoor Sandstone) form a broadly conformable package which dips southwest off the western slope of the Lickey and Clent Hills. These formations have the steepest dip in the area of up to 10 degrees. Together with older rocks that include the Ordovician Lickey Quartzite Formation, these Permo-Triassic strata form the footwall of the Lickey Fault (Figure 6).
- D. Block D, which is penetrated by the Webheath PWS boreholes, is a narrow, triangular sliver of Bromsgrove Sandstone which dips 5 degrees to the south-east beneath a cover of Mercia Mudstone. The block is a horst, bounded on the west by the Lickey Fault and on the east by the Longbridge Fault. The Longbridge Fault downthrows the base of the Mercia Mudstone to the east by around 200 m.
- E. This is a relatively steeply dipping (approximately 10 degrees) Kidderminster Formation to Mercia Mudstone succession faulted against the eastern slope of the Lickey Hills and bounded to the east by the Longbridge Fault. The 1:50000 map sheet shows a structurally complex, faulted outcrop pattern here which is greatly simplified in the bedrock geology model.
- F. This is a large area of Mercia Mudstone on the eastern, downthrown side of the north-south trending Longbridge Fault. The Bromsgrove Sandstone aquifer is concealed by around 200 m of Mercia Mudstone and is approximately half the thickness of equivalent strata west of the Lickey High.

This structural subdivision is also shown in the surfaces of the bedrock geology model (Figure 5).

4 Conclusions and Recommendations

The bedrock geology modelling process integrates a range of 2D and 3D geological information to produce an interpretation of the subsurface geology of the project area. Every effort has been made to ensure that the bedrock geology model is consistent with this information and the conceptual geological model for the project area. However, this bedrock geology model represents one interpretation of the available data, and it is important to recognise that other equally valid interpretations may be possible. Groundwater modelling within the Bromsgrove aquifer may highlight areas where the bedrock geology model requires modification.

Some stratigraphical observations and correlations in the report are new and would benefit from more lithological and particularly geophysical log correlation. This should be undertaken if it becomes apparent during the groundwater modelling phase, that the internal stratigraphy of the Bromsgrove Sandstone and Wildmoor Sandstone formations is influencing the storage and movement of groundwater.

References

British Geological Survey holds most of the references listed below, and copies may be obtained via the library service subject to copyright legislation (contact libuser@bgs.ac.uk for details). The library catalogue is available at: <http://geolib.bgs.ac.uk>.

Barclay, W.J., Ambrose, K., Cahdwick, R.A. and Pharoah, T.C. 1997. Geology of the country around Worcester. *Memoir of the British Geological Survey, Sheet 199 (England and Wales)*.

Chadwick, R. 1985. Seismic reflection investigations into the stratigraphy and structural evolution of the Worcester Basin. *Journal fo the Geological Society of London*, Vol. 142, 187-202

Old, R.A., Hamblin, R.J.O., Ambrose, K. and Warrington, G., 1991. Geology of the country around Redditch. *Memoir of the British Geological Survey, Sheet 183 (England and Wales)*

Old, R.A., Sumbler, M.G. and Ambrose, K., 1987. Geology of the country around Warwick. *Memoir of the British Geological Survey, Sheet 184 (England and Wales)*.

Poole E.G. and Williams, B.J. 1980. The Keuper Saliferous Beds of the Droitwich area. *Report of the Institute of Geological Sciences*, No. 81/2

Powell, J.H., Glover, B.W. and Waters, C.N., 2000. Geology of the Birmingham. *Memoir of the British Geological Survey, Sheet 168 (England and Wales)*.

Warrington, G., 1970. The stratigraphy and palaeontology of the "Keuper" Series of the central Midlands of England. *Quarterly Journal of the Geological Society of London*, Vol.126, 183-223.

Warrington, G. et al., 1980. A correlation of Triassic rocks in the British Isles. *Special Report of the Geological Society of London*, No. 13.

Whittaker, A. 1980. Kempsey No.1 Geological Well Completion Report. *Report Deep Geology Unit Institute of Geological Sciences*, No. 80/1.

Appendix 1 Borehole Stratigraphy

Boreholes in the table contain a stratigraphical contact. Boreholes not in the table which are contained within one unit are: Fairfield No.1, Meadow Farm = WRS; Fockbury Mill, New Road Shallow, Burcott Lane, Tardebigge, Hewell Grange ALF/OBH = BMS. Z is depth in metres above or below ordnance datum. MD is measured downhole depth. HGZ in the MarkerName column stands for High Gamma Zone and represents an informal muddier subdivision of the formation that is prefixed. Dip and azimuth represent the dip and direction of dip of the surface in the borehole.

WellName	X	Y	Z	MD	MarkerName	Dip	Azimuth
Battlefield_Brook_SP	394800	270500	29.0	59.0	5 2 BMS HGZ TOP	2.6	137.2
Brockhill_No1	400100	270160	98.1	36.5	6 1 WRS HGZ TOP	2.5	225.6
Brockhill_No1	400100	270160	76.6	57.9	6 2 WRS HGZ BASE	2.5	225.6
Brockhill_No1	400100	270160	13.0	121.6	6 WRS	2.5	225.6
Brockhill_No2	400180	270140	100.5	32.8	6 1 WRS HGZ TOP	3.1	307.5
Brockhill_No2	400180	270140	79.1	54.3	6 2 WRS HGZ BASE	3.1	308.1
Brockhill_No2	400180	270140	15.4	117.9	6 WRS	2.9	303.9
Burcot_No1	398470	271610	81.5	50.2	6 1 WRS HGZ TOP	10.0	236.7
Burcot_No1	398470	271610	54.1	77.6	6 2 WRS HGZ BASE	9.9	236.6
Burcot_No1	398470	271610	2.0	129.7	6 WRS	9.9	236.6
Burcot_No2	398480	271630	84.7	47.0	6 1 WRS HGZ TOP	6.9	234.3
Burcot_No2	398480	271630	57.3	74.5	6 2 WRS HGZ BASE	6.9	234.3
Burcot_No3	398470	271710	89.4	44.1	6 1 WRS HGZ TOP	6.9	234.3
Burcot_No3	398470	271710	61.9	71.6	6 2 WRS HGZ BASE	6.9	234.3
Burcot_No3	398470	271710	9.9	123.6	6 WRS	6.9	234.3
Burcot_No3	398470	271710	-96.2	229.7	7 KDM	5.4	234.2
Burcot_No4	398460	271720	89.1	42.6	6 1 WRS HGZ TOP	6.9	234.3
Burcot_No4	398460	271720	61.7	70.1	6 2 WRS HGZ BASE	6.9	234.3
Burcot_No4	398460	271720	9.6	122.1	6 WRS	6.9	234.3
Catshill	395760	273860	113.3	8.1	5 2 BMS HGZ TOP	1.4	158.8
Fairfield_No2	395650	275670	132.6	14.0	6 2 WRS HGZ BASE	8.3	233.9
Meadow_Farm	395680	274800	62.3	70.2	6 1 WRS HGZ TOP	8.8	216.3
Meadow_Farm	395680	274800	40.8	91.7	6 2 WRS HGZ BASE	7.7	220.7
Meadow_Farm	395680	274800	-13.8	146.2	6 WRS	6.2	230.7
New_Road_Deep	396290	270260	-25.4	107.9	5 2 BMS HGZ TOP	3.4	142.5
Sanders_Park_OBH	395310	270770	25.3	59.5	5 2 BMS HGZ TOP	2.5	149.0
Sugarbrook_No1	396020	268150	52.0	16.9	4 MMG LOWER	3.9	169.5
Sugarbrook_No1	396020	268150	-123.7	192.6	5 2 BMS HGZ TOP	4.1	160.8
Sugarbrook_No1	396020	268150	-168.9	237.8	5 1 BMS HGZ BASE	4.0	162.9
Sugarbrook_No2	396100	268210	54.7	15.1	4 MMG LOWER	3.3	167.7
Sugarbrook_No2	396100	268210	-121.5	191.3	5 2 BMS HGZ TOP	4.1	160.8
Sugarbrook_No2	396100	268210	-166.6	236.4	5 1 BMS HGZ BASE	4.0	162.9
Sugarbrook_No3	396190	268160	50.7	22.5	4 MMG LOWER	3.5	175.9
Sugarbrook_No3	396190	268160	-125.3	198.5	5 2 BMS HGZ TOP	3.3	177.7
Sugarbrook_No3	396190	268160	-170.3	243.4	5 1 BMS HGZ BASE	3.3	177.7
Sugarbrook_No4	396100	268070	45.6	24.8	4 MMG LOWER	3.5	175.9
Sugarbrook_No4	396100	268070	-130.3	200.7	5 2 BMS HGZ TOP	3.3	177.7
Sugarbrook_No4	396100	268070	-175.2	245.6	5 1 BMS HGZ BASE	3.3	177.7
Washingstocks_No1	395930	273240	93.4	22.4	5 2 BMS HGZ TOP	1.7	153.9
Washingstocks_No1	395930	273240	49.1	66.7	5 1 BMS HGZ BASE	1.7	153.9
Washingstocks_No2	395800	273190	93.8	17.9	5 2 BMS HGZ TOP	1.7	153.9
Washingstocks_No2	395800	273190	49.5	62.2	5 1 BMS HGZ BASE	1.7	153.9
Washingstocks_No3	395930	273220	92.9	22.3	5 2 BMS HGZ TOP	1.7	153.9
Washingstocks_No3	395930	273220	48.6	66.6	5 1 BMS HGZ BASE	1.7	153.9
Webheath	400930	266870	77.9	35.1	4 MMG LOWER	3.1	281.4
Webheath_No1	400980	266930	79.9	33.2	4 MMG LOWER	3.4	266.8
Webheath_No1	400980	266930	-253.6	366.7	5 BMS	3.7	241.3
Webheath_No2	400800	267000	71.2	50.0	4 MMG LOWER	2.3	294.8
Webheath_No3	401180	266890	-243.6	365.5	5 BMS	3.7	241.3
Whitford	394410	270690	49.0	41.8	5 2 BMS HGZ TOP	2.9	135.9
Whitford	394410	270690	-11.3	102.1	5 1 BMS HGZ BASE	3.1	137.1
Wildmoor	395961	274641	104.5	20.5	6 1 WRS HGZ TOP	19.5	244.4
Wildmoor_No1	395490	275180	110.9	28.3	6 1 WRS HGZ TOP	9.6	181.0
Wildmoor_No1	395490	275180	79.9	59.4	6 2 WRS HGZ BASE	8.8	185.8
Wildmoor_No1	395490	275180	-3.6	142.8	6 WRS	6.2	230.6
Wildmoor_No2	395450	275190	108.7	30.5	6 1 WRS HGZ TOP	7.2	262.3
Wildmoor_No2	395450	275190	77.4	61.9	6 2 WRS HGZ BASE	7.2	262.3
Wildmoor_No2	395450	275190	-7.5	146.8	6 WRS	7.2	262.3

Appendix 2 Figures as Separate PDF file

The following figures are bundled in the pdf file which accompanies this report.

Figure 1 Maps showing the geographical features and elevation of the area of interest (blue polyline).

Figure 2 Distribution of boreholes used in the investigation area that are held in the BGS Single Onshore Borehole Index (SOBI).

Figure 3 Diagram showing the workflow employed in the generation of surfaces

Figure 4 Perspective view of the topography and geology of the Bromsgrove area

Figure 5 Structural blocks of the Bromsgrove area.

Figure 6 Cross-sections derived from the bedrock geology model

Figure 7 Geophysical and lithological logs from the Wildmoor No. 2 Borehole.



Figure 8 Correlation of gamma-ray logs from the Wildmoor No. 2, Wildmoor (1914) Burcot No. 4 and Brockhill No. 1 boreholes. The distinctive high gamma zone appears to be related to an interval of clayey sandstones within the formation (depths in red are downhole depth in feet).



Figure 9 Correlation of lithological and geophysical logs in the Bromsgrove Sandstone Formation in the Bromsgrove structural block.

Figure 10 Correlation of lithological and geophysical logs in the Bromsgrove Sandstone Formation across the wider Bromsgrove area.

Figure 11 Distribution of the Droitwich Halite.

Figure 12 A-G Contour maps on the bases of the main formations and thickness maps of the Mercia Mudstone Group and Sherwood Sandstone Formation.









1.07 (71.57)	S007SE21 (N4776)	SISST,BMS,Fill,null
1.52 (71.12)		SISST,BMS,Soft sandstone,
9.3 (63.34)		SISST,BMS,Hard brown sandstone,
14.33 (68.31)		MDST,BMS,"Grey sandstone, siltstone and shale",
22.86 (49.78)		SISST,BMS,null,
24.26 (46.26)		SISST,BMS,Gray sandstone with plants,PLANT,
33.91 (39.64)		SISST,BMS,Coarse sandstone with mudstone,
42.46 (29.66)		SISST,BMS,"Grey sandstone, coarse with plants",PLANT,
45.72 (26.91)		MDST,BMS,"Grey, micaceous",
48.62 (24.00)		SISST,BMS,null,
49.07 (23.57)		SISST,BMS,null,
51.97 (20.66)		MDST,BMS,null,
54.1 (19.54)		SISST,BMS,Gray sandstone with salt pseudomorphs,
63.09 (9.54)		SISST,BMS,Gray sandstone and shale,
74.07 (-1.49)		SISST,BMS,Frequent chocolate shales,
86.26 (-13.62)		SISST,BMS,Light grey sandstone with abundant plants,PLANT,
89.92 (-17.27)		MDST,BMS,Some grey shale and thin sandstones,
94.38 (-21.54)		SISST,BMS,Light grey sandstone,
95.74 (-23.07)		MDST,BMS,Brown and grey shale,
97.23 (-24.59)		SISST,BMS,Gray sandstone,
101.19 (-28.55)		SISST,BMS,Brown and grey sandstone,
103.33 (-30.69)		SISST,BMS,Gray siltstone,
104.55 (-31.91)		SISST,BMS,Hard brown sandstone,
106.07 (-33.43)		SISST,BMS,Hard grey sandstone,
113.64 (-40.93)		SISST,BMS,Hard red sandstone,
114 (-41.36)		MDST,BMS,Chocolate shale,
115 (-42.38)		SISST,BMS,Brown sandstone,
117.35 (-44.71)		MDST,BMS,Chocolate shale,
119.46 (-46.94)		SISST,BMS,Brown siltstone sandstone,
119.79 (-47.15)		MDST,BMS,null,
123.14 (-50.51)		SISST,BMS,Brown sandstone,
123.75 (-51.11)		MDST,BMS,null,
124.97 (-52.33)		SISST,BMS,null,
126.19 (-53.55)		MDST,BMS,Purple mudstone,
132.59 (-59.94)		SISST,BMS,null,
133.66 (-61.00)		MDST,BMS,null,
137.36 (-64.52)		SISST,BMS,null,
137.77 (-65.13)		MDST,BMS,null,
138.68 (-66.03)		SISST,BMS,null,
140.21 (-67.57)		MDST,BMS,null,
141.12 (-68.47)		SISST,BMS,null,
143.26 (-70.61)		MDST,BMS,null,
146.59 (-75.94)		SISST,BMS,"Brown, micaceous, occasional shale",
150.11 (-77.47)		SISST,BMS,Hard sandstone,
161.85 (-89.21)		MDST,BMS,Chocolate shale,
162.46 (-89.82)		MDST,BMS,Chocolate shale,
165.81 (-93.17)		MDST,BMS,null,
170.89 (-98.35)		SISST,BMS,Coarse brown sandstone,
174.86 (-102.32)		MDST,BMS,null,
177.7 (-105.06)		SISST,BMS,Coarse sandstone,
178 (-105.36)		MDST,BMS,null,

1. Convert paper borehole records to database records

2. Construct geological cross-sections in GSI3d




3. Create fence diagram in GSI3d and refine cross-sections




4. Export cross-section XYZ points to GOCAD and generate faulted triangulated mesh



Use 3d visualisation to refine borehole interpretations and repeat loop multiple times





- Mercia Mudstone
- Bromsgrove Sandstone
- Wildmoor Sandstone
- Kidderminster Conglomerate
- Clent Breccia






Structural Blocks


- A:** Wildmoor Sandstone to Jurassic sequence dipping at around five degrees SE. Forms the northern limit of the Worcester Basin lying west of the Western Boundary-Lickey Fault system
- B:** Fault block of Bromsgrove Sandstone to Mercia Mudstone dipping around 3 degrees SE
- C:** Clent Breccia to Wildmoor Sandstone dipping at around 8 degrees SW
- D:** Steeply dipping faulted block of Kidderminster Conglomerate to Mercia Mudstone
- E:** Fault block of Bromsgrove Sandstone to Mercia Mudstone dipping around five degrees SE
- F:** Downfaulted block of Mercia mudstone over Bromsgrove Sandstone to the east of the Longbridge Fault system.



Section 1





Section 2



Scales in metres

PROJECT Bromsgrove Aquifer Project
 WELL Wildmoor ABH 2
 LOCATION
 ELEVATION
 LOGGER
 DATE





Wildmoor2

Wildmoor1914

Burcott4

BrockhillPS1




North

Boreholes are not adjusted to true elevation


British Geological Survey
 BGS
 BRITISH GEOLOGICAL SURVEY
 Natural Resources Research Centre

PROJECT: Birmingham Aquifer Study
 WELL: VIMBOROUGH SOUTH 17/5
 LOCATION: VIMBOROUGH SOUTH
 ELEVATION: 100.0
 LOGGERS: WYBORN 03/03/15
 DATE: 2015




British Geological Survey
 BGS
 BRITISH GEOLOGICAL SURVEY
 Natural Resources Research Centre

PROJECT: Birmingham Aquifer Study
 WELL: SANDHILL PARK
 LOCATION: SANDHILL PARK
 ELEVATION: 100.0
 LOGGERS: WYBORN 03/03/15
 DATE: 2015




British Geological Survey
 BGS
 BRITISH GEOLOGICAL SURVEY
 Natural Resources Research Centre

PROJECT: Birmingham Aquifer Study
 WELL: NEW ROAD DIRT
 LOCATION: NEW ROAD DIRT
 ELEVATION: 100.0
 LOGGERS: WYBORN 03/03/15
 DATE: 2015




British Geological Survey
 BGS
 BRITISH GEOLOGICAL SURVEY
 Natural Resources Research Centre

PROJECT: Birmingham Aquifer Study
 WELL: HAZELTON FARM 098/E22
 LOCATION: HAZELTON FARM 098/E22
 ELEVATION: 100.0
 LOGGERS: WYBORN 03/03/15
 DATE: 2015




British Geological Survey
 BGS
 BRITISH GEOLOGICAL SURVEY
 Natural Resources Research Centre

PROJECT: Birmingham Aquifer Study
 WELL: SANDHILL PARK
 LOCATION: SANDHILL PARK
 ELEVATION: 100.0
 LOGGERS: WYBORN 03/03/15
 DATE: 2015




British Geological Survey
 BGS
 BRITISH GEOLOGICAL SURVEY
 Natural Resources Research Centre

PROJECT: Birmingham Aquifer Study
 WELL: SANDHILL PARK
 LOCATION: SANDHILL PARK
 ELEVATION: 100.0
 LOGGERS: WYBORN 03/03/15
 DATE: 2015



British Geological Survey
 BGS
 BRITISH GEOLOGICAL SURVEY
 Natural Resources Research Centre

PROJECT: Birmingham Aquifer Study
 WELL: SANDHILL PARK
 LOCATION: SANDHILL PARK
 ELEVATION: 100.0
 LOGGERS: WYBORN 03/03/15
 DATE: 2015



South



Base of Mercia Mudstone

Sandstones with plant with remains


Mud-rich high-gamma zone

Gravelly sandstone with calcretes

Presumed base of Bromsgrove Sandstone




British Geological Survey
 PROJECT: Bromsgrove Aquifer Study
 WELL: Bromsgrove Aquifer Study
 LOCATION: Bromsgrove Aquifer Study
 ELEVATION: Bromsgrove Aquifer Study
 LOGGING: Bromsgrove Aquifer Study
 DATE: Bromsgrove Aquifer Study




**Mercia Mudstone Group
 Bromsgrove Sandstone**


British Geological Survey
 PROJECT: Bromsgrove Aquifer Study
 WELL: Bromsgrove Aquifer Study
 LOCATION: Bromsgrove Aquifer Study
 ELEVATION: Bromsgrove Aquifer Study
 LOGGING: Bromsgrove Aquifer Study
 DATE: Bromsgrove Aquifer Study




British Geological Survey
 PROJECT: Bromsgrove Aquifer Study
 WELL: Bromsgrove Aquifer Study
 LOCATION: Bromsgrove Aquifer Study
 ELEVATION: Bromsgrove Aquifer Study
 LOGGING: Bromsgrove Aquifer Study
 DATE: Bromsgrove Aquifer Study



British Geological Survey
 PROJECT: Bromsgrove Aquifer Study
 WELL: Bromsgrove Aquifer Study
 LOCATION: Bromsgrove Aquifer Study
 ELEVATION: Bromsgrove Aquifer Study
 LOGGING: Bromsgrove Aquifer Study
 DATE: Bromsgrove Aquifer Study




British Geological Survey
 PROJECT: Bromsgrove Aquifer Study
 WELL: Bromsgrove Aquifer Study
 LOCATION: Bromsgrove Aquifer Study
 ELEVATION: Bromsgrove Aquifer Study
 LOGGING: Bromsgrove Aquifer Study
 DATE: Bromsgrove Aquifer Study




**Mud
 Zone**

British Geological Survey
 PROJECT: Bromsgrove Aquifer Study
 WELL: Bromsgrove Aquifer Study
 LOCATION: Bromsgrove Aquifer Study
 ELEVATION: Bromsgrove Aquifer Study
 LOGGING: Bromsgrove Aquifer Study
 DATE: Bromsgrove Aquifer Study




British Geological Survey
 PROJECT: Bromsgrove Aquifer Study
 WELL: Bromsgrove Aquifer Study
 LOCATION: Bromsgrove Aquifer Study
 ELEVATION: Bromsgrove Aquifer Study
 LOGGING: Bromsgrove Aquifer Study
 DATE: Bromsgrove Aquifer Study




**Plant-rich
 sandstones**

British Geological Survey
 PROJECT: Bromsgrove Aquifer Study
 WELL: Bromsgrove Aquifer Study
 LOCATION: Bromsgrove Aquifer Study
 ELEVATION: Bromsgrove Aquifer Study
 LOGGING: Bromsgrove Aquifer Study
 DATE: Bromsgrove Aquifer Study




Gravelly sandstones with calcrete



Bromsgrove Sandstone

Wildmoor Sandstone

Bromsgrove Sandstone



No control on salt distribution

Droitwich Halite




390000

400000

280000

270000

260000



Scale (m)





390000

400000


280000

270000

260000



Scale (m)





390000 400000

280000

270000

260000



Jurassic cover



