



Drivers of long-term variability in CO₂ net ecosystem exchange in a temperate peatland

C. Helfter¹, C. Campbell², K. J. Dinsmore¹, J. Drewer¹, M. Coyle¹, M. Anderson¹, U. Skiba¹, E. Nemitz¹, M. F. Billett^{1,3}, and M. A. Sutton¹

¹Centre for Ecology and Hydrology (Edinburgh), Penicuik, Scotland, UK

²Scottish Environmental Protection Agency, Stirling, Scotland, UK

³University of Stirling, Stirling, Scotland, UK

Correspondence to: C. Helfter (caro2@ceh.ac.uk)

Received: 15 July 2014 – Published in Biogeosciences Discuss.: 22 October 2014

Revised: 26 February 2015 – Accepted: 2 March 2015 – Published: 18 March 2015

Abstract. Land–atmosphere exchange of carbon dioxide (CO₂) in peatlands exhibits marked seasonal and inter-annual variability, which subsequently affects the carbon (C) sink strength of catchments across multiple temporal scales. Long-term studies are needed to fully capture the natural variability and therefore identify the key hydrometeorological drivers in the net ecosystem exchange (NEE) of CO₂. Since 2002, NEE has been measured continuously by eddy-covariance at Auchencorth Moss, a temperate lowland peatland in central Scotland. Hence this is one of the longest peatland NEE studies to date. For 11 years, the site was a consistent, yet variable, atmospheric CO₂ sink ranging from -5.2 to -135.9 g CO₂-C m⁻² yr⁻¹ (mean of -64.1 ± 33.6 g CO₂-C m⁻² yr⁻¹). Inter-annual variability in NEE was positively correlated to the length of the growing season. Mean winter air temperature explained 87 % of the inter-annual variability in the sink strength of the following summer, indicating an effect of winter climate on local phenology. Ecosystem respiration (R_{eco}) was enhanced by drought, which also depressed gross primary productivity (GPP). The CO₂ uptake rate during the growing season was comparable to three other sites with long-term NEE records; however, the emission rate during the dormant season was significantly higher. To summarise, the NEE of the peatland studied is modulated by two dominant factors:

- phenology of the plant community, which is driven by winter air temperature and impacts photosynthetic potential and net CO₂ uptake during the growing season (colder winters are linked to lower summer NEE),

- water table level, which enhanced soil respiration and decreased GPP during dry spells.

Although summer dry spells were sporadic during the study period, the positive effects of the current climatic trend towards milder winters on the site's CO₂ sink strength could be offset by changes in precipitation patterns especially during the growing season.

1 Introduction

Northern peatlands are one of the most important global sinks of atmospheric CO₂; with their ability to sequester C controlled by hydrometeorological variables such as precipitation, temperature, length of growing season and period of snow cover, they also potentially represent an important climatic feedback mechanism (Aurela et al., 2001; Frohling et al., 2001; Lafleur et al., 2003). Peatland carbon models generally suggest a decline in net sink strength in a warming climate, although the magnitude of the decline predicted by individual models is variable (Clark et al., 2010). UK peatlands are predicted to become a net source of carbon in response to climate change (Worrall et al., 2007), with climate models predicting a rise in global temperature of ca. 3 °C between 1980–1999 and 2100 (IPCC, 2007; scenario A1B which considers a balanced distribution between fossil fuel intensive and non-fossil fuel energy sources). A greater understanding of drivers and feedback mechanisms, across a range of temporal scales, is therefore a current research priority.

Eddy covariance measurements using fixed flux towers provide the best method for assessing inter-annual changes in land–atmosphere exchange of CO₂ at the catchment scale (Dinsmore et al., 2010; Nilsson et al., 2008; Roulet et al., 2007). In most years and in most peatlands, net ecosystem exchange (NEE) is the largest and most variable of the C flux terms (Roulet et al., 2007). In combination with aquatic fluxes (downstream and evasion losses) and CH₄ emissions, it is a key component of C and greenhouse gas (GHG) budgets for peatland systems (Billett et al., 2010; Dinsmore et al., 2010). Although more sites are now being established globally, there are still relatively few peatland sites (< 10) with published NEE measurements for periods of 3 years or more.

Including the Auchencorth Moss site, there are to our knowledge only six peatland sites in the Northern Hemisphere for which long-term (≥ 3 years) data sets of NEE are now available and all show that peatlands operate as a sink for atmospheric CO₂, albeit with different annual sink strengths. The 6-year mean NEE for Mer Bleue peatland (Ontario, Canada) was $-40.2 \text{ g C m}^{-2} \text{ yr}^{-1}$ (negative values signify uptake), varying year-to-year from a minor (-2) to a major (-112) CO₂ sink (Roulet et al., 2007). Similarly, McVeigh et al. (2014) found that a blanket bog in south-west Ireland had a mean 9-year NEE of $-55.7 \text{ g C m}^{-2} \text{ yr}^{-1}$ and exhibited significant inter-annual variability (-32.1 to $-79.2 \text{ g C m}^{-2} \text{ yr}^{-1}$). Degerö Stormyr in northern Sweden showed consistent yet variable CO₂ uptake over 12 consecutive years (12-year mean $-58 \pm 21 \text{ g C m}^{-2} \text{ yr}^{-1}$, range -18 to $-105 \text{ g C m}^{-2} \text{ yr}^{-1}$) (Peichl et al., 2014). Eddy covariance measurements at Lompolojänkka, a nutrient-rich fen in northern Finland, again showed that the site operated as a weak ($-3 \text{ g C m}^{-2} \text{ yr}^{-1}$) to strong ($-59 \text{ g C m}^{-2} \text{ yr}^{-1}$) CO₂ sink over a 3-year period (Aurela et al., 2009). In contrast to the variability exhibited by these sites, a sub-arctic permafrost mire in Northern Sweden was relatively stable over the period 2001–2008 ($-46 \text{ g C m}^{-2} \text{ yr}^{-1}$) (Christensen et al., 2012).

Quantifying inter-annual variability in NEE is a prerequisite for detecting longer term trends or step changes in flux magnitude in response to climatic or anthropogenic influences and identifying its drivers. In the UK, there have been significant regional changes in precipitation and temperature since the beginning of the 20th century, with the most rapid changes occurring over the last 50 years (Jenkins et al., 2009). During the period 1961–2006 annual precipitation increased by 2.5–23.2 %, with the largest increases occurring in the winter (particularly in Scotland and northern England); summer months were typically characterised by a decrease in precipitation. Mean annual temperature during the same period increased in parts of the UK by 1.05–1.64 °C (Jenkins et al., 2009), with winter months (January–February) warming much faster than the other months of the year in some parts (Holden and Rose, 2011). These data show that significant changes are taking place in seasonal climatic

patterns, which are likely to have a major impact on annual net CO₂ uptake by peatland systems.

Meteorological conditions such as rainfall, temperature and levels of photosynthetic active radiation (PAR) control NEE and its components, total ecosystem respiration (R_{eco}) and gross primary productivity (GPP). R_{eco} is composed of a plant respiration term (autotrophic respiration, R_{A}), which quantifies metabolic respiration from both above- and below-ground biomass, and a heterotrophic respiration term (R_{H}) resulting from microbial decomposition of organic matter in the soil. Autotrophic respiration can account for up to 60 % or R_{eco} whilst total belowground respiration can account for up to 70 % (van der Molen et al., 2011). R_{eco} and GPP have been shown to be tightly linked in a range of ecosystems on both short-term and annual timescales (Irvine et al., 2008; Law, 2005; Ryan and Law, 2005) and respond similarly, although not necessarily with the same magnitude, to extreme events such as drought. For example, short-term dynamics of R_{eco} can be more sensitive to the availability of labile C compounds produced by photosynthesis than to the effects of varying soil moisture on soil microbial activity (Irvine et al., 2008). On a global scale, drought is the main cause of decreased GPP alongside continent-specific secondary drivers such as cold spells and precipitation (Zscheischler et al., 2014a, b). Although less well understood and modelled than GPP, R_{eco} plays a major role in ecosystem C exchange dynamics, and increases in R_{eco} have been shown to turn a sink of C into a source (Lund et al., 2012). In order to interpret inter-annual variability in NEE, it is therefore crucial to partition NEE into GPP and R_{eco} and study their dynamics with respect to meteorology. We have done this on Auchencorth Moss, an ombrotrophic peatland in south-east Scotland.

The first eddy covariance measurements of CO₂ exchange at Auchencorth Moss took place in 1995–1996 (Hargreaves et al., 2003), with continuous measurements starting in 2002. Previous measurements of NEE have been published for specific 2–3 year time periods and suggest that inter-annual variability is high. Dinsmore et al. (2010) and Drewer et al. (2010) reported that over a 3-year period (2006–2008) the peatland acted as a very strong CO₂ sink (-88 to $-136 \text{ g C m}^{-2} \text{ yr}^{-1}$), whereas Billett et al. (2004) reported that between 1995–1996 it was acting as a weaker CO₂ sink (-36 and $-8 \text{ g C m}^{-2} \text{ yr}^{-1}$). In comparison to NEE, CH₄ emissions at Auchencorth Moss are small (average of $0.32 \text{ g CH}_4\text{-C m}^{-2} \text{ yr}^{-1}$ in 2007 and 2008; Dinsmore et al., 2010). Although these individual studies highlight significant inter-annual variability at Auchencorth Moss, they are for relatively short periods of time and are insufficient to investigate the drivers of inter-annual variability in NEE. Here we present the first complete analysis of the 2002–2013 data set in terms of monthly, seasonal and annual fluxes and explore the drivers of temporal variability in NEE. We use our data to test the following hypotheses:

- Colder than average winter temperatures affect the ecosystem's phenology and reduce summer GPP and NEE.
- Ecosystem respiration is related to water table depth and the peatland releases more CO₂ to the atmosphere during dry spells.
- Annual NEE is positively correlated with the length of the growing season.

2 Materials and methods

2.1 Site description

Auchencorth Moss (55°47'32" N, 3°14'35" W; 267 m a.s.l.) is a low-lying ombrotrophic peatland situated 17 km south-west of Edinburgh (Scotland, UK). Parent material comprises Upper Carboniferous/Lower Devonian sedimentary rocks overlain by fluvio-glacial till; peat depth range from <0.5 to >5 m. Long-term research (e.g. Billett et al., 2004; Dinsmore et al., 2010) on C fluxes is focussed on the 3.4 km² upper part of the catchment (elevation range 249–300 m) where the soils comprise peats (85 %), Gleysols (9 %), Humic Gleysols (3 %) and Cambisols (3 %). The open moorland site has an extensive uniform fetch over blanket bog to the south, west and north with a dominant wind direction from the south-west; winds from the north-east are the second most important wind direction. The terrain is relatively flat with a complex micro-topography consisting of hummocks and hollows. Hummocks are relatively small in size (typically 40 cm in diameter and ~30 cm in height) and covered by either a mix of *Deschampsia flexuosa* and *Eriophorum vaginatum*, or *Juncus effusus*. In contrast, hollows are dominated by mosses (*Sphagnum papillosum* and *Polytrichum commune*) and a layer of grasses (Dinsmore et al., 2009).

The site was drained more than 100 years ago (Leith et al., 2014); the drains have become progressively less effective and re-vegetated over time, leading to slow and progressive rewetting of the site. Over the last 20 years the site has been used for seasonal low intensity sheep grazing; areas of peat extraction occur at the margins of the catchment outside the footprint of the flux tower measurements.

2.2 Instrumentation and data processing

Fluxes of carbon dioxide (CO₂) have been measured continuously by eddy covariance (EC) at Auchencorth Moss since May 2002. The principles of operation and flux calculation methods using the eddy-covariance technique have been extensively discussed elsewhere (Aubinet et al., 2000; Baldocchi et al., 2001). The EC system at Auchencorth Moss consists of a LI-COR 7000 closed-path infrared gas analyser operating at 10 Hz for the simultaneous measurement of carbon

dioxide and water vapour. Turbulence measurements were made with an ultrasonic anemometer (initially model Solent R1012A R2 operating at 20.8 Hz; from 2009 Gill Windmaster Pro operating at 20 Hz; both Gill Instruments, Lymington, UK), mounted atop a 3 m mast. The effective measurement height is 3.5 m with a vertical separation of 20 cm between the anemometer and the inlet of the sampling line. Air is sampled at 20 L min⁻¹ through a 40 m long Dekabon line (internal diameter 4 mm). In addition to eddy-covariance measurements, the site is equipped with a Campbell Scientific 23X data logger for the automated acquisition of a comprehensive suite of meteorological parameters which include net radiation (Skye instruments SKS1110), PAR (Skye instruments SKP215), air temperature (fine wire type-E thermocouple), air pressure (Vaisala PTB101C), wind speed and direction (Gill Instruments WindSonic), soil water content (Campbell Scientific CS616 TDR probes), soil temperature (Campbell Scientific 107 thermistors at 10, 20, 30, and 40 cm), rainfall (tipping bucket rain gauge) and, since April 2007, water table depth (Druck PDCR 1830).

High-frequency eddy-covariance data is acquired by in-house software written in LabView (National Instruments) and processed offline into half-hourly fluxes.

Half-hourly data points were excluded from further analysis if any of the criteria listed below were not met:

- The total number of “raw” (high-frequency) data points per notional half-hour period was less than 90 % of the maximum possible number of points (36 000), i.e. below a minimum averaging period of 27 min.
- The number of spikes in raw w (vertical wind velocity component), CO₂ (CO₂ mole fraction) and H₂O mole fraction exceeded 1 % of the total number of points per half-hour period.
- The stationarity test devised by Foken and Wichura (1996), which compares half-hourly fluxes to the average of six 5 min averaging periods within the half hour, did not fulfil the quality criterion.
- Turbulence was insufficient for reliable EC measurements ($u_* < 0.1 \text{ m s}^{-1}$).
- CO₂ mole fractions < 330 ppm.
- Half-hourly CO₂ fluxes (F_{CO2}) fell outside the (–50 μmol m⁻² s⁻¹, +120 μmol m⁻² s⁻¹) interval.
- Half-hourly latent fluxes (LE) fell outside the (–250 W m⁻², +600 W m⁻²) interval.

After quality control, the number of good data points ranged from 45 % (in 2005) to 78 % (in both 2004 and 2008), with an annual mean of 65 ± 11 %.

Due to technical difficulties with the sampling pump (gradual decline in pumping performance), which were not detected immediately, most of the flux data for the summer period of 2011 were discarded as a precautionary measure.

2.3 Calculations of ecosystem respiration, Q_{10} and GPP

Gap filling of net ecosystem exchange (NEE) measured by eddy-covariance and partitioning of the gap-filled half-hourly fluxes into ecosystem respiration (R_{eco}) and gross primary production (GPP) was achieved using an online tool developed at the Max Planck Institute for Biogeochemistry, Jena, Germany¹ (Reichstein et al., 2005). In this flux partitioning approach, daytime R_{eco} is obtained by extrapolation of the night time parameterisation of NEE on air temperature (using an exponential relationship of the form given in Eq. 1).

$$R_{\text{eco}} = a \exp(bT) \quad (1)$$

where T is air temperature and a and b are fitting coefficients. GPP was then calculated as the difference between R_{eco} and measured NEE.

The growth rate (Q_{10}) for ecosystem respiration for a change of 10 °C was determined using the relationship:

$$Q_{10} = \left(\frac{R_2}{R_1} \right)^{\frac{10}{(T_2 - T_1)}} \quad (2)$$

T_1 and T_2 are reference temperatures (5 and 15 °C, respectively), and R_1 and R_2 are the corresponding respiration rates. R_1 and R_2 for each calendar year of the study were calculated from Eq. (1) using 24 h averages of measured night time T_{air} and NEE (see Supplement for non-linear regression statistics).

GPP was parameterised with respect to PAR using the following rectangular hyperbolic regression function:

$$\text{GPP} = \frac{\alpha \times \text{GPP}_{\text{sat}} \times \text{PAR}}{\text{GPP}_{\text{sat}} + \alpha \times \text{PAR}} \quad (3)$$

where GPP_{sat} (GPP at light saturation) and α (quantum efficiency) are fitting parameters.

2.4 Statistical tests

Statistical dependence between ecosystem response and hydro-meteorological variables was tested using Spearman's rank correlation. This allows testing for monotony between pairs of variables without making assumptions as to the nature of the function linking them. The independent variables winter air temperature, length of growing season (LGS) and annual water table depth (WTD), were tested for rank correlation against the dependent variables summertime NEE, R_{eco} , GPP, α and GPP_{sat} , annual NEE and annual GPP_{sat} .

The Spearman's correlation coefficient (ρ) is calculated using Eq. (4):

$$\rho = 1 - \frac{6 \sum d_i^2}{n(n^2 - 1)} \quad (4)$$

In Eq. (4), d_i is the difference between ranked variables and n the sample size.

Potential dependence between daily growing season (March–September) water table depth and ecosystem response (R_{eco} , GPP and NEE) was further investigated using one-way analysis of variance (ANOVA). The assumptions made were that (a) the 10 WTD classes (> 0 cm to < -45 cm in increments of 5 cm) constitute different treatments and (b) that the plant community has reached a steady state in terms of growth. The null hypothesis tested using this ANOVA is that WTD has no influence on ecosystem response.

3 Results

3.1 Site meteorology

During the study period (2002–2013) the site received a mean annual precipitation of 1018 ± 166 mm (\pm values denote standard deviation). Autumn (September–November) was the wettest season with 96 ± 11 mm of rain per month, and spring (March–May) was the driest with 64 ± 17 mm per month. Rainfall is highly variable year on year but records from a weather station of the UK Met Office (UK Meteorological Office, 2013) located 3.5 km north of the study site indicated a slight upward trend since the early 1970s (average annual precipitation 899 ± 166 mm for the period 1961–2001).

Mean annual air temperatures were 8.3 ± 4.6 °C for the study period 2002–2013 compared to 7.7 ± 4.5 °C for 1961–2001. Despite year-on-year variability there are indications of a steady increase of the order of 0.019 °C yr⁻¹ since records began in 1961 at the nearby Met Office station, which is consistent with UK and global trends (Jenkins et al., 2009). All seasons were warmer in 2002–2013 than in 1961–2001, albeit not significantly. Summer (June–August) was the warmest season with an average temperature of 13.6 ± 1.1 °C, and winter (December–February) the coldest with 3.7 ± 1.0 °C (Fig. 1).

Over the period April 2007–December 2013, water table depth (WTD) was within 4 cm of the peat surface for 50 out of 81 months (62 %). During dry periods, however, the water table could fall quickly to depths < -35 cm (Table 2).

3.2 Seasonal and inter-annual variability of R_{eco} , GPP and NEE

Ecosystem respiration typically peaked in July/August and was distributed asymmetrically around its peak value (Fig. 2), following the annual cycle of temperature. Plotting monthly GPP as a function of photosynthetically active radiation (PAR) reveals two separate plant productivity regimes culminating around mid-summer (Fig. 3). The hysteresis of GPP vs. PAR is characterised by an exponential growth phase from March to June/July followed by a logarithmic decline in photosynthetic efficiency. The ratio of GPP to R_{eco} showed

¹<http://www.bgc-jena.mpg.de/~MDIwork/eddyproc/upload.php>




Figure 1. Monthly air temperature, rainfall and photosynthetically active radiation (PAR) for the study period 2002–2013.




Figure 2. 10-year monthly averages of ecosystem respiration and, inset, ratio of gross primary production (GPP) to ecosystem respiration (horizontal dashed lines are 10-year annual mean, and error bars are the standard deviations).

that on average carbon uptake by vegetative growth exceeded losses to the atmosphere through respiration for six months of the year, from April to September (Fig. 2, inset).

A negative correlation was established between mean annual values of GPP_{sat} (GPP at light saturation, Eq. 3) and WT (Spearman $\rho = -0.63$, $p < 0.05$, Table 1) indicating that the photosynthetic capacity of the plant community tended to decrease as WT deepened. Furthermore, GPP_{sat} was positively correlated to the average temperature during the preceding winter ($\rho = 0.73$, $p < 0.01$, Table 1).

Both GPP and R_{eco} exhibited significant inter-annual variability with peak summer time values ranging from 96 to

245 g CO₂-C m⁻² month⁻¹ for GPP and 76 to 198 g CO₂-C m⁻² month⁻¹ for R_{eco} (August 2010 and July 2006, for minima and maxima, respectively). The site was consistently a sink for CO₂, however inter-annual variability was large. NEE (mean -64.1 ± 33.6 g CO₂-C m⁻²) ranged from -5.2 to -135.9 g CO₂-C m⁻² yr⁻¹ with minimum and maximum CO₂ uptake in 2013 and 2007, respectively (Fig. 4). As observed at other sites (Christensen et al., 2012), annual values of NEE were well correlated to the length of the growing seasons (LGS from here onward; $R^2 = 0.64$; Fig. 5). Furthermore, whilst mean spring/summer (April–September) NEE (NEE_{SS}) at Auchencorth Moss was not significantly corre-




Figure 3. Hysteresis in gross primary production (GPP) as a function of photosynthetically active radiation (PAR) (10-year monthly means).

Table 1. Spearman's rank correlation coefficients (ρ) and associated p values for all statistically significant inter-annual correlations between ecosystem response and hydro-meteorological parameters observed at Auchencorth Moss during the study period 2003–2013. The suffix SS denotes spring/summer means and LGS is the length of the growing season. GPP_{sat} and α are GPP at light saturation and quantum efficiency, respectively, obtained by non-linear regression between GPP and PAR using Eq. (3).

Ecosystem response	Parameter	ρ	p value
NEE_{SS}	Winter T_{air}	-0.96	< 0.01
NEE (annual)	LGS	-0.80	< 0.01
GPP_{SS}	Winter T_{air}	0.73	< 0.01
R_{ecoSS}	Winter T_{air}	0.61	0.02
GPP_{sat}	Winter T_{air}	0.68	0.02
α	Winter T_{air}	0.68	0.02
GPP_{sat} (annual)	WT (annual)	-0.63	< 0.05

lated to summer temperature, a strong negative correlation (i.e. net uptake increased with increasing winter T_{air}) was observed between mean NEE_{SS} and the mean air temperature of the preceding winter (December–March) ($R^2 = 0.87$, Fig. 6; $p < 0.01$). Comparable correlations to winter T_{air} were observed for GPP_{SS} and R_{ecoSS} ($R^2 = 0.43$, $p < 0.01$; $R^2 = 0.28$, $p = 0.02$, respectively, Fig. 6).

3.3 Effects of dry periods on CO₂ exchange

Throughout most years and most seasons Auchencorth Moss can be considered a wet site, with mean water table depth (WTD) -3.5 ± 6.8 cm and monthly range $+3.8$ cm (flooded; positive values denote water table levels above the peat surface) to -36 cm (WTD measurements commenced in




Figure 4. Annual NEE for 2003–2013 (no data for 2011 due to instrument failure during the growing season); the horizontal line is the mean NEE for the study period.




Figure 5. Annual NEE as a function of the length of the growing season.

April 2007). The site was generally waterlogged during the autumn and winter months. During dry spells, which we arbitrarily define as any period lasting 1 week or longer with $WTD < -5$ cm, the water table can drop quickly at rates up to 3 cm day^{-1} (Table 2).

Three notable dry spells occurred during the summer of 2010 and two during the summer of 2008, characterised by cycles of rapid fall and rise of the water table. Meanwhile, air temperatures exhibited little variation. Details of the drainage rates (water table drawdown) and minimum water table depths are given in Table 2. Under normal hydrological conditions (water table typically within 3–5 cm of the peat surface), R_{eco} at Auchencorth Moss did not exhibit a significant correlation with WTD. In contrast, during the dry spells of 2008 and 2010, daily R_{eco} was non-linearly correlated to WTD (Fig. 7). The response of R_{eco} to changes in WTD occurred with time lags ranging from 0 to 5 days (Table 2).

Table 2. Water table drainage rates and minimum water table depths (WTD) observed during the summer dry spells of 2008, 2010 and 2013. The time lag is the number of days elapsed between the start of the dry period and the onset of a response from the ecosystem respiration (R_{eco}); the time lag was determined by optimising the polynomial fit between R_{eco} and WTD. The minimum value of R_{eco} for each dry spell and the water table depth corresponding to each minimum value of R_{eco} were calculated using a second degree polynomial regression function between R_{eco} and WTD. No parabolic relationship was observed in 2013 between R_{eco} and WTD; for this reason, time lag, minimum R_{eco} and WTD for minimum R_{eco} could not be calculated.

Period	Drainage rate (cm day ⁻¹)	Minimum WTD (cm)	Time lag (days)	Minimum R_{eco} ($\mu\text{mol m}^{-2} \text{s}^{-1}$)	WTD for minimum R_{eco} (cm)	Mean T_{air} (°C)	Wind direction (°)
05–29/05/2008	1.2	–20.4	2	0.03	–1.5	10.1	70
22/07–01/08/2008	3.0	–19.1	3	2.31	–4.5	16.1	100
15–26/05/2010	1.6	–30.7	2	1.05	–15.6	12.9	181
09–24/06/2010	2.0	–36.1	0	1.58	–12.5	13.0	176
21/07–08/08/2010	2.0	–22.1	5	2.01	–2.9	11.4	191
26/05–06/09/2013	1.4	–48.5	–	–	–	14.5	222




Figure 6. Spring/summer fluxes of NEE, GPP and R_{eco} (mean from April to September) as a function of the preceding winter's mean air temperature (mean from December to March).

During the first two dry spells of 2010 the relationship between R_{eco} and WTD was of clear parabolic form, with R_{eco} reaching a minimum a few days after the onset of the dry period. Except for the second dry period of 2010, the residuals of the regressions between R_{eco} and WTD were not correlated with air nor soil temperature. The two dry spells of 2008 exhibited similar parabolic relationships between R_{eco} and WTD but differed in magnitude.

Such parabolic relationships between R_{eco} and WTD were not observed during the summer of 2013, which was the second driest in the 2002–2013 study period (the driest was 2003 with 346 mm rain between April and September compared to 361 mm in 2013); 2013 also had the longest winter of the study period (start of the growing season at day 103 in 2013 compared to day 77 ± 21 for the entire study period) as well




Figure 7. Daily ecosystem respiration as a function of water table depth during five dry spells (two in summer 2008 and three in 2010).

as the lowest soil temperatures. Soil temperature at -5 cm increased by 3°C in the 10 days prior to the start of the thermal growing season; T_{soil} increased steadily until mid-July and reached 15°C , the highest value of the 11-year study period, on 26 July. The dry period began on 25 May, culminating on 22 July (WTD = -48 cm), and WTD was ≤ -5 cm until early September. In 2013, the relationship between R_{eco} and WTD was linear across the 6 temperature classes considered (Fig. 8). Above 16°C , the positive correlation between R_{eco} and WTD was less pronounced and was even found to be negative for the 16 – 18°C temperature class. Above 18°C , the positive linear correlation was no longer statistically significant.

For all years for which WTD data was available, the sensitivity of R_{eco} to air temperature (Q_{10}) decreased with a drop in water table; in contrast, the theoretical values of R_{eco} at $T_{\text{air}} = 0^\circ\text{C}$ (obtained by extrapolation to the origin of




Figure 8. Ecosystem respiration as a function of water table depth and air temperature (daily means for May–September 2013).

the temperature-dependent functions fitted to monthly R_{eco} and averaged to annual values) were found to increase with WTD. One-way analysis of variance (ANOVA) on GPP, NEE and R_{eco} with respect to 10 WTD classes (Table 3) demonstrated that the position of the WT does have a statistically significant impact on R_{eco} for all years between 2007 and 2013, except 2012. For GPP, the correlation with WTD was significant in 2008 and 2010, and for NEE in 2013 only.

4 Discussion

The following sections discuss the effects of winter meteorology and water table depth on ecosystem response during the growing season and place the Auchencorth Moss peatland into a broader context by comparing it to other sites in the Northern Hemisphere with published NEE records ≥ 3 years.

4.1 Effect of winter meteorology on ecosystem response

Mean winter T_{air} explained 87 % of inter-annual variability in NEE during the following summer (NEE_{SS}), 43 % of GPP_{SS} and 28 % of R_{ecoSS} (Fig. 6), which is consistent with observations over a 12-year period at a boreal fen in northern Sweden (Peichl et al., 2014). A number of studies have reported correlations between winter meteorological conditions and the development of plant populations later in the year. Weltzin (2000) reported increased total net primary productivity (TNPP) in shrubs, a decrease in graminoids and no effect on bryophytes exposed to a gradient of infrared loading (i.e. continuous heating by infrared lamps). Individual species of bryophytes at a temperate UK site have been shown to respond to winter warming and/or summer drought in opposite ways, but this was not reflected at the

Table 3. Results (p value) of 1-way analysis of variance (ANOVA) on daily GPP, R_{eco} and NEE with respect to 10 water table depth (WTD) classes (>0 to <-45 cm in increments of 5 cm). Missing values denote failure of the equal variance test.

Year	NEE	R_{eco}	GPP
2007	–	0.02	–
2008	0.14	<0.01	0.02
2009	0.72	0.04	0.80
2010	0.93	<0.01	<0.01
2012	–	0.06	0.48
2013	0.03	<0.01	0.05

community level, whose mean cover did not exhibit significant differences between treatments (Bates et al., 2005). Kreyling (2008) demonstrated enhancement of aboveground net primary productivity (ANPP) in grasses as a result of freeze-thaw cycles the preceding winter, whilst belowground net primary productivity (BNPP) was adversely affected. Eddy-covariance measurements provide spatially integrated fluxes representative of the entire plant community within the footprint of the flux tower. The contributions of individual species, whose productivity can vary from year to year (Bates et al., 2005; Kreyling et al., 2010, 2008; Weltzin et al., 2000) cannot be assessed by EC. However, based on the knowledge that Sphagnum mosses are capable of photosynthesising as soon as the snow cover disappears and daily air temperature reaches >0 °C (Loisel et al., 2012), we speculate that the sensitivity of GPP, GPP_{sat} and α to winter air temperature is predominantly caused by graminoids and other non-moss species.

4.2 Effect of water table level on GPP and R_{eco}

WTD had a statistically significant negative effect on GPP indicating a decrease in plant productivity caused by the onset of drought stress. This has previously been shown to be important at other sites, particularly in moss species (Aurela et al., 2009; Lafleur et al., 2003; van der Molen et al., 2011); furthermore, a negative linear relationship between leaf area index (LAI) and WTD has been reported for a grassland established on drained organic soil in Ireland (Renou-Wilson et al., 2014), which illustrates the effect of water availability on graminoid productivity. It must however be noted that the WTD range in the Renou-Wilson (2014) study was significantly deeper (typically 20–60 cm below peat surface) than at our study site. Wet-adapted moss species growing in hollows are known to have large variability in growth rate directly linked to WTD (faster growth than hummock and lawn species under wet conditions, but susceptible to desiccation under dry conditions; Gunnarsson, 2005; Loisel et al., 2012). Weltzin et al. (2000) showed that, along a gradient of decreasing WTD of range consistent with our study site, TNPP increased in bryophytes, decreased in shrubs and was un-

changed in graminoids. As graminoids and bryophytes were the dominant species in the EC footprint, the sensitivity of GPP to WTD observed at our study site was likely to be mainly due to mosses.

The parabolic trend seen in the relationship between R_{eco} and WTD during dry spells (Fig. 7) may help understand the mechanistic drivers of R_{eco} at Auchencorth Moss. The parabolic trends were especially strong during the two first dry spells of 2010 (15 May–9 June 2010 and 10 June–10 July 2010) during which the prevailing wind direction was from the south. The WTD measurements might not be representative of the entire flux footprint, which could perhaps explain the markedly different trends observed in 2008 when wind was blowing from the east. We postulate that the initial decline in respiration was caused by a reduction in plant metabolic activity as water availability decreased (Lund, 2012). Drought has been shown to decrease C assimilation, slow the translocation of photosynthates between above- and belowground biomass, and reduce root-mediated respiration within days (Ruehr et al., 2009). Meanwhile, the lowering of the WT also favours aerobic processes, increases microbial activity and decomposition of organic matter (Hendriks et al., 2007; Moyano et al., 2013), and facilitates CO₂ diffusion within the peat profile (Moldrup et al., 1999; Tang et al., 2005) causing a net increase in CO₂ efflux from the soil. Minimum R_{eco} could then correspond to equilibrium between declining autotrophic and increasing heterotrophic respiration. The decrease of the sensitivity of R_{eco} with respect to T_{air} (Q_{10}) at our site is consistent with findings at other hydric sites where soil respiration (in particular heterotrophic respiration) has been shown to be enhanced by drought (Wang et al., 2014). The subsequent net increase in R_{eco} with deepening WTD could then be explained by a gradual increase in the ratio of heterotrophic to autotrophic respiration.

Based on these observations, we attribute the differences in respiration patterns during the dry spells to water table dynamics, which differs from drier sites where temperature (not WT) was found to be the dominant control of R_{eco} (Lafleur et al., 2005; Updegraff et al., 2001). This is further supported by the result of one-way ANOVA, which demonstrates a statistically significant correlation between R_{eco} and WTD for all growing seasons (except for 2012 which had a wetter than average growing season with WT near or above the peat surface for the entire growing season). The linear (rather than parabolic) response of R_{eco} to WTD in 2013 could perhaps be linked to the long winter of 2013 (the thermal growing season began 69 days later than in 2008, and 10 days later than in 2010) and the fact that the dry spell, which lasted most of the summer, began less than a month after the start of the growing season. Under these conditions, the moss population could have switched from relatively low metabolic activity to desiccation while active growth had just begun in the graminoid community. Hence, the $R_{\text{H}}/R_{\text{A}}$ ratio could have been smaller than in previous years. In contrast to other

years, GPP during summer 2013 was positively correlated to WTD ($p < 0.001$), which suggests growth in species less susceptible to drought-stress than mosses.

Disentangling the effects of lower than average winter air temperature and summer dry spells on annual NEE is not straightforward, but the former seems to be the dominant driver based on our results (Table 1). The combined effects of a long, relatively cold winter and warm, dry summer which could have slowed plant growth, disturbed the normal phenological cycle and enhanced carbon losses from the peatland through enhanced heterotrophic respiration, were illustrated in 2013 when the sink strength of Auchencorth Moss was significantly weakened ($-5.2 \text{ g C-CO}_2 \text{ m}^{-2} \text{ yr}^{-1}$) compared to the long-term mean of $-64.1 \pm 33.6 \text{ g C-CO}_2 \text{ m}^{-2} \text{ yr}^{-1}$ (2003–2012).

4.3 NEE in Northern Hemisphere peatland C budgets

Compared to other peatlands in the Northern Hemisphere, annual values of NEE at Auchencorth Moss are at the high end of both the mean ($-64.1 \pm 33.6 \text{ g CO}_2\text{-C m}^{-2} \text{ yr}^{-1}$) and inter-annual range (-5.2 to $-135.9 \text{ g CO}_2\text{-C m}^{-2} \text{ yr}^{-1}$). However, when the length of the growing season (LGS; the start of the growing season was defined as the first day of the year when mean diurnal air temperature exceeded 5°C for 5 consecutive days. Conversely, the end of the growing season was defined as the first day of the year when mean diurnal air temperature fell below 5°C for 5 consecutive days) was accounted for, the mean daily growing season NEE (NEE_{GS}) at Auchencorth Moss ($-0.57 \text{ g CO}_2\text{-C m}^{-2} \text{ day}^{-1}$) was remarkably similar to that found at both Mer Bleue (cool temperate bog; $-0.58 \text{ g CO}_2\text{-C m}^{-2} \text{ day}^{-1}$; Roulet et al., 2007) and Degerö Stormyr (boreal mire; $-0.48 \text{ g CO}_2\text{-C m}^{-2} \text{ day}^{-1}$; Peichl et al., 2014). By contrast, mean daily NEE_{GS} at Glencar (maritime blanket bog; Koehler et al., 2011, McVeigh et al., 2014) was slightly lower ($-0.39 \text{ g CO}_2\text{-C m}^{-2} \text{ day}^{-1}$), whilst the two sub-arctic Scandinavian peatlands Lompolojänkä (nutrient-rich sedge fen; Aurela et al., 2009) and Stordalen (sub-arctic palsa mire; Christensen et al., 2012) stand out with mean daily growing season NEE rates 2 to 2.5 times higher than the values found for Auchencorth Moss, Degerö Stormyr and Mer Bleue, and over 3 times higher than the value found at Glencar (Table 4).

Auchencorth Moss had a mean daily NEE during the dormant season (NEE_{DS}) of $0.61 \text{ g CO}_2\text{-C m}^{-2} \text{ day}^{-1}$, the highest amongst the aforementioned catchments (10, 5, 3 and 2 times higher than Glencar, Degerö Stormyr, Stordalen and Mer Bleue, respectively). Mean daily NEE_{DS} at Lompolojänkä was only slightly lower than at Auchencorth Moss ($0.52 \text{ g CO}_2\text{-C m}^{-2} \text{ day}^{-1}$).

Despite the lower daily mean NEE, the long growing season at Auchencorth Moss made its total NEE_{GS} comparable to that of Lompolojänkä and Stordalen. The vigorous net uptake at Lompolojänkä during the growing season was offset by relatively high carbon losses during the rest of the year.

Table 4. Annual minimum, maximum and mean values of NEE at several long-term peatland monitoring sites in the Northern Hemisphere. LGS and LDS are the length of growing and dormant season respectively, and subscripts GS and DS denote growing and dormant season. The length of the growing season for the study site Auchencorth Moss was bounded by the first and last day for which mean daily air temperatures exceeded 5 °C for 5 consecutive days. For the other sites, LGS was estimated from data available in the respective articles.

Site	Auchencorth Moss (this study)	Stordalen ¹	Mer Bleue ²	Glencar ³	Lompolojännkä ⁴	Degerö Stormyr ⁵
Latitude	55°47′	68°20′	45°23′	51°55′	68°0′	64°11′
Duration (years)	11	8	6	9	3	12
Minimum NEE (g CO ₂ -C m ⁻²)	-5.2	-20	-2	-32.1	-3.3	-18
Maximum NEE (g CO ₂ -C m ⁻²)	-135.9	-95	-112	-79.7	-58.9	-105
Mean NEE (g CO ₂ -C m ⁻²)	-64.1 ± 33.6	-66 ± 29.1	-40.2 ± 40.5	-55.7 ± 30.0	-31.9 ± 27.8	-58.0 ± 21.0
Mean NEE _{GS} (g CO ₂ -C m ⁻²)	-142 ± 55.0	-133 ± 28.0	-97.1 ± 38.7	-60 ± 15.0	-160 ± 13.0	-84.8 ± 23.6
Length of growing season (LGS) (days)	247	117 ^c	168 ^a	153 ^b	119	120
NEE _{GS} /LGS (g CO ₂ -C m ⁻² day ⁻¹)	-0.57	-1.14	-0.58	-0.39	-1.34	-0.48
NEE _{DS} /LDS (g CO ₂ -C m ⁻² day ⁻¹)	0.61	0.27	0.29	0.06	0.52	0.11

References: ¹ Christensen et al. (2012); ² Roulet et al. (2007); ³ McVeigh et al. (2014); ⁴ Aurela et al. (2009); ⁵ Peichl et al. (2014). ^a Estimated from Lafleur et al. (2003): growing season from May to September (1998–2002); ^b mean growing season lengths 2002–2007 (Sottocornola and Kiely, 2010); ^c use of NEE and LGS for the years 2006–2008 only, as winter measurements of NEE during the other years of the study were deemed unreliable by the authors (Christensen et al., 2012).

Auchencorth Moss, Lompolojännkä and Stordalen therefore had comparable NEE but for very different reasons: Auchencorth Moss had long growing seasons but also relatively high carbon losses the rest of the year, which could be due to milder winters with minimal snow cover. Lompolojännkä and Stordalen had vigorous carbon uptake rates, their LGS were comparable to one another, but were half that of Auchencorth Moss, whilst Lompolojännkä had high carbon losses during the dormant season which strongly reduced the site's sink strength.

Carbon uptake rates at Degerö Stormyr and Mer Bleue were very similar to Auchencorth Moss but their carbon loss rates, which were comparable to Stordalen's, were half or less than that of Auchencorth Moss. This could be explained by cooler climate and prolonged periods of snow cover compared to Auchencorth Moss.

Considering the differences in latitude, climate, hydrology and vegetation, these sites (with the exception of Stordalen and Lompolojännkä) are remarkably similar in terms of their daily mean NEE_{GS}.

NEE represents only one flux pathway within the full net ecosystem C budget (NECB). When terrestrial CH₄ emissions (2007–2008; Dinsmore et al., 2010), downstream aquatic flux losses and water surface evasion (2007–2011; Dinsmore et al., 2013) are accounted for, the total long-term sink strength of Auchencorth Moss is reduced to approximately -28 g C m⁻² yr⁻¹ (whilst recognising un-

certainty as the fluxes are not measured over the same time period). Using literature values of CH₄ (Roulet et al., 2007) and aquatic C losses for Mer Bleue (Billett and Moore, 2008) results in an approximate total C sink strength of -17 g C m⁻² yr⁻¹; for Degerö Stormyr the total C sink strength is -24 g C m⁻² yr⁻¹ (Nilsson et al., 2008), -30 g C m⁻² yr⁻¹ for Glencar (Koehler et al., 2011) and -34 g C m⁻² yr⁻¹ for Stordalen (Christensen et al., 2012; Lundin et al., 2013; Olefeldt et al., 2013); data for Lompolojännkä could not be found. Hence when all flux pathways are accounted for the C balances of the different peatlands appear to converge.

5 Summary

Eleven years of continuous monitoring of net ecosystem exchange of carbon dioxide at a temperate Scottish peatland revealed highly variable inter-annual dynamics despite little or no change in land management. Variation in climate and especially winter air temperature is thought to be the dominant control at the study site. The latter explained 87% of inter-annual changes in NEE and a modest rise of 1 °C above average winter air temperature for the 2002–2013 study period was accompanied by a 20% increase in CO₂ uptake. Colder winters appear to have an adverse effect on the peatland CO₂ sink strength possibly due to disturbances to the phenological cycle of the graminoid species at the site. Dry

spells have been linked to enhanced ecosystem respiration and depressed GPP and it is thought that (a) heterotrophic respiration can become the dominant term as water availability decreases, and (b) mosses are more sensitive to WTD than other species at the site. Cold winters and dry summers both have negative effects on the CO₂ sink strength of the bog; these two factors converged in 2013 and led to a significant reduction in net CO₂ uptake (−90 % compared to the 11-year mean). Auchencorth Moss, although always a sink of CO₂ during the study period, is highly sensitive to even modest changes in hydro-meteorological conditions at relatively short timescales. The large inter-annual variability of NEE observed to date makes future trends difficult to predict and quantify. Changes in seasonal hydro-meteorological conditions, especially changes in precipitation patterns and intensity, could however be pivotal for the CO₂ cycling of this peatland. Drier summers could lead to a reduction in net CO₂ uptake but this could be offset by milder temperatures, particularly in winter, and longer growing seasons. Mean annual temperatures at the study site have risen by 0.019 °C yr^{−1} since 1961, which could, in theory, benefit C uptake by the peatland in the long-term since NEE was found to be closely linked to the length of the growing season.

The Supplement related to this article is available online at doi:10.5194/bg-12-1799-2015-supplement.

Acknowledgements. The authors acknowledge a succession of European projects (FP5 CarboMont, FP6 NitroEurope, FP7 GHG Europe and ÉCLAIRE) and CEH carbon catchments for support of the measurements and data analysis. Measurements were supported by CEH colleagues Ian Leith, Ivan Simmons and Sarah Leeson.

Edited by: E. Pendall

References

- Aubinet, M., Grelle, A., Ibrom, A., Rannik, U., Moncrieff, J., Foken, T., Kowalski, A. S., Martin, P. H., Berbigier, P., Bernhofer, C., Clement, R., Elbers, J., Granier, A., Grunwald, T., Morgenstern, K., Pilegaard, K., Rebmann, C., Snijders, W., Valentini, R., and Vesala, T.: Estimates of the annual net carbon and water exchange of forests: The Euroflux methodology, *Adv. Ecol. Res.*, 30, 113–175, 2000.
- Aurela, M., Laurila, T., and Tuovinen, J. P.: Seasonal CO₂ balances of a subarctic mire, *J. Geophys. Res.-Atmos.*, 106, 1623–1637, doi:10.1029/2000jd900481, 2001.
- Aurela, M., Lohila, A., Tuovinen, J.-P., Hatakka, J., Riutta, T., and Laurila, T.: Carbon dioxide exchange on a northern boreal fen, *Boreal Environ. Res.*, 14, 699–710, 2009.
- Baldocchi, D., Falge, E., Gu, L. H., Olson, R., Hollinger, D., Running, S., Anthoni, P., Bernhofer, C., Davis, K., Evans, R., Fuentes, J., Goldstein, A., Katul, G., Law, B., Lee, X. H., Malhi, Y., Meyers, T., Munger, W., Oechel, W., Paw, K. T., Pilegaard, K., Schmid, H. P., Valentini, R., Verma, S., Vesala, T., Wilson, K., and Wofsy, S.: Fluxnet: A new tool to study the temporal and spatial variability of ecosystem-scale carbon dioxide, water vapor, and energy flux densities, *B. Am. Meteorol. Soc.*, 82, 2415–2434, 2001.
- Bates, J. W., Thompson, K., and Grime, J. P.: Effects of simulated long-term climatic change on the bryophytes of a limestone grassland community, *Glob. Change Biol.*, 11, 757–769, doi:10.1111/j.1365-2486.2005.00953.x, 2005.
- Billett, M. F. and Moore, T. R.: Supersaturation and evasion of CO₂ and CH₄ in surface waters at Mer Bleue peatland, Canada, *Hydro. Process.*, 22, 2044–2054, doi:10.1002/hyp.6805, 2008.
- Billett, M. F., Palmer, S. M., Hope, D., Deacon, C., Storeton-West, R., Hargreaves, K. J., Flechard, C., and Fowler, D.: Linking land-atmosphere-stream carbon fluxes in a lowland peatland system, *Global Biogeochem. Cy.*, 18, GB1024, doi:10.1029/2003gb002058, 2004.
- Billett, M. F., Charman, D. J., Clark, J. M., Evans, C. D., Evans, M. G., Ostle, N. J., Worrall, F., Burden, A., Dinsmore, K. J., Jones, T., McNamara, N. P., Parry, L., Rowson, J. G., and Rose, R.: Carbon balance of UK peatlands: Current state of knowledge and future research challenges, *Clim. Res.*, 45, 13–29, doi:10.3354/cr00903, 2010.
- Christensen, T. R., Jackowicz-Korczynski, M., Aurela, M., Crill, P., Heliasz, M., Mastepanov, M., and Friborg, T.: Monitoring the multi-year carbon balance of a subarctic peatland mire with micrometeorological techniques, *Ambio*, 41, 207–217, doi:10.1007/s13280-012-0302-5, 2012.
- Clark, J. M., Billett, M. F., Coyle, M., Croft, S., Daniels, S., Evans, C. D., Evans, M., Freeman, C., Gallego-Sala, A. V., Heinemeyer, A., House, J. I., Monteith, D. T., Nayak, D., Orr, H. G., Prentice, I. C., Rose, R., Rowson, J., Smith, J. U., Smith, P., Tun, Y. M., Vanguelova, E., Wetterhall, F., and Worrall, F.: Model inter-comparison between statistical and dynamic model assessments of the long-term stability of blanket peat in Great Britain (1940–2099), *Clim. Res.*, 45, 227–281, doi:10.3354/cr00974, 2010.
- Dinsmore, K. J., Skiba, U. M., Billett, M. F., and Rees, R. M.: Effect of water table on greenhouse gas emissions from peatland mesocosms, *Plant Soil*, 318, 229–242, doi:10.1007/s11104-008-9832-9, 2009.
- Dinsmore, K. J., Billett, M. F., Skiba, U. M., Rees, R. M., Drewer, J., and Helfter, C.: Role of the aquatic pathway in the carbon and greenhouse gas budgets of a peatland catchment, *Glob. Change Biol.*, 16, 2750–2762, doi:10.1111/j.1365-2486.2009.02119.x, 2010.
- Dinsmore, K. J., Billett, M. F., and Dyson, K. E.: Temperature and precipitation drive temporal variability in aquatic carbon and GHG concentrations and fluxes in a peatland catchment, *Glob. Change Biol.*, 19, 2133–2148, doi:10.1111/gcb.12209, 2013.
- Drewer, J., Lohila, A., Aurela, M., Laurila, T., Minkkinen, K., Penttila, T., Dinsmore, K. J., McKenzie, R. M., Helfter, C., Flechard, C., Sutton, M. A., and Skiba, U. M.: Comparison of greenhouse gas fluxes and nitrogen budgets from an ombrotrophic bog in Scotland and a minerotrophic sedge fen in Finland, *Eur. J. Soil Sci.*, 61, 640–650, doi:10.1111/j.1365-2389.2010.01267.x, 2010.

- Foken, T. and Wichura, B.: Tools for quality assessment of surface-based flux measurements, *Agr. Forest Meteorol.*, 78, 83–105, doi:10.1016/0168-1923(95)02248-1, 1996.
- Frolking, S., Roulet, N. T., Moore, T. R., Richard, P. J. H., Lavoie, M., and Muller, S. D.: Modeling northern peatland decomposition and peat accumulation, *Ecosystems*, 4, 479–498, doi:10.1007/s10021-001-0105-1, 2001.
- Gunnarsson, U.: Global patterns of sphagnum productivity, *J. Bryol.*, 27, 269–279, doi:10.1179/174328205x70029, 2005.
- Hargreaves, K. J., Milne, R., and Cannell, M. G. R.: Carbon balance of afforested peatland in Scotland, *Forestry*, 76, 299–317, doi:10.1093/forestry/76.3.299, 2003.
- Hendriks, D. M. D., van Huissteden, J., Dolman, A. J., and van der Molen, M. K.: The full greenhouse gas balance of an abandoned peat meadow, *Biogeosciences*, 4, 411–424, doi:10.5194/bg-4-411-2007, 2007.
- Holden, J. and Rose, R.: Temperature and surface lapse rate change: A study of the UK's longest upland instrumental record, *Int. J. Climatol.*, 31, 907–919, doi:10.1002/joc.2136, 2011.
- IPCC: Technical summary, Cambridge, UK, 19–92, 2007.
- Irvine, J., Law, B. E., Martin, J. G., and Vickers, D.: Interannual variation in soil CO₂ efflux and the response of root respiration to climate and canopy gas exchange in mature ponderosa pine, *Glob. Change Biol.*, 14, 2848–2859, doi:10.1111/j.1365-2486.2008.01682.x, 2008.
- Jenkins, G., Perry, M., and Prior, J.: The climate of the United Kingdom and recent trends, Hadley Centre, Met Office, Exeter, 2009.
- Koehler, A.-K., Sottocornola, M., and Kiely, G.: How strong is the current carbon sequestration of an Atlantic blanket bog?, *Glob. Change Biol.*, 17, 309–319, doi:10.1111/j.1365-2486.2010.02180.x, 2011.
- Kreyling, J., Beierkuhnlein, C., Pritsch, K., Schloter, M., and Jentsch, A.: Recurrent soil freeze-thaw cycles enhance grassland productivity, *New Phytol.*, 177, 938–945, doi:10.1111/j.1469-8137.2007.02309.x, 2008.
- Kreyling, J., Beierkuhnlein, C., and Jentsch, A.: Effects of soil freeze-thaw cycles differ between experimental plant communities, *Basic Appl. Ecol.*, 11, 65–75, doi:10.1016/j.baec.2009.07.008, 2010.
- Lafleur, P. M., Roulet, N. T., Bubier, J. L., Frolking, S., and Moore, T. R.: Interannual variability in the peatland-atmosphere carbon dioxide exchange at an ombrotrophic bog, *Global Biogeochem. Cy.*, 17, 1036, doi:10.1029/2002gb001983, 2003.
- Lafleur, P. M., Moore, T. R., Roulet, N. T., and Frolking, S.: Ecosystem respiration in a cool temperate bog depends on peat temperature but not water table, *Ecosystems*, 8, 619–629, doi:10.1007/s10021-003-0131-2, 2005.
- Law, B.: Carbon dynamics in response to climate and disturbance: Recent progress from multi-scale measurements and modeling in Ameriflux, in: Plant responses to air pollution and global change, edited by: Omasa, K., Nouchi, I., and DeKok, L. J., 205–213, 2005.
- Leith F. I., Garnett M. H., Dinsmore, K. J., Billett, M. F., and Heal, K. V.: Source and age of dissolved and gaseous carbon in a peatland-riparian-stream continuum: a dual isotope (¹⁴C and ^δ¹³C) analysis, *Biogeochemistry*, 119, 415–433, doi:10.1007/s10533-014-9977-y, 2014.
- Loisel, J., Gallego-Sala, A. V., and Yu, Z.: Global-scale pattern of peatland *Sphagnum* growth driven by photosynthetically active radiation and growing season length, *Biogeosciences*, 9, 2737–2746, doi:10.5194/bg-9-2737-2012, 2012.
- Lund, M., Christensen, T. R., Lindroth, A., and Schubert, P.: Effects of drought conditions on the carbon dioxide dynamics in a temperate peatland, *Environ. Res. Lett.*, 7, 045704, doi:10.1088/1748-9326/7/4/045704, 2012.
- Lundin, E. J., Giesler, R., Persson, A., Thompson, M. S., and Karlsson, J.: Integrating carbon emissions from lakes and streams in a subarctic catchment, *J. Geophys. Res.-Biogeo.*, 118, 1200–1207, 10.1002/jgrg.20092, 2013.
- McVeigh, P., Sottocornola, M., Foley, N., Leahy, P., and Kiely, G.: Meteorological and functional response partitioning to explain interannual variability of CO₂ exchange at an Irish Atlantic blanket bog, *Agr. Forest Meteorol.*, 194, 8–19, doi:10.1016/j.agrformet.2014.01.017, 2014.
- Moldrup, P., Olesen, T., Yamaguchi, T., Schjonning, P., and Rolston, D. E.: Modeling diffusion and reaction in soils: IX. The Buckingham-Burdine-Campbell equation for gas diffusivity in undisturbed soil, *Soil Science*, 164, 542–551, doi:10.1097/00010694-199908000-00002, 1999.
- Moyano, F. E., Manzoni, S., and Chenu, C.: Responses of soil heterotrophic respiration to moisture availability: An exploration of processes and models, *Soil Biol. Biochem.*, 59, 72–85, doi:10.1016/j.soilbio.2013.01.002, 2013.
- Nilsson, M., Sagerfors, J., Buffam, I., Laudon, H., Eriksson, T., Grelle, A., Klemedtsson, L., Weslien, P., and Lindroth, A.: Contemporary carbon accumulation in a boreal oligotrophic minerogenic mire – a significant sink after accounting for all C-fluxes, *Glob. Change Biol.*, 14, 2317–2332, doi:10.1111/j.1365-2486.2008.01654.x, 2008.
- Olefeldt, D., Roulet, N., Giesler, R., and Persson, A.: Total waterborne carbon export and DOC composition from ten nested subarctic peatland catchments – importance of peatland cover, groundwater influence, and inter-annual variability of precipitation patterns, *Hydrol. Process.*, 27, 2280–2294, doi:10.1002/hyp.9358, 2013.
- Peichl, M., Oquist, M., Lofvenius, M. O., Ilstedt, U., Sagerfors, J., Grelle, A., Lindroth, A., and Nilsson, M. B.: A 12-year record reveals pre-growing season temperature and water table level threshold effects on the net carbon dioxide exchange in a boreal fen, *Environ. Res. Lett.*, 9, 055006, doi:10.1088/1748-9326/9/5/055006, 2014.
- Reichstein, M., Falge, E., Baldocchi, D., Papale, D., Aubinet, M., Berbigier, P., Bernhofer, C., Buchmann, N., Gilmanov, T., Granier, A., Grunwald, T., Havrankova, K., Ilvesniemi, H., Janous, D., Knohl, A., Laurila, T., Lohila, A., Loustau, D., Matteucci, G., Meyers, T., Miglietta, F., Ourcival, J. M., Pumpanen, J., Rambal, S., Rotenberg, E., Sanz, M., Tenhunen, J., Seufert, G., Vaccari, F., Vesala, T., Yakir, D., and Valentini, R.: On the separation of net ecosystem exchange into assimilation and ecosystem respiration: Review and improved algorithm, *Glob. Change Biol.*, 11, 1424–1439, doi:10.1111/j.1365-2486.2005.001002.x, 2005.
- Renou-Wilson, F., Barry, C., Müller, C., and Wilson, D.: The impacts of drainage, nutrient status and management practice on the full carbon balance of grasslands on organic soils in a maritime temperate zone, *Biogeosciences*, 11, 4361–4379, doi:10.5194/bg-11-4361-2014, 2014.

- Roulet, N. T., Lafleur, P. M., Richard, P. J. H., Moore, T. R., Humphreys, E. R., and Bubier, J.: Contemporary carbon balance and late Holocene carbon accumulation in a northern peatland, *Glob. Change Biol.*, 13, 397–411, doi:10.1111/j.1365-2486.2006.01292.x, 2007.
- Ruehr, N. K., Offermann, C. A., Gessler, A., Winkler, J. B., Ferrio, J. P., Buchmann, N., and Barnard, R. L.: Drought effects on allocation of recent carbon: From beech leaves to soil CO₂ efflux, *New Phytol.*, 184, 950–961, doi:10.1111/j.1469-8137.2009.03044.x, 2009.
- Ryan, M. G. and Law, B. E.: Interpreting, measuring, and modeling soil respiration, *Biogeochemistry*, 73, 3–27, doi:10.1007/s10533-004-5167-7, 2005.
- Sottocornola, M. and Kiely, G.: Hydro-meteorological controls on the CO₂ exchange variation in an Irish blanket bog, *Agr. Forest Meteorol.*, 150, 287–297, doi:10.1016/j.agrformet.2009.11.013, 2010.
- Tang, J. W., Misson, L., Gershenson, A., Cheng, W. X., and Goldstein, A. H.: Continuous measurements of soil respiration with and without roots in a ponderosa pine plantation in the Sierra Nevada mountains, *Agr. Forest Meteorol.*, 132, 212–227, doi:10.1016/j.agrformet.2005.07.011, 2005.
- UK Meteorological Office: Met office integrated data archive system (MIDAS) land and marine surface stations data (1853–current), in: NCAS British Atmospheric Data Centre, available at: http://badc.nerc.ac.uk/view/badc.nerc.ac.uk__ATOM__dataent_ukmo-midas, (last access: January 2014), 2013.
- Updegraff, K., Bridgman, S. D., Pastor, J., Weishampel, P., and Harth, C.: Response of CO₂ and CH₄ emissions from peatlands to warming and water table manipulation, *Ecol. Appl.*, 11, 311–326, doi:10.2307/3060891, 2001.
- van der Molen, M. K., Dolman, A. J., Ciais, P., Eglin, T., Geron, N., Law, B. E., Meir, P., Peters, W., Phillips, O. L., Reichstein, M., Chen, T., Dekker, S. C., Doubkova, M., Friedl, M. A., Jung, M., van den Hurk, B. J. J. M., de Jeu, R. A. M., Kruijft, B., Ohta, T., Rebel, K. T., Plummer, S., Seneviratne, S. I., Sitch, S., Teuling, A. J., van der Werf, G. R., and Wang, G.: Drought and ecosystem carbon cycling, *Agr. Forest Meteorol.*, 151, 765–773, doi:10.1016/j.agrformet.2011.01.018, 2011.
- Wang, Y., Hao, Y., Cui, X. Y., Zhao, H., Xu, C., Zhou, X., and Xu, Z.: Responses of soil respiration and its components to drought stress, *J. Soil. Sediment.*, 14, 99–109, doi:10.1007/s11368-013-0799-7, 2014.
- Weltzin, J. F., Pastor, J., Harth, C., Bridgman, S. D., Updegraff, K., and Chapin, C. T.: Response of bog and fen plant communities to warming and water-table manipulations, *Ecology*, 81, 3464–3478, doi:10.1890/0012-9658(2000)081[3464:robafp]2.0.co;2, 2000.
- Worrall, F., Burt, T., Adamson, J., Reed, M., Warburton, J., Armstrong, A., and Evans, M.: Predicting the future carbon budget of an upland peat catchment, *Climatic Change*, 85, 139–158, doi:10.1007/s10584-007-9300-1, 2007.
- Zscheischler, J., Mahecha, M. D., von Buttlar, J., Harmeling, S., Jung, M., Rammig, A., Randerson, J. T., Schoelkopf, B., Seneviratne, S. I., Tomelleri, E., Zaehle, S., and Reichstein, M.: A few extreme events dominate global interannual variability in gross primary production, *Environ. Res. Lett.*, 9, 035001, doi:10.1088/1748-9326/9/3/035001, 2014a.
- Zscheischler, J., Reichstein, M., Harmeling, S., Rammig, A., Tomelleri, E., and Mahecha, M. D.: Extreme events in gross primary production: a characterization across continents, *Biogeosciences*, 11, 2909–2924, doi:10.5194/bg-11-2909-2014, 2014b.