

Article (refereed) - postprint

Lange, Markus; Eisenhauer, Nico; Sierra, Carlos A.; Bessler, Holger; Engels, Christoph; Griffiths, Robert I.; Mellado-Vázquez, Perla G.; Malik, Ashish A.; Roy, Jacques; Scheu, Stefan; Steinbeiss, Sibylle; Thomson, Bruce C.; Trumbore, Susan E.; Gleixner, Gerd. 2015. **Plant diversity increases soil microbial activity and soil carbon storage.** *Nature Communications*, 6, 6707. 8, pp. <https://doi.org/10.1038/ncomms7707>

© 2015 Macmillan Publishers Limited.

This version available <http://nora.nerc.ac.uk/id/eprint/511229/>
NERC has developed NORA to enable users to access research outputs wholly or partially funded by NERC. Copyright and other rights for material on this site are retained by the rights owners. Users should read the terms and conditions of use of this material at <http://nora.nerc.ac.uk/policies.html#access>

This document is the author's final manuscript version of the journal article following the peer review process. Some differences between this and the publisher's version may remain. You are advised to consult the publisher's version if you wish to cite from this article.

<https://www.nature.com/>

Contact CEH NORA team at
noraceh@ceh.ac.uk

Plant diversity increases soil microbial activity and at the same time soil carbon storage

Markus Lange¹, Nico Eisenhauer^{2,3,4}, Carlos A. Sierra¹, Holger Bessler⁵, Christoph Engels⁵, Robert I. Griffiths⁶, Perla G. Mellado-Vázquez¹, Ashish Malik¹, Jacques Roy⁷, Stefan Scheu⁸, Sibylle Steinbeiss¹, Bruce C. Thomson⁶, Susan E. Trumbore¹, Gerd Gleixner¹

¹ Max Planck Institute for Biogeochemistry, POB 100164, 07701 Jena, Germany

² Friedrich Schiller University Jena, Institute of Ecology, Dornburger Str. 159, 07743 Jena, Germany

³ German Centre for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig, Deutscher Platz 5e, 04103 Leipzig, Germany

⁴ Institute for Biology, University of Leipzig, Johannisallee 21, 04103 Leipzig, Germany

⁵ Department of Plant Nutrition, Faculty of Agriculture and Horticulture, Humboldt-Universität zu Berlin, Unter den Linden 6, 10099 Berlin, Germany

⁶ Centre for Ecology and Hydrology, Maclean Building, Benson Lane, Crowmarsh Gifford, Wallingford, OX10 8BB, UK

⁷ Centre National de la Recherche Scientifique, Ecotron - UPS 3248, Campus Baillarguet, 34980, Montferrier-sur-Lez, France

⁸ J.F. Blumenbach Institute of Zoology and Anthropology, Georg August University Göttingen, Berliner Str. 28, 37073 Göttingen, Germany

Corresponding authors: M.L. (email: mlange@bgc-jena.mpg.de) or G.G. (email: ggleix@bgc-jena.mpg.de)

Abstract:

Plant diversity strongly influences ecosystem functions and services, such as soil carbon storage. However, the mechanisms underlying the positive plant diversity effects on soil carbon storage are poorly understood. We explored this relationship using long-term data from a grassland biodiversity experiment (The Jena Experiment) and radiocarbon (^{14}C) modelling. Here we show that higher plant diversity increases rhizosphere carbon inputs into the microbial community resulting in both increased microbial activity and carbon storage. Increases in soil carbon were related to the enhanced accumulation of recently fixed carbon in high diversity plots, while plant diversity had less pronounced effects on the decomposition rate of existing carbon. The present study shows that elevated carbon storage at high plant diversity is a direct function of the soil microbial community, indicating that the increase in carbon storage is mainly limited by the integration of new carbon into soil and less by the decomposition of existing soil carbon.

Introduction

Climate and land use change can cause alterations in biodiversity^{1,2} with significant consequences for ecosystem functions and services³. Quantifying the impact of biodiversity loss on ecosystem services requires long-term investigations in replicated experimental systems⁴⁻⁶. This is especially true for soil carbon storage as changes in stocks are slow⁷, the spatial heterogeneity is large⁸ and the involved processes are complex^{9,10}. Soils store the vast majority of terrestrial carbon¹¹, and increased soil carbon storage may counteract the increase in atmospheric carbon dioxide. Traditional explanatory theories focus on the recalcitrance of less reactive compounds and physical protection as controls of soil carbon stocks, while more recent theories highlight the

importance of soil microorganisms for the persistence of soil organic matter. With respect to microbial regulation, emerging theories focus on either: spatial factors which constrain the accessibility of substrates to decomposer organisms; or microbial carbon *re-cycling* and the consequent microbial production of soil organic matter through microbially derived products^{9,10,12-15}.

In general, the amount of carbon stored in soil represents the balance between plant shoot and root litter production, root exudates and their microbial decomposition¹⁶. The observed increase in carbon storage with plant diversity¹⁷ therefore either reflects higher primary production^{18,19} or longer persistence of plant-derived organic materials due to slower decomposition^{10,16}. Increased plant residue inputs provide also more substrate for soil microorganisms, resulting in a more active and more abundant microbial community²⁰. However, different mechanisms are proposed, how the microbial community affects soil carbon. Higher amounts of plant residue inputs might either accelerate decomposition and reduce carbon storage through decreased metabolic efficiency (microbial carbon use efficiency²¹) or enhance decomposition of existing soil carbon (positive priming effect²²). Alternatively or in parallel, higher amounts of plant residue inputs could enhance carbon storage through increased microbial necromass accumulation over time²³. Therefore although soil carbon storage increased with plant diversity in a number of long-term grassland experiments^{24,25}, the underlying mechanisms are yet to be understood.

Here, we analysed comprehensive data collected over nine years from The Jena Experiment, a grassland biodiversity experiment manipulating plant species and plant functional group richness²⁶. Differences in plant carbon input between treatments likely originate from differences in root exudates and dead roots, as the plots were mown twice a year and aboveground plant biomass was

removed as is typical for Central European hay meadows. We applied path analysis to identify the relationships between explanatory variables affecting soil carbon storage. Furthermore, to determine whether soil carbon accumulation resulted from increased root-derived carbon inputs or reduced decomposition, we modelled the carbon flow in the system using observations of soil carbon stocks and radiocarbon (^{14}C) in soil organic matter (SOM) as a double constraint²⁷. The combined results of path analysis and soil carbon modelling suggest a unique cause for enhanced carbon storage with plant diversity in our experimental plots. Although both approaches are based on data from the top, biologically most active soil layer (0–5 cm), we were able to generalise the proposed mechanism of plant diversity effects on soil carbon storage to deeper soil layers (0–30 cm).

Results

Effects of plant diversity on soil organic carbon changes

In 2011, nine years after establishment of The Jena Experiment, soil carbon concentration in the topsoil layer (0–5 cm, averaged across all diversity treatments) had increased by 27% to $4.74 \pm 0.28 \text{ g kg}^{-1}$ (mean \pm s.e.m.). An increase in SOM was expected because the experiment had been established on a ploughed arable field turned into permanent grassland. The increase in soil carbon concentration was highly correlated with sown plant species richness ($P < 0.001$, Fig. 1). The presence of legumes negatively affected soil carbon concentration ($P = 0.040$, Fig. 1), while other plant functional groups or their richness did not significantly influence carbon concentration (Table 1; Supplementary Table 1).

Mechanisms of soil organic carbon storage

In this strictly confirmatory approach²⁸, we tested different hypotheses (Table 2; Fig. 2), resulting in set of different path models (Supplementary Fig. 1), assessing if the models deviate or not from the observed data structure. All individual models were tested for their adequacy in reflecting the observed data. Path analyses revealed that plant species richness had direct positive effects on root carbon inputs as well as metabolic activity of soil microorganisms, measured as basal respiration (Fig. 3a and b; Methods). There was no direct relationship between plant species richness or the presence of legumes and soil carbon storage, showing that fine root carbon inputs and microbial activity are the main drivers explaining the positive effect of plant diversity on soil carbon storage. Low root biomass on plots with legumes²⁹ may have caused the negative effect of legumes on microbial activity²⁰ via reduced carbon inputs. Importantly, the increased soil carbon storage was directly only related to higher microbial activity. This potential chain of causality was the best model that adequately explained the data (Supplementary Table 2; Supplementary Notes 1–4), also when soil carbon sequestration was set up in a path model as a time series (Supplementary Fig. 2), which increased the likelihood of the model.

Dynamics of soil organic carbon

Radiocarbon based modelling of SOM storage (Methods), suggested that the increased carbon storage in plots with higher plant species richness was mainly due to the accumulation of ‘new’ carbon (Fig. 4a). Furthermore, the sequestration of ‘new’ carbon and the loss of existing ‘old’ carbon (positive priming) were positively correlated ($R^2 = 0.863$, $P = 0.007$, Fig. 4b) and depended on plant species richness. High diversity plots (richness levels 8, 16 and 60) accumulated 21.8% more carbon compared to low diversity plots (richness levels 1, 2 and 4). At the same time, high diversity plots primed on average only 9.6% of ‘old’ carbon. Interestingly, carbon losses were mainly observed in the fast cycling carbon (Fig. 4c), whereas in the more sustainable slow cycling

carbon these losses were reduced at higher plant diversity, and more ‘new’ carbon was stored in the slow cycling carbon (Fig. 4d). An explanation for these opposite patterns is the transfer or processing rate (conversion efficiency, Methods) of carbon from fast to slow cycling carbon, which is – in accordance with our path model – proportional to soil microbial biomass and activity (Fig. 3c). Consequently, the slow cycling carbon (turnover time = 47.7 years, Supplementary Table 3) dominates the longer-term dynamics of the total stock, whereas priming and other effects of the fast cycling carbon (turnover time = 4.0 years) become less important over longer time scales.

Discussion

The results reported in this study indicate that storage of soil carbon is governed by the metabolic activity of soil microbes, which is mediated by plant diversity via higher root inputs and other yet unidentified mechanisms, as suggested by the direct relation in the path model (Fig. 3a). A number of processes influence microbial activity: First, a denser vegetation in highly diverse plant communities reduces evaporation from the topsoil, which in turn promotes higher soil microbial activity and growth³⁰. Second, soil carbon storage is linked to root inputs^{31,32} including root exudation, which is known to change the activity and composition of the microbial community³³. Indeed, carbon uptake in rhizospheric microorganisms at high plant diversity was increased compared to low plant diversity (Fig. 5) as indicated by a complementary ¹³CO₂ labelling experiment in the Ecotron facility in Montpellier (Methods). This showed higher microbial label uptake was most likely attributed to higher levels of root exudation with increasing plant diversity, as the soil microbial community was sampled immediately after a three week label application and the label enrichment in roots and root sugars was not affected by diversity (data not shown). Third, increasing diversity of soil microbial communities increases microbial respiration³⁴. In fact, bacterial and fungal diversity increased with higher plant species richness in The Jena Experiment

(Table 1; Fig. 6). And finally, a shift in the metabolic activity of soil microorganisms towards anabolic activity with plant diversity resulting in increased necromass accumulation over time²³, indicated by slightly but significantly more microbial carbon per unit root carbon in high diversity plots ($R^2 = 0.06$, $P = 0.025$).

The missing direct path from legumes to microbial activity indicates minor importance of improvement of soil nitrogen status (measured as changes in ^{15}N of bulk soil, Table 1) for soil microbial activity, and thus for carbon storage. This was partly unexpected as legumes have been reported to increase the availability of nitrogen in soils^{35,36}, and are therefore suggested to increase plant biomass production (above and below ground)²⁴, resulting in increased metabolic activity of soil microorganisms³⁷. However, the presence of legumes did not increase root biomass production and the metabolic activity of soil microorganisms in The Jena Experiment^{20,38}. Although the increase in total nitrogen content was highly correlated with the increase in soil organic carbon ($R^2 = 0.79$, $P < 0.001$), the lower root biomass on legume plots²⁹ likely caused the negative legume effect on soil carbon storage. However, the legume effect might be most pronounced in the topsoil as it was not significant when considering the top 30 cm of the soil (Table 1). In contrast, the positive plant diversity effect on soil carbon storage (Table 1), root biomass³⁹, and soil microbial biomass⁴⁰ was also found for deeper soil layers in The Jena Experiment. Despite the consistency of the plant diversity effect, it may be strongest in the topsoil and decrease with soil depth^{25,39}, which may be due to the fact that root carbon inputs^{24,39} and microbial activity⁴¹ decrease with soil depth. Therefore, the mechanism proposed for the topsoil is very likely to also be relevant for deeper soil layers, but the decreased biological activity at deeper soil layers has to be taken into account. Moreover, other mechanisms like spatial separation from decomposers^{9,12} might become more important with soil depth.

The positive plant diversity effect on carbon storage is driven by root carbon inputs, but mediated by the soil microbial community. Longer term, we expect this link to become even more important through both an increase of the positive plant diversity effect on plant (root) biomass production^{6,39} and an increased microbial contribution to soil carbon storage by elevated soil microbial activity²⁰. Both mechanisms in concert might be responsible for an increase of the plant diversity effect on carbon storage, as indicated by the time series path model (Supplementary Fig. 2).

In summary, higher levels of carbon inputs to the soil and more favourable microclimatic conditions linked to more diverse plant communities result in more active, more abundant and more diverse soil microbial communities. Microbial activity increases the turnover rates of root litter and exudates as indicated by increased microbial respiration. Thus, microbial-products associated with increased microbial respiration such as microbial necromass end up in slow-cycling SOM pools in the form of reduced organic material^{10,14}. The importance of microbial residues for soil carbon storage is suggested to even increase over time²³. Thus, in line with the recent debate⁴²⁻⁴⁴, our findings challenge previous views underestimating or even negating the influence of size and activity of soil microorganisms on soil carbon storage.

Our data from long-term field experiments emphasize the importance of plant diversity and its effects on soil microorganisms for ecosystem carbon storage. Although we cannot rule out that the proposed mechanisms are particularly pronounced at the field site of The Jena Experiment, the underlying relationships between plant diversity and soil carbon storage^{17,24}, soil microbial biomass and activity^{37,45,46}, and root biomass^{24,47} have been reported in several independent experimental settings providing some evidence that the proposed links are of general significance. This leads us to reconsider the role of soil microorganisms as sources rather than sinks for slow cycling organic matter. Thus, our results suggest that the activity and composition of soil microbial communities

can serve as a proxy for carbon transfer into sustainable slow cycling forms of soil carbon, and that plant diversity and associated soil microbial communities can significantly contribute to sequestration of atmospheric carbon dioxide.

Methods

Study site of The Jena Experiment

Our study was carried out as a part of The Jena Experiment, a large-scale grassland diversity experiment²⁶. The experiment is located on the floodplain of the Saale River near the city of Jena (Thuringia, Germany; 50°57'N, 11°35'E). The soil of the field site is classified as Eutric Fluvisol⁴⁸. In spring 2002, eighty-two experimental grassland plots of 20 × 20 m were established. One monoculture plot was abandoned due to poor establishment, making the total number of plots 81 instead of the 82 originally planned. Plots are arranged in four blocks to account for changes in abiotic factors with distance from the river. Soil texture in the upper soil ranges from sandy loam to silty clay with increasing distance to the river. Sand content declines from 40% near the river to 7% at the furthest plot, while silt content increases from 44% to 69%, respectively. The clay content ranges from 16–24%, but is not related to distance from the river. For the 40 years prior to establishing the experiment, the field site was a cultivated field with mineral fertilizer input. The initial physico-chemical properties, such as pH (7.1–8.4), soil organic carbon (5–33 g C kg⁻¹), and soil nitrogen concentrations (1.0–2.7 g N kg⁻¹), varied across the field site and are considered in the block design of The Jena Experiment²⁶.

The plots differed in both the number of plant species (1, 2, 4, 8, 16, 60 species), randomly chosen from a common pool of 60 plant species, and the number of plant functional groups, *i.e.*

grasses, legumes, small herbs, and tall herbs, which were classified into four functional groups based on morphological, phenological and physiological traits; for details see Roscher *et al.*²⁶. Experimental communities are weeded manually twice a year to maintain the levels of diversity and mown twice yearly in June and September as is typical for hay meadows in Central Europe.

Sampling and laboratory analyses

Data included in this paper originated from soil samples taken at a depth of 0–5 cm, where biological activity is at a maximum. Fine root biomass in some cases had to be extrapolated to obtain estimates for the topsoil (0–5 cm), as not all sampling campaigns were set up for stratified sampling (see below).

Soil organic carbon

Soil sampling was performed before sowing in April 2002 and was repeated in April in the years 2004, 2006, 2008 and 2011, as described in more detail in Steinbeiss *et al.*²⁵. In short, three soil samples were taken per plot (4.8 cm in diameter, 0–30 cm deep) using a split tube sampler (Eijkelkamp Agrisearch Equipment, Giesbeek, the Netherlands). In the year 2002, a total of five soil samples were taken per plot. The soil was dried, sieved (2 mm mesh) and milled (4 min., frequency 30 s⁻¹). Total carbon of ground samples was determined by an elemental analyzer after combustion at 1150 °C (Elementaranalysator vario Max CN, Elementar Analysensysteme GmbH, Hanau, Germany). Inorganic carbon concentration was measured by elemental analysis after removing organic carbon for 16 h at 450 °C in a muffle furnace by oxidation. Organic carbon concentration was calculated from the difference between total and inorganic carbon concentrations. Carbon stocks were calculated as the product of bulk soil density and its

concentration of carbon⁸. In order to exclude physical aspects of bulk density (*e.g.* inter-annual differences caused by degree of winter freezing) statistical analyses were done with carbon concentrations [g kg^{-1}] and stocks were estimated using the mean of bulk soil density across all years.

Periodic sampling of other data used in this study did not always occur over the same time interval (annual: microbial activity, biannual: soil and root sampling). Thus, the mean of the other variables (basal respiration fine root carbon input) was calculated referring to each biannual soil carbon sampling campaigns.

Fine root standing biomass

Root standing biomass was sampled in 2003, 2004, 2006, and 2008 and separated in fine and coarse roots based on root diameter (fine roots ≤ 2 mm; coarse roots > 2 mm). Roots were calculated as grams dry mass per square meter. In the first three sampling campaigns fine roots were determined to a depth of 30 cm. In 2008, stratified sampling was carried out with a finer resolution of 5 cm increments. Based on the fine root ratio of the topsoil (0–5 cm) and entire sampling depth (0–30 cm) in 2008, fine root standing biomass in the topsoil was estimated for each prior sampling year. The carbon content of fine roots was determined using root material from ingrowth cores sampled in 2004²⁹. Fine root biomass comprised on average 80.9% (± 16.8 s.d.) of the total root biomass and was highly correlated to the total root biomass ($R^2 = 0.66$, $P < 0.001$).

Basal respiration

Basal respiration was measured annually from 2003 to 2011, except in 2005. Five soil samples per plot were taken to a depth of 5 cm, pooled and stored at 5°C. The samples were homogenized, sieved (2 mm) to remove larger roots, animals, and stones, and adjusted to a

gravimetric soil water content of 25%. Basal respiration was measured using an O₂ microcompensation apparatus⁴⁹. The microbial respiratory response was measured at hourly intervals for 24 h at 22°C. Basal respiration ($\mu\text{L O}_2 \text{ h}^{-1} \text{ g soil dry mass}^{-1}$) was determined without addition of substrate and measured as the mean of the O₂ consumption rates 14 to 24 hours after the start of the measurements (for details see Eisenhauer *et al.*²⁰).

Radiocarbon and soil nitrogen isotope ratios

Radiocarbon concentrations (¹⁴C) were measured in soil samples of the years 2002 and 2011 as described in detail by Tefs and Gleixner⁵⁰. Briefly, after decalcification with HCl and graphite production, the ¹⁴C content of the samples was measured with accelerator mass spectroscopy (3 MV Tandetron 4130 AMS, High Voltage Engineering Europa, Amersfoort, The Netherlands). Soil nitrogen isotope ratios ($\delta^{15}\text{N}$) were measured from 50 mg of dried ground soil with an elemental analyser coupled to an isotope-ratio mass spectrometer (EA-IRMS; Delta C prototype IRMS, Finnigan MAT).

Root exudation

Within the experimental setup of The Jena Experiment it is not possible to determine effects of plant diversity on the amount of root exudation. Therefore we took advantage of the Ecotron facility in Montpellier (France) to overcome this limitation. Twelve monoliths (2.0 m in depth and 1.6 m in diameter) were excavated from the field site of The Jena Experiment and installed in the macrocosms in the Ecotron in spring 2012. These twelve monoliths reflected the diversity gradient of The Jena Experiment on a smaller scale⁵¹. Monoliths with high (16 plant species) and low plant diversity (4 plant species) were labelled with ¹³CO₂ for four weeks. We analysed root exudation and the turnover of the exudates by root associated microbes using the phospholipid fatty acids

(PLFA) method^{52,53}. Particular specific monounsaturated fatty acids like PLFA 16:1 ω 5, 16:1 ω 7, 17:1 ω 8 and 18:1 ω 7 were shown to be highly enriched by root-derived organic carbon in label experiments^{54,55}. We measured the concentration and enrichment in $\delta^{13}\text{C}$ of these PLFA markers in the Ecotron to assess the effect of plant diversity on root exudation and the incorporation of the exudates in soil microbes.

Genetic diversity of soil bacteria and fungi

The genetic diversity of soil microbial communities was determined using terminal-restriction fragment length polymorphism (TRFLP). Briefly, total nucleic acids were extracted from 0.25 g fresh weight of soil using the MoBio Powersoil 96 kit. Bacterial 16S rRNA genes were amplified using forward primer 63F (5'-CAGGCCTAACACATGCAAGTC-3')⁵⁶, fluorescently labelled with 6-FAM at the 5' end, and reverse primer 519r (5'-GTATTACCGCGGCTGCTG-3')⁵⁷. For fungi we used the primer pair ITS1F (5'-CTTGGTCATTTAGAGGAAGTAA-3')⁵⁸ and ITS4 (5'-TCCTCCGCTTATTGATATGC-3')⁵⁹, spanning the ITS1 and ITS2 regions. Amplicons were purified by gel filtration using Sephadex G50 (Sigma-Aldrich, Gillingham, UK) prior to digestion with restriction endonucleases Msp 1 (CCGG) and Taq 1 (TCGA) (Promega, Southampton, UK) for bacteria and fungi respectively. A 3730 DNA analyser (Applied Biosystems, CA, USA) was used for fragment analysis, prior to binning of individual terminal restriction fragments (T-RFs) using Genemarker software (SoftGenetics, PA, USA). The relative abundance of individual T-RFs was calculated by dividing the intensity of each fragment by the total intensity of all fragments prior to calculation of Simpson's diversity index.

Statistical analyses of main effects of plant diversity

The Jena Experiment is based on a factorial design with different combinations of sown plant species richness (PSR) and number of functional groups (FG). To assess for the design variables, analyses of variance (ANOVA) with sequential sum of squares (type I SS) were applied using R⁶⁰. The ‘block effect’, mainly accounting for differences in soil texture among the blocks²⁶, was included as random factor and fitted first, followed by PSR and FG. Finally, the presence of each plant functional group (legumes, grasses, small herbs and tall herbs) was included into the model in a series of alternative models.

Path analysis

We applied path analyses (maximum likelihood)²⁸ to explore the potentially causal relationships between plant diversity and community composition and soil carbon storage using the lavaan package in R⁶¹. Depending on the results of the ANOVA approach, all variables with significant effect on changes of soil carbon content were included as exogenous (predictor) variables. As variables potentially mediating plant community effects we included fine root carbon as measure of carbon input^{31,32} and basal respiration as measure of microbial metabolic activity^{20,34,37}. We tested four different ways how plant diversity can affect carbon storage: due to carbon input from roots, due to microbial activity, due to both root inputs as well as microbial activity, or directly via plant diversity, indicating that other mechanisms than root carbon input and microbial activity underlie the plant diversity effect. Each of the four storage ways comprised a set of models (Supplementary Fig. 1), differing in the relations from plant diversity to basal respiration and fine root carbon to basal respiration. For all models, the adequacy of the hypothesized structural relationships and the data was verified using Chi-square tests and the residual mean square error of approximation (RMSEA). To test if carbon storage is a function of the microbial activity or *vice versa*, we tested all models twice, first with the path from basal

respiration towards soil carbon content, as ‘sink models’ (as described above, Supplementary Note 1) and second with a path from soil carbon content towards basal respiration as ‘source models’ (Supplementary Note 2). In the final step we identified the minimal adequate (most parsimonious) model based on the Bayes Information Criterion (BIC). Therefore, all adequate models were compared (Supplementary Note 3), including non-significant pathways and when only significant pathways are contained.

Eisenhauer *et al.*²⁰ and Ravenek *et al.*³⁹ showed that plant species richness was not closely linked to basal respiration and root standing biomass in the first years of the experiment; therefore, we did not use data from the first two years in the present analyses. Missing values were replaced by the mean. All variables were standardized to [0, 1]. The total sample size was $N = 81$.

Three-pool ^{14}C model with serial structure

To describe soil organic carbon dynamics in The Jena Experiment, we implemented a dynamic model of the form

$$\begin{aligned}\frac{dC_f}{dt} &= I - k_1 C_f \\ \frac{dC_s}{dt} &= \alpha_{2,1} k_1 C_f - k_2 C_s \\ \frac{dC_p}{dt} &= \alpha_{3,2} k_2 C_s - k_3 C_p\end{aligned}\tag{1}$$

where C_f , C_s , and C_p represent carbon storage in the fast, slow, and passive pool, respectively. I represents the total amount of carbon inputs to the first 5 cm of the soil, and k_1 , k_2 , k_3 represent first-order decomposition rates for the fast, slow, and passive pools, respectively. The transfer coefficients $\alpha_{2,1}$ and $\alpha_{3,2}$ represent the proportion of the decomposed carbon from each pool that is transferred to a slower cycling pool^{62,63}.

One particular aspect of this model, suggested from the path analysis, is that we represent the effects of the microbial community on decomposition as a control on the transfer rate. That is, we assume

$$\alpha_{2,1}k_1 = f \cdot MB, \quad (2)$$

where MB is the microbial biomass and f a scaling factor. Notice that we do not represent the microbial biomass as an additional pool in our dynamical system because it leads to unrealistic nonlinear behaviour⁶⁴ not supported by our own observations.

To account for radiocarbon dynamics, we implemented a model similar to equation (1) representing each pool as a fraction of radiocarbon with respect to an internationally-agreed standard^{27,63}. The model has similar form as equation (1) with an additional term accounting for radioactive decay. The model was implemented in the R environment for computing using the SoilR package⁶⁵. The passive pool is mainly needed to account for the low ^{14}C values in the bulk soil organic matter. As described by other authors⁶⁶ such low ^{14}C values are common for arable fields in Europe.

Based on our findings in the path analysis, we implemented a measure of soil microbial community activity (basal respiration) as a proxy for the transfer between the fast and slow carbon pool. Unfortunately, no data on basal respiration were available for arable control plots, which would have been needed for the spin up (please see section “Spin up and parameter estimation for the spin up”). Thus, we took advantage of the microbial biomass (based on the chloroform fumigation extraction (CFE) analysis⁶⁷) instead. Basal respiration and microbial biomass were highly correlated (Fig. 3c). For each level of plant species richness we applied a model with the level specific parameter estimation (Table 2; Fig. 3b). However, we could neither evaluate the real

carbon input (carbon of standing fine root biomass and root exudates, root tips *etc.*) nor the extent of the transfer coefficient. Thus, we applied a set of different models for each level of plant diversity, modified in the carbon input and transfer coefficient. Implementing the transfer coefficient between the fast and the slow pool depending on soil microbial biomass resulted in good agreement between model predictions and measured data (Table 3; Supplementary Table 3; and Supplementary Fig. 3).

Calculation of new carbon accumulated and loss of old carbon

Based on the mass balance and the isotopic mass balance for the isotopic mass ratio (equation 3) we calculated the yearly (a) amount of new carbon sequestered (C_{new}) and loss of the old carbon ($C_{\text{loss old}}$).

$$\begin{aligned}\Delta C &= C_{\text{new}} - C_{\text{loss old}} = C_a - C_{a+1} \\ \delta_{\Delta C} \Delta C &= \delta_{\text{new}} C_{\text{new}} - \delta_{\text{loss old}} C_{\text{loss old}} = \delta_{a_n} C_{a_n} - \delta_{a_n+1} C_{a_n+1}\end{aligned}\tag{3}$$

Spin up and parameter estimation for the spin up

The objective of the spin up simulations is to match the carbon stocks and ^{14}C data for the pre-experimental conditions. We let the spin up simulations run for 10.000 years, to assure the modelled pre-experimental conditions are in equilibrium. As there are no comparable data for fine root biomass on the arable control plots, we used the mean fine root carbon of all experimental plots from of all fine root sampling campaigns (mean = 149.77 [g m⁻²], SD = 98.05). Due to ploughing of the soil before the implementation of the experimental plots, a homogeneous

distribution of fine root carbon can be assumed as 25 g m^{-2} in each soil segment of 5 cm. In contrast, after establishment of experimental plots, the fine-root biomass distribution was similar to meadows with the main amount of roots in the topsoil, strongly decreasing towards deeper depths. Thus, a fine root carbon input of 67.04 g m^{-2} (on average) was assumed on experimental plots. The ratio of the assumed fine root inputs between experimental plots and arable land is similar to the ratio of carbon input on meadows and arable lands reported in Schulze *et al.*⁶⁸ Furthermore, we applied a fine root turnover of 1.7 years on experimental grassland plots⁶⁹. To meet the measured and the modelled data, a fine root turnover of 1.2 years for arable lands was applied. In the spin up, we set the input factor to 1.7 (Supplementary Table 3), the transfer coefficient to 0.1 reflecting C_{mic} of $0.22 \text{ [ng g}^{-1}\text{]}$ on the arable control plots (*cf.* Table 1). This value of the transfer coefficient is within the expected range. The resulting sizes of the three pools and their ages are shown in Supplementary Table 3.

Acknowledgments:

We thank Uta Gerighausen, Steffen Rühlow and Maike Habekost for their help during soil sampling, the RoMA and ^{14}C Analysis Facility at the MPI for Biogeochemistry provided the data on carbon and radiocarbon respectively. Markus Reichstein and Winfried Voigt gave valuable comments on earlier versions of the manuscript. Lisa Geesink created the artwork. We gratefully acknowledge Wolfgang W. Weisser, Ernst-Detlef Schulze, Christiane Roscher, Alex Weigelt and Anne Ebeling and all the people that were involved in planning, set up and maintaining of the experiment and the German Research Foundation (DFG) for financial support of FOR 456 and Gl 262-14. We thank Alex Milcu and the Ecotron team for organising coordinating and running the Ecotron experiment respectively. The Ecotron benefited from the CNRS human and technical resources and from the state 'Investissement d'Avenir' ANR-11-INBS-0001. Further support came from the joint International Max Planck Research School of the Friedrich Schiller University Jena and the Max Planck Institute for Biogeochemistry Jena, and from the German Centre for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig, funded by the German Research Foundation (FZT 118).

Author Contributions:

ML, ST and GG designed the study, ML and GG wrote the manuscript, ML and NE performed statistical analyses, ML and CS modelled soil carbon, PM, ML, GG and JR performed the Ecotron experiment, AM, BT and RIG performed TRFLP analyses, NE and SS measured microbial activity, HB and CE measured root biomass. All authors reviewed and improved the manuscript.

Supplementary Information accompanies this paper at www.nature.com/naturecommunications

Competing financial interests: The authors declare no competing financial interests.

Figure Legends:

Figure 1: The relationship between plant diversity and soil organic carbon storage. Soil carbon changes [g kg^{-1}] between 2002 and 2011 as affected by (log) plant species richness ($F_{1,73} = 41.29$, $P < 0.001$) and the presence (orange circles) and absence of legumes (blue circles) ($F_{1,73} = 4.37$, $P = 0.040$).

Figure 2: Conceptual model on the expected causal relationships between plant diversity and soil carbon storage through soil carbon inputs and soil microbial activity based on hypotheses derived from previous experiments (see Table 2 for hypotheses and references).

Figure 3: Mechanisms of soil carbon storage. a) Most parsimonious path model explaining the underlying mechanisms of the positive relationship between plant diversity and soil organic carbon storage ($\chi^2_4 = 5.35$, $P = 0.253$; RMSEA = 0.065 CI90 [0.000; 0.191]). Numbers next to the endogenous variables indicate their explained variance (R^2). Numbers next to the arrows indicate standardized path coefficients and asterisks mark their significance: < 0.05 (*), < 0.01 (**) or < 0.001 (***). b) Impact of (log) plant species richness on mean fine root carbon and mean soil microbial biomass. These mean values were used as specific parameters for SOM storage modelling (three pool model) for each level of plant species richness. c) Regression of mean metabolic activity and mean biomass of the microbial community at plant species richness (PSR) levels ($R^2 = 0.73$, $P = 0.031$). Metabolic community activity of soil microbes was measured as basal respiration and microbial community biomass was measured as microbial carbon (using chloroform fumigation analysis (CFE)). Both measures are specified per g dry soil. Error bars are defined by s.e.m.

Figure 4: Sequestration of new and loss of old soil organic carbon at The Jena Experiment.

Effect of plant species richness (PSR) on a) mean annual loss of old carbon ($C_{\text{loss old}}$) and gain of

new carbon (C_{new}). Relation of new carbon sequestered and the loss of old carbon of the b) total stock, c) fast pool and d) slow pool. Calculations are based on the pool-model and the isotopic mass balance (Methods). Error bars are given as s.e.m.

Figure 5: Enrichment of root associated microorganisms in long-term ^{13}C labelling experiment. Influence of plant species richness on the label uptake into phospholipid fatty acids (PLFA) of the root associated microbial community ($F_{1,9} = 6.99$, $P = 0.027$) studied in a three week labelling experiment under fully controlled conditions (ECOTRON facilities, Montpellier, France³⁴) Error bars are given as s.e.m.

Figure 6: Genetic diversity of the soil microbial community. Terminal-restriction fragment length polymorphism (TRFLP) separately analysed for fungal (orange circles) and bacterial communities (blue circles) as affected by plant species richness (Table 1).

Tables:

Table 1: Results of ANOVAs on the effects of the experimental variables. Impact of plant diversity (plant species richness (PSR, log transformed) and number plant functional groups (FG)) and the presence of distinct plant functional groups (legumes, grasses, small herbs, tall herbs) on a) changes in soil organic carbon (ΔC_{org} 0–5 and 0–30 cm), genetic diversity (determined using terminal-restriction fragment length polymorphism (TRFLP)) of the soil microbial community (fungi and bacteria) and changes in $\delta^{15}N$ (0–5 cm). Numbers give *F*-values, asterisks mark their significance: < 0.05 (*), < 0.01 (**) or < 0.001 (***). Detailed ANOVA results are given in Supplementary Table 1.

	df	ΔC_{org} (0–5cm)		ΔC_{org} (0–30cm)		TRFLP fungi		TRFLP bacteria		$\Delta^{15}N$ (0–5cm)	
Block	3	0.08		1.46		1.05		1.60		5.00	**
PSR	1	41.29	***	21.82	***	12.06	***	19.37	***	16.69	***
FG	1	3.69		0.72		0.36		3.36		10.79	**
Legumes	1	4.38	*	0.23		0.06		0.04		6.44	*
Grasses	1	1.65		0.16		0.01		0.81		3.76	
Small herbs	1	1.96		0.15		1.74		0.00		0.06	
Tall herbs	1	0.33		0.08		1.52		1.03		0.13	

Table 2: Potential relations between the considered variables. Hypothesised mechanisms of the individual paths in the conceptual model (Fig. 2). Arrows indicate the direction of the paths.

Path #	Pathway	Hypothesised mechanism
1	Plant species richness → Carbon inputs	Plant diversity increases root biomass or root carbon concentration ^{24,29,39}
2	Plant species richness → Microbial activity	Plant diversity increases microbial community activity ^{20,30}
3	Legumes → Carbon inputs	Plant community composition, such as the presence of legumes, influences the amount of root carbon residues ²⁴
4	Legumes → Microbial activity	The presence of legumes decreases soil microbial activity ²⁰
5	Carbon inputs → Microbial activity	Positive effects of root carbon residues on soil microbial activity ^{20,37}
6	Carbon inputs → Carbon storage	Carbon storage increases with higher inputs of root carbon residues ¹⁵ or with more plant derived recalcitrant compounds ⁷⁰
7	Microbial activity → Carbon storage	Microbial community activity influences soil carbon storage ^{14,42-44}
8	Carbon storage → Microbial activity	Higher soil organic carbon content increases soil microbial community activity ²²
9	Plant species richness → Carbon storage	Plant species richness influences carbon storage by other mechanisms than root carbon inputs and microbial activity

10	Legumes → Carbon storage	The presence of legumes influences carbon storage by other mechanisms than through root carbon inputs and microbial activity
----	--------------------------	--

Table 3: Variations of soil carbon properties and their drivers in The Jena Experiment.

Means and standard deviation (s.d.) for each level of plant species richness (PSR) in changes of soil carbon content (ΔC_{org} [g kg⁻¹], Methods) and radiocarbon ($\Delta^{14}C$ [‰]) from the pre-establishing phase (2002) to 2011. Mean root carbon inputs (C_{root} [g m⁻²], calculated as the product of fine root standing biomass and carbon concentration of fine root, Methods) and mean microbial metabolic activity (measured as basal respiration (BR) [μ L O₂ h⁻¹ g soil dry mass], Methods) were not considered throughout the first two years of the (2002 and 2003) as both measures did not respond to the established plant diversity gradient during this time (Methods²⁰). Microbial biomass (C_{mic} [mg g⁻¹ dw]) was determined in 2007 using chloroform fumigation extraction (CFE, Methods).

PSR	N plots	ΔC_{org}	ΔC_{org} s.d.	$\Delta^{14}C$	$\Delta^{14}C$ s.d.	C_{root}	C_{root} s.d.	BR	BR s.d.	C_{mic}	C_{mic} s.d.
1	15*	155.64	77.44	-10.41	19.12	63.98	49.86	2.57	0.300	0.328	0.096
2	16	162.62	121.48	-0.07	18.31	60.71	30.42	3.05	0.402	0.343	0.082
4	16	246.20	123.96	1.14	17.39	62.17	25.82	3.24	0.421	0.358	0.080
8	16	306.27	71.99	3.60	18.01	71.77	46.95	3.22	0.339	0.374	0.119
16	14	323.66	112.76	4.97	10.19	77.32	32.81	3.44	0.350	0.406	0.096
60	4	390.23	114.68	11.69	8.75	68.48	15.53	3.68	0.201	0.485	0.050

* One monoculture plot was abandoned due to poor establishment, making the total number of plots 81 instead of the 82, which were originally set up.



References


- 1 Loreau, M. *et al.* Ecology - Biodiversity and ecosystem functioning: Current knowledge and future challenges. *Science* 294, 804-808, doi:10.1126/science.1064088 (2001).
- 2 Sala, O. E. *et al.* Global biodiversity scenarios for year 2100. *Science* 287, 1770-1774 (2000).
- 3 Hooper, D. U. *et al.* A global synthesis reveals biodiversity loss as a major driver of ecosystem change. *Nature* 486, 105-U129, doi:10.1038/nature11118 (2012).
- 4 Allan, E. *et al.* A comparison of the strength of biodiversity effects across multiple functions. *Oecologia* 173, 223-237, doi:10.1007/s00442-012-2589-0 (2013).
- 5 Cardinale, B. J. *et al.* Biodiversity loss and its impact on humanity. *Nature* 486, 59-67, doi:10.1038/nature11148 (2012).
- 6 Reich, P. B. *et al.* Impacts of Biodiversity Loss Escalate Through Time as Redundancy Fades. *Science* 336, 589-592, doi:10.1126/science.1217909 (2012).
- 7 Poeplau, C. *et al.* Temporal dynamics of soil organic carbon after land-use change in the temperate zone - carbon response functions as a model approach. *Global Change Biology* 17, 2415-2427, doi:10.1111/j.1365-2486.2011.02408.x (2011).
- 8 Schrumpf, M., Schulze, E. D., Kaiser, K. & Schumacher, J. How accurately can soil organic carbon stocks and stock changes be quantified by soil inventories? *Biogeosciences* 8, 1193-1212, doi:10.5194/bg-8-1193-2011 (2011).
- 9 Gleixner, G. Soil organic matter dynamics: a biological perspective derived from the use of compound-specific isotopes studies. *Ecological Research* 28, 683-695, doi:10.1007/s11284-012-1022-9 (2013).
- 10 Schmidt, M. W. I. *et al.* Persistence of soil organic matter as an ecosystem property. *Nature* 478, 49-56, doi:10.1038/nature10386 (2011).
- 11 Jobbagy, E. G. & Jackson, R. B. The vertical distribution of soil organic carbon and its relation to climate and vegetation. *Ecological Applications* 10, 423-436, doi:10.2307/2641104 (2000).
- 12 Dungait, J. A. J., Hopkins, D. W., Gregory, A. S. & Whitmore, A. P. Soil organic matter turnover is governed by accessibility not recalcitrance. *Global Change Biology* 18, 1781-1796, doi:10.1111/j.1365-2486.2012.02665.x (2012).
- 13 Gleixner, G., Poirier, N., Bol, R. & Balesdent, J. Molecular dynamics of organic matter in a cultivated soil. *Organic Geochemistry* 33, 357-366, doi:10.1016/s0146-6380(01)00166-8 (2002).
- 14 Miltner, A., Bombach, P., Schmidt-Bruecken, B. & Kaestner, M. SOM genesis: microbial biomass as a significant source. *Biogeochemistry* 111, 41-55, doi:10.1007/s10533-011-9658-z (2012).
- 15 Don, A., Roedenbeck, C. & Gleixner, G. Unexpected control of soil carbon turnover by soil carbon concentration. *Environmental Chemistry Letters* 11, 407-413, doi:10.1007/s10311-013-0433-3 (2013).
- 16 Jastrow, J. D., Amonette, J. E. & Bailey, V. L. Mechanisms controlling soil carbon turnover and their potential application for enhancing carbon sequestration. *Clim. Change* 80, 5-23, doi:10.1007/s10584-006-9178-3 (2007).
- 17 De Deyn, G. B. *et al.* Additional carbon sequestration benefits of grassland diversity restoration. *Journal of Applied Ecology* 48, 600-608, doi:10.1111/j.1365-2664.2010.01925.x (2011).
- 18 Marquard, E. *et al.* Plant species richness and functional composition drive overyielding in a six-year grassland experiment. *Ecology* 90, 3290-3302, doi:10.1890/09-0069.1 (2009).

- 19 Tilman, D. *et al.* Diversity and productivity in a long-term grassland experiment. *Science* 294, 843-845, doi:10.1126/science.1060391 (2001).
- 20 Eisenhauer, N. *et al.* Plant diversity effects on soil microorganisms support the singular hypothesis. *Ecology* 91, 485-496, doi:10.1890/08-2338.1 (2010).
- 21 Manzoni, S., Taylor, P., Richter, A., Porporato, A. & Agren, G. I. Environmental and stoichiometric controls on microbial carbon-use efficiency in soils. *New Phytol.* 196, 79-91, doi:10.1111/j.1469-8137.2012.04225.x (2012).
- 22 Fontaine, S. *et al.* Stability of organic carbon in deep soil layers controlled by fresh carbon supply. *Nature* 450, 277-U210, doi:10.1038/nature06275 (2007).
- 23 Liang, C., Cheng, G., Wixon, D. L. & Balser, T. C. An Absorbing Markov Chain approach to understanding the microbial role in soil carbon stabilization. *Biogeochemistry* 106, 303-309, doi:10.1007/s10533-010-9525-3 (2011).
- 24 Fornara, D. A. & Tilman, D. Plant functional composition influences rates of soil carbon and nitrogen accumulation. *Journal of Ecology* 96, 314-322, doi:10.1111/j.1365-2745.2007.01345.x (2008).
- 25 Steinbeiss, S. *et al.* Plant diversity positively affects short-term soil carbon storage in experimental grasslands. *Global Change Biology* 14, 2937-2949, doi:10.1111/j.1365-2486.2008.01697.x (2008).
- 26 Roscher, C. *et al.* The role of biodiversity for element cycling and trophic interactions: an experimental approach in a grassland community. *Basic and Applied Ecology* 5, 107-121, doi:10.1078/1439-1791-00216 (2004).
- 27 Trumbore, S. Radiocarbon and Soil Carbon Dynamics. *Annual Review of Earth and Planetary Sciences* 37, 47-66, doi:10.1146/annurev.earth.36.031207.124300 (2009).
- 28 Grace, J. B. *Structural Equation Modeling and Natural Systems*. (Cambridge University Press, 2006).
- 29 Bessler, H. *et al.* Aboveground overyielding in grassland mixtures is associated with reduced biomass partitioning to belowground organs. *Ecology* 90, 1520-1530, doi:10.1890/08-0867.1 (2009).
- 30 Lange, M. *et al.* Biotic and Abiotic Properties Mediating Plant Diversity Effects on Soil Microbial Communities in an Experimental Grassland. *PLoS ONE* 9(5), doi:10.1371/journal.pone.0096182 (2014).
- 31 Kramer, C. *et al.* Recent (< 4 year old) leaf litter is not a major source of microbial carbon in a temperate forest mineral soil. *Soil Biology & Biochemistry* 42, 1028-1037, doi:10.1016/j.soilbio.2010.02.021 (2010).
- 32 Rasse, D. P., Rumpel, C. & Dignac, M. F. Is soil carbon mostly root carbon? Mechanisms for a specific stabilisation. *Plant and Soil* 269, 341-356, doi:10.1007/s11104-004-0907-y (2005).
- 33 Bais, H. P., Weir, T. L., Perry, L. G., Gilroy, S. & Vivanco, J. M. The role of root exudates in rhizosphere interactions with plants and other organisms. *Annu Rev Plant Biol* 57, 233-266, doi:DOI 10.1146/annurev.arplant.57.032905.105159 (2006).
- 34 Bell, T., Newman, J. A., Silverman, B. W., Turner, S. L. & Lilley, A. K. The contribution of species richness and composition to bacterial services. *Nature* 436, 1157-1160, doi:10.1038/nature03891 (2005).
- 35 Oelmann, Y. *et al.* Soil and plant nitrogen pools as related to plant diversity in an experimental grassland. *Soil Science Society of America Journal* 71, 720-729, doi:10.2136/sssaj2006.0205 (2007).
- 36 Spehn, E. M. *et al.* The role of legumes as a component of biodiversity in a cross-European study of grassland biomass nitrogen. *Oikos* 98, 205-218, doi:10.1034/j.1600-0706.2002.980203.x (2002).


- 37 Zak, D. R., Holmes, W. E., White, D. C., Aaron, D. P. & Tilman, D. Plant Diversity, Soil Microbial
Communities, and Ecosystem Function: Are There Any Links? *Ecology* 84, 2042-2050 (2003).
- 38 Bessler, H. *et al.* Nitrogen uptake by grassland communities: contribution of N-2 fixation,
facilitation, complementarity, and species dominance. *Plant and Soil* 358, 301-322,
doi:10.1007/s11104-012-1181-z (2012).
- 39 Ravenek, J. M. *et al.* Long-term study of root biomass in a biodiversity experiment reveals
shifts in diversity effects over time. *Oikos* 123, 1528-1536, doi:10.1111/oik.01502 (2014).
- 40 Guenay, Y., Ebeling, A., Steinauer, K., Weisser, W. W. & Eisenhauer, N. Transgressive
overyielding of soil microbial biomass in a grassland plant diversity gradient. *Soil Biology &
Biochemistry* 60, 122-124, doi:10.1016/j.soilbio.2013.01.015 (2013).
- 41 Kramer, C. & Gleixner, G. Soil organic matter in soil depth profiles: Distinct carbon preferences
of microbial groups during carbon transformation. *Soil Biology & Biochemistry* 40, 425-433,
doi:10.1016/j.soilbio.2007.09.016 (2008).
- 42 Liang, C. & Balser, T. C. Microbial production of recalcitrant organic matter in global soils:
implications for productivity and climate policy. *Nature Reviews Microbiology* 9,
doi:10.1038/nrmicro2386-c1 (2011).
- 43 Schimel, J. P. & Schaeffer, S. M. Microbial control over carbon cycling in soil. *Frontiers in
microbiology* 3, doi:10.3389/fmicb.2012.00348 (2012).
- 44 Simpson, A. J., Simpson, M. J., Smith, E. & Kelleher, B. P. Microbially derived inputs to soil
organic matter: Are current estimates too low? *Environmental Science & Technology* 41, 8070-
8076, doi:10.1021/es071217x (2007).
- 45 Eisenhauer, N. *et al.* Plant diversity effects on soil food webs are stronger than those of
elevated CO₂ and N deposition in a long-term grassland experiment. *Proc Natl Acad Sci U S A*
110, 6889-6894, doi:10.1073/pnas.1217382110 (2013).
- 46 Spehn, E. M., Joshi, J., Schmid, B., Alphei, J. & Körner, C. Plant diversity effects on soil
heterotrophic activity in experimental grassland ecosystems. *Plant and Soil* 224, 217-230
(2000).
- 47 Mueller, K. E., Tilman, D., Fornara, D. A. & Hobbie, S. E. Root depth distribution and the
diversity-productivity relationship in a long-term grassland experiment. *Ecology* 94, 787-793
(2013).
- 48 FAO-UNESCO. *Soil map of the world. Revised legend with corrections and update.*, 1997).
- 49 Scheu, S. Automated measurement of the respiratory response of soil microcompartments -
active microbial biomass in earthworm feces. *Soil Biology & Biochemistry* 24, 1113-1118,
doi:10.1016/0038-0717(92)90061-2 (1992).
- 50 Tefs, C. & Gleixner, G. Importance of root derived carbon for soil organic matter storage in a
temperate old-growth beech forest - Evidence from C, N and C-14 content. *Forest Ecology and
Management* 263, 131-137, doi:10.1016/j.foreco.2011.09.010 (2012).
- 51 Milcu, A. *et al.* Functional diversity of leaf nitrogen concentrations drives grassland carbon
fluxes. *Ecology Letters* 17, 435-444, doi:10.1111/ele.12243 (2014).
- 52 Kramer, C. & Gleixner, G. Variable use of plant- and soil-derived carbon by microorganisms in
agricultural soils. *Soil Biology & Biochemistry* 38, 3267-3278, doi:10.1016/j.soilbio.2006.04.006
(2006).
- 53 Zelles, L. Phospholipid fatty acid profiles in selected members of soil microbial communities.
Chemosphere 35, 275-294, doi:10.1016/s0045-6535(97)00155-0 (1997).
- 54 Esperschuetz, J. *et al.* Microbial response to exudates in the rhizosphere of young beech trees
(*Fagus sylvatica* L.) after dormancy. *Soil Biology & Biochemistry* 41, 1976-1985,
doi:10.1016/j.soilbio.2009.07.002 (2009).

- 55 Treonis, A. M. *et al.* Identification of groups of metabolically-active rhizosphere microorganisms by stable isotope probing of PLFAs. *Soil Biology & Biochemistry* 36, 533-537, doi:10.1016/j.soilbio.2003.10.015 (2004).
- 56 Marchesi, J. R. *et al.* Design and evaluation of useful bacterium-specific PCR primers that amplify genes coding for bacterial 16S rRNA. *Applied and Environmental Microbiology* 64, 795-799 (1998).
- 57 Lane, D. J. in *Nucleic acid techniques in bacterial systematics* (eds E. Stackebrandt & M. Goodfellow) 115 - 175 (John Wiley and Sons, 1991).
- 58 Gardes, M. & Bruns, T. D. ITS primers with enhanced specificity for basidiomycetes - application to the identification of mycorrhizae and rusts. *Molecular Ecology* 2, 113-118, doi:10.1111/j.1365-294X.1993.tb00005.x (1993).
- 59 White, D. C., Davis, W. M., Nickels, J. S., King, J. D. & Bobbie, R. J. Determination of the Sedimentary Microbial Biomass by Extractable Lipid Phosphate. *Oecologia* 40, 51-62 (1979).
- 60 R: A language and environment for statistical computing v. 3.0.3 (R Foundation for Statistical Computing, Vienna, Austria, 2014).
- 61 Rosseel, Y. lavaan: An R Package for Structural Equation Modeling. *Journal of Statistical Software* 48, 1-36 (2012).
- 62 Andren, O. & Katterer, T. ICBM: The introductory carbon balance model for exploration of soil carbon balances. *Ecological Applications* 7, 1226-1236 (1997).
- 63 Sierra, C. A. *et al.* Predicting decadal trends and transient responses of radiocarbon storage and fluxes in a temperate forest soil. *Biogeosciences* 9, 3013-3028, doi:10.5194/bg-9-3013-2012 (2012).
- 64 Wang, Y. P. *et al.* Oscillatory behavior of two nonlinear microbial models of soil carbon decomposition. *Biogeosciences Discuss.* 10, 19661-19700, doi:10.5194/bgd-10-19661-2013 (2013).
- 65 Sierra, C. A., Müller, M. & Trumbore, S. E. Modeling radiocarbon dynamics in soils: SoilR version 1.1. *Geoscientific Model Development Discussions* 7 (2014).
- 66 Brodowski, S., Amelung, W., Haumaier, L. & Zech, W. Black carbon contribution to stable humus in German arable soils. *Geoderma* 139, 220-228, doi:10.1016/j.geoderma.2007.02.004 (2007).
- 67 Vance, E. D., Brookes, P. C. & Jenkinson, D. S. An Extraction Method for Measuring Soil Microbial Biomass-C. *Soil Biology & Biochemistry* 19, 703-707 (1987).
- 68 Schulze, E. D. *et al.* Importance of methane and nitrous oxide for Europe's terrestrial greenhouse-gas balance. *Nature Geoscience* 2, 842-850, doi:10.1038/ngeo686 (2009).
- 69 Solly, E. *et al.* Mean age of carbon in fine roots from temperate forests and grasslands with different management. *Biogeosciences* 10, 4833-4843, doi:10.5194/bg-10-4833-2013 (2013).
- 70 Sollins, P., Homann, P. & Caldwell, B. A. Stabilization and destabilization of soil organic matter: Mechanisms and controls. *Geoderma* 74, 65-105, doi:10.1016/s0016-7061(96)00036-5 (1996).







a)







b)



c)



a) Annual Stock Changes



Plant species richness

