



Making the most of what we have: application of extrapolation approaches in radioecological wildlife transfer models



Nicholas A. Beresford^{a, b, *}, Michael D. Wood^b, Jordi Vives i Batlle^c, Tamara L. Yankovich^d, Clare Bradshaw^e, Neil Willey^f

^a NERC Centre for Ecology & Hydrology, Lancaster Environment Center, Library Av., Bailrigg, Lancaster LA14AP, UK

^b School of Environment & Life Sciences, University of Salford, Manchester M4 4WT, UK

^c Belgian Nuclear Research Centre, Boeretang 200, 2400 Mol, Belgium

^d International Atomic Energy Agency, Vienna International Centre, 1400 Vienna, Austria

^e Department of Ecology, Environment and Plant Sciences, Stockholm University, SE-10691, Sweden

^f Centre for Research in Biosciences, University of the West of England, Coldharbour Lane, Frenchay, Bristol BS16 1QY, UK

ARTICLE INFO

Article history:

Received 22 December 2014

Received in revised form

4 March 2015

Accepted 7 March 2015

Available online 4 April 2015

Keywords:

Allometry

REML

Concentration ratio

Stoichiometry

Extrapolation

Biological half-life

ABSTRACT

We will never have data to populate all of the potential radioecological modelling parameters required for wildlife assessments. Therefore, we need robust extrapolation approaches which allow us to make best use of our available knowledge. This paper reviews and, in some cases, develops, tests and validates some of the suggested extrapolation approaches.

The concentration ratio ($CR_{\text{product-diet}}$ or $CR_{\text{wo-diet}}$) is shown to be a generic (trans-species) parameter which should enable the more abundant data for farm animals to be applied to wild species.

An allometric model for predicting the biological half-life of radionuclides in vertebrates is further tested and generally shown to perform acceptably. However, to fully exploit allometry we need to understand why some elements do not scale to expected values.

For aquatic ecosystems, the relationship between $\log_{10}(a)$ (a parameter from the allometric relationship for the organism–water concentration ratio) and $\log(K_d)$ presents a potential opportunity to estimate concentration ratios using K_d values.

An alternative approach to the $CR_{\text{wo-media}}$ model proposed for estimating the transfer of radionuclides to freshwater fish is used to satisfactorily predict activity concentrations in fish of different species from three lakes. We recommend that this approach (REML modelling) be further investigated and developed for other radionuclides and across a wider range of organisms and ecosystems.

Ecological stoichiometry shows potential as an extrapolation method in radioecology, either from one element to another or from one species to another.

Although some of the approaches considered require further development and testing, we demonstrate the potential to significantly improve predictions of radionuclide transfer to wildlife by making better use of available data.

© 2015 The Authors. Published by Elsevier Ltd. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

1. Introduction

The assessment of the exposure of wildlife to ionising radiation for planned, existing and emergency scenarios requires predictions to be made of the transfer of a wide range of radionuclides to a diversity of species. Most models assessing the exposure of wildlife

for regulatory purposes use a simple concentration ratio ($CR_{\text{wo-media}}$) relating the whole organism activity concentration to that in an environmental medium (i.e. soil, air or water) (Beresford et al., 2008a). Other models use radionuclide biological half-lives and transfer from the diet (e.g. USDOE, 2002).

For many of the radionuclide-species combinations that require assessment, there are no empirical data. When empirical data are lacking, predictions are often made using transfer parameter values derived using extrapolation approaches, though clarity on the use of such approaches varies between publications/models. For instance, the $CR_{\text{wo-media}}$ value for an organism of ‘similar taxonomy’

* Corresponding author. NERC Centre for Ecology & Hydrology, Lancaster Environment Center, Library Av., Bailrigg, Lancaster LA14AP, UK.

E-mail address: nab@ceh.ac.uk (N.A. Beresford).

may be assumed (e.g. a mammal value may be used to model birds if data for the latter are lacking) (e.g. Beresford et al., 2008b). Brown et al. (2013) recently evaluated how the most commonly used approaches performed against additional data now available

intake, between animals of differing age or species, rather than any difference in radionuclide transfer (e.g. Smith and Beresford, 2005; Galeriu et al., 2007; Beresford et al., 2007). The rationale for this is that the transfer coefficient is defined as:

$$F_f = \frac{\text{Radionuclide activity concentration in meat (Bq kg}^{-1}\text{)}}{\text{Daily dry matter intake (kg d}^{-1}\text{)} \times \text{Radionuclide activity concentration of the diet (Bq kg}^{-1}\text{ DM)}}$$

(Coppelstone et al., 2013). They concluded that the extrapolation approaches commonly used to date have under-predicted the empirical 95th percentile $CR_{\text{wo-media}}$ value as often as they have over-predicted. This highlights the need to treat results from these previous extrapolations with caution, but there have been various recent advances that may bring greater confidence in the application of extrapolation methods. In this paper, we assess our ability to extrapolate radioecological data of relevance to wildlife radiological assessments considering these recent advances and future potential.

2. Transfer from the diet to terrestrial vertebrates is a constant across species

The transfer coefficient was first proposed by Ward et al. (1965) to describe the transfer of radiocaesium from the diet to the milk of dairy cattle. The authors defined the transfer coefficient as the ratio between the radiocaesium activity concentration in milk and the daily dietary radionuclide intake. Ward et al. (1965) reported that this parameter exhibited less variability between individual animals within their experimental herd than when transfer was expressed as the total amount of Cs excreted in milk (represented as a percentage of intake). Ward and Johnson (1965) subsequently defined the meat transfer coefficient as the ratio of the ^{137}Cs activity concentration in boneless meat to the dietary daily ^{137}Cs intake.

Following the publications of Ward and co-workers in the 1960s, the transfer coefficient was adopted as the basis for quantifying transfer to milk (F_m , d l^{-1} or d kg^{-1}), and meat and eggs (F_f , d kg^{-1}) for all radionuclides. By the late 1970s to early 1980s, transfer coefficient values were being recommended for most radionuclide-animal product combinations (e.g. Ng, 1982; Ng et al., 1977, 1979, 1982) and the International Atomic Energy Agency (IAEA) recommended their use (IAEA, 1994). These recommended values have been incorporated into many predictive food chain models (e.g. Brown and Simmonds, 1995; Müller and Pröhl, 1993; USNRC, 1977; Yu et al., 2001).

On the basis of the many studies conducted over the approximately 50 years since the transfer coefficient concept was introduced, it has generally been accepted that transfer coefficients for smaller animals are higher than those for larger animals, and that those for adults are lower than those for young (and hence smaller) livestock. For instance, F_f values for sheep meat recommended by IAEA (2010) for many radionuclides are *circa* one order of magnitude higher than those recommended for beef. Similarly, F_m values for goat milk tend to be one order of magnitude higher than those recommended for cow milk. The use of transfer coefficients has also been suggested for wildlife (e.g. Thomas et al., 1994; MacDonald, 1996; Moss and Horrill, 1996) and some models use simple food chains to estimate radionuclide concentrations in wildlife (e.g. USDOE, 2002).

However, it has been suggested that much of the observed difference in $F_{f,m}$ values is a consequence of differences in dry matter

Consequently, the concentration ratio ($CR_{\text{meat-diet}}$) is equal to:

$$CR_{\text{meat-diet}} = F_f (\text{d kg}^{-1}) \times \text{Daily dry matter intake (kg d}^{-1}\text{)}$$

The above equations are for meat but a similar derivation can be performed for milk or eggs.

A between species similarity in CR values for animal derived food products should not be surprising, given that the concentrations of many elements in meat, or milk, are similar across species (Mertz, 1986, 1987). A particular advantage of being able to assume that the milk or meat CR for many radionuclides varies little between species is that generic values can be derived for animals for which no data are currently available. Recognising this, the IAEA (2010) summarised milk and meat CR values, as well as transfer coefficients.

We would also expect that CR values for wildlife would vary little between species (Beresford et al., 2004) and would be similar to those of farm animals. To test this hypothesis, Table 1 presents $CR_{\text{meat-diet}}$ values for seven herbivorous species of wild mammals and birds. The $CR_{\text{meat-diet}}$ values for these species are similar to those for the meat of farm animals in IAEA (2010), which presents a generic value of 0.39 based on data for four farm animal species. Since Cs is relatively homogeneously distributed throughout the body tissues (Yankovich et al., 2010a), it can then be assumed that, for Cs $CR_{\text{meat-diet}} \approx CR_{\text{org-diet}}$ (where $CR_{\text{org-diet}}$ is the ratio of the radionuclide activity concentration in the whole organism to that in its diet). Currently there are few data with which to test our hypothesis for the transfer of other radionuclides to wild animals. For many elements other than Cs, distribution is not homogenous throughout the body tissues. However, for such elements an assumption that the distribution within the body was similar across animal species would be reasonable (e.g. Sr accumulates in the bone of all vertebrates).

3. Allometry

Size affects rates of biological processes from cellular metabolism to population dynamics (Peters, 1983; Hoppeler and Weibel, 2005). The dependence of a biological variable (Y) on body mass (M) is typically characterised by an allometric scaling law. There are several allometric equations that can be proposed, the simplest being to assume that:

$$Y = aM^b$$

where a and b (the allometric exponent) are constants, b is dimensionless and a has the units of the variable, Y , per mass to the power of $-b$.

Kleiber (1932) found that basal metabolic rate (measured as heat production) across 13 groups of mature animals, ranging from

Table 1
Caesium $CR_{\text{meat-diet}}$ values for wildlife; for both source references 'diet' concentration is determined from dried stomach contents.

Species Latin	Species common	Mean \pm SD ^a	n	Reference
<i>Cervus elaphus</i>	Red deer	0.40 \pm 0.20	56	Chaplow et al., 2015
<i>Odocoileus virginianus</i>	Whitetail deer	0.53 ^b	11	Sheppard 2013
<i>Lepus europaeus</i>	Brown hare	0.46	1	Chaplow et al., 2015
<i>Lepus timidus</i>	Blue hare	0.48 \pm 0.06	5	Chaplow et al., 2015
<i>Oryctolagus cuniculus</i>	Rabbit	0.27 \pm 0.11	20	Chaplow et al., 2015
<i>Tetrao tetrix</i>	Black grouse	0.40–0.85	2	Chaplow et al., 2015
<i>Lagopus lagopus scotica</i>	Red grouse	0.73 \pm 0.40	10	Chaplow et al., 2015

^a Where $n = 2$ the range is presented.

^b Geometric mean as presented in source publication.

<0.2 to circa 680 kg body mass, was proportional to mass to the power 0.74 (i.e. $M^{0.74}$). Kleiber subsequently suggested that 'metabolic body size' (now generally referred to as metabolic live-weight) could be determined as $M^{0.75}$ (Kleiber, 1947); this has since become known as *Kleiber's law*.

There is considerable debate with regard to the numerical values for the allometric exponent, in particular whether it should be 0.75 or perhaps 0.67 for basal metabolic rate (e.g. West et al., 1997; Hoppeler and Weibel, 2005; Isaac and Carbone, 2010; Agutter and Tuszynski, 2011). The debate hinges on theoretical considerations, such as whether allometry of the metabolic rate relates to the fractal nature of energy distribution networks (West et al., 1999; Brown et al., 2002) with an exponent of 0.75, or whether it is based on a consideration of surface area-to-volume ratios with an exponent of 0.67 (Farrell-Gray and Gotelli, 2005).

In discussing this issue with respect to radioecological models, Higley and Bytwerk (2007) suggested that, given other uncertainties in radioecological modelling, the exact value of the allometric scaling exponent 'may not be of critical importance' for practical (rather than theoretical) purposes. We support this suggestion and Fig. 1 demonstrates the relatively small effect of assuming $M^{0.67}$ rather than $M^{0.75}$ over a likely range of vertebrate live masses.

3.1. Allometry in radioecology

3.1.1. Homeothermic vertebrates

Many of the reported allometric relationships, such as dry matter food ingestion rates, water ingestion rates and inhalation




Fig. 1. A comparison of predictions assuming allometric exponents of 0.67 and 0.75 across a mass range appropriate for terrestrial mammals (predictions are M^b). Predictions using an exponent of 0.037, appropriate to reptiles, are also shown.

rates, have been used in models of the radionuclide transfer to wildlife (e.g. Beresford et al., 2008a; Johansen et al., 2012), including the US Department of Energy's 'Graded Approach' as implemented in the RESRAD-Biota model (USDOE, 2002, 2004). However, in the 1970s, radionuclide biological half-life ($T_{1/2B}$) was also shown to scale allometrically (e.g. Stara et al., 1971; Kitchings et al., 1976). The application of allometry to radioecological parameters has received revived attention during the development of models to predict the exposure of wildlife to radionuclides in terrestrial (Higley et al., 2003; Higley, 2010; Beresford et al., 2004; Sheppard, 2001) and aquatic ecosystems (Vives i Batlle et al., 2007, 2009; Brown et al., 2004).

USDOE (2002) presents $T_{1/2B}$ allometric relationships for 16 elements in terrestrial/riparian vertebrates. When used in model inter-comparison exercises, allometric relationships from USDOE (2002), and similar models, produced results comparable to models using $CR_{\text{wo-media}}$ values (Beresford et al., 2009; Wood et al., 2009a). Of the allometric expressions describing radionuclide $T_{1/2B}$ from USDOE (2002), a number have an exponent of approximately 0.25 (Cs, Co, Ra, Sb, Sr, U, Zn and Zr). Iodine has an exponent of 0.13 and H of 0.55, but other literature suggests these too should approximate to 0.25 (Galeriu et al., 2003; MacDonald, 1996). Therefore, it appears that for most radionuclides, $T_{1/2B}$ is driven by metabolic rate and Beresford and Vives i Batlle (2013) demonstrated that this was logical by considering a simple radionuclide retention model (Sazykina, 2000) and Kleibers Law. Notable exceptions are Pu, Am, Ce and Eu, which USDOE (2002) suggest scale to circa 0.8.

Sheppard (2001) proposed that, if it is accepted that there is an approximation of the exponent applicable for all elements (i.e. in the case of $T_{1/2B}$, circa 0.25), then only an estimation of the multiplicand is needed for any given element. Through algebraic derivation, Beresford and Vives i Batlle (2013) proposed a method of determining the multiplicand for the $T_{1/2B}$ relationship:

$$a_B = \frac{\ln 2}{a_1 f_1} CR_{\text{org-diet}}$$

where a_B is the multiplicand for the allometric relationship describing $T_{1/2B}$, a_1 is the multiplicand for the allometric relationship describing dry matter intake and f_1 is the fractional gastrointestinal absorption coefficient. By substituting this expression for a_B in the equation describing the $T_{1/2B}$ allometric relationship, Beresford and Vives i Batlle (2013) suggested that an estimate of $T_{1/2B}$ could be derived as:

$$T_{1/2B} = \frac{\ln 2}{a_1 f_1} CR_{\text{org-diet}} M^{0.25}$$

Values of a_1 are relatively well documented for terrestrial vertebrates (e.g. Nagy, 2001), f_1 values are collated in publications (e.g. IAEA, 2010), and, if we accept that $CR_{\text{org-diet}}$ is a constant across

species (see above), then this parameter can also be sourced from IAEA (2010) for many radionuclides.

Beresford & Vives i Batlle tested their hypothesis using data primarily summarised by Whicker and Schultz (1982), which tabulates $T_{1/2B}$ estimates for a number of radionuclides and terrestrial organisms. Data were for mammals ranging from 0.01 kg (harvest mouse) to 80 kg (reindeer) and radionuclides of Co, Cs, I and Sr. A comparison of predicted $T_{1/2B}$ values across all radionuclides made using a_1 values appropriate to the feeding type of each species with measured data yielded a linear regression ($p < 0.001$) with an R^2 value of 0.58, a slope of 1.4 and an intercept which was not significantly different from zero (Beresford and Vives i Batlle, 2013).

Subsequent to Beresford and Vives i Batlle (2013), a review of radionuclide $T_{1/2B}$ values has been conducted (this is contributing to IAEA Modelling and Data for Radiological Impact Assessments (MODARIA) programme activities, see: <http://bit.ly/1e9Nxxq>). Data identified in this review enable us to further test the Beresford and Vives i Batlle (2013) model and to consider a wider range of elements (Ag, Co, Cs, I, Na, Nb, Ru, Se, Sr and Zn) and species. The data used had to meet various criteria:

- animal mass had to be presented in the source reference;
- only data for adults were used;
- $T_{1/2B}$ values had to be for the whole body except in the cases of I and Cs, for which thyroid and muscle data were also used, respectively, assuming these tissues reflected whole body loss rates;
- with the exception of I, Sr and Cs, only data for studies where the radionuclide had been ingested or administered orally were considered; for I, Sr and Cs, data from studies using intravenous administration were also used, as the weight of evidence suggested that these elements behave in a similar manner in the circulatory system after either oral or intravenous administration (Mayes et al., 1996).

A total of 123 $T_{1/2B}$ values have been considered in this expanded model testing (source references are listed in Supplementary Materials); these included data originally considered by Beresford and Vives i Batlle (2013), though source references were consulted rather than relying on the Whicker and Schultz (1982) compilation. Data were largely for mammals (ranging from 8 g to 70 kg body mass), although limited data for ducks were also identified. When multiple components of loss were cited in the source reference, the longest $T_{1/2B}$ value was used for comparison with predictions.

Radionuclide-specific parameter values used to make predictions are presented in Table 2. Estimates of f_1 were taken from IAEA (2010), which cites values from ICRP (2006) for monogastric

animals and additionally presents f_1 values for ruminants (the ruminant values were only used here if no monogastric f_1 value was available; Beresford and Vives i Batlle (2013) compared the effect of using ruminant vs monogastric f_1 values). Values of $CR_{org-diet}$ were estimated from $CR_{meat-diet}$ values presented by IAEA (2010). Whilst IAEA (2010) presents $CR_{meat-diet}$ values for Co, Cs and I, it does not include a $CR_{meat-diet}$ value for Sr. Dietary transfer coefficients presented in IAEA (2010) were, therefore, used together with typical dry matter intake rates from IAEA (1994) to estimate an average Sr $CR_{org-diet}$ across all five species for which F_f data were available (cattle, goat, sheep, poultry and pig) in IAEA (2010). To convert $CR_{meat-diet}$ values to $CR_{org-diet}$ values, conversion factors from Yankovich et al. (2010a) were used; if Yankovich et al. contained no data for a given element, then conversion factors were estimated from Coughtrey et al. (1983) or the data of Barnett et al. (2014).

Values of a_1 were sourced from Nagy (2001), and both the 'all mammals' value ($a_1 = 0.057 \text{ d}^{-1} \text{ kg}^{0.25}$) and values for the most appropriate taxonomic or feeding strategy group were used: carnivorous mammals ($a_1 = 0.027 \text{ d}^{-1} \text{ kg}^{0.25}$); herbivorous mammals ($a_1 = 0.15 \text{ d}^{-1} \text{ kg}^{0.25}$); omnivorous mammals ($a_1 = 0.077 \text{ d}^{-1} \text{ kg}^{0.25}$); omnivorous birds ($a_1 = 0.119 \text{ d}^{-1} \text{ kg}^{0.25}$). Values presented by Nagy, in grammes, were converted to kilograms for application here, using the multiplicative factor: $(a_1 \times 1000^{0.75})/1000$.

The majority of predictions were within an order of magnitude of the observed data: 97 of 110 predictions using the a_1 for all mammals and 107 of 123 predictions made using the most appropriate feeding group a_1 (Table 3). Allometry is a broad indicator only (Higley and Bytwerk, 2007) and hence predictions within an order of magnitude are considered acceptable here.

Using the a_1 for 'all mammals', nine predictions were more than an order of magnitude lower than the observed data; with the exception of one prediction (Cs in rabbit), these large under-predictions were for Nb ($n = 4$) and Ru ($n = 4$). These eight comparisons comprised all of the available data for Ru and Nb. Only four predictions were more than an order of magnitude higher than the observed data. All of these over predictions were for Sr; these data had not previously been considered in the initial evaluation of the model by Beresford and Vives i Batlle (2013). These observations are discussed further below.

When predictions were made using a_1 values for specific feeding strategies (see Table 3), the following were observed:

Carnivorous mammals – Using the a_1 value for carnivorous mammals increased the predicted $T_{1/2B}$ value compared to that predicted using the 'all mammals' a_1 value (Fig. 2). The predictions using the carnivorous mammal a_1 value were generally in better agreement with the observed data.

Herbivorous mammals – Using the a_1 value for herbivorous mammals decreased the predicted $T_{1/2B}$ value compared to that predicted using the 'all mammals' a_1 value (Fig. 3). There was not a consistent trend across all of the elements to improve, or not, the level of agreement between predictions and observed data (e.g. for I predictions using the a_1 value for herbivorous mammals resulted in better agreement with the observed data than those using the 'all mammals' a_1 value, whereas the opposite was the case for Zn).

Omnivorous mammals – Using the a_1 value for omnivorous mammals decreased the predicted $T_{1/2B}$ value compared to that predicted using the 'all mammals' a_1 value, but only by circa 25%.

Predictions for I were in good agreement for herbivorous and carnivorous mammals. For omnivores, predicted values tended to be <20% of the observed data values. Caesium data were the most numerous ($n = 55$). If we omit an outlying value for rabbit, which was under-predicted by more than an order of magnitude, a linear regression of measured Cs $T_{1/2B}$ values to predictions made using

Table 2

Parameter values used to predict the biological half-life of radionuclides in mammals and birds (data sources are cited in text).

Element	f_1	$CR_{meat-diet}$	Wholebody to muscle correction factor	$CR_{org-diet}$
Ag	5E-2	4.3E-4	1.2E+2	5.2E-2
Co	1E-1	3.1E-1	3E+0	9.3E-1
Cs	1E+0	3.9E-1	1E+0	3.9E-1
I	1E+0	9.4E-2	5E+0	4.7E-1
Po	5E-1	1.4E-1	2E+0	2.8E-1
Sr	3E-1	2.2E-2	4.E+2	8.8E+0
Zn	5E-1	1.9E+0	1.8E+0	3.42E+0
Na	9E-1	9.7E-1	1E+0	9.7E-1
Nb	1E-2	6.5E-6	1.1E+1	7.2E-5
Ru	5E-2	5.4E-4	1E+0	5.4E-4
Se	8E-1	1.1E+0	1E+0	1.1E+0

Table 3

A summary comparison of predicted with observed $T_{1/2B}$ values.

Element	Animal type compared ^a	N ^b	Mean (range) predicted:observed using 'all mammal' a_1	Mean (range) predicted:observed using feeding group a_1
Ag	Mouse species, rat, monkey, dog	4	3.1 (0.79–6.8)	2.5 (1.2–5.0)
Co	Mouse, rat	2	(1.2–8.5)	0.85–6.3
Cs	Mouse species, rat, monkey, dog, wild boar, rabbit, guinea pig, deer species, fox species, coyote, wolf	55	0.44 (0.08–2.2)	0.34 (0.03–1.0)
I	Mouse, rat, monkey, dog, guinea pig, deer	11	0.98 (0.14–2.5)	0.50 (0.07–1.3)
Na	Mouse, rat species, monkey	4	0.21 (0.13–0.31)	0.16 (0.10–0.23)
Nb	Mouse, rat, monkey, dog	4	0.04 (0.002–0.10)	0.04 (0.002–0.08)
Ru	Mouse, rat, monkey, dog	4	0.008 (0.001–0.02)	0.01 (0.0008–0.03)
Se	Mouse, rat, monkey, dog	4	0.49 (0.40–0.56)	0.55 (0.29–1.2)
Sr	Mouse, rat, rabbit, dog, deer	6	22 (1.2–50)	21 (1.7–74)
Zn	Mouse species, rat species, vole, monkey, dog, duck	29	1.9 (0.21–5.9) ^c	1.0 (0.15–4.4) ^d

^a Where 'species' is used data were available for more than one species falling into the broad animal types listed.

^b Number of comparisons.

^c Mammals only ($n = 16$).

^d Includes ducks ($n = 29$).

the appropriate feeding group a_1 value yields an R^2 of 0.52, a slope of 0.43, and an intercept which is not significantly different to zero. The slope demonstrates a tendency to under-predict the measured values.

There was a tendency to over-predict Sr $T_{1/2B}$ values, some by more than an order of magnitude (Table 3). Those four values which were over-predicted by more than an order of magnitude were all from studies using intravenous administration or radiostrontium. The reported half-lives (see DiGregorio et al., 1978) were

comparatively short compared to the two studies reporting Sr $T_{1/2B}$ values following oral administration and also data from farm animal studies (Fesenko et al., 2015).

Predictions for Ag, Co, Se and Zn all had an acceptable level of agreement. Those for Na were within an order of magnitude of the observed $T_{1/2B}$ values, but were all comparatively low.

Predictions for both Ru and Nb were poor, both elements being considerably under-predicted. In the cases of Ru and Nb, it is worth noting that the $CR_{meat-diet}$ values in IAEA (2010) are each based




Fig. 2. The effect of a_1 value on the predicted $T_{1/2B}$ values for carnivorous mammals; note three outlying data points (one each for Nb, Ru and Sr) are not shown. Data represent individual studies/animals for which live-mass and biological half-life values were available.




Fig. 3. The effect of a_1 value on the predicted $T_{1/2B}$ values for herbivorous mammals; note two outlying data points (one each for Cs and Sr) are not shown. Data represent individual studies/animals for which live-mass and biological half-life values were available.

upon one observation only, although this is also the case for Ag and Se. To improve the predictions such that they are within an order of magnitude of the observed values, the $CR_{\text{meat-diet}}$ for Nb would need to increase by approximately 1–2 orders of magnitude and that for Ru by about two-orders of magnitude. This is plausible as IAEA (2010) presents $CR_{\text{meat-diet}}$ values for other elements which vary by 1–2 orders of magnitude between studies. The f_1 values used also impact on the predicted $T_{1/2B}$ values for both Nb and Ru. There are reported f_1 values for these elements, which are lower than those we have used in this assessment (Coughtrey and Thorne, 1983; Beresford et al., 1998a); application of a lower f_1 would improve the predicted values. However, the Nb and Ru $T_{1/2B}$ used here are from Furchner and Drake (1971) and Furchner et al. (1971) respectively; the f_1 values estimated by these authors from experimental data were similar to those used to make our predictions.

There was reasonable agreement between the predicted and measured Zn $T_{1/2B}$ values for ducks; the ratio of predicted to measured values ranged from 0.2 to 1.5. However, $T_{1/2B}$ values varied considerably more for the observed data (28–250 d; $n = 13$) than the predicted values (41–42 d). The low variation in predictions is not surprising given that the mass of the ducks varied little (1.1–1.3 kg).

Overall, our ability to obtain reasonable predictions is, in part, dependent upon the quality of data available for the required input parameters. For many elements in IAEA (2010), $CR_{\text{meat-diet}}$ values are based upon few observations. Similarly, the correction factors to convert from $CR_{\text{meat-diet}}$ to $CR_{\text{org-diet}}$ are based on relatively few data, as exemplified by the need to derive them for Co, I and Sr. Investigation of the data used by Nagy (2001) to derive the allometric dry matter intake relationship for herbivores shows that they are dominated by relatively small species, with many of the larger species being marsupials.

It should also be acknowledged that the dry matter intake relationships presented by Nagy (2001) are for animals under field and not laboratory conditions, and that field metabolic rates are generally higher than basal metabolic rates determined for housed animals (Nagy, 2005). This may result in a tendency to under-predict $T_{1/2B}$ for housed (i.e. experimental) animals. Overall, in the above assessment, there was a tendency to under- rather than over-predict, 103 of the 123 predictions using feeding group a_1 values were less than the observed value.

An assumption of the approach described here is that $T_{1/2B}$ scales to the power of 0.25. For five elements in USDOE (2002) (Am, Ce, Eu, Pu and Th), $T_{1/2B}$ scales to the power of 0.8. For all of these elements, International Commission on Radiological Protection publications (ICRP, 1979, 1981, 1988) are quoted as the data source, but unfortunately, we have not been able to find the data within these publications to independently verify the allometric equations presented by USDOE (2002). None of the elements scaling to *circa* 0.8 play an important biological role, and hence, it could, perhaps, be suggested that there is no reason for them to follow a metabolically driven uptake process (i.e. as signified by a mass scaling function of 0.25). However, we acknowledge that some of the elements which do scale as mass to the power of 0.25 are neither essential elements nor their analogues (e.g. U). Therefore, it is recommended that the reasons for the deviation of some elements from the mass scaling function of 0.25 be explored and/or the relationships presented by USDOE (2002) be independently verified.

For comparison with our predictions, we have used the longest $T_{1/2B}$ value reported where multiple components of loss were observed. In doing so, we made no differentiation based on the magnitude of this loss component, even though in many instances, the long component of loss contributes relatively little to the total loss (<5%). It is questionable whether the longest $T_{1/2B}$ always best

represents metabolic turnover when three and four component loss equations are reported.

The assumption of a single long-component of loss, as currently used in most wildlife assessment models (e.g. USDOE, 2002; Avila et al., 2004), should, if the models are correctly parameterised, yield estimates of the equilibrium activity concentration in organisms which are conservative (i.e. they should be over-estimated compared to a model that assumes more than one loss component). However, if used in dynamic models, this assumption will predict slower changes in organism activity concentrations than would be observed in reality as a result of changes in activity concentrations in environmental media.

3.1.2. Application to reptiles

Although only proposed for homeothermic vertebrates, allometric models for $T_{1/2B}$, such as USDOE (2002), have been used to make predictions of radionuclide activity concentrations in (poikilothermic) reptiles and amphibians (e.g. Wood et al., 2009a; Beresford et al., 2010; Yankovich et al., 2010b; Johansen et al., 2012).

Beresford and Wood (2014) evaluated to what extent the Beresford and Vives i Batlle (2013) model could be applied to reptiles. Similar criteria as used above for mammal and bird data were applied to identify $T_{1/2B}$ values against which to compare predictions. In total 28 $T_{1/2B}$ values for Cs, three for Sr and two for Ra were identified (see Beresford and Wood (2014)). Data for Cs were predominantly for different species of snake (Staton et al., 1974), with one value for a turtle species (*Trachemys scripta scripta*) (Scott et al., 1986). Data identified for Sr and Ra were all for *T. scripta scripta* (Scott et al., 1986; Hinton et al., 1992). Data covered reptiles with masses ranging from 0.02 to 1.5 kg. Most of the estimated $T_{1/2B}$ values available were longer than the length of the studies from which they were derived. For instance, the maximum radiocaesium half-life determined by Staton et al. (1974) for snakes from their 63 d long study was 430 d. This will undoubtedly add some uncertainty to the reported $T_{1/2B}$ values.

Using parameter values for mammals, all but one prediction was under-estimated; in the case of Cs, the under-estimates were by 1–2 orders of magnitude.

The exponent in the allometric biological half-life expressions is, in effect, defined by the exponent for the allometric model of metabolic rate (B_r) of *circa* 0.75 for mammals; the exponent for $T_{1/2B}$ is '(1 – the exponent for B_r)'. However, the exponent for the allometric model describing B_r for reptiles has been shown to be in the range *circa* 0.80–0.92 (Nagy, 2005; Isaac and Carbone, 2010). Therefore, the exponent describing $T_{1/2B}$ for reptiles should be in the range 0.08–0.20. From Nagy (2001), Beresford and Wood (2014) estimated exponents of 0.037 for snakes, derived from information for carnivorous reptiles, and 0.08 for turtles, derived from information for 'all reptiles'. These exponents, together with the relevant values of a_1 for reptiles (Nagy, 2001) and reptile-specific values of $CR_{\text{org-diet}}$ and f_1 (see Beresford and Wood (2014) for details of derivation), were used to estimate $T_{1/2B}$ values to compare with the available data (Fig. 4). Only one of the 33 predictions deviates by more than a factor of six from the measured value; the one 'outlier', a prediction of the Sr $T_{1/2B}$ in *T. scripta scripta*, was predicted to be 14 times higher than the measured value. However, $T_{1/2B}$ allometric exponents applicable to reptiles result in a relatively low dependence of $T_{1/2B}$ with mass (see Figs. 1 and 4). Over a mass range 0.1 g–1000 kg, the predicted biological half-lives for reptiles vary by less than a factor of four using an exponent of 0.08, and by less than a factor of two using an exponent of 0.037. For homeotherms, $M^{0.25}$ predicts $T_{1/2B}$ values varying over *circa* 60-fold (Fig. 1). Given the small influence of mass on $T_{1/2B}$ predictions for reptiles, Beresford and Wood (2014) suggested that if sufficient reported $T_{1/2B}$ values are available for a given element,




Fig. 4. A comparison of measured radionuclide biological half-life (d) in reptiles with predictions using reptile specific parameter values in the model of Beresford and Vives i Batlle (2013) (from Beresford and Wood, 2014).

then it is likely that these would be applicable to any reptile. For instance all of the 28 reported values of Cs $T_{1/2B}$ for reptiles considered by Beresford and Wood (2014), which covered a 50-fold mass range, were within a factor of 5 of the mean.

However, the relatively good agreement between predicted and measured $T_{1/2B}$ in Fig. 4 demonstrates that if no reptile data are available for a given radionuclide, then the Beresford and Vives i Batlle model (2013), populated with reptile-specific parameter values, will give reasonable estimates.

There is some evidence that the dry matter intake rate of herbivorous reptiles (namely tortoises), scales to mass with an exponent of circa 0.75 (Franz et al., 2011). Therefore, we may expect the allometric relationship for $T_{1/2B}$ for tortoise species to scale to circa 0.25, but we currently have no data to test this.

3.1.3. Aquatic organisms

For marine mammals, the allometric models developed by USDOE (2002) for terrestrial and riparian mammals and birds have been used within radioecological transfer models (e.g. Brown et al., 2004). As discussed above for reptiles, for most other aquatic organisms, this would not be a valid approach. However, the application of allometry to derive both biological half-life and CR values for marine organisms has been the subject of some studies (see review by Vives i Batlle et al., 2009).

Table 4 presents allometric parameters describing $T_{1/2B}$ for Tc, Cs, Pu and Am in marine organisms. To parameterise these models, data were available for fish, crustaceans, molluscs, macroalgae/seaweed and plankton (Vives i Batlle et al., 2007, 2008). The mean value of b for the $T_{1/2B}$ relationships was 0.16 ± 0.03 . Vives i Batlle et al. (2009) presented allometric $T_{1/2B}$ expressions for Sr and I from previous studies. However, these expressions were taken from USDOE (2002) and were for terrestrial animals; they are not valid for most marine organisms (although they may be applicable to marine birds and mammals).

Table 4

Allometric parameters describing the biological half-life (d) of radionuclides in marine organisms (adapted from Vives i Batlle et al., 2009).

	Tc	Cs	Pu	Am
a	98	54	631	251
b	0.15	0.17	0.20	0.13
R^2	0.72	0.92	0.91	0.76
p	0.03	0.04	0.2	0.05
n	6	4	3	5

For CR, the mean (\pm SD) value of b for marine organisms, calculated across all those elements (Ac, Am, Ce, Cm, Mn, Pa, Pm/Eu, Po, Pu, Ra, Ru, Th and Zr), was -0.26 ± 0.09 and the fitted relationship had an R^2 value >0.7 (Vives i Batlle et al., 2009) (see Table 5). For the individual radionuclides with an R^2 value in Table 5 greater than 0.7 (i.e. Am, Ce, Cm, Pm/Eu, Pu, Ra, Ru, and Th), Vives i Batlle et al. (2009) also found a significant linear relationship between $\log_{10}(a)$ and $\log_{10}(K_d)$ (i.e. the sediment-water distribution coefficient) (Fig. 5), showing particle-reactive (or sediment seeking) radionuclides to have the highest $\log_{10}(a)$ values. The relationship between $\log_{10}(a)$ and $\log(K_d)$ improved when Ru and Pm/Eu were removed though the authors had no scientific justification for doing this. With the addition of Ac, Mn, Pa, Po and Zr (which have R^2 values <0.7 in Table 5), the trend became weaker ($\log_{10}(a) = 0.38 \times \log_{10}(K_d) - 1.6$; $R^2 = 0.50$; $n = 13$), but the statistical significance remained strong ($p = 0.007$) (Vives i Batlle et al., 2009).

For CR, if we accept that the exponent will approximate to -0.25 for cations, then the relationship between $\log_{10}(a)$ and $\log(K_d)$ observed by Vives i Batlle et al. (2009) presents a potential opportunity to estimate CR values when data are lacking, assuming K_d is known.

An anomaly when considering the allometric relationships available for marine organisms with those derived for terrestrial organisms is that the actinide and lanthanide elements scale with an exponent similar to other elements for marine organisms, whereas this is not the case for terrestrial organisms. A potential reason for this with respect to CR for marine organisms is the relationship between $\log_{10}(K_d)$ and $\log_{10}(a)$ which implies that the more particle-reactive the nuclide is, the more strongly it attaches to organic matter (e.g. food). As the ingestion of food is related to metabolism, and hence, to body mass via Kleiber's law, the resultant CR will scale allometrically. Obviously, this argument applies to animals, but not to plants.

Pan and Wang (2008) have previously made similar suggestions with respect to metabolically driven allometry of the ingestion rate and the uptake of Cd and Zn by marine invertebrates. However, Vives i Batlle et al. (2009) suggest that for some elements, the process could also be surface-area driven (e.g. passive sorption of radionuclides on body surfaces, such as mollusc shells). Supporting this suggestion, it has been shown that dead cells of phytoplankton had the same uptake of transuranic elements as live cells, indicating that the uptake process is passive (Fisher et al., 1983). Similarly, Ginn and Fein (2008) observed that metal adsorption on bacteria was the same for many different bacteria species, suggesting that the surface area rather than the biology of the bacteria is important in metal transfer. For a perfectly isometrically scaled organism, all surface area-based properties change with mass to the power of $2/3$ (Galilei, 1638), this would result in the CR scaling to $M^{-0.33}$ instead of $M^{-0.25}$.

4. An alternative to the CR_{wo-media} model for wildlife?

There are many cases where empirical data to derive CR_{wo-media} are lacking. For instance, in the revised version of the ERICA Tool (released 2014 <http://www.ERICA-tool.com/news/news-articles/2014/ERICA-assessment-tool-version-1.2/>) (Brown et al., 2008), data were available for only 622 of 1521 required values. Furthermore, this parameter is highly variable as it incorporates many processes and will be largely determined by site-specific characteristics (e.g. Beresford et al., 2008a; Johansen et al., 2012; Wood et al., 2009b, 2013; Yankovich et al., 2010b).

Soil-to-plant transfer of elements of radiological interest has been related to plant evolutionary history, or phylogeny (Willey, 2010), including for Cs (Broadley et al., 1999; Willey et al., 2005),

Table 5
Allometric parameters for CR ($L\ kg^{-1}$) in marine organisms (adapted from Vives i Batlle et al., 2009); a and b are the constant and exponent for the allometric model. The authors describe those relationships with R^2 values of <0.7 as 'potentially approaching statistical significance'. For all elements $n = 9$.

	Pu	Am	Ru	Ce	Pm/Eu	Ra	Th	Cm	Mn	Zr	Po	Ac	Pa
a	209	251	29	468	1549	85	562	363	4365	269	5495	380	58
b	-0.30	-0.28	-0.46	-0.25	-0.18	-0.11	-0.27	-0.27	-0.13	-0.30	-0.17	-0.18	-0.17
R^2	0.80	0.78	0.75	0.72	0.72	0.79	0.89	0.85	0.46	0.68	0.64	0.58	0.53
p	0.001	0.002	0.002	0.004	0.004	0.001	0.0002	0.0004	0.04	0.007	0.01	0.02	0.03

Note: $^{154,155}\text{Eu}$ and ^{147}Pm were assumed to be biological analogues.




Fig. 5. Linear trend between $\log_{10}(a)$ for the CR and sediment $\log_{10}(K_d)$ for marine ecosystems, both with (left) and without (right) Ru, Eu and Pm. Error bars represent the standard error of $\log_{10}(a)$ (adapted from Vives i Batlle et al. (2009)).

Sr (Willey and Fawcett, 2005a), Ru (Willey and Fawcett, 2006), Cl (Willey and Fawcett, 2005b) and Co (Willey and Wilkins, 2008) (Fig. 6). Such phylogenetic relationships present a potential approach to enable predictions of transfer, with some scientific justification, for taxonomic groups for which there are no data either at the generic or site-specific level (Willey, 2010). The potential to derive phylogenetic relationships for radionuclide transfer to other organism types was demonstrated by Jeffree et al. (2010, 2013). Jeffree et al. suggested that the transfer of a number of radionuclides to marine teleost and chondrichthyan fishes, and to the amphioxus (fish like chordate) species, *Branchiostoma lanceolatum*, was influenced by phylogeny. However, the work of Jeffree et al. was based upon the results of laboratory studies that only considered the exposure pathway from radionuclide contaminated seawater. Although this usefully removes the influences of many confounding factors, it is not directly applicable to environmental conditions as food chain transfer was excluded.

Beresford et al. (2013) used the approach of Willey (2010) to analyse data of Cs transfer to freshwater fish species. Starting with the database on radionuclide transfer to freshwater organisms as described by Yankovich et al. (2013) (see also Coplestone et al., 2013), they compiled a data set of 597 entries covering 53 species of freshwater fish from 67 sites. The 53 fish species all belonged to the class Actinopterygii (ray-finned fishes), with 10 taxonomic orders, 14 families and 33 genera being represented in the dataset.

The Residual Maximum Likelihood (REML) fitting of a mixed-model regression (as described by Willey, 2010) was used to analyse the dataset. The output consists of a mean value for each species on a common scale after REML adjustment (the fixed factor), taking account of the effect of the random factor (i.e. inter-site variation). This provides a method for statistically accounting for as much of the effect of site as possible within the collated data. The mean value output for each species provides a relative scaling value. The REML-adjusted means for different taxonomic groups of freshwater fish are presented in Table 6.

The results demonstrated differences in Cs transfer to freshwater fish based upon phylogenetically-derived taxonomic




Fig. 6. Extrapolated concentration ratios for radionuclides of chlorine, caesium, ruthenium, cobalt and strontium. Published REML-adjusted data (Broadley et al., 1999; Willey and Fawcett, 2005a,b, 2006; Willey et al., 2005; Willey and Wilkins, 2008), have been transformed to have a geometric mean of IAEA recommended transfer value for 'grass and herbs' (IAEA, 2014) then divided into phylogenetic groups of the Angiosperm Phylogeny Group III phylogeny for flowering plants (1 = Monocotyledonous plants, 2 = Rosids, 3 = Caryophyllids, 4 = Asterids. Geomeans with 95% confidence intervals. Cl $n = 106$ species, Ru $n = 114$, Sr $n = 155$, Cs $n = 256$, Co $n = 241$).

groupings, though they did not indicate a phylogenetic trend as previously seen for plants (e.g. Fig. 6). However, although a relatively large dataset was considered, data were only available for 53 of the total 11,952 freshwater species (Nelson, 2006), representing only 10 orders and one class. Earlier analyses, which have suggested phylogenetic relationships for the transfer of radionuclides to plants (Willey, 2010) and marine fish (Jeffree et al., 2010, 2013), have included species encompassing much wider evolutionary time scales (e.g. >500 million years in the case of marine fish).

Beresford et al. (2013) suggested that the REML-adjusted means potentially provide a more refined approach than the CR_{wo-media} model. By taking into account inter-site variation, the REML-adjusted means in effect provide a mechanism of accounting for site-specific variables (e.g. K concentrations in water in the case of Cs transfer to fish). To independently test this hypothesis, data from

27 Finnish lakes (STUK, 2012; <http://bit.ly/1xDJQu4>), which had not been used in the model calibration, were analysed. Data were available for four or more fish species at each lake with a total of 11 fish species being represented. As *Perca fluviatilis* was present at all 27 sites and was also well represented within the dataset used for the REML analysis, it was used as the ‘known species’ from which to calculate activity concentrations for the other species (treated as ‘unknowns’). This allowed 100 predictions to be made.

There was relatively good agreement between predicted and measured values, with a linear regression fit yielding an R^2 of 0.83 ($p < 0.001$) and a slope (\pm standard error) of 0.98 ± 0.04 ($p < 0.001$) (Beresford et al., 2013). The intercept was not significantly different to zero.

Predictions were considerably better than if the appropriate feeding group (i.e. benthic feeding, piscivorous or forage feeding),

Table 6
REML-adjusted means for different taxonomic groups; these are relative values and not absolute values of CR_{wo-water} (reproduced from Beresford et al., 2013).

Order		Family		Genus		Species	
Lepisosteiformes	6.8	Lepisosteidae	6.9	<i>Lepisosteus</i>	7.6	<i>osseus</i>	7.4
Amiiformes	3.2	Amiidae	2.9	<i>Amia</i>	3.6	<i>calva</i>	3.0
Anguilliformes	1.8	Anguillidae	1.9	<i>Anguilla</i>	1.9	<i>anguilla</i>	2.1
Clupeiformes	5.0	Clupeidae	4.9	<i>Dorosoma</i>	4.7	<i>cepedianum</i>	4.3
				<i>Alosa</i>	4.0	<i>pseudoharengus</i>	3.7
Osmeriformes	5.6	Osmeridae	5.5	<i>Osmerus</i>	4.5	<i>mordax</i>	4.2
Salmoniformes	5.5	Salmonidae	5.5	<i>Coregonus</i>	3.8	<i>clupeaformis</i>	3.9
						<i>hoyi</i>	5.0
						<i>artedi</i>	3.5
						<i>spp.</i>	3.2
				<i>Oncorhynchus</i>	8.1	<i>kisutch</i>	9.3
						<i>mykiss</i>	6.5
						<i>tschawytscha</i>	8.9
				<i>Salmo</i>	6.0	<i>trutta</i>	5.8
				<i>Salvelinus</i>	7.8	<i>alpinus</i>	7.9
						<i>fontinalis x namaycush</i>	5.6
						<i>namaycush</i>	8.0
						<i>siscowet</i>	10.8
				<i>Stenodus</i>	5.4	<i>leucichthys</i>	5.6
Esociformes	8.1	Esocidae	8.3	<i>Esox</i>	8.5	<i>lucius</i>	8.8
						<i>niger</i>	3.0
Cypriniformes	4.6	Catostomidae	5.3	<i>Catostomus</i>	5.0	<i>catostomus</i>	4.5
						<i>commersoni</i>	4.4
				<i>Moxostoma</i>	4.7	<i>aureolum</i>	4.2
		Cyprinidae	4.5	<i>Abramis</i>	4.5	<i>brama</i>	4.8
				<i>Carassius</i>	4.4	<i>auratus</i>	4.3
						<i>carassius</i>	4.9
				<i>Cyprinus</i>	1.2	<i>carpio</i>	1.2
					6.2	<i>crysoleucas</i>	5.7
				<i>Rutilus</i>	4.8	<i>rutilus</i>	5.0
				<i>Scardinius</i>	4.4	<i>erythrophthaimus</i>	4.7
				<i>Notropis</i>	3.9	<i>hudsonius</i>	3.6
				<i>Gobio</i>	6.2	<i>gobio</i>	6.5
				<i>Tinca</i>	3.1	<i>tinca</i>	3.2
Siluriformes	7.6	Ictaluridae	7.6	<i>Ictalurus</i>	6.2	<i>punctatus</i>	5.7
						<i>spp.</i>	5.1
Perciformes	8.6	Centrarchidae	7.0	<i>Ambloplites</i>	14.2	<i>rupestris</i>	13.8
				<i>Lepomis</i>	4.1	<i>gulosus</i>	5.3
						<i>macrochirus</i>	3.7
						<i>gibbosus</i>	3.7
				<i>Micropterus</i>	9.6	<i>microlophus</i>	2.9
						<i>dolomieu</i>	8.7
						<i>salmoides</i>	8.5
				<i>Pomoxis</i>	8.3	<i>annularis</i>	7.3
						<i>nigromaculatus</i>	9.3
		Moronidae	7.5	<i>Morone</i>	9.1	<i>chrysops</i>	8.8
		Percidae	9.0	<i>Perca</i>	9.4	<i>flavescens</i>	7.3
						<i>fluviatilis</i>	10.5
				<i>Sander</i>	10.0	<i>lucioperca</i>	7.8
						<i>canadensis</i>	12.3
						<i>vitreus</i>	11.8
				<i>Gymnocephalus</i>	1.9	<i>cernuus</i>	2.0
		Sciaenidae	15.9	<i>Aplodinotus</i>	11.2	<i>grunniens</i>	10.3

geometric mean $CR_{\text{wo-water}}$ values from Yankovich et al. (2013) are used to predict the ^{137}Cs activity concentrations in the fish. The $CR_{\text{wo-water}}$ values generally resulted in under-predictions, with a linear regression of predicted-to-measured activity concentrations yielding a slope of only 0.31.

In Fig. 7, we further test the model of Beresford et al. (2013) against recent data for three lakes in northern England (Barnett et al., 2015). Across the three lakes, data were available for vendace (*Coregonus albula*), Northern pike (*Esox lucius*), ruffe (*Gymnocephalus cernua*), perch (*Perca fluviatilis*), roach (*Rutilus rutilus*) and brown trout (*Salmo trutta*). Replication for each species ranged from 6 to 20, although not all species were sampled in each lake. Perch was present at all sites and was hence used as our ‘known’ species. Activity concentrations in the other species were then predicted as:

Cs concentration in ‘unknown’ species

$$= \frac{\text{REML} - \text{adjusted mean for unknown species}}{\text{REML} - \text{adjusted mean for perch}} \times \text{concentration in perch}$$

The results of this comparison were again encouraging, yielding a significant linear regression ($p < 0.03$) with a slope of 0.82, an R^2 value of 0.58, and an intercept not significantly different from zero. Water data were not available for these lakes so a comparison of predictions with those using CR values was not possible. However, this demonstrates a potential advantage of the alternative (‘REML’) approach, in that water concentrations, which are prone to comparatively large temporal variations, are not required.

5. Stoichiometry and ionomics

The terms, ‘ecological stoichiometry’ and ‘ionomics’, are closely related. Ecological stoichiometry (or ecostochiometry) is ‘the study of the balance of chemical elements in components, interactions, and processes in ecosystems’ (Sternler and Elser, 2002) and has its origins in aquatic ecology. Ionomics is the study of the total elemental composition of an organism, the ionome (Salt, 2004), and the term has been used mostly within terrestrial botany.

5.1. Previous and current uses

Early studies in aquatic ecosystems mainly used ecological stoichiometry to investigate food web dynamics (Lindeman, 1942; Lotka, 1925), and carbon cycling and energy flows (Odum, 1960). Redfield (1958) recognised that the elemental composition of seawater and plankton was fairly constant across wide oceanographic areas and that biological processes could be controlled by elements other than C, such as N and P and trace elements. Since then, most studies have continued to focus on C, N and P and primary producers. More recently, ionomics studies focussing on macro- and micronutrients have been used to explore plant phylogenetics and physiology (Broadley et al., 2004; White et al., 2012). Elements other than C, N and P have also been used in the natural sciences to trace the origins of organisms or materials in the environment. For example, elemental signatures have been used to identify the biological origin of raw materials of biodiesels (Paredes et al., 2014) and the geographic origin of a range of food products (Gonzalez et al., 2009). Multivariate methods (e.g. principal component analysis (PCA) and discriminant analysis), based on elemental composition, have been used to distinguish between ecosystem components and trophic groups in a coastal ecosystem (Kumblad and Bradshaw, 2008), different angiosperm families in grassland habitats (White et al., 2012) and freshwater invertebrate taxa (Karimi and Folt, 2006). However, such approaches have not been widely used in radioecology.

In radioecology, it is known from field and experimental data that concentrations of certain elements/radionuclides are closely correlated to each other, to such a degree that they may be used as analogues for each other. The most common examples are radio-Sr/Ca and radio-Cs/K. The concentration of Sr or Cs in fish can be estimated from the water concentrations of Ca or K, respectively (e.g. Kryshev and Ryabov, 2000; Smith et al., 2009) and the Sr concentration in milk can be predicted from Ca concentrations in the diet and milk (Beresford et al., 1998b). Similar relationships between Ca and P (e.g. Hendrixson et al., 2007) or Ra and Ba (e.g. Gonnee et al., 2013) offer potential for further exploration.

Another approach that has been used in radioecology is based on the earlier ecological theory of organic matter (carbon) and




Fig. 7. A comparison of measured mean Cs concentrations in fish of different species from three different lakes in northern England with values predicted using measured data for perch and the REML-adjusted means (see Table 6). Error bars are minimum and maximum values, and the solid line represent the 1:1 relationship between predicted and measured values.

energy flows in ecosystems. Radionuclide or element concentrations in ecosystem components are normalised to their C content, i.e. element:C ratios are calculated, based on the assumption that many elements are stoichiometrically related to the carbon content because of their role in metabolism and structural components of the organism (Elser et al., 2000; Bradshaw et al., 2012). Ecosystem models based on carbon flows are thus constructed as the basis for radionuclide/element transfer models, using CR values based on C-normalised element concentrations (Kumblad et al., 2006; Bradshaw et al., 2012; Konovalenko et al., 2014).

5.2. Natural variation in element composition

A key issue in ecostochiometry is whether an element is essential or non-essential. The ability of organisms to maintain constant body concentrations, despite changing concentrations in the environment and/or their resource supply, (homeostasis) is generally assumed to be weak for autotrophs and strong for heterotrophs (Sterner and Elser, 2002). Therefore, plant and algae stoichiometry is thought to more closely reflect that of the environment than animals, though this is much debated (e.g. Persson et al., 2010). The degree of homeostasis has been shown to vary depending on whether elements are macronutrients, essential micronutrients, or non-essential elements (Karimi and Folt, 2006; Bradshaw et al., 2012). Organisms often take up the necessary amounts of trace elements from their food within the 'window of essentiality' (Hopkin, 1989) to ensure essential levels but avoid toxic concentrations in the body.

These principles could be made use of when using ecostochiometry in extrapolation. For example, for autotrophs, extrapolation from environmental concentrations or ratios is probably more appropriate, whereas for heterotrophs, extrapolation within taxonomic groups may be more relevant (Karimi and Folt, 2006; White et al., 2012). Higher trophic levels may have a more similar elemental composition to their food than lower trophic levels, and stoichiometric ratios may shift between abiotic-biotic components and from primary producers to primary consumers (Bradshaw et al., 2012). In cases of homeostasis, element ratios will be constrained by the biology and ecology of the organism/ecosystem. A better understanding of these natural ranges would greatly help in our ability to predict element concentrations.

6. Discussion

We will never have data to populate all of the potential radioecological modelling parameters for wildlife (or indeed human) assessments. Therefore, we need robust extrapolation approaches which allow us to make best use of our available knowledge. In this study, we have reviewed, developed, tested and validated some of these extrapolation approaches.

Application of the transfer coefficient concept has been widely adopted in radioecology to estimate activity concentrations in farm animal products; applicability to wildlife assessments has also been proposed. However, this parameter is, in part, dependent upon the dry matter intake of animals and this gives a false impression of differences in transfer between different species. The concentration ratio ($CR_{\text{product-diet}}$ or $CR_{\text{wo-diet}}$) is a more generic parameter, and should enable the more abundant data for farm animal species to be applied to wild species.

Allometric expressions describing radionuclide biological half-life have previously been proposed for some elements. Beresford and Vives i Batlle (2013) present an approach, based on allometry, to predict acceptable $T_{1/2B}$ for a number of radionuclides and over orders of magnitude range in animal live-weight. However, we acknowledge that for some radionuclides (i.e. Am, Ce, Eu, Pu and

Th), $T_{1/2B}$ has been reported to scale to *circa* $M^{0.8}$ and not the *circa* 0.25, as assumed in Beresford and Vives i Batlle (2013). To fully exploit the potential of allometric models, the reasons for this need to be explored.

Beresford and Wood (2014) demonstrated that the relationship derived by Beresford and Vives i Batlle (2013) for homoeothermic vertebrates is not directly applicable to poikilothermic reptiles. However, it is possible to parameterise the Beresford & Vives i Batlle model using reptile-specific values to obtain reasonable agreement between measured and predicted $T_{1/2B}$ values. Similarly it is possible to adapt the model parameters for other animal groupings using published information (e.g. as we have demonstrated above for different categories of mammals).

For aquatic ecosystems, the relationship between $\log_{10}(a)$ (on the allometric relationship for CR) and $\log(K_d)$ observed by Vives i Batlle et al. (2009) presents a potential opportunity to estimate CR values when data are lacking, assuming K_d is known.

The commonly used $CR_{\text{wo-media}}$ approach to estimating the radionuclide activity concentrations in wildlife is open to criticism, as $CR_{\text{wo-media}}$ values can be highly variable, largely due to site-specific factors. The analyses of available data using the REML analysis, as demonstrated by Beresford et al. (2013), should compensate for inter-site variation, assuming sufficient data are available for the analysis. For freshwater fish the outputs of the REML analysis predicted ^{137}Cs activity concentrations in a new independent dataset well. We recommend that this approach of producing relative values be further investigated and developed for other radionuclides and across a wider range of organisms and ecosystems. A disadvantage of the approach is that it requires relatively large datasets which must meet specific criteria. However, recent data compilations (Howard et al., 2013; Yankovich et al., 2013; Copplestone et al., 2013) should enable similar analysis to be conducted for a number of elements for terrestrial, marine and freshwater species.

The ICRP (2009) suggested identifying a series of terrestrial, freshwater and marine sites from which samples of their Reference Animals and Plants (RAPs) could be sampled and analysed to serve as 'points of reference'. Such studies have been initiated and results are starting to be published (Barnett et al., 2014). However, such data are highly site-specific, potentially limiting their wider applicability. The application of the REML approach to data, such as those presented by Barnett et al. (2014), to derive relative values for different organisms should provide a more generic set of 'reference data'. A collaboration of the TREE (<http://www.ceh.ac.uk/tree>) and COMET (www.comet-radioecology.org) projects is taking this area of research forward, including the focused sampling of species falling within the RAP definitions at a number of terrestrial sites in different countries (Japan, Norway, Spain, UK and Ukraine).

Ecological stoichiometry shows potential as an extrapolation method in radioecology, either from one element to another or from one species to another. This will be most successful when using data from elements, organisms (taxa, functional groups) and/or ecosystems that are as similar as possible to each other. At the very least, stoichiometry could be used to set limits on the range of possible element concentrations; there are biological and ecological limits to many element concentrations/ratios, even if there is variability.

Multi-element datasets are becoming more common as analytical techniques (e.g. ICP-MS) become easier, faster and cheaper. In the near future we will have access to a much larger amount of data (e.g. Baxter et al., 2007; <http://www.ionomicshub.org/home/PiiMS>), which can be used to test stoichiometric assumptions and theories, and to develop extrapolation methods. Such multi-element datasets could be analysed to identify similarities in element profiles between or within taxa, trophic level,

functional group, habitat, geographical region, season or year, depending on the temporal and spatial resolution of the data. This would identify the most promising groupings for which extrapolation methods could subsequently be tested, using simple element-specific ratios or predictive multivariate methods, such as those developed by the food industry. There, multi-element profiles are usually compared against known profiles using a range of pattern recognition methods (mostly multivariate techniques). Neural networks and regression trees have also been used to separate samples by fingerprinting techniques (see Gonzalez et al., 2009) and many of these can also be used in a predictive rather than a descriptive or analytical way. The TREE project (<http://www.ceh.ac.uk/tree>) is taking forward this area of research, primarily for crop plants, but with some consideration of wildlife (and potentially farm animals).

Acknowledgements

Much of the work described here was conducted within the EC funded STAR Network of Excellence (<http://www.star-radioecology.org/>) and is, in part, based upon one of the projects deliverable reports (Beresford et al., 2014). We are grateful to all our colleagues who contributed to this wider report, including Justin Brown (NRPA), Ali Hosseini (NRPA) and Claire Wells (NERC-CEH). The biological half-life database, in part used here, was compiled within framework of the work programme of the IAEA MODARIA Working Group 8 (<http://www-ns.iaea.org/projects/modaria/default.asp?l=116>); the database will subsequently be published in its entirety. The inputs of N. Willey, M.D. Wood, T.L. Yankovich and N.A. Beresford to this paper were, in part, supported by the TREE project (funded by the NERC, Environment Agency and Radioactive Waste Management Ltd.) (<http://www.ceh.ac.uk/tree>). C. Bradshaw was partly funded by the Baltic Sea Adaptive Management programme at Stockholm University.

Appendix A. Supplementary data

Supplementary data (see review by Beresford et al. 2015) related to this article can be found at <http://dx.doi.org/10.1016/j.jenvrad.2015.03.022>.

References

- Agutter, P.S., Tuszynski, J.A., 2011. Analytic theories of allometric scaling. *J. Exp. Biol.* 214, 1055–1062.
- Avila, R., Beresford, N.A., Agüero, A., Broed, R., Brown, J., Iospje, M., Robles, B., Suárez, A., 2004. Study of the uncertainty in estimation of the exposure of non-human biota to ionizing. *J. Radiol. Prot.* 24, A105–A122.
- Barnett, C.L., Beresford, N.A., Walker, L.A., Baxter, M., Wells, C., Copplestone, D., 2014. Transfer parameters for ICRP reference animals and plants collected from a forest ecosystem. *Radiat. Environ. Biophys.* 53, 125–149.
- Barnett, C.L., Wells, C., Thacker, S., Guyatt, H.J., Fletcher, J.M., Lawlor, A., Beresford, N.A., 2015. Elemental Concentrations in Fish from Lakes in Northwest England. NERC-Environmental Information Data Centre. <http://doi.org/10.5285/ed90df1b-462c-46bb-afb-59794fb03f6b>.
- Baxter, I., Ouzzani, M., Orcun, S., Kennedy, B., Jandhyala, S.S., Salt, D.E., 2007. Purdue ionomics information management system. An integrated functional Genomics Platform. *Plant Physiol.* 143, 600–611.
- Beresford, N.A., Crout, N.M.J., Mayes, R.W., Howard, B.J., Lamb, C.S., 1998a. Dynamic distribution of radioisotopes of cerium, ruthenium and silver in sheep tissues. *J. Environ. Radioact.* 38, 317–338.
- Beresford, N.A., Mayes, R.W., Hansen, H.S., Crout, N.M.J., Hove, K., Howard, B.J., 1998b. Generic relationship between calcium intake and radiostrontium transfer to milk of dairy ruminants. *Radiat. Environ. Biophys.* 37, 129–131.
- Beresford, N.A., Broadley, M.R., Howard, B.J., Barnett, C.L., White, P.J., 2004. Estimating radionuclide transfer to wild species - data requirements and availability for terrestrial ecosystems. *J. Radiol. Prot.* 24, A89–A103.
- Beresford, N.A., Mayes, R.W., Barnett, C.L., Howard, B.J., 2007. The transfer of radionuclides to ewes through a breeding cycle - an illustration of the pitfalls of the transfer coefficient. *J. Environ. Radioact.* 98, 24–35.
- Beresford, N.A., Barnett, C.L., Brown, J., Cheng, J.-J., Copplestone, D., Filistovic, V., Hosseini, A., Howard, B.J., Jones, S.R., Kamboj, S., Kryshev, A., Nedveckaite, T., Olyslaegers, G., Saxén, R., Sazykina, T., Vives i Batlle, J., Vives-Lynch, S., Yankovich, T., Yu, C., 2008a. Inter-comparison of models to estimate radionuclide activity concentrations in non-human biota. *Radiat. Environ. Biophys.* 47, 491–514.
- Beresford, N.A., Barnett, C.L., Howard, B.J., Scott, W.A., Brown, J.E., Copplestone, D., 2008b. Derivation of transfer parameters for use within the ERICA Tool and the default concentration ratios for terrestrial biota. *J. Environ. Radioact.* 99, 1393–1407.
- Beresford, N.A., Barnett, C.L., Beaugelin-Seiller, K., Brown, J.E., Cheng, J.-J., Copplestone, D., Gaschak, S., Hingston, J.L., Horyna, J., Hosseini, A., Howard, B.J., Kamboj, S., Kryshev, A., Nedveckaite, T., Olyslaegers, G., Sazykina, T., Smith, J.T., Telleria, D., Vives i Batlle, J., Yankovich, T.L., Heling, R., Wood, M.D., Yu, C., 2009. Findings and recommendations from an international comparison of models and approaches for the estimation of radiological exposure to non-human biota. *Radioprotection* 44, 565–570.
- Beresford, N.A., Barnett, C.L., Brown, J.E., Cheng, J.-J., Copplestone, D., Gaschak, S., Hosseini, A., Howard, B.J., Kamboj, S., Nedveckaite, T., Olyslaegers, G., Smith, J.T., Vives i Batlle, J., Vives-Lynch, S., Yu, C., 2010. Predicting the radiation exposure of terrestrial wildlife in the Chernobyl exclusion zone: an international comparison of approaches. *J. Radiol. Prot.* 30, 341–373.
- Beresford, N.A., Vives i Batlle, J., 2013. Estimating the biological half-life for radionuclides in homeothermic vertebrates: a simplified allometric approach. *Radiat. Environ. Biophys.* 52, 505–511.
- Beresford, N.A., Yankovich, T.L., Wood, M.D., Fesenko, S., Andersson, P., Muikku, M., Willey, N.J., 2013. A new approach to predicting environmental transfer of radionuclides to wildlife taking account of inter-site variation using residual maximum likelihood mixed-model regression: a demonstration for freshwater fish and caesium. *Sci. Total Environ.* 463–464, 284–292.
- Beresford, N.A., Brown, J.E., Vives i Batlle, J., Hosseini, A., Bradshaw, C. (Eds.), 2014. Evaluation of Extrapolation Approaches to Provide Radioecological Parameters. STAR Deliverable (D-N°3.2). (Contract Number: Fission-2010-3.5.1-269672). Available from: <http://bit.ly/1zisRbf>.
- Beresford, N.A., Wood, M.D., 2014. A new simplified allometric approach for predicting the biological half-life of radionuclides in reptiles. *J. Environ. Radioact.* 138, 116–121.
- Beresford, N.A., Beaugelin-Seiller, K., Burgos, J., Cujic, M., Fesenko, S., Kryshev, A., Pachal, N., Real, A., Su, B.S., Tagami, K., Vives i Batlle, J., Vives-Lynch, S., Wells, C., Wood, M.D., December 2015. Radionuclide biological half-life values for terrestrial and aquatic wildlife. *J. Environ. Radioact.* 150, 270–276. <http://dx.doi.org/10.1016/j.jenvrad.2015.08.018>. ISSN 0265-931X.
- Bradshaw, C., Kautsky, U., Kumblad, L., 2012. Ecological stoichiometry and multi-element transfer in a coastal ecosystem. *Ecosystems* 15, 591–603.
- Broadley, M.R., Willey, N.J., Meade, A., 1999. The effect of taxonomic position on radiocaesium uptake by flowering plants. *Environ. Pollut.* 106, 341–349.
- Broadley, M.R., Bowen, H.C., Cotterill, H.L., Hammond, J.P., Meacham, M.C., Mead, A., White, P.J., 2004. Phylogenetic variation in the shoot mineral concentration of angiosperms. *J. Exp. Bot.* 55, 321–336.
- Brown, J., Simmonds, J., 1995. FARMLAND a Dynamic Model for the Transfer of Radionuclides through Terrestrial Foodchains. NRPB-R273, Chilton, UK.
- Brown, J., Børretzen, P., Dowdall, M., Sazykina, T., Kryshev, I., 2004. The derivation of transfer parameters in the assessment of radiological impacts on arctic marine biota. *Arctic* 57, 279–289.
- Brown, J.E., Alfonso, B., Avila, R., Beresford, N.A., Copplestone, D., Pröhl, G., Ulanovsky, A., 2008. The ERICA tool. *J. Environ. Radioact.* 99, 1371–1383.
- Brown, J.E., Beresford, N.A., Hosseini, A., 2013. Approaches to providing missing transfer parameter values in the ERICA Tool - how well do they work? *J. Environ. Radioact.* 126, 399–411.
- Brown, J.H., Gupta, V.K., Li, B.-L., Milne, B.T., Restrepo, C., West, G.B., 2002. The fractal nature of nature: power laws, ecological complexity and biodiversity. *Philos. Trans. Roy. Soc. B* 357, 619–626.
- Chaplow, J.S., Beresford, N.A., Barnett, C.L., 2015. Post Chernobyl surveys of radionuclides in soil, vegetation, wildlife and fungi in Great Britain. *Earth Syst. Sci. Data* 7, 215–221. <http://www.earth-syst-sci-data.net/7/215/2015/essd-7-215-2015.pdf>.
- Copplestone, D., Beresford, N.A., Brown, J.E., Yankovich, T., 2013. An international database of radionuclide concentration ratios for wildlife: development and uses. *J. Environ. Radioact.* 126, 288–298.
- Coughtrey, P.J., Thorne, M., 1983. Radionuclide Distribution and Transport in Terrestrial and Aquatic Ecosystems e a Critical Review of Data, vol. 1. A.A. Balkema, Rotterdam.
- Coughtrey, P.J., Jackson, D., Thorne, M., 1983. Radionuclide Distribution and Transport in Terrestrial and Aquatic Ecosystems a Critical Review of Data, vol. 3. AA Balkema, Rotterdam.
- DiGregorio, D., Kitchings, T., Van Voris, P., 1978. Radionuclide transfer in terrestrial animals. *Health Phys.* 34, 3–31.
- Elser, J.J., Sterner, R.W., Gorokhova, E., Fagan, W.F., Markow, T.A., Cotner, J.B., Harrison, J.F., Hobbie, S.E., Odell, G.M., Weider, L.J., 2000. Biological stoichiometry from genes to ecosystems. *Ecol. Lett.* 3, 540–550.
- Fesenko, S., Isamov, N., Barnett, C.L., Beresford, N.A., Howard, B.J., Sanzharova, N., Fesenko, E., 2015. Review of Russian language studies on radionuclide behaviour in agricultural animals: biological half-lives. *J. Environ. Radioact.* 142, 136–151.

- Farrell-Gray, C.C., Gotelli, N.J., 2005. Allometric exponents support a 3/4-power scaling law. *Ecology* 86, 2083–2087.
- Fisher, N.S., Bjerregaard, P., Fowler, S.W., 1983. Interactions of marine plankton with transuranic elements. I. Biokinetics of neptunium, plutonium americium and californium in phytoplankton. *Limnol. Oceanogr.* 28, 432–447.
- Franz, R., Hummel, J., Müller, D.W.H., Bauert, M., Hatt, J.-M., Clauss, M., 2011. Herbivorous reptiles and body mass: effect on food intake, digesta retention, digestibility and gut capacity, and a comparison with mammals. *Comp. Biochem. Physiol., Part A* 158, 94–101.
- Furchner, J., Drake, A., 1971. Comparative metabolism of radionuclides in mammals-VI. Retention of Nb-95 in the mouse, rat, monkey and dog. *Health Phys.* 21, 173–180.
- Furchner, J., Richmond, C., Drake, A., 1971. Comparative metabolism of radionuclides in mammals-VII. Retention of Ru-106 in the mouse, rat, monkey and dog. *Health Phys.* 21, 355–365.
- Galeriu, D., Beresford, N.A., Takeda, H., Melintescu, A., Crout, N.M., 2003. Towards a model for the dynamic transfer of tritium and carbon in mammals. *Radiat. Prot. Dosim.* 105, 387–390.
- Galeriu, D., Melintescu, A., Beresford, N.A., Crout, N.M.J., Peterson, R., Takeda, H., 2007. Modelling ^3H and ^{14}C transfer to farm animals and their products under steady state conditions. *J. Environ. Radioact.* 98, 205–217.
- Galilei, G., 1638. *Discorsi e dimostrazione matematiche intorno à due nuove scienze*. Elsevier, Leiden.
- Ginn, B.R., Fein, J.B., 2008. The effect of species diversity on metal adsorption onto bacteria. *Geochim. Cosmochim. Acta* 72, 3939–3948.
- Gonneea, M.E., Mulligan, A.E., Charette, M.A., 2013. Seasonal cycles in radium and barium within a subterranean estuary: implications for groundwater derived chemical fluxes to surface waters. *Geochim. Cosmochim. Acta* 119, 164–177.
- Gonzalez, A., Armenta, S., de la Guardia, M., 2009. Trace-element composition and stable-isotope ratio for discrimination of foods with Protected Designation of Origin. *Trends Anal. Chem.* 28, 1295–1311.
- Hendrixson, H.A., Sterner, R.W., Kay, A.D., 2007. Elemental stoichiometry of freshwater fishes in relation to phylogeny, allometry and ecology. *J. Fish. Biol.* 70, 121–140.
- Higley, K.A., Domotor, S.L., Antonio, E.J., 2003. A kinetic-allometric approach to predicting tissue radionuclide concentrations for biota. *J. Environ. Radioact.* 66, 61–74.
- Higley, K.A., Bytwerk, D.P., 2007. Generic approaches to transfer. *J. Environ. Radioact.* 98, 4–23.
- Higley, K.A., 2010. Estimating transfer parameters in the absence of data. *Radiat. Environ. Biophys.* 49, 645–656.
- Hinton, T.G., Whicker, F.W., Pinder III, J.E., Ibrahim, S.A., 1992. Comparative Kinetics of ^{47}Ca , ^{85}Sr and ^{226}Ra in the Freshwater Turtle, *Trachemys scripta*. *J. Environ. Radioact.* 16, 25–47.
- Hopkin, S.P., 1989. *Ecophysiology of Metals in Terrestrial Invertebrates*. Elsevier Applied Science, London.
- Hoppeler, H., Weibel, E.R., 2005. Editorial Scaling functions to body size: theories and facts. *J. Exp. Biol.* 208, 1573–1574.
- Howard, B.J., Beresford, N.A., Copplestone, D., Telleria, D., Proehl, G., Fesenko, S., Jeffree, R., Yankovich, T., Brown, J., Higley, K., Johansen, M., Mulye, H., Vandenhove, H., Gashchak, S., Wood, M.D., Takata, H., Andersson, P., Dale, P., Ryan, J., Bollhöfer, A., Doering, C., Barnett, C.L., Wells, C., 2013. The IAEA handbook on radionuclide transfer to wildlife. *J. Environ. Radioact.* 121, 55–74.
- IAEA, 1994. Handbook of Transfer Parameter Values for the Prediction of Radionuclide Transfer in Temperate Environments. IAEA-TRS-364. IAEA, Vienna.
- IAEA, 2010. Handbook of Parameter Values for the Prediction of Radionuclide Transfer in Terrestrial and Freshwater Environments. IAEA-TRS-472. IAEA, Vienna.
- IAEA, 2014. Handbook of Parameter Values for the Prediction of Radionuclide Transfer to Wildlife. IAEA-TRS-479. IAEA, Vienna.
- ICRP, 1979. Limits for intakes of radionuclides by workers. ICRP Publication 30 (Part 1). *Ann. ICRP* 2 (3–4).
- ICRP, 1981. Limits for intakes of radionuclides by workers. ICRP Publication 30 (Part 3). *Ann. ICRP* 6 (2–3).
- ICRP, 1988. Limits for intakes of radionuclides by workers: an addendum. ICRP Publication 30 (Part 4). *Ann. ICRP* 19 (4).
- ICRP, 2006. Human alimentary tract model for radiological protection. ICRP Publication 100. *Ann. ICRP* 36 (1–2).
- ICRP, 2009. Environmental protection: transfer parameters for Reference Animals and Plants. ICRP Publication 114. *Ann. ICRP* 39 (6).
- Isaac, N.J.B., Carbone, C., 2010. Why are metabolic scaling exponents so controversial? Quantifying variance and testing hypotheses. *Ecol. Lett.* 13, 728–735.
- Jeffree, R.A., Oberhansli, F., Teyssie, J.-L., 2010. Phylogenetic consistencies among chondrichthyan and teleost fishes in their bioaccumulation of multiple trace elements from seawater. *Sci. Tot. Environ.* 408, 3200–3210.
- Jeffree, R.A., Oberhansli, F., Teyssie, J.-L., 2013. Marine radionuclide transfer factors and phylogeny. *J. Environ. Radioact.* 126, 388–398.
- Johansen, M.P., Barnett, C.L., Beresford, N.A., Brown, J.E., Černe, M., Howard, B.J., Kamboj, S., Keum, D.-K., Smodiš, B., Twining, J.R., Vandenhove, H., Vives i Batlle, J., Wood, M.D., Yu, C., 2012. Assessing doses to terrestrial wildlife at a radioactive waste disposal site: inter-comparison of modelling approaches. *Sci. Tot. Environ.* 427–428, 238–246.
- Karimi, R., Folt, C.L., 2006. Beyond macronutrients: element variability and multi-element stoichiometry in freshwater invertebrates. *Ecol. Lett.* 9, 1273–1283.
- Kitchings, T., DiGregorio, D., Van Voris, P., 1976. A review of the ecological parameters of radionuclide turnover in vertebrate food chains. In: Cushing, C.E., Cutshall, N.H., Fraley, L.F., French, N.R., Murphy, P.G., Sharitz, R.R., Trabalka, J.R., Turner, F.R., Whicker, F.W., Wolfe, D.A. (Eds.), *Radioecology and Energy Resources*. Dowden, Hutchinson & Ross, Inc, Stroudsburg, Pennsylvania, pp. 304–313.
- Kleiber, M., 1932. Body size and metabolism. *Hilgardia* 6, 315–353.
- Kleiber, M., 1947. Body size and metabolic rate. *Physiol. Rev.* 27, 511–541.
- Konovalenko, L., Bradshaw, C., Kumblad, L., Kautsky, U., 2014. Radionuclide transfer in marine coastal ecosystems, a modelling study using metabolic processes and site data. *J. Environ. Radioact.* 133, 48–59.
- Kryshev, A.I., Ryabov, I.N., 2000. A dynamic model of ^{137}Cs accumulation by fish of different age classes. *J. Environ. Radioact.* 50, 221–233.
- Kumblad, L., Kautsky, U., Næslund, B., 2006. Transport and fate of radionuclides in aquatic environments - the use of ecosystem modelling for exposure assessments of nuclear facilities. *J. Environ. Radioact.* 87, 107–129.
- Kumblad, L., Bradshaw, C., 2008. *Element Composition of Biota, Water and Sediment in the Forsmark Area, Baltic Sea: Concentrations, Bioconcentration Factors and Partitioning Coefficients (Kd) of 48 Elements*. Technical report, SKB TR-08-09. Available from: www.skb.se.
- Lindeman, R.L., 1942. The trophic-dynamic aspect of ecology. *Ecology* 23, 399–417.
- Lotka, A.J., 1925. *Elements of Physical Biology*. Williams and Wilkins, Baltimore.
- MacDonald, C.R., 1996. Ingestion Rates and Radionuclide Transfer in Birds and Mammals on the Canadian Shield. Report TR-722 COG-95–551. Atomic Energy of Canada Limited, Ontario.
- Mayes, R.W., Beresford, N.A., Howard, B.J., Vandecasteele, C.M., Stakelum, G., 1996. Use of the true absorption coefficient as a measure of the bioavailability of radiocaesium in ruminants. *Radiat. Environ. Biophys.* 35, 101–109.
- Mertz, W., 1986. *Trace Elements in Human and Animal Nutrition*, fifth ed. Academic Press Inc, London.
- Mertz, W., 1987. *Trace Elements in Human and Animal Nutrition*, fifth ed. Academic Press Inc, London.
- Moss, R.W., Horrill, A.D., 1996. Metabolism of radiocaesium in Red Grouse. *J. Environ. Radioact.* 33, 49–62.
- Müller, H., Pröhl, G., 1993. A dynamic model for assessing radiological consequences of nuclear accidents. *Health Phys.* 64, 232–252.
- Nagy, K.A., 2001. Food Requirements of Wild Animals: Predictive Equations for Free-living Mammals, Reptiles and Birds. *Nutrition Abstracts and Reviews*. In: Series B Livestock Feeds and Feeding, vol. 71, pp. 21R–31R.
- Nagy, K.A., 2005. Field metabolic rate and body size. *J. Exp. Biol.* 208, 1627–1634.
- Nelson, J.S., 2006. *Fishes of the World*, fourth ed. John Wiley & Sons, Inc, Hoboken, New Jersey.
- Ng, Y.C., 1982. A review of transfer factors for assessing the dose from radionuclides in agricultural products. *Nucl. Saf.* 23 (1), 57–71.
- Ng, Y.C., Colsher, C.S., Quinn, D.J., Thompson, S.E., 1977. Transfer Co-efficients for the Prediction of the Dose to Man via the Forage-cow-milk Pathway from Radionuclides Released to the Biosphere. Lawrence Livermore Laboratory, Livermore.
- Ng, Y.C., Colsher, C.S., Thompson, S.E., 1979. Transfer factors for assessing the dose from radionuclides in agricultural products. In: *Proceedings of an International Symposium on the Biological Implications of Radionuclides Released from Nuclear Industries*. IAEA, Vienna.
- Ng, Y.C., Colsher, C.S., Thompson, S.E., 1982. Transfer Coefficients for Assessing the Dose from Radionuclides in Meat and Eggs. Final report, NUREG/CR-2976. US Nuclear Regulatory Commission, Washington, DC.
- Odum, H.P., 1960. Ecological potential and analogue circuits for the ecosystem. *Am. Sci.* 48, 1–8.
- Pan, K., Wang, W.-X., 2008. Allometry of cadmium and zinc concentrations and bioaccumulation in the scallop *Chlamys nobilis*. *Mar. Ecol. Prog. Ser.* 365, 115–126. Prepress Abstract.
- Paredes, E., Can, S.Z., Quélet, C.R., 2014. Examining the possibility of using ratios of element mass fractions as markers for origin determination of soya and canola based biodiesels. Measurement procedure validation to underpin this study. *Fuel* 123, 248–255.
- Persson, J., Fink, P., Goto, A., Hood, J.M., Jonas, J., Kato, S., 2010. To be or not to be what you eat: regulation of stoichiometric homeostasis among autotrophs and heterotrophs. *Oikos* 119, 741–751.
- Peters, R.H., 1983. *The Ecological Implications of Body Size*. Cambridge University Press, Cambridge.
- Redfield, A.C., 1958. The biological control of chemical factors in the environment. *Am. Sci.* 46, 205–221.
- Salt, D.E., 2004. Update on plant ionomics. *Plant Physiol.* 136, 2451–2456.
- Sazykina, T.G., 2000. ECOMOD — An ecological approach to radioecological modelling. *J. Environ. Radioact.* 50, 207–220.
- Scott, D.E., Whicker, F.W., Gibbons, J.W., 1986. Effect of season on the retention of ^{137}Cs and ^{90}Sr by the yellow-bellied slider turtle (*Pseudemys scripta*). *Can. J. Zool.* 64, 2850–2853.
- Sheppard, S.C., 2001. Toxicants in the environment: bringing radioecology and ecotoxicology together. In: Bréchnignac, F., Howard, B.J. (Eds.), *Radioactive Pollutants Impact on the Environment*. EDP Sciences, France, pp. 63–74.
- Sheppard, S.C., 2013. Transfer factors to Whitetail deer: comparison of stomach-content, plant-sample and soil-sample concentrations as the denominator. *J. Environ. Radioact.* 126, 434–437.

- Smith, J., Beresford, N.A., 2005. Chernobyl Catastrophe and Consequences. Praxis Publishing/Springer, Chichester. <http://nora.nerc.ac.uk/id/eprint/18096>.
- Smith, J.T., Sasina, N.V., Kryshev, A.I., Belova, N.V., Kudelsky, A.V., 2009. A review and test of predictive models for the bioaccumulation of radiostrotrium in fish. *J. Environ. Radioact.* 100, 950–954.
- Stara, J.F., Nelson, N.S., Dellarosa, R.J., Bustad, L.K., 1971. Comparative metabolism of radionuclides in mammals: a review. *Health Phys.* 20, 113–137.
- Staton, M.A., Brisbin, I.L., Geiger, R.A., 1974. Some aspects of radiocesium retention in naturally contaminated captive snakes. *Herpetologica* 30, 204–211.
- Sturner, R.W., Elser, J.J., 2002. Ecological Stoichiometry: the Biology of Elements from Molecules to the Biosphere. Princeton University Press, Princeton, New Jersey, p. 584.
- STUK (Radiation and Nuclear Safety Authority), 2012. Cs-137 in Freshwater Fish in Finland after Chernobyl Accident Metadata Record. <http://datacehacuk/metadata/c806d247-bb4e-47d3-b34f-acfd24af0bfb> (last accessed 24.09.12.).
- Thomas, P.A., Sheard, J.W., Swanson, S., 1994. Transfer of ^{210}Po and ^{210}Pb through the lichen-caribou-wolf food chain of Northern Canada. *Health Phys.* 66, 666–677.
- United States Department of Energy (USDOE), 2002. A Graded Approach for Evaluating Radiation Doses to Aquatic and Terrestrial Biota. Technical Standard DOE-STD-1153–2002, Module 1–3. US. Department of Energy, Washington, DC.
- United States Department of Energy (USDOE), 2004. RESRAD-Biota: a Tool for Implementing a Graded Approach to Biota Dose Evaluation. User's Guide, Version 1. United States Department of Energy Standard, pp. 2004–22102. DOE/EH-0676. ISCORS Technical Report.
- United States Nuclear Regulatory Commission (USNRC), 1977. Calculation of Annual Doses to Man from Routine Releases of Reactor Effluents for the Purpose of Evaluating Compliance with 10 CFR 50. Appendix I, Regulatory Guide 1.109. Nuclear Regulatory Commission, Office of Standards Development, Washington, DC.
- Vives i Batlle, J., Wilson, R.C., McDonald, P., 2007. Allometric methodology for the calculation of biokinetic parameters for marine biota. *Sci. Tot. Environ.* 388, 256–269.
- Vives i Batlle, J., Wilson, R.C., Watts, S.J., Jones, S.R., McDonald, P., Vives-Lynch, S., 2008. Dynamic model for the assessment of radiological exposure to marine biota. *J. Environ. Radioact.* 99, 1711–1730.
- Vives i Batlle, J., Wilson, R.C., Watts, S.J., McDonald, P., Craze, A., 2009. Derivation of allometric relationships for radionuclides in marine phyla. *Radioprotection* 44, 7–52.
- Ward, G.M., Johnson, J.E., 1965. The cesium-137 content of beef from dairy and feed lot cattle. *Health Phys.* 11, 95–100.
- Ward, G.M., Johnson, J.E., Stewart, H.F., 1965. Cesium-137 passage from precipitation to milk. In: Klement, A.W. (Ed.), Proceedings of the Second Conference on Radioactive Fallout from Nuclear Weapons Tests. National Technical Information Service, Springfield, pp. 703–710.
- West, G.B., Brown, J.H., Enquist, B.J., 1997. A general model for the origin of allometric scaling laws in biology. *Science* 276, 122–126.
- West, G.B., Brown, J.H., Enquist, B.J., 1999. The fourth dimension of life: fractal geometry and allometric scaling of organisms. *Science* 284, 1677–1679.
- Whicker, F.W., Schultz, V., 1982. Radioecology: Nuclear Energy and the Environment, vol. II. CRC Press, Boca Raton, Florida.
- White, P.J., Broadley, M.R., Thompson, J.A., McNicol, J.W., Crawley, M.J., Poulton, P.R., Johnston, A.E., 2012. Testing the distinctness of shoot ionomes of angiosperm families using the Rothamsted Park Grass Continuous Hay Experiment. *New Phytol.* 196, 101–109.
- Willey, N.J., Fawcett, K., 2005a. A phylogenetic effect on strontium concentrations in angiosperms. *Environ. Exp. Bot.* 57, 258–269.
- Willey, N.J., Fawcett, K., 2005b. Species selection for phytoremediation of $^{36}\text{Cl}/^{35}\text{Cl}$ using angiosperm phylogeny and inter-taxa differences in uptake. *Int. J. Phytotrem.* 7, 295–306.
- Willey, N.J., Tang, S., Watt, N., 2005. Predicting inter-taxa differences in plant uptake of $^{134}/^{137}\text{Cs}$. *J. Environ. Qual.* 34, 1478–1489.
- Willey, N.J., Fawcett, K., 2006. Inter-taxa differences in root uptake of $^{103}/^{106}\text{Ru}$ by plants. *J. Environ. Radioact.* 86, 227–240.
- Willey, N.J., Wilkins, J., 2008. Phylogeny and growth strategy as predictors of differences in cobalt concentrations between plant species. *Environ. Sci. Technol.* 42, 2162–2167.
- Willey, N.J., 2010. Phylogeny can be used to make useful predictions of soil-to-plant transfer factors for radionuclides. *Radiat. Environ. Biophys.* 49, 613–623.
- Wood, M.D., Beresford, N.A., Barnett, C.L., Copplestone, D., Leah, R.T., 2009a. Assessing radiation impact at a protected coastal sand dune site: an inter-comparison of models for estimating the radiological exposure of non-human biota. *J. Environ. Radioact.* 100, 1034–1052.
- Wood, M.D., Leah, R.T., Jones, S.R., Copplestone, D., 2009b. Radionuclide transfer to invertebrates and small mammals in a coastal sand dune ecosystem. *Sci. Total Environ.* 407 (13), 4062–4074.
- Wood, M.D., Beresford, N.A., Howard, B.J., Copplestone, D., 2013. What are the limitations of equilibrium transfer parameters? *J. Environ. Radioact.* 126, 299–313.
- Yankovich, T.L., Beresford, N.A., Wood, M., Aono, T., Andersson, P., Barnett, C.L., Bennett, P., Brown, J., Fesenko, S., Hosseini, A., Howard, B.J., Johansen, M., Phaneuf, M., Tagami, K., Takata, H., Twining, J., Uchida, S., 2010a. Whole-body to tissue concentration ratios for use in biota dose assessments for animals. *Radiat. Environ. Biophys.* 49, 549–565.
- Yankovich, T.L., Vives i Batlle, J., Vives-Lynch, S., Beresford, N.A., Barnett, C.L., Beaugelin-Seiller, K., Brown, J.E., Cheng, J.-J., Copplestone, D., Heling, R., Hosseini, A., Howard, B.J., Kamboj, S., Kryshev, A.I., Nedveckaite, T., Smith, J.T., Wood, M.D., 2010b. An international model validation exercise on radionuclide transfer and doses to freshwater biota. *J. Radiol. Prot.* 30, 299–340.
- Yankovich, T.L., Beresford, N.A., Fesenko, S., Fesenko, J., Phaneuf, M., Dagher, E., Outola, I., Andersson, P., Thiessen, K., Ryan, J., Wood, M.D., Bollhöfer, A., Barnett, C.L., Copplestone, D., 2013. Establishing a database of radionuclide transfer parameters for freshwater wildlife. *J. Environ. Radioact.* 126, 299–313.
- Yu, C., Zielen, A.J., Cheng, J.J., LePoire, D.J., Gnanapragasam, E., Kamboj, S., Arnish, J., Wallo III, A., Williams, W.A., Peterson, H., 2001. User's Manual for RESRAD Version 6. Aragonne National Laboratory, Ilanosis.