

**Regional modelling of permafrost thicknesses over the past 130 ka: implications for permafrost development in Great Britain**

JONATHAN P. BUSBY, JONATHAN R. LEE, SEV KENDER, J. PAUL WILLIAMSON AND SIMON NORRIS

Busby, J. P., Lee, J. R., Kender, S., Williamson, J. P. & Norris, S. 2015: Regional modelling of permafrost thicknesses over the past 130 ka: implications for permafrost development in Great Britain

The greatest thicknesses of permafrost in Great Britain most likely occurred during the last glacial-interglacial cycle, since this is when some of the coldest conditions occurred during the last 1 M years. The regional development of permafrost across Great Britain during the last glacial-interglacial cycle has been modelled from a ground surface temperature history based on mean annual temperatures and the presence of glacier ice. To quantify the growth and decay of permafrost, modelling was undertaken at six locations across Great Britain that represents upland glaciated, lowland glaciated, upland unglaciated and lowland unglaciated conditions. Maximum predicted permafrost depths derived in this academic study range between several tens of metres to over a 100 m depending on various factors including elevation, glacier ice cover, geothermal heat flux and air temperature. In general, the greatest maximum permafrost thicknesses occur at upland glaciated locations with minimum thickness at lowland sites. Current direct geological evidence for permafrost is from surface or shallow processes, mainly associated with the active layer. Further research is recommended to identify the imprint of freeze/thaw conditions in permanently frozen porous rocks from beneath the active layer.

*Jonathan P. Busby (jpbu@bgs.ac.uk), Jonathan R. Lee, Sev Kender, and J. Paul Williamson, British Geological Survey, Keyworth, Nottingham, NG12 5GG, UK; Simon Norris, Radioactive Waste Management Limited., Curie Avenue, Harwell, Didcot, OX11 0RA, UK.*

Modern permafrost has been recognised in several mid-high latitude regions that are subjected to High Arctic (Smith *et al.* 2007), Continental (Halsey *et al.* 1995; Bockheim & Hall 2002) and High-altitude (Gorbunov 1978; Guodong & Shaoling 1982) climatic regimes. It has also been widely interpreted from the geological record within mid- to high-latitude areas where it developed repeatedly during successive Quaternary cold stages (Ballantyne & Harris 1994; Rozenbaum & Shpolyanskaya 1998; Clayton *et al.* 2001; French *et al.* 2007; French 2008; Candy *et al.* 2011; French & Millar 2014; Vandenberghe *et al.* 2014).

Preserved evidence for past permafrost development typically relates to phenomenon produced by 'seasonal' frost action (French 2008). Recognising evidence for 'perennial' permafrost is by contrast more challenging. This is because 'perennial' permafrost is likely to have occurred largely at depth and the potential subtlety and ambiguity of the geological evidence. Specific challenges relate to the identification of geological evidence for the growth of ice within various materials, the effects of groundwater chemistry upon freezing at depth and the recognition of materials that have remained dry, but were 'thermally-frozen'. Recent studies from deep mines and boreholes have demonstrated modern 'thermally-frozen' permafrost thicknesses of 500-600 m in northern Canada (Ruskeeniemi *et al.* 2002; Ruskeeniemi *et al.* 2004), 1500 m in central Siberia (Fotiev 1997) and nearly 100 m of permafrost buried at depth in Poland (Szewczyk & Nawrocki 2011). What is less clear are the thicknesses of 'perennial' permafrost that developed during the Quaternary.

Understanding past permafrost thicknesses is important from several geological and applied angles. Firstly, it provides a direct insight into the depth to which cold climate processes have operated and this has major implications for weathering rates and the development of regolith/saprolite. From an applied perspective, the depth to which permafrost may have developed is directly relevant to establishing a safety case for a geological disposal facility (GDF) for radioactive waste (McEwen & de Marsily 1991;

Ershov *et al.* 2003; Chapman & Hooper 2012). The safety case for a GDF relies on complementary natural and engineered barriers to provide the necessary level of containment in the post-closure period. Post-closure safety case studies typically consider natural changes over the first one million years following GDF closure.

There are no definitive indicators of the maximum depth of permafrost development in Great Britain during the Quaternary. Within this paper, we utilise a numerical modelling approach linked to the past climate, thermal conductivity and geothermal heat flux to reconstruct permafrost thicknesses for six localities in Great Britain during the last 130 ka. Each locality was carefully chosen to exhibit variations in inferred glacier cover during the Late Devensian, latitude, elevation and geology.

## Background

### *Definition of permafrost*

Historically, several definitions of the term *permafrost* exist. The term was originally employed as an abbreviated form of ‘permanently frozen’ (Muller 1945). However, this definition is somewhat restrictive because it relies upon the availability of water within the substrate for the ground to freeze. Within modern literature, *permafrost* is widely defined as ground which remains at or below 0 °C for at least two consecutive years and perennially frozen ground as ground that keeps frozen for at least two consecutive years (French 2007). Hence, permafrost is solely based on temperature, thus disregarding the texture, degree of compaction, water content, and lithologic character of the material, whereas perennially frozen ground is defined on the basis of freezing of water. The freezing of water is itself dependent on pressure, salinity and by the adsorptive and capillary properties of the ground matter. In the study here *permafrost* (i.e. ground which remains at or below the 0 °C (32 °F)

isotherm) has been predicted by one dimensional heat conduction modelling that does not take into account the freezing of water, the effect of groundwater movement or the change in ground properties due to freezing. Therefore this modelling identifies the potential for ice formation.

### *The Quaternary record of permafrost within Great Britain*

Mid-latitude regions including Great Britain have proven particularly sensitive to global-scale climatic change during the Quaternary with marked climatic variability recognised at both Milankovitch (Bridgland 2000; Bridgland 2006; Bates *et al.* 2010; White *et al.* 2010; Penkman *et al.* 2011) and sub-Milankovitch time-scales (Lewis *et al.* 2001; McDermott *et al.* 2001; Walker *et al.* 2003; Briant *et al.* 2005; Candy & Schreve 2007; Gao *et al.* 2007).

Evidence demonstrates the cyclical occurrence of cold and temperate (interglacial) climatic conditions in Great Britain during the Quaternary although the geological record is highly-fragmented and far from complete. Based upon the marine oxygen isotope record, Great Britain has experienced at least 50 major cold climatic stages (Lisiecki & Raymo 2005) and at least 30 distinct episodes of glaciation of varying scale and extent during the Quaternary (Lee *et al.* 2011; Böse *et al.* 2012; Lee *et al.* 2012; Thierens *et al.* 2012). Likewise, multiple episodes of permafrost development have been recognised during the Early-Middle (Rose & Allen 1977; West 1980; Kemp *et al.* 1993; Murton *et al.* 1995; Lee *et al.* 2003; Candy *et al.* 2011) and Late Pleistocene (Watson & Morgan 1977; Seddon & Holyoak 1985; Allen 1987; Ballantyne & Harris 1994; Bateman 1995; Murton *et al.* 2003; Boardman 2011; Bateman *et al.* 2014). Palaeontological evidence from the last two cold periods – the Younger Dryas (12.9-11.7 ka BP) and Late Devensian glaciation (30-18 ka BP), reveal that mean winter air temperatures were at least -25 °C (Atkinson *et al.* 1987; Huijzer &

Vandenberghe 1998). From this climate record it might be expected that depths of periglacial disturbance might extend to 10s, or even 100s, of metres. However, despite evidence for permafrost being both temporally and spatially common, there is no currently known geological evidence that refers to thicknesses of it beyond several tens of metres.

*Permafrost modelling approach*

The modelling is based on the periodic heating at the surface of a column of infinite depth and the propagation of the heat into the ground in the vertical (depth) dimension. Only the variation of thermal properties in the vertical (z) dimension is considered. In situations where the geology is horizontally layered this 1D assumption is reasonable but it will lead to a more approximate solution if the geology varies substantially in the horizontal (x, y) plane. For the purposes of a regional investigation to assess the potential for permafrost development this, static, 1D modelling approach is appropriate, but it does not lend itself to a full sensitivity analysis whereby all the input modelling parameters are varied across their permissible limits.

A surface temperature history consists of a series of cyclical swings in temperature between warm and cold conditions. Each cycle can last from a few hundred to several thousand years. An example of a surface temperature history for Prudhoe Bay, Alaska, based on an East Siberia model, is shown in Fig. 1a (Osterkamp & Gosink 1991). The change in surface temperature will be propagated into the ground, but the amplitude of the change will decay exponentially with depth and there is a time-lag between the temperature perturbation at the surface and at depth. The rate of the exponential decay and the time-lag are both dependent on the thermal diffusivity of the geological strata. A surface temperature history can also be represented as a series of step changes in temperature. The surface temperature history for Prudhoe Bay, shown in Fig. 1a, is shown as a series of step changes in Fig. 1b.

The depth to which the effect is noticeable depends on the magnitude of the temperature step, the time since the event and the thermal diffusivity of the geological strata. Carslaw & Jaeger (1959) have shown that:

$$T_{\theta} = T_0 \times \operatorname{erfc} [z/2\sqrt{\kappa t}], \quad (1)$$

where  $T_{\theta}$  is the departure from original equilibrium temperature at depth  $z$  and time  $t$  after an instantaneous change in surface temperature of  $T_0$ ;  $\kappa$  is the average thermal diffusivity of the geological strata down to depth  $z$  and  $\operatorname{erfc}(x)$  is the complementary error function. Noting that the change in surface temperature is the difference in temperature between successive steps, the effect of more than one temperature step is found by addition of all the steps, i.e.

$$T_{\theta} = \sum T_{\theta i}, \quad (2)$$

where  $T_{\theta i}$  is the temperature deviation due to the  $i$ th event.

Therefore to model the evolution of sub-surface temperatures through time, an initial sub-surface temperature profile is perturbed by any step changes in surface temperature that have occurred from the initial time up to the time being considered. The initial time has been taken as 126 ka BP and sub-surface temperatures have been calculated at 5 m depth intervals to 1 km every 250 years up to present day. The derivation of the step changes in surface temperature, i.e. the palaeoclimate of the last glacial-interglacial cycle and its variations across Great Britain, are discussed below.

## Temperature and ice sheet evolution over the last glacial-interglacial cycle

For the permafrost modelling a surface temperature history is required for each locality considered. This has been achieved by constructing a generic record of past mean annual air

temperature (MAAT) over the last glacial-interglacial cycle (0-130 ka BP) which has then been modified for latitude and periods of ice cover.

There are several proxies for past land-surface temperature, which include fossil pollen and beetle assemblages, and although each has its uncertainties, using multi-proxy data can provide reasonably robust reconstructions. Estimating cold winter temperatures from fossil beetle assemblages is thought to be particularly problematic. This is due to a lack of proxy data in the calibration curve (Coope *et al.* 1998); the lack of abundance data in the transfer function used (Bray *et al.* 2006); plus the ability of beetles to bury themselves in the ground during winter months – beetles therefore may only respond to summer temperatures (Westaway & Younger 2013), which is not suitable for MAAT reconstruction.

Last Glacial-Interglacial cycle MAAT curve

Westaway & Younger (2013) attempted to reproduce the MAAT curve for southern and northern UK over the last 150 ka, based on a combination of climate proxy records including sea-surface temperature (SST), derived from planktonic foraminifera, and ice-core temperature data. Here, we produce a different MAAT record by utilising NE Atlantic SST records (Fig. 2B-D), and pollen-based MAAT records (Fig. 2A), to reconstruct an idealized curve (Fig. 3A) which looks different from Greenland ice core temperature records (Fig. 3C). This approach is necessary as there are no land-surface MAAT pollen records from the UK during the Last Glacial (c. 30-19 ka BP) (Davis *et al.* 2003; Bartlein *et al.* 2011).

Time interval 12–0 ka BP. - For this interval, which spans the Younger Dryas and current Holocene interglacial, pollen records are available across Europe, including the UK, which have been quantitatively analysed in Davis *et al.* (2003). Based on this large dataset (500

1  
2  
3 sites), these authors produced MAAT records for 6 different regions of Europe. The ‘central  
4  
5 west’ (which is the approximate region, N-S, from northern Scotland to northern Spain and  
6  
7 W-E from western Ireland to Poland) MAAT curve has been followed for the last 12 ka (Fig.  
8  
9 2A), as this is the proximity region that includes Great Britain. In this reconstruction the  
10  
11 temperature anomaly (relative to present day temperature) for the Younger Dryas is  $-4^{\circ}\text{C}$ .  
12  
13  
14  
15  
16

17 Time interval 22–12 ka BP. - This important period encompasses the progressive break-up of  
18  
19 the Last British-Irish Ice Sheet, encompassing the global climatic minimum (c. 21 ka), the  
20  
21 complicated phase of climatic amelioration commonly referred to as the Lateglacial, and the  
22  
23 onset of the Younger Dryas stadial. The most comprehensive analysis of the MAAT history  
24  
25 for this interval is provided by the study of Shakun *et al.* (2012) that summarizes 80 proxy  
26  
27 records of both sea surface and land surface proxies to reconstruct global latitudinal  
28  
29 temperature curves. Their temperature curve corresponding to  $30\text{--}60^{\circ}\text{N}$  (Fig. 2B) is used  
30  
31 here.  
32  
33  
34  
35  
36  
37

38 Time interval 40–22 ka BP. - The time interval between 40–22 ka BP includes the build-up of  
39  
40 the Last British-Irish Ice Sheet to its maximum-known extent at approximately 27 ka BP and  
41  
42 then its gradual decline (Clark *et al.* 2012). For the time interval before 22 ka ago there are no  
43  
44 available regional time-series compilations of MAAT, and therefore proximal NE Atlantic  
45  
46 SST records have been used to estimate the shape of the UK MAAT curve. For 22–40 ka the  
47  
48 proximal SST record from BOFS (Biogeochemical Oceanic Flux Study) ocean core 5K was  
49  
50 referenced (Fig. 2C), which is located west of the Porcupine Seabight at a similar latitude to  
51  
52 southern Britain (Maslin *et al.* 1995). This is believed to be a reasonably accurate estimate  
53  
54 due to the proximity of the site, and similarity to the Davis *et al.* (2003) and Shakun *et al.*  
55  
56  
57  
58  
59  
60

(2012) compilation curves from the 0–22 ka interval (Fig. 2). For BOFS 5K, both summer and winter absolute temperatures are reconstructed by using planktonic foraminiferal assemblage data and different transfer functions which all show similar trends (Fig. 2).

Time interval 140–40 ka BP. - The final time-interval spans the Last Interglacial (Ipswichian/Eemian/MIS 5e) and the complex transition into the Last Glaciation (Late Devensian). For the longer, oldest time interval the NE Atlantic SST reconstruction from ODP (Ocean Drilling Program) ocean core 980 (Fig. 2D) has been used, which is located in the Rockall Trough at a similar latitude to central Great Britain (McManus *et al.* 1999). This record has some similarities to BOFS 5K, but is at a far lower temporal resolution and is therefore only applied for the older section.

*Last Glaciation – absolute temperature*

In order to scale the idealised MAAT curve (Fig. 3B) to each of the six Great British localities, modern MAAT is applied as the Holocene average (maximum temperature), and the Annan & Hargreaves (2013) global estimate of MAAT during the global peak of the last glaciation (19–23 ka BP) as a guide for the minimum temperature at each locality, with a linear scaling of temperature between. The Annan & Hargreaves (2013) dataset is based on a combination of computer modelling and pollen MAAT proxies, where the southern England and Wales localities are estimated to be 8–12 °C below present, and the northern England and Scotland sites 12–20 °C below present. Therefore the estimated climate range for the last glacial-interglacial cycle is 12 to 17 °C below modern, between southern English and northern Scottish sites respectively.

The average global estimate for the climatic peak of the last glaciation fits reasonably well with the NE Atlantic SST records, which show  $\sim 11^{\circ}\text{C}$  difference between Holocene and global climatic minimum values at BOFS 5K (Maslin *et al.* 1995), and  $\sim 8^{\circ}\text{C}$  difference at ODP 980 (McManus *et al.* 1999). These values are slightly greater than the SST estimates of the oceans surrounding the British Isles by Waelbroeck *et al.* (2009), which was based on a synthesis of global integrated fossil and geochemical proxies. It also fits well with pollen based reconstructions at 21 ka BP from central Europe (Bartlein *et al.* 2011) and NE France (Busschers *et al.* 2007), showing MAAT  $\sim 10^{\circ}\text{C}$  below present, and the pollen-based European regional reconstructions for the Younger Dryas (12.9-11.7 ka) at  $\sim 4^{\circ}\text{C}$  below present (Davis *et al.* 2003).

#### *Insulating effect of ice cover*

Due to the insulating capacity of ice sheets and glaciers, for the final land-surface temperature estimates, it is necessary to reconstruct the time-intervals that each location was covered by ice and the temperature at the base of the ice. Ice sheet dynamics in Great Britain have been modelled by several studies (Siegert & Dowdeswell 2004; Boulton & Hagdorn 2006; Evans *et al.* 2009; Hubbard *et al.* 2009; Clark *et al.* 2012; Patton *et al.* 2013) with the reconstructions of Hubbard *et al.* (2009) providing a series of time-slices with reconstructed ice sheet extent from 0-35 ka ago and these have been used here for the period  $>35$  ka BP. To reconstruct the presence of ice cover in the older part of the record, the sensitivity of the ice sheet or glacier presence at each location for the 0–35 ka BP record, derived from Hubbard *et al.* (2009), is used to extend back using the reconstructed temperature record; i.e. for each locality, when the temperature in the older part of the record falls below that in the recent record ( $<35$  ka BP) when ice is predicted, ice is assumed in that part of the older record. The modelled reconstructions of Hubbard *et al.* (2009) appear to correlate well with the proxy

data, and hence the ice cover presence/absence data over the interval 0–35 ka BP are far less uncertain than those from the interval 35–130 ka BP. Periods of ice cover at each of the six localities are given in the description of the localities in Table 1.

The theory behind the reconstruction of temperatures beneath glaciers in Scotland has been discussed by Glasser & Siegert (2002) and Hall & Glasser (2003). Warm-based ice that is often found under thick ice in topographical troughs is warmer than cold-based ice frozen to its bed. Temperatures at the base of the ice will also be affected by the mean annual air temperature and ice thickness. Sliding ice may be due to the pressure at the base of the ice being close to the pressure melting point; a sufficiently high pressure that the ice melts even though its temperature is below 0 °C. Hall & Glasser (2003) have shown that during basal freezing conditions, basal temperatures can range from -12 to -6 °C (absolute). However, when basal melting occurs, basal temperatures are more likely to be in the range of -1 to +1 °C (absolute).

The basal ice thermal regime may also change with time and space as the ice mass thins or thickens. However, for simplicity and for each surface temperature history, a single basal ice temperature has been assigned for periods of ice cover. Hubbard *et al.* (2009) have modelled the development of the Last British-Irish Ice Sheet and have predicted the cumulative time that the ice sheet bed was at pressure melting point. At each of the localities, if Hubbard *et al.* (2009) predict predominantly persistent frozen basal conditions then cold-based ice is assumed, however, if predominantly pressure melting conditions are predicted then warm-based ice is assumed, for the entire period from 0-130 ka BP. Basal absolute temperatures for cold-based frozen conditions are therefore estimated as -3 °C (absolute). This temperature is not as low as modelled by Hall & Glasser (2003) as it is assumed that warm-based conditions would have also occurred as the ice sheet grew and declined. For warm-based ice conditions, absolute temperatures are assumed to be +1 °C.

Two examples of surface temperature histories are shown in Fig. 4. Central East Anglia was ice-free during the Late Devensian (Clark *et al.* 2004; Bateman *et al.* 2014). Therefore, the land surface temperature estimates infer a minimum temperature of 12 °C below present, which occurred at the global peak of the last glaciation (19-23 ka BP) (Annan & Hargreaves 2013). By contrast, within the Stainmore Trough, the land-surface was insulated from the minimum MAAT of 14 °C below present by the extensive periods of ice cover (Livingstone *et al.* 2012). These occur between 110-102, 76-37 and 34-14.5 ka BP and the temperature during these periods is constant at +1 °C absolute (8.2 °C below present). The minimum temperature of 9.8 °C below present occurred between 13-12 ka BP after the ice had retreated.

## Modelling results

### *Permafrost modelling*

In order to assess how permafrost development may vary across Great Britain, six locations have been selected which present a variety of latitude, elevation, substrate geology and periods of ice cover. These are shown in Fig. 5. For each location, the vertical geological bedrock section, at the 1:625 000 scale, has been extracted to 1 km depth from the BGS GB3D national geological model (Mathers *et al.* 2014). In summary these locations comprise;

#### Upland glaciated

Mid-Wales	Silurian mudstone, sandstone, siltstone and conglomerate
Northwest Highlands	Precambrian psammite

#### Lowland glaciated

Stainmore Trough	Carboniferous sedimentary rocks of mixed lithologies
------------------	--

Midland Valley	Carboniferous Coal Measures and Clackmannan Group limestone, sandstone and mudstone
Upland unglaciated	
Dartmoor	Permian granite pluton
Lowland unglaciated	
East Anglia	Chalk and Mesozoic argillaceous rocks overlying Lower Palaeozoic basement

Each geological unit has been assigned a thermal conductivity after (Rollin 2002), which is required for the calculation of thermal diffusivity that is used in equation (1). (Note that specific heat and density that are also required for the calculation of thermal diffusivity, were taken as standard values for the geological lithologies.) Sketch diagrams of the geological succession at each of the glaciated locations are shown in Fig. 6 and the unglaciated locations in Fig. 7. In addition, for each location, a mean annual present day surface temperature and heat flow (Downing & Gray 1986; Lee *et al.* 1987; Rollin 1995; Rollin *et al.* 1995; Barker *et al.* 2000) were assigned, which are required for the estimation of the initial vertical sub-surface temperature profile (see below). All of these parameters are presented in Table 1.

Permafrost models have been run with the repeated application of equation (1) with the surface temperature histories. The initial temperature profile at 130 ka BP has been estimated by application of Fourier’s Law

$$\text{grad } T_z = -\frac{q}{\lambda_z}, \tag{3}$$

where grad  $T_z$  is the temperature gradient ( $\text{K m}^{-1}$ ) between depths  $z_i$  and  $z_{i+1}$ ,  $q$  is heat flow ( $\text{W m}^{-2}$ ) and  $\lambda_z$  is thermal conductivity ( $\text{W m}^{-1} \text{K}^{-1}$ ) between depths  $z_i$  and  $z_{i+1}$ . This is applied

from the ground surface downwards where, from the derivation of the surface temperature histories discussed above, at 130 ka BP the ground surface temperatures have been taken to be 5 °C less than present day surface temperature at all locations (see Fig. 4).

The evolution of sub-surface temperatures at the six localities as a result of the surface temperature histories has been modelled with sub-surface temperatures generated every 250 years from 126 ka BP to present day. Classed plots of the sub-surface temperatures for each locality are shown in Figs. 8 to 10 where permafrost ( $T \leq 0$  °C) is shown as shades of blue/cyan. For the upland glaciated sites periods of ice cover are associated with cold based ice leading to extensive permafrost development with maximum depths of 105 and 180 m for mid-Wales and the Northwest Highlands respectively. Within the Northwest Highlands, permafrost is modelled as continuous from 110 to 12 ka BP due to the sub-zero temperatures imposed by the cold-based glacial ice. Permafrost development is very different between the lowland glaciated sites. For the Midland Valley where cold-based ice has been assumed, the maximum permafrost depth is 110 m, similar to the upland sites. For the Stainmore Trough, despite three periods of ice cover, permafrost only develops within the model after the ice has retreated and then to a maximum depth of only 20 m. Dartmoor has been classified as an upland unglaciated locality although there is an ongoing debate regarding the presence of localised glaciers during the Quaternary (Evans *et al.* 2012; Straw 2013; Evans *et al.* 2014; Straw 2014). However, the small localised glaciers advocated by Evans *et al.* (2012) are unlikely to exert a major influence on the regional permafrost development. Within the model, permafrost develops on Dartmoor whenever the MAAT is less than 9.0 °C below present day (0 °C absolute). This occurs in intervals from 62 to 14.5 ka BP. The maximum permafrost thickness of 80 m occurs during the peak of the Late Devensian cold stage. At the lowland unglaciated location, East Anglia, the pattern of permafrost development is similar to Dartmoor except there is no permafrost development until after 40 ka BP and the maximum

permafrost thickness is 65 m. In this case, the lowland location results in a 1 °C rise in MAAT. Maximum thicknesses of permafrost at the six locations are summarised in Table 2.

Discussion

*Controls on permafrost development*

In the current modelling, maximum predicted permafrost depths in Great Britain range between several tens of metres to over a 100 m depending on various factors including elevation, glacier ice cover, geothermal heat flux and air temperature. The results comprise 1D heat conduction models that have not considered the effects of freezing or groundwater flow. In addition, due to the static modelling approach, our best estimate for each input parameter (e.g. thermal conductivity, periods of ice cover etc.) has been applied rather than considering the full permissible range for each parameter. The evolution through time of sub-surface temperatures has been modelled when the surface temperature has been allowed to vary based on the climate from the last glacial-interglacial cycle. It is not possible to make any statements about ice formation or the depth to which partially or completely frozen ground might penetrate. However, as shown in other studies, perennially frozen ground is likely to correlate with the extent of the 0 °C isotherm (Hartikainen *et al.* 2010; Govaerts *et al.* 2011). Freezing will depend upon other factors such as the porosity of the geological strata and salinity of the groundwater (French 2007; Stotler *et al.* 2009).

The subglacial thermal regime plays an important role in permafrost development and specifically the speed and depth to which permafrost can penetrate. The deepest permafrost development was modelled in the Northwest Highlands where extensive, but thin cold-based ice cover has been assumed for much of the last 110 ka up to the beginning of the Holocene. The predominantly frozen basal conditions would have led to colder basal-ice, assumed to

be  $-3^{\circ}\text{C}$  (absolute) enabling propagation of sub-zero temperatures into the substrate for an extensive period of time. However, cold basal-ice still had an insulating effect on the substrate because without ice cover, the MAAT during the last glaciation is predicted to have fallen to  $-9.8^{\circ}\text{C}$  (absolute) which would have led to deeper and more rapid permafrost development. By contrast, the least and shallowest permafrost development corresponds to the Stainmore Trough which was a major zone of fast-flowing warm-based ice during the Late Devensian (Evans *et al.* 2009; Livingstone *et al.* 2012). A warm-based ice temperature of  $+1^{\circ}\text{C}$  (absolute) has been assumed, limiting MAAT transfer into the substrate and restricting shallow permafrost development to a short period of ice-free conditions during deglaciation. At this northerly latitude the surface ground temperature could have fallen to  $-4.8^{\circ}\text{C}$  (absolute) had there been no glacial ice cover. Hence, in both of these examples, ice cover has had a major insulating effect protecting the ground from the extremes of air temperature. In general, ice cover has an insulating effect because the temperature at the base of the ice, even for cold-based ice, is warmer than the extremes of the cold temperatures, especially during the peak of the Late Devensian.

The results obtained from the modelling can be compared to similar studies undertaken as part of international studies related to implementing geological disposal of radioactive waste. The geology at Mol (Belgium) comprises an alternating sedimentary succession of sandstones and mudstones. For a climate model with a minimum MAAT of  $-19^{\circ}\text{C}$  compared to present day and length of cold periods comparable to those used here, maximum permafrost thicknesses (defined as the  $0^{\circ}\text{C}$  isotherm combined with 50% frozen ground) of 160 m and 215 m for the climate models with and without surface cover respectively were modelled (Govaerts *et al.* 2011). The minimum MAAT applied here was  $-17^{\circ}\text{C}$  compared to present day in the Northwest Highlands where the maximum depth of permafrost was 180 m, which is comparable to the results of Govaerts *et al.* (2011).

The Forsmark site in Sweden is located over crystalline bedrock and maximum modelled permafrost depths (0 °C isotherm) were 260 m for the repetition of the last glacial-interglacial cycle and 390 m for a severe permafrost case (Hartikainen *et al.* 2010). The Dartmoor and Northwest Highlands sites here are analogous in terms of geology and the range in permafrost thicknesses between the three locations are compatible, given the differences in the surface temperature history resulting from the geographical location of Forsmark compared to Great Britain, and the extent and nature of glacial ice cover.

The principal factors that determine the depth of permafrost are the long-term surface temperature history, the thermal properties of the geological strata and the geothermal heat flux. Geological strata with high thermal conductivities will produce smaller temperature gradients and will therefore support thicker permafrost. Conversely, strata with low thermal conductivities act as insulators, creating a blanketing effect that leads to high temperature gradients and thinner permafrost. A geological sensitivity analysis was run where permafrost development was modelled at the six localities using the same surface temperature history and an initial temperature profile calculated with the same heat flow (0.065 W m<sup>-2</sup>) and surface temperature (5 °C), i.e. the only variable was the change in geology. This indicated a linear relationship between average thermal conductivity and permafrost depth where a 100% increase in thermal conductivity produced an 88% increase in permafrost depth. Of the two locations with the highest average thermal conductivities (Northwest Highlands and Dartmoor), Dartmoor is associated with the highest heat flow region in Great Britain, which will reduce permafrost thickness.

The surface temperature histories used in the modelling were derived from a review of recent literature. There is general agreement on the timing of major cold stages, but authors disagree on the minimum temperatures that may have been reached. For instance, Westaway & Younger (2013) have suggested that during the last glacial cycle temperatures

in southern England reached  $-20^{\circ}\text{C}$  (relative to present day) between 65-75 ka BP and at the peak of the last glaciation. Govaerts *et al.* (2011) in their 'realistic glacial cycle' indicated a minimum temperature in northern Belgium (which would be expected to be similar to southern England) of  $-19^{\circ}\text{C}$  (relative to present day). The surface temperature history applied here for southern England has a minimum temperature of  $-12^{\circ}\text{C}$  (relative to present day) during the last glaciation. Adoption of lower temperatures in the modelling would have led to the development of thicker permafrost, especially at the unglaciated locations.

Clearly, all the parameters in the modelling have a range of values that will influence permafrost depth and these could be tested with a full sensitivity analysis. The modelling has adopted a, static, 1D approach to defining the parameters and these have been chosen to be reasonable (and defensible) to produce a first pass assessment of permafrost thicknesses. The result therefore, provides an insight into regional differences across Great Britain of permafrost thicknesses, rather than a detailed analysis at each location of the range of potential permafrost depths.

#### *Implications for major landscape change*

Recent developments in the understanding of the scale and frequency of Quaternary climate change reveal that mid-latitude regions such as Great Britain have been subjected cyclically to cold stage conditions and presumably periglacial climatic regimes (Candy *et al.* 2011; Böse *et al.* 2012). Outputs from the modelling presented within this paper indicate the potential for permafrost to extend through the substrate to depths of between several tens of metres to, in-excess of, 100 m depending on location and a range of geological and climatic factors. This implies that over the duration of the Quaternary, surface and shallow sub-surface materials to considerable depths have been 'thermally stressed' by repeated cycles of permafrost development. Potential evidence for this 'thermal stressing' is the development of extensive weathered regolith (saprolite) capping interfluvial areas, its downslope transportation

via colluvium-mantled slopes (Booth *et al.* 2015) and transportation and deposition by rivers forming widespread aggradational river terrace sequences (Gibbard & Allen 1994; Rose 1994; Maddy *et al.* 1995; Rose *et al.* 1999; Westaway *et al.* 2006; Howard *et al.* 2007; Boreham *et al.* 2010; White *et al.* 2010). These deposits are particularly well-preserved beyond the southern-most limits of successive Quaternary glaciations, which acted to erode and remove the products of weathering (Booth *et al.* 2015).

*Evolution of the modelling approach*

The regional permafrost modelling presented here provides a valuable insight into the temporal and spatial controls of permafrost development. However, the approach is 1D in nature and could be enhanced by consideration of further variables. For instance, latent heat produced during freezing will result in thinner perennially frozen ground compared to permafrost. In addition groundwater flow and composition, groundwater recharge and discharge, porosity and saturation of the rock strata will all influence the development of frozen ground. Such modelling should also aim to explore the change in groundwater chemistry such as salinity and its vertical variability. The insulating effect of ice cover has been shown to be an important factor in shielding the ground surface from air temperature and limiting permafrost development. The temperature at the base of the ice can vary over a large temperature range and these temperatures will not be constant, but will change depending on ice thickness, prevailing climate (if the ice is thin enough) and glacier dynamics. Modelling schemes for estimating basal ice temperatures are available (Glasser & Siegert 2002; Forsström 2005), but they require the evolution of the ice sheet through time to be estimated. Hence it is recommended that further modelling studies of ice sheet evolution, particularly for the period prior to 35 ka BP should be undertaken.

Most experimental studies and direct geological evidence for permafrost is the result of surface or shallow processes, particularly those that occur within the active layer (Hallet *et al.* 1991; Murton *et al.* 2000; Murton *et al.* 2006; French 2008). By comparison, little evidence has been published relating to the geological signature of permanently frozen ground. Hutchinson & Thomas-Betts (1990) have reported depths of periglacial disturbance in southern England to be less than 20 m, except for one, unreferenced, site to 50 m depth. Some studies have been carried out for Finnish and Swedish programmes related to implementing geological disposal of radioactive waste by studying conditions at depth at Lupin in northern Canada (Ruskeeniemi *et al.* 2002; Ruskeeniemi *et al.* 2004; Stotler *et al.* 2009). However, these sites are all located on metamorphosed crystalline rocks and so differ significantly from those that underlie much of Great Britain. Therefore, there is a general lack of knowledge on the imprint that freeze/thaw conditions might have on saturated porous rocks in permanently frozen ground. Laboratory studies to simulate in-situ conditions of frozen ground might detect imprints such as damaged pore structures and changes to the geochemistry of pore fluids and associated precipitates. Such imprints could then be searched for from deep core material for evidence of permafrost during the last glacial-interglacial cycle.

## Conclusions

- One-dimensional heat conduction modelling across Great Britain has demonstrated that permafrost development during the last glacial-interglacial cycle was likely at all locations and in some cases was extensive to over 100 m in depth.
- The surface temperature history at any location has been derived from a general reconstruction of the UK mean annual air temperature modified by latitude and

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

periods of ice cover. Simplifying assumptions have been made to temperatures beneath glacial ice but a distinction has been made between warm-based basal conditions and cold, frozen conditions.

- In general, the greatest maximum permafrost thicknesses occur at upland glaciated locations (cold-based ice) and localities that remained largely ice-free, with minimum thickness at insulated lowland sites.

*Acknowledgments* - Radioactive Waste Management Limited. (RWM) is thanked for their financial support and scientific interest in this study, which also benefitted from review comments from RWM staff. Dr Andrew Finlayson is thanked for his helpful discussions. This paper is published by permission of the Executive Director of the British Geological Survey (NERC).

References

Allen, J. 1987: Dimlington Stadial (Late Devensian) ice-wedge casts and involutions in the Severn Estuary, southwest Britain. *Geological Journal* 22, 109-118.

Annan, J. & Hargreaves, J. 2013: A new global reconstruction of temperature changes at the Last Glacial Maximum. *Climate of the Past* 9, 367-376.

Atkinson, T. C., Briffa, K. R. & Coope, G. R. 1987: Seasonal Temperatures in Britain during the Past 22,000 Years, Reconstructed Using Beetle Remains. *Nature* 325, 587-592.

Ballantyne, C. K. & Harris, C. 1994: *The periglaciation of Great Britain*. 330 pp. Cambridge University Press, Cambridge.

- 1  
2  
3 Barker, J., Downing, R., Gray, D., Findlay, J., Kellaway, G., Parker, R. & Rollin, K. 2000:  
4  
5 Hydrogeothermal studies in the United Kingdom. *Quarterly Journal of Engineering*  
6  
7 *Geology and Hydrogeology* 33, 41-58.  
8  
9  
10 Bartlein, P. J., Harrison, S. P., Brewer, S., Connor, S., Davis, B. A. S., Gajewski, K., Guiot,  
11  
12 J., Harrison-Prentice, T. I., Henderson, A., Peyron, O., Prentice, I. C., Scholze, M.,  
13  
14 Seppä, H., Shuman, B., Sugita, S., Thompson, R. S., Viau, A. E., Williams, J. & Wu,  
15  
16 H. 2011: Pollen-based continental climate reconstructions at 6 and 21 ka: a global  
17  
18 synthesis. *Climate Dynamics* 37, 775-802.  
19  
20  
21 Bateman, M. 1995: Thermoluminescence dating of the British coversand deposits.  
22  
23 *Quaternary Science Reviews* 14, 791-798.  
24  
25  
26 Bateman, M. D., Hitchens, S., Murton, J. B., Lee, J. R. & Gibbard, P. L. 2014: The evolution  
27  
28 of periglacial patterned ground in East Anglia, UK. *Journal of Quaternary Science*  
29  
30 29, 301-317.  
31  
32  
33 Bates, M. R., Briant, R. M., Rhodes, E. J., Schwenninger, J.-L. & Whittaker, J. E. 2010: A  
34  
35 new chronological framework for Middle and Upper Pleistocene landscape evolution  
36  
37 in the Sussex/Hampshire Coastal Corridor, UK. *Proceedings of the Geologists'*  
38  
39 *Association* 121, 369-392.  
40  
41  
42 Boardman, J. E. 2011: *Periglacial processes and landforms in Britain and Ireland*. 308 pp.  
43  
44 Cambridge University Press, Cambridge.  
45  
46  
47 Bockheim, J. G. & Hall, K. J. 2002: Permafrost, active-layer dynamics and periglacial  
48  
49 environments of continental Antarctica: Periglacial and Permafrost Research in the  
50  
51 Southern Hemisphere. *South African Journal of Science* 98, 82-90.  
52  
53  
54 Booth, S., Merritt, J. & Rose, J. 2015: Quaternary Provinces and Domains – a quantitative  
55  
56 and qualitative description of British landscape types. *Proceedings of the Geologists'*  
57  
58 *Association, in press*.  
59  
60

- 1  
2  
3 Boreham, S., White, T. S., Bridgland, D. R., Howard, A. J. & White, M. J. 2010: The  
4  
5 Quaternary history of the Wash fluvial network, UK. *Proceedings of the Geologists'*  
6  
7 *Association 121*, 393-409.  
8  
9  
10 Böse, M., Lüthgens, C., Lee, J. R. & Rose, J. 2012: Quaternary glaciations of northern  
11  
12 Europe. *Quaternary Science Reviews 44*, 1-25.  
13  
14 Boulton, G. & Hagdorn, M. 2006: Glaciology of the British Isles Ice Sheet during the last  
15  
16 glacial cycle: form, flow, streams and lobes. *Quaternary Science Reviews 25*, 3359-  
17  
18 3390.  
19  
20  
21 Bray, P. J., Blockley, S. P. E., Coope, G. R., Dadswell, L. F., Elias, S. A., Lowe, J. J. &  
22  
23 Pollard, A. M. 2006: Refining mutual climatic range (MCR) quantitative estimates of  
24  
25 palaeotemperature using ubiquity analysis. *Quaternary Science Reviews 25*, 1865-  
26  
27 1876.  
28  
29  
30 Briant, R. M., Bateman, M. D., Russell Coope, G. & Gibbard, P. L. 2005: Climatic control on  
31  
32 Quaternary fluvial sedimentology of a Fenland Basin river, England. *Sedimentology*  
33  
34 *52*, 1397-1423.  
35  
36  
37 Bridgland, D. 2000: River terrace systems in north-west Europe: an archive of environmental  
38  
39 change, uplift and early human occupation. *Quaternary Science Reviews 19*, 1293-  
40  
41 1303.  
42  
43 Bridgland, D. R. 2006: The Middle and Upper Pleistocene sequence in the Lower Thames: a  
44  
45 record of Milankovitch climatic fluctuation and early human occupation of southern  
46  
47 Britain: Henry Stopes Memorial Lecture 2004. *Proceedings of the Geologists'*  
48  
49 *Association 117*, 281-305.  
50  
51  
52 Busschers, F., Kasse, C., Van Balen, R., Vandenberghe, J., Cohen, K., Weerts, H., Wallinga,  
53  
54 J., Johns, C., Cleveringa, P. & Bunnik, F. 2007: Late Pleistocene evolution of the  
55  
56  
57  
58  
59  
60

- Rhine-Meuse system in the southern North Sea basin: imprints of climate change, sea-level oscillation and glacio-isostasy. *Quaternary Science Reviews* 26, 3216-3248.
- Candy, I. & Schreve, D. 2007: Land–sea correlation of Middle Pleistocene temperate sub-stages using high-precision uranium-series dating of tufa deposits from southern England. *Quaternary Science Reviews* 26, 1223-1235.
- Candy, I., Silva, B. & Lee, J. 2011: Climates of the early Middle Pleistocene in Britain : environments of the earliest humans in northern Europe. In Ashton, N., Lewis, S. & Stringer, C. (eds.): *The ancient human occupation of Britain*, 11-22, Elsevier, Amsterdam.
- Carlshaw, H. A. & Jaeger, J. C. 1959: *Conduction of heat in solids. 2nd Edition*, Oxford University Press, Oxford.
- Chapman, N. & Hooper, A. 2012: The disposal of radioactive wastes underground. *Proceedings of the Geologist's Association* 123, 46-63.
- Clark, C. D., Gibbard, P. L. & Rose, J. 2004: Pleistocene glacial limits in England, Scotland and Wales. In Ehlers, J. & Gibbard, P. L. (eds.): *Developments in Quaternary Science*, 47-82, Elsevier, Amsterdam.
- Clark, C. D., Hughes, A. L., Greenwood, S. L., Jordan, C. & Sejrup, H. P. 2012: Pattern and timing of retreat of the last British-Irish Ice Sheet. *Quaternary Science Reviews* 44, 112-146.
- Clayton, L., Attig, J. W. & Mickelson, D. M. 2001: Effects of late Pleistocene permafrost on the landscape of Wisconsin, USA. *Boreas* 30, 173-188.
- Coope, G. R., Lemdahl, G., Lowe, J. J. & Walkling, A. 1998: Temperature gradients in northern Europe during the last glacial–Holocene transition (14-9 14C ka BP) interpreted from coleopteran assemblages. *Journal of Quaternary Science* 13, 419-433.

- Davis, B. A. S., Brewer, S., Stevenson, A. C. & Guiot, J. 2003: The temperature of Europe during the Holocene reconstructed from pollen data. *Quaternary Science Reviews* 22, 1701-1716.
- Downing, R. & Gray, D. 1986: Geothermal resources of the United Kingdom. *Journal of the Geological Society* 143, 499-507.
- Ershov, E., Parmuzin, S., Lobanov, N. & Lopatin, V. 2003: Problems of radioactive waste burial in perennally frozen ground. In Phillips, M., Springman, S. M. & Arenson, L. U. (eds.): *Permafrost*, 660 pp. Swets and Zeitlinger, Lisse, ISBN 90 5809 582 7.
- Evans, D. J. A., Harrison, S., Vieli, A. & Anderson, E. 2012: The glaciation of Dartmoor: the southernmost indepent Pleistocene ice cap in the British Isles. *Quaternary Science Reviews* 45, 31-53.
- Evans, D. J. A., Harrison, S., Vieli, A. & Anderson, E. 2014: Dartmoor Glaciation - fact or fiction? - a comment on Allan Straw's article. *Quaternary Newsletter* 130, 11-19.
- Evans, D. J. A., Livingstone, S. J., Vieli, A. & Ó Cofaigh, C. 2009: The palaeoglaciology of the central sector of the British and Irish Ice Sheet: reconciling glacial geomorphology and preliminary ice sheet modelling. *Quaternary Science Reviews* 28, 739-757.
- Forsström, P.-L. 2005: Through a glacial cycle: simulation of the Eurasian ice sheet dynamics during the last glaciation. *Annales Academiae Scientiarum Fennicae. Geologica – Geographica* 168, 94 pp.
- Fotiev, S. M. 1997: Permafrost groundwater: Russian literature review. . In Haldorsen, S., Liebman, M., Nelson, G., van Everdingen, R. O. & Boike, J. (eds.): *State-of-the-art report on saturated water movement in permafrost areas. Report 4/97 (lnr 54)*, Agricultural University of Norway.
- French, H. 2008: Recent contributions to the study of past permafrost. *Permafrost and Periglacial Processes* 19, 179-194.

- French, H. M. 2007: *The Periglacial Environment*. 478 pp. John Wiley, London.
- French, H. M., Demitroff, M., Forman, S. L. & Newell, W. L. 2007: A chronology of Late-Pleistocene permafrost events in southern New Jersey, Eastern USA. *Permafrost and Periglacial Processes* 18, 49-59.
- French, H. M. & Millar, S. W. S. 2014: Permafrost at the time of the Last Glacial Maximum (LGM) in North America. *Boreas* 43, 667-677.
- Gao, C., Boreham, S., Preece, R., Gibbard, P. & Briant, R. 2007: Fluvial response to rapid climate change during the Devensian (Weichselian) Lateglacial in the River Great Ouse, southern England, UK. *Sedimentary Geology* 202, 193-210.
- Gibbard, P. L. & Allen, L. G. 1994: Drainage Evolution in South and East England During the Pleistocene. *Terra Nova* 6, 444-452.
- Glasser, N. F. & Siegert, M. J. 2002: Calculating basal temperatures in ice sheets: an Excel spreadsheet method. *Earth Surface Processes and Landforms* 27, 673-680.
- Gorbunov, A. 1978: Permafrost investigations in high-mountain regions. *Arctic and Alpine Research* 10, 283-294.
- Govaerts, J., Weetjens, E. & Beerten, K. 2011: Numerical simulation of permafrost depth at the Mol site. *External Report of the Belgian Nuclear Research Centre, SCK-CEN-ER-148*, 24 pp. Mol, Belgium.
- Guodong, C. & Shaoling, W. 1982: On the zonation of high-altitude permafrost in China. *Journal of Glaciology and Geocryology* 4, 1-17.
- Hall, A. M. & Glasser, N. F. 2003: Reconstructing the basal thermal regime of an ice stream in a landscape of selective linear erosion: Glen Avon, Cairngorm Mountains, Scotland. *Boreas* 32, 191-207.

- 1  
2  
3 Hallet, B., Walder, J. & Stubbs, C. 1991: Weathering by segregation ice growth in  
4 microcracks at sustained subzero temperatures: Verification from an experimental  
5 study using acoustic emissions. *Permafrost and Periglacial Processes* 2, 283-300.  
6  
7  
8  
9  
10 Halsey, L. A., Vitt, D. H. & Zoltai, S. C. 1995: Disequilibrium response of permafrost in  
11 boreal continental western Canada to climate change. *Climatic Change* 30, 57-73.  
12  
13  
14 Hartikainen, J., Kouhia, R. & Wallroth, T. 2010: *Permafrost simulations at Forsmark using a*  
15 *numerical 2D thermo-hydro-chemical model*. SKB TR-09-17, 142 pp. Svensk  
16 Kärnbränslehantering AB, Swedish Nuclear Fuel and Waste Management Company.  
17  
18  
19  
20  
21 Howard, A. J., Bridgland, D., Knight, D., McNabb, J., Rose, J., Schreve, D., Westaway, R.,  
22 White, M. J. & White, T. S. 2007: The British Pleistocene fluvial archive: East  
23 Midlands drainage evolution and human occupation in the context of the British and  
24 NW European record. *Quaternary Science Reviews* 26, 2724-2737.  
25  
26  
27  
28  
29  
30 Hubbard, A., Bradwell, T., Golledge, N., Hall, A., Patton, H., Sugden, D., Cooper, R. &  
31 Stoker, M. 2009: Dynamic cycles, ice streams and their impact on the extent,  
32 chronology and deglaciation of the British-Irish ice sheet. *Quaternary Science*  
33 *Reviews* 28, 758-776.  
34  
35  
36  
37  
38  
39 Huijzer, B. & Vandenberghe, J. 1998: Climate reconstruction of the Weichselian Pleniglacial  
40 in northwestern and central Europe. *Journal of Quaternary Science* 13, 391-417.  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50 Hutchinson, J. N. & Thomas-Betts, A. 1990. Extent of permafrost in southern Britain in  
51 relation to geothermal flux. *Quarterly Journal of Engineering Geology, London* 23,  
52 387-390.  
53  
54  
55  
56  
57  
58  
59  
60 Kemp, R. A., Whiteman, C. A. & Rose, J. 1993: Paleoenvironmental and Stratigraphic  
Significance of the Valley Farm and Barham Soils in Eastern England. *Quaternary*  
*Science Reviews* 12, 833-848.

- Lee, J. R., Brown, E. J., Rose, J., Moorlock, B. S. P. & Hamblin, R. J. O. 2003: A Reply to 'Implications of a Middle Pleistocene Ice-Wedge Cast at Trimingham, Norfolk, Eastern England'. *Permafrost and Periglacial Processes* 14, 75-78.
- Lee, J. R., Busschers, F. S. & Sejrup, H. P. 2012: Pre-Weichselian Quaternary glaciations of the British Isles, The Netherlands, Norway and adjacent marine areas south of 68°N: implications for long-term ice sheet development in northern Europe. *Quaternary Science Reviews* 44, 213-228.
- Lee, J. R., Rose, J., Hamblin, R. J., Moorlock, B. S., Riding, J. B., Phillips, E., Barendregt, R. W. & Candy, I. 2011: The Glacial History of the British Isles during the Early and Middle Pleistocene: Implications for the long-term development of the British Ice Sheet. In Ehlers, J., Gibbard, P. L. & Hughes, P. D. (eds.): *Quaternary Glaciations—Extent and Chronology, A Closer look. Developments in Quaternary Science*, 59-74, Elsevier, Amsterdam.
- Lee, M., Brown, G., Webb, P., Wheildon, J. & Rollin, K. 1987: Heat flow, heat production and thermo-tectonic setting in mainland UK. *Journal of the Geological Society* 144, 35-42.
- Lewis, S. G., Maddy, D. & Scaife, R. G. 2001: The fluvial system response to abrupt climate change during the last cold stage: the Upper Pleistocene River Thames fluvial succession at Ashton Keynes, UK. *Global and Planetary Change* 28, 341-359.
- Lisiecki, L. E. & Raymo, M. E. 2005: A Pliocene-Pleistocene stack of 57 globally distributed benthic  $\delta^{18}\text{O}$  records. *Paleoceanography* 20, PA1003.
- Livingstone, S. J., Evans, D. J., Ó Cofaigh, C., Davies, B. J., Merritt, J. W., Huddart, D., Mitchell, W. A., Roberts, D. H. & Yorke, L. 2012: Glaciodynamics of the central sector of the last British–Irish Ice Sheet in Northern England. *Earth-Science Reviews* 111, 25-55.

Maddy, D., Green, C. P., Lewis, S. G. & Bowen, D. Q. 1995: Pleistocene Geology of the Lower Severn Valley, Uk. *Quaternary Science Reviews* 14, 209-222.

Maslin, M., Shackleton, N. & Pflaumann, U. 1995: Surface water temperature, salinity, and density changes in the northeast Atlantic during the last 45,000 years: Heinrich events, deep water formation, and climatic rebounds. *Paleoceanography* 10, 527-544.

Mathers, S. J., Terrington, R. L., Waters, C. N. & Leslie, A. G. 2014: GB3D – a framework for the bedrock geology of Great Britain. *Geoscience Data Journal* 1, 30-42.

McDermott, F., Matthey, D. P. & Hawkesworth, C. 2001: Centennial-scale Holocene climate variability revealed by a high-resolution speleothem  $\delta^{18}\text{O}$  record from SW Ireland. *Science* 294, 1328-1331.

McEwen, T. & de Marsily, G. 1991: *The potential significance of permafrost to the behaviour of a deep radioactive waste repository*. SKI TR 91:8, 69 pp. Statens kärnkraftinspektion, Stockholm.

McManus, J. F., Oppo, D. W. & Cullen, J. L. 1999: A 0.5-million-year record of millennial-scale climate variability in the North Atlantic. *Science* 283, 971-975.

Muller, S. W. 1945: *Permafrost or permanently frozen ground and related engineering problems*. U.S. Geological Survey, Military Geology Unit, Strategic Engineering Study Special Report 62, pp. 135 (2nd printing, with corrections, 1945: pp. 231; reprint, 1947: Ann Arbor, Mich., J.W. Edwards).

Murton, J. B., Bateman, M. D., Baker, C. A., Knox, R. & Whiteman, C. A. 2003: The Devensian periglacial record on Thanet, Kent, UK. *Permafrost and Periglacial Processes* 14, 217-246.

Murton, J. B., Coutard, J. P., Lautridou, J. P., Ozouf, J. C., Robinson, D. A., Williams, R. B., Guillemet, G. & Simmons, P. 2000: Experimental design for a pilot study on bedrock

- weathering near the permafrost table. *Earth Surface Processes and Landforms* 25, 1281-1294.
- Murton, J. B., Peterson, R. & Ozouf, J.-C. 2006: Bedrock fracture by ice segregation in cold regions. *Science* 314, 1127-1129.
- Murton, J. B., Whiteman, C. A. & Allen, P. 1995: Involutions in the Middle Pleistocene (Anglian) Barham Soil, Eastern England - a Comparison With Thermokarst Involutions From Arctic Canada. *Boreas* 24, 269-280.
- Osterkamp, T. E. & Gosink, J. P. 1991: Variations in permafrost thickness in response to changes in paleoclimate. *Journal of Geophysical Research: Solid Earth* 96, 4423-4434.
- Patton, H., Hubbard, A., Glasser, N. F., Bradwell, T. & Golledge, N. R. 2013: The last Welsh Ice Cap: Part 1 – Modelling its evolution, sensitivity and associated climate. *Boreas* 42, 471-490.
- Penkman, K. E. H., Preece, R. C., Bridgland, D. R., Keen, D. H., Meijer, T., Parfitt, S. A., White, T. S. & Collins, M. J. 2011: A chronological framework for the British Quaternary based on Bithynia opercula. *Nature* 476, 446-449.
- Rollin, K. 1995: A simple heat-flow quality function and appraisal of heat-flow measurements and heat-flow estimates from the UK Geothermal Catalogue. *Tectonophysics* 244, 185-196.
- Rollin, K. 2002: Assessment of BGS data for ground source heat pump installations in the UK. *British Geological Survey Internal Report, IR/02/196* 50 pp.
- Rollin, K., Kirby, G. & Rowley, W. 1995: Atlas of geothermal resources in Europe: UK revision. *British Geological Survey Technical Report WK/95/07C*, 160 pp.
- Rose, J. 1994: Major river systems of central and southern Britain during the Early and Middle Pleistocene. *Terra Nova* 6, 435-443.

- Rose, J. & Allen, P. 1977: Middle Pleistocene stratigraphy in southeast Suffolk. *Journal of the Geological Society of London* 133, 83-102.
- Rose, J., Whiteman, C. A., Allen, P. & Kemp, R. A. 1999: The Kesgrave Sands and Gravels: 'pre-glacial' quaternary deposits of the River Thames in East Anglia and the Thames Valley. *Proceedings of the Geologists Association* 110, 93-116.
- Rozenbaum, G. & Shpolyanskaya, N. 1998: Late cenozoic permafrost history of the Russian Arctic. *Permafrost and Periglacial Processes* 9, 247-273.
- Ruskeenieniemi, T., Ahonen, L., Paananen, M., Frape, S., Stotler, R. L., Hobbs, M., Kaija, J., Degnan, P., Blomqvist, R., Jensen, M. R., Lehto, Jensen, M. R., Moren, L., Puigdomenech, I. & Snelling, M. 2004: Permafrost at Lupin: Report of Phase 2. *Geological Survey of Finland, YST-119*, 89 pp.
- Ruskeenieniemi, T., Paananen, M., Ahonen, L., Kaija, J., Kuivamäki, A., Frape, S., Moren, L. & Degnan, P. 2002: Permafrost at Lupin: Report of Phase 1. *Geological Survey of Finland, YST-112*, 59 pp.
- Seddon, M. B. & Holyoak, D. T. 1985: Evidence of sustained regional permafrost during deposition of fossiliferous Late Pleistocene river sediments at Stanton Harcourt (Oxfordshire, England). *Proceedings of the Geologists' Association* 96, 53-71.
- Shakun, J. D., Clark, P. U., He, F., Marcott, S. A., Mix, A. C., Liu, Z., Otto-Bliesner, B., Schmittner, A. & Bard, E. 2012: Global warming preceded by increasing carbon dioxide concentrations during the last deglaciation. *Nature* 484, 49-54.
- Siegert, M. J. & Dowdeswell, J. A. 2004: Numerical reconstructions of the Eurasian Ice Sheet and climate during the Late Weichselian. *Quaternary Science Reviews* 23, 1273-1283.
- Smith, L. C., Sheng, Y. & MacDonald, G. M. 2007: A first pan-Arctic assessment of the influence of glaciation, permafrost, topography and peatlands on northern hemisphere lake distribution. *Permafrost and Periglacial Processes* 18, 201-208.

- Stotler, R. L., Frape, S. K., Ruskeeniemä, T., Ahonen, L., Onstott, T. C. & Hobbs, M. Y. 2009: Hydrogeochemistry of groundwaters in and below the base of thick permafrost at Lupin, Nunavut, Canada. *Journal of Hydrology* 373, 80-95.
- Straw, A. 2013: Dartmoor Glaciation - fact or fiction. *Quaternary Newsletter* 129, 46-51.
- Straw, A. 2014: Dartmoor Glaciation - response to comments by David Evans *et al.* *Quaternary Newsletter* 130, 20-22.
- Szewczyk, J. A. N. & Nawrocki, J. 2011: Deep-seated relict permafrost in northeastern Poland. *Boreas* 40, 385-388.
- Thierens, M., Pirlet, H., Colin, C., Latruwe, K., Vanhaecke, F., Lee, J. R., Stuut, J. B., Titschack, J., Huvenne, V. A. I., Dorschel, B., Wheeler, A. J. & Henriot, J. P. 2012: Ice-rafting from the British-Irish ice sheet since the earliest Pleistocene (2.6 million years ago): implications for long-term mid-latitudinal ice-sheet growth in the North Atlantic region. *Quaternary Science Reviews* 44, 229-240.
- Vandenberghe, J., French, H. M., Gorbunov, A., Marchenko, S., Velichko, A. A., Jin, H., Cui, Z., Zhang, T. & Wan, X. 2014: The Last Permafrost Maximum (LPM) map of the Northern Hemisphere: permafrost extent and mean annual air temperatures, 25-17 ka BP. *Boreas* 43, 652-666.
- Waelbroeck, C., Paul, A., Kucera, M., Rosell-Melé, A., Weinelt, M., Schneider, R., Mix, A., Abelmann, A., Armand, L. & Bard, E. 2009: Constraints on the magnitude and patterns of ocean cooling at the Last Glacial Maximum. *Nature Geoscience* 2, 127-132.
- Walker, M., Coope, G., Sheldrick, C., Turney, C., Lowe, J., Blockley, S. & Harkness, D. 2003: Devensian Lateglacial environmental changes in Britain: a multi-proxy environmental record from Llanilid, South Wales, UK. *Quaternary Science Reviews* 22, 475-520.

Watson, E. & Morgan, A. 1977: The Periglacial Environment of Great Britain During the Devensian [and Discussion]. *Philosophical Transactions of the Royal Society of London. B, Biological Sciences* 280, 183-198.

West, R. G. 1980: *The Pre-glacial Pleistocene of the Norfolk and Suffolk Coasts*. 204 pp. Cambridge University Press, Cambridge.

Westaway, R., Bridgland, D. & White, M. 2006: The Quaternary uplift history of central southern England: evidence from the terraces of the Solent River system and nearby raised beaches. *Quaternary Science Reviews* 25, 2212-2250.

Westaway, R. & Younger, P. L. 2013: Accounting for palaeoclimate and topography: A rigorous approach to correction of the British geothermal dataset. *Geothermics* 48, 31-51.

White, T. S., Bridgland, D. R., Westaway, R., Howard, A. J. & White, M. J. 2010: Evidence from the Trent terrace archive, Lincolnshire, UK, for lowland glaciation of Britain during the Middle and Late Pleistocene. *Proceedings of the Geologists' Association* 121, 141-153.

- Figure 1. A) A surface temperature history for Prudhoe Bay, Alaska, based on an East Siberia model (after Osterkamp & Gosink 1991). B) The same surface temperature history represented as a series of step changes in temperature.
- Figure 2. Selected temperature proxy records used to construct the average temperature trends over the UK during the past 140 ka. A) European pollen records from Davis *et al.* (2003). B) Sea surface temperature and land proxy records from Shakun *et al.* (2012). C) Sea surface temperature records from Maslin *et al.* (1995). D) Sea surface temperature records from McManus *et al.* (1999). Note that curves A, B and C are drawn against the top time axis and curve D against the bottom time axis. The dashed lines are to facilitate comparison between the curves, with the shaded area representing time greater than 40 ka BP.
- Figure 3. The reconstructed temperature trend for Great Britain over the last 130 ka (B), aside the appended data used from several sources and regions to create the record (A; see also Fig. 2 where the four elements used to construct curve A are shown). Ice core temperature proxy shown for comparison (C). This reconstructed temperature trend was used as a basis for the surface temperature histories at each of the six localities, where the maximum and minimum MAAT were adjusted, along with the effect of ice sheet presence, at each location.
- Figure 4. Surface temperature histories for A) East Anglia which was ice free and the minimum temperature of 12 °C below present occurred at the peak of the last glaciation. B) The Stainmore Trough, where ice cover between 110-102, 76-37 and 34-14.5 ka BP insulated the land surface from the extremes of air temperature. Hence, despite a minimum MAAT of 14 °C below present, the minimum temperature of 9.8 °C below present occurred after the ice had retreated. Periods of ice cover are indicated by the blue bars.
- Figure 5. The six selected locations across Great Britain for permafrost modelling overlaid on the 1:625 000 scale bedrock geology. The geological key for this map can be viewed at <http://mapapps.bgs.ac.uk/geologyofbritain/home.html>.
- Figure 6. Sketch diagrams (not to scale) of the geological succession at each of the glaciated locations. A) Mid-Wales. B) The Northwest Highlands. C) The Stainmore Trough. D) The Midland Valley. Depths (m), relative to Ordnance Datum (+ve above sea level), are shown to the left of each sketch and the geology (with, in brackets, the thermal conductivity in  $\text{W m}^{-1} \text{K}^{-1}$ ) on the sketch. The rock unit name (left part of geology code) can be found at <http://www.bgs.ac.uk/lexicon/> and the rock description (right part of geology code) can be found at [http://www.bgs.ac.uk/data/vocabularies/dictionary.cfm?name=DIC\\_ROCK\\_SI\\_GMA](http://www.bgs.ac.uk/data/vocabularies/dictionary.cfm?name=DIC_ROCK_SI_GMA).
- Figure 7. Sketch diagrams (not to scale) of the geological succession at each of the unglaciated locations. A) Dartmoor. B) East Anglia. Depths (m), relative to Ordnance Datum (+ve above sea level), are shown to the left of each sketch and the geology (with, in brackets, the thermal conductivity in  $\text{W m}^{-1} \text{K}^{-1}$ ) on or to the right of the sketch. The rock unit name (left part of geology code) can be found at <http://www.bgs.ac.uk/lexicon/> and the rock description (right part of geology code) can be found at

[http://www.bgs.ac.uk/data/vocabularies/dictionary.cfm?name=DIC\\_ROCK\\_SI\\_GMA](http://www.bgs.ac.uk/data/vocabularies/dictionary.cfm?name=DIC_ROCK_SI_GMA).

- Figure 8. Permafrost evolution modelling for the upland glaciated locations presented as classed plots of ground temperatures from 126 ka BP to the present. Mid-Wales is shown in the top panel and the Northwest Highlands in the lower panel.
- Figure 9. Permafrost evolution modelling for the lowland glaciated locations presented as classed plots of ground temperatures from 126 ka BP to the present. The Stainmore Trough is shown in the top panel and the Midland Valley in the lower panel.
- Figure 10. Permafrost evolution modelling for the unglaciated locations presented as classed plots of ground temperatures from 126 ka BP to the present. Dartmoor, which is an upland unglaciated location, is shown in the top panel and East Anglia, which is a lowland unglaciated location, in the lower panel.
- Table 1. Descriptions of the permafrost modelling locations. Coordinates are British National Grid and elevation is relative to Ordnance Datum (OD, +ve above mean sea level). The minimum mean annual air temperature (MAAT) relative to the present, predicted by the surface temperature histories, is shown along with periods of ice cover and assumed temperatures at the base of the ice (relative temperature, and in brackets, the absolute temperature).
- Table 2. Maximum regional, modelled thicknesses of permafrost at the six locations.






Figure 1  
195x304mm (72 x 72 DPI)








Figure 4  
180x367mm (200 x 200 DPI)




Figure 5  
70x122mm (300 x 300 DPI)




Figure 6




Figure 7  
99x71mm (220 x 220 DPI)




Figure 8  
668x554mm (100 x 100 DPI)




Figure 9  
668x554mm (100 x 100 DPI)




Figure 10  
668x554mm (100 x 100 DPI)

	Mid-Wales	NW Highlands	Stainmore Trough	Midland Valley	Dartmoor	East Anglia
Easting (m)	310453	230850	444128	290971	278000	603000
Northing (m)	292587	861650	523367	686182	84000	287294
Elevation (m OD)	160	165	28	66	257	42
Present day MAAT (°C)	8.8	7.2	9.2	9.0	9.0	10.0
Heat flow (W m <sup>-2</sup> )	0.055	0.058	0.067	0.069	0.105	0.035
Min MAAT relative to present (°C)	-12	-17	-14	-14	-12	-12
Periods of ice cover (ka BP)	40-37; 29-26; 24-15.5	110-12	110-102; 76-37; 34-14.5	62-37; 34-14.5	No ice cover	No ice cover
Temp. below ice relative to present (and absolute) (°C)	-11.8 (-3)	-10.2 (-3)	-8.2 (+1)	-12.0 (-3)		

Table 1

Location		Maximum thickness of permafrost (m)
Upland glaciated	Mid-Wales	105
	Northwest Highlands	180
Lowland glaciated	Stainmore Trough	20
	Midland Valley	110
Upland unglaciated	Dartmoor	80
Lowland unglaciated	East Anglia	65

Table 2

For Review Only