

Article (refereed) - postprint

Medinets, Sergiy; Skiba, Ute; Rennenberg, Heinz; Butterbach-Bahl, Klaus.
2015. **A review of soil NO transformation: associated processes and possible physiological significance on organisms.**

© 2014 Elsevier Ltd.

This manuscript version is made available under the CC-BY-NC-ND 4.0 license <http://creativecommons.org/licenses/by-nc-nd/4.0/>



This version available <http://nora.nerc.ac.uk/512955/>

NERC has developed NORA to enable users to access research outputs wholly or partially funded by NERC. Copyright and other rights for material on this site are retained by the rights owners. Users should read the terms and conditions of use of this material at <http://nora.nerc.ac.uk/policies.html#access>

NOTICE: this is the author's version of a work that was accepted for publication in *Soil Biology and Biochemistry*. Changes resulting from the publishing process, such as peer review, editing, corrections, structural formatting, and other quality control mechanisms may not be reflected in this document. Changes may have been made to this work since it was submitted for publication. A definitive version was subsequently published in *Soil Biology and Biochemistry* (2015), 80. 92-117. [10.1016/j.soilbio.2014.09.025](https://doi.org/10.1016/j.soilbio.2014.09.025)

www.elsevier.com/

Contact CEH NORA team at
noraceh@ceh.ac.uk

Title page

**A review of soil NO transformation: associated processes and
possible physiological significance on organisms**

Sergiy Medinets^{a,b,c}, Ute Skiba^d, Heinz Rennenberg^c, Klaus Butterbach-
Bahl^{b,e*}

^aRegional Centre for Integrated Environmental Monitoring, Odessa
National I. I. Mechnikov University (ONU), Mayakovskogo lane 7, 65082
Odessa, Ukraine, s.medinets@gmail.com

^bInstitute for Meteorology and Climate Research (IMK), Karlsruhe
Institute of Technology (KIT), Kreuzeckbahnstraße 19, D-82467 Garmisch-
Partenkirchen, Germany, klaus.butterbach-bahl@kit.edu

^cInstitute of Forest Sciences, Chair of Tree Physiology, University of
Freiburg, Georges-Koehler-Allee 53/54, D-79110 Freiburg, Germany,
heinz.rennenberg@ctp.uni-freiburg.de

^dCentre for Ecology and Hydrology (CEH) Edinburgh, Bush Estate,
Penicuik, Midlothian, EH26 0QB, United Kingdom, ums@ceh.ac.uk

^eInternational Livestock Research Institute (ILRI), Old Naivasha Road,
Nairobi, Kenya

**Corresponding author:*
Klaus Butterbach-Bahl
Institute for Meteorology and Climate Research (IMK), Karlsruhe
Institute of Technology (KIT)
Kreuzeckbahnstraße 19, D-82467 Garmisch-Partenkirchen, Germany
tel.: +49 8821 183136
e-mail: klaus.butterbach-bahl@kit.edu

**A review of soil NO transformation: associated processes and
possible physiological significance on organisms**

Sergiy Medinets^{a,b,c}, Ute Skiba^d, Heinz Rennenberg^c, Klaus Butterbach-
Bahl^{b,e*}

^aRegional Centre for Integrated Environmental Monitoring, Odessa National I. I.
Mechnikov University (ONU), Mayakovskogo lane 7, 65082 Odessa, Ukraine

^bInstitute for Meteorology and Climate Research (IMK), Karlsruhe Institute of
Technology (KIT), Kreuzeckbahnstraße 19, D-82467 Garmisch-Partenkirchen, Germany,

^cInstitute of Forest Sciences, Chair of Tree Physiology, University of Freiburg,
Georges-Koehler-Allee 53/54, D-79110 Freiburg, Germany

^dCentre for Ecology and Hydrology (CEH) Edinburgh, Bush Estate, Penicuik,
Midlothian, EH26 0QB, United Kingdom

^eInternational Livestock Research Institute (ILRI), Old Naivasha Road, Nairobi, Kenya

Abstract

NO emissions from soils and ecosystems are of outstanding importance
for atmospheric chemistry. Here we review the current knowledge on
processes involved in the formation and consumption of NO in soils, the
importance of NO for the physiological functioning of different organisms,
and for inter- and intra-species signalling and competition, e.g. in the
rooting zone between microbes and plants. We also show that prokaryotes
and eukaryotes are able to produce NO by multiple pathways and that
unspecific enzymo-oxidative mechanisms of NO production are likely to
occur in soils. Nitric oxide production in soils is not only linked to NO
production by nitrifying and denitrifying microorganisms, but also linked to
extracellular enzymes from a wide range of microorganisms.

Further investigations are needed to clarify molecular mechanisms of NO production and consumption, its controlling factors, and the significance of NO as a regulator for microbial, animal and plant processes. Such process understanding is required to elucidate the importance of soils as sources (and sinks) for atmospheric NO.

Key words: nitric oxide, nitrification, denitrification, unspecific enzymo-oxidative mechanisms, NOS, SOD, dissimilatory nitrate reduction to ammonium, chemodenitrification, NO signalling, NO consumption, NO production, archaea, protists, invertebrates

Highlights

- Abiotic and biotic pathways of NO transformation are discussed
- Interrelation between NO transformation processes is discussed
- Unspecific enzymo-oxidative mechanisms of NO transformation are proposed
- Physiological NO functions/effects in/for various groups of organisms are shown
- Importance of bacterial NO as signalling substance for others organisms is highlighted

1. Introduction

Nitric oxide (nitrogen monoxide, NO) is a highly reactive constituent of the troposphere (Fowler et al., 2009) and is considered to be the main

precursor of ground-level tropospheric ozone in rural areas (Chameides et al., 1994; Laville et al., 2011), impacting human health and plant productivity (Staffelbach et al., 1997; Ludwig et al., 2001). The main sources of NO in the troposphere are fossil fuel combustion, biomass burning, soil emissions and lightning (Delmas et al., 1997). Nevertheless agricultural soils can be the predominant NO source in rural regions, where the contribution of fossil fuel combustion is low (Bouwman et al., 2002; Butterbach-Bahl et al., 2009).

The global soil NO production is estimated at $\sim 8.9 \text{ Tg N a}^{-1}$, of which $\sim 15\%$ is produced in Europe (IPCC, 2007). However, an earlier estimate by Davidson and Kinglerlee's (1997) provided much higher values ranging from 13 to 21 Tg N a^{-1} . Such large divergence between estimates results from insufficient knowledge of the full range of soil microbial processes involved in NO production and consumption and the interactions of these processes with environmental variables.

Biological N transformation processes in soils, namely nitrification and denitrification, are usually considered the dominant sources of soil NO production. However, also abiotic chemical N transformations can be an important source (Ludwig et al., 2001; Butterbach-Bahl et al., 2011, 2013). Linking NO production, consumption and emission to the source and sink processes of nitrification and denitrification *in situ* still remains challenging, as they can occur simultaneously and in the same soil aggregates (Arah, 1997). Moreover, they can be spatially or temporally linked to each other, using products (Garrido et al., 2002) and/or intermediates from one process by the other (Butterbach-Bahl et al., 2013). That is why *in vitro*

experiments, applying molecular methods, stable isotopes and inhibitors are important to disentangle processes and mechanisms involved in soil NO production and emission. Simulating field conditions for different soil types from a wide range of climate zones will allow us to crack open the veil of soil NO transformations and reveal potential mechanisms and drivers. Better process understanding is the basis to develop mitigation strategies for reducing soil NO emissions.

Endogenous NO is generally considered as a freely diffusible molecule in cells with a significant importance as a signaling substance. Thus, NO acts as a short-lived messenger molecule with numerous molecular targets, playing numerous physiological roles at organelle, intra- and inter-cellular levels in both prokaryotes and eucaryotes (Jacklet, 1997; Gusarov et al., 2008; Johnson et al., 2008; Leitner et al., 2009; Velayutham and Zweier, 2013).

The main purpose of this review is to present recent advances from field and laboratory studies focusing on NO transformation and underlying processes as well as investigating the potential of other processes not yet associated with NO production or consumption, and to highlight the physiological and ecological significance of such processes.

2. Reactivity of NO

NO is a stable free radical with an ionization potential of 9.26 eV and an electron affinity of 0.024 eV (Natalis et al., 1979). Its high reactivity is due to its electronic configuration, i.e. the existence of an unpaired electron

residing in a π^* molecular orbital (Wong et al., 1989). Consequently, NO can be easily oxidized to the nitrosonium ion (NO^+), reduced to the nitroxide ion (NO^-), or converted to nitrogen dioxide (NO_2) by oxygen (O_2) (McCleverty, 2004 and reference therein). NO and its ions share isoelectronic properties with other molecule and ions. For example, NO is isoelectronic with O_2^+ , meanwhile NO^- is isoelectronic with O_2 and NO^+ with CO and CN^- (McCleverty, 2004 and reference therein). A very important property of nitric oxide related to its redox-activity in solution is its ability to form nitrosyl as well as multi nitrosyl complexes with transitional metals (e.g., Fe, Mn, Co, Ru) and metal-containing enzymes (e.g., copper-containing nitrite reductase (NIR)) (Ruggiero et al., 1993; Ford and Lorkovic, 2002; Lee et al., 2002 and references therein). It has been shown that the reversible process $\text{NO} \leftrightarrow \text{NO}^+$ in water is strongly pH-dependent (Lee et al., 1990; Kim and Kochi, 1991) and NO can be produced from nitrite NO_2^- under strongly basic conditions (Stanbury, 1989).

3. Soil processes associated with NO production and consumption

The main microbiological processes of N transformation in soils, such as nitrification, nitrifier and heterotrophic denitrification, as well as abiotic chemodenitrification are classically considered as important pathways of both soil NO production and consumption under different environmental condition (Firestone and Davidson, 1989; Conrad, 1996; Yamulki et al., 1997; Skiba et al., 1997; Zumft, 1997; Gasche and Papen, 1999; Ludwig et al., 2001; Wrage et al., 2001; Garrido et al., 2002; Venterea et al., 2005;

170 Kesik et al., 2006; Robertson and Groffman, 2007; Skiba, 2008; Kool et al.,
 171 2009a, 2009b; Bru et al., 2010; Wu et al., 2010; Baggs, 2011; Ju et al.,
 172 2011; Butterbach-Bahl et al., 2011, 2013; Bakken et al., 2012; Luo et al.,
 173 2012; Schreiber et al., 2012; Barton et al., 2013; Pilegaard, 2013 and many
 174 others). In a recent review Schreiber et al. (2012) provided an overview of
 175 microbial and chemical NO and N₂O production processes and innovative
 176 experimental approaches, but did not include the role of NO in higher
 177 organisms. Another recent review by Pilegaard (2013) focused on soil NO
 178 emission and its regulating factors, but did not include process description at
 179 the organism level. To fill these gaps we have considered additional
 180 processes associated with NO exchange, for example codenitrification (e.g.,
 181 Shoun et al., 1992; Tanimoto et al., 1992; Spott et al., 2011), dissimilatory
 182 nitrate reduction to ammonium (e.g., Bengtsson and Bergwall, 2000; Silver
 183 et al., 2001, 2005; Rütting et al., 2008; Templer et al., 2008; Wan et al.,
 184 2009; Schmidt et al., 2011), anaerobic ammonium oxidation (e.g., Strous et
 185 al., 1996; Humbert et al., 2010; Kartal et al., 2011), nitrite-dependant
 186 anaerobic oxidation of methane (e.g., Raghoebarsingetal, 2006; Ettwig et
 187 al., 2010; Harron et al., 2013), nitric oxide synthase mediated NO
 188 production (e.g., Fritz-Laylin et al., 2009; Messner et al., 2009; Chen et al.,
 189 2010; Forstermann and Sessa, 2012) and the theoretically feasible,
 190 unspecific enzyme mediated mechanisms of oxidation of soil N described
 191 for the first time in this review in detail. We also provide a brief overview of
 192 the physiological functions of NO in different groups of organisms living in
 193 and on soil (e.g., Jacklet 1997; Gusarov et al., 2008, 2009; Johnson et al.,

2008; Fritz-Laylin et al., 2009; Schreiber et al., 2011; Forstermann and Sessa, 2012).


3.1. Abiotic processes

3.1.1. Chemodenitrification

The term chemodenitrification describes the strictly chemical, non-enzymatic conversion of nitrite (NO_2^-) or nitrate (NO_3^-) to gaseous nitrogen species at low pH (below 5). This process normally requires the presence of ammonium (NH_4^+), amines or reduced metals (e.g. Fe^{2+}), as well as high soil organic matter (Clark, 1962; Broadbent and Clark, 1965; Wullstein and Gilmour, 1966; Chalk and Smith, 1983; Zumft, 1997) and soil water contents (Venterea et al., 2005). The most important reaction of chemodenitrification (Equation (1)) is the formation of NO via nitrous acid (HNO_2 (aqueous phase), HONO (gas phase)) decomposition (Van Cleemput and Baert, 1976; Chalk and Smith, 1983; Zumft, 1997; Venterea et al., 2005):




If reduced metals are available (e.g. Fe^{2+}) the Equation (2) can be presented as:



pH is the major controlling factor for chemodenitrification in soils (Chalk and Smith, 1983; Zumft, 1997), while NO_2^- concentrations (Ludwig et al., 2001), temperature (Kesik et al., 2005, 2006) and soil water content (Venterea et al., 2005) have been identified as additional controllers. The chemical decomposition of NO_2^- mainly occurs under acidic soil conditions ($\text{pH} < 4.5$), and Yamulki et al. (1997) detected NO emissions from sterile acidic soil. However, also at more neutral pH (5 – 7) ranges, NO may be produced chemically or react with humic substances producing N_2O and CO_2 (Porter, 1969; , Stevenson et al., 1970). As for every chemical reaction, reaction rates increase with rising temperature (Kesik et al., 2006) and high rates of soil NO emissions during warm periods from acidic soils were attributed partially to chemodenitrification in agricultural (Cheng et al., 2004) and N-affected temperate forest soils (Kesik et al., 2006; Luo et al., 2012).

Another soil related source of atmospheric NO is the emission of HONO from acidic soils (Su et al., 2011):



Air concentrations of HONO determine the sink and source function of soils. If air HONO concentrations are lower than in the soil aqueous or gaseous phase, a net emission is observed, while otherwise soils function as a sink for atmospheric HONO (Su et al., 2011). For instance, in typical acidic ($\text{pH} 4\text{--}5$) tropical forest and boreal soils even small soil NO_2^-

concentrations (ca. 0.001-0.01 $\mu\text{g g}^{-1}$) can lead to significant HONO emissions into the atmosphere (Su et al., 2011). Therefore, this process seems to be important at least for some natural ecosystems and may be an additional source of atmospheric NO and OH⁻ (Su et al., 2011).

3.1.2. Chemical consumption

It is widely known that abiotic nitrosation reactions via NO₂⁻ can result in N immobilization or ‘chemodenitrification’ including the production of NO, N₂O or N₂ (e.g., Bremner and Fúhr, 1966; Stevenson et al., 1970; Williams, 2004). Since both NO₂⁻ and NO can be considered as nitroso donors and since this reaction is likely to be reversible (Spott et al., 2011 and references there in), it can be assumed that under observed soil NO concentrations of 60-180 ppbv (Dong, Simon and Rennenberg, unpublished data), not only NO₂⁻ but also NO should be involved in abiotic nitrosation reactions. In particular, the nitrosation reactions of NO₂⁻ (and thus also of NO) with humic substances (e.g., secondary aliphates, aromates, amides) have been widely reported and proposed to be considered as an abiotic pathway of N incorporation into soil organic matter (SOM) (Bremner and Fúhr, 1966; Smith and Chalk, 1980; van Cleemput and Samater, 1996; Thorn and Mikita, 2000). Azhar et al. (1986a,b) provide evidence that during nitrification the NO₂⁻ (and likely NO) formed contributes to the nitrosation of organic matter under neutral or weak acidic soil pH conditions. Comparable results for the reaction of NO and organic matter have been reported by Stephenson (1970). In addition, metal-nitrosyl

complexes as formed e.g. during denitrification can function as a nitrosyl donor to a variety of N-, O-, S- and C-nucleophilic organic matter constituents (Garber and Hollocher, 1982b). This seems to be a significant process for SOM nitrosation in fertilized soils with high $\text{NH}_4^+/\text{NH}_3$ concentrations (Thorn and Mikita, 2000) where NO_2^- accumulates due to the inhibition of *Nitrobacter* spp. - driving the conversion of NO_2^- to NO_3^- in neutral to high pH soils –by increased levels of NH_3 .

Chemical reactions of NO in aqueous solution are well documented (e.g. Williams, 2004) and should occur in soils too. According to Williams (2004) NO in aqueous solution (irrespective of the pH) can react with: i) amides to produce N-nitrosamides, ii) alcohols to give alkyl nitrites, iii) hydrogen peroxide to generate peroxyxynitrous acid, and iv) thiols to form S-nitrosothiols. Moreover, in aerated water NO may react with O_2 to produce NO_2 , which can further react with NO to form the nitrosating agent N_2O_3 , which then hydrolyzes to NO_2^- (Williams, 2004 and references therein).

The main pathway of consumption of soil emitted NO in surface air and/or inside the canopy is its rapid reaction with O_3 or R-OO^* (derived from the reaction of mostly biogenic volatile organic carbon (VOC) with OH^*) to form NO_2 . Plant leaves can take up NO_2 and further metabolize it. Several studies (Geßler et al., 2000; Butterbach-Bahl et al., 2004; Sparks, 2009) have suggested that soil NO emission and in-canopy conversion to NO_2 results in re-deposition onto plant leaves and uptake as NO_2 . Thus, soil NO emissions can be an important process of nutrient dispersal and recycling at ecosystem scale. Also direct diffusive uptake of atmospheric NO by leaves constitutes a canopy sink. However, due to the low solubility

of NO in the aqueous solution of the apoplastic space, this process is less important than plant leaf uptake of NO₂ (Hanson and Lindberg, 1991). A second possible pathway of atmospheric consumption of soil NO is the reversible reaction with OH⁻ to form HONO (Su et al., 2011).


In the troposphere, NO can react with hydroperoxy radicals (HO₂^{*}) (Hertel et al., 2011) and organic peroxy radicals (RO₂^{*}) (Finlayson-Pitts and Pitts, 1986; Pringlecombe, 1996) to produce NO₂. In sunlight (hν = 200-420 nm) NO₂ photo-dissociates to form NO and the very short-lived O(³P) radical, which in most cases combines with O₂ to form O₃; during night-time NO₂ can react with O₃ to form the NO₃^{*} radical and O₂ (Pringlecombe, 1996, Hertel et al., 2011). After aldehydes are photo-dissociated or react with OH⁻, an alkyl radical is formed and can be converted to a peroxy acetyl radical, which can react with NO₂ to form peroxy acetyl nitrate (PAN) (Pringlecombe, 1996, Fowler et al., 2009). Alternatively, NO₂ can react with OH⁻ to form HNO₃ at an average rate of ca. 5% per hour (Pringlecombe, 1996).

3.2. Biotic processes

3.2.1. Nitrification

Nitrification is the biological oxidation of ammonium (NH₄⁺) via hydroxylamine (NH₂OH) to nitrite (NO₂⁻) and further on to nitrate (NO₃⁻) (Equation (4)) (Wrage et al., 2001; Butterbach-Bahl et al., 2011, 2013). It is

one of the most important processes of ecosystem N-cycling, both in agricultural and natural soils (Ludwig et al., 2001).



Nitrification can be performed by heterotrophic and autotrophic nitrifiers. Autotrophic nitrifiers use the oxidation of NH_4^+ or NO_2^- as an energy source for CO_2 fixation, while heterotrophic nitrifiers use N-containing organic substances as energy and C source (Prosser, 1989; Wrage et al., 2001; Arp et al., 2002; Conrad, 2002; Costa et al., 2006; Butterbach-Bahl et al., 2011). Heterotrophic nitrifiers (e.g. *Arthrobacter*) can oxidize both NH_4^+ and organic N with similar intermediates, but use different enzymes for the transformation of these substrates (Wrage et al., 2001; Conrad, 2002). Ammonium oxidizing bacteria (AOB) are very specific organisms, e.g. *Nitrosomonas*, *Nitrosospira*, *Nitrosococcus* spp., that oxidize NH_4^+ to NH_2OH catalysed by ammonia monooxygenase (AMO) and NH_2OH to NO_2^- catalysed by hydroxylamine oxidoreductase (HAO). Ammonium can also be oxidized by autotrophic ammonium oxidizing archaea (AOA), belonging to the phylum *Thaumarchaeota* (Könneke et al., 2005; Brochier-Armanet et al., 2008; Tourna et al., 2008; Martens-Habbena et al., 2009; Spang et al., 2010). AOA may even dominate NH_4^+ oxidation in soils (Leininger et al., 2006; Prosser and Nicol, 2008, 2012). Nitrite-oxidizing bacteria (NOB), e.g. *Nitrobacter*, *Nitrospira*, *Nitrococcus*, *Nitrospina* spp.,

perform further oxidation of NO_2^- to NO_3^- , catalysed by nitrite oxidoreductase (NXR).

Whilst heterotrophic nitrifying bacteria can use ammonia as well as organic N forms as substrate (Papen et al., 1989), fungal nitrification seems to exclusively rely on organic pathways (Robertson and Groffman, 2007):



Typically this process involves oxidation of amines or amides, is not coupled to ATP production and, therefore, is not involved in heterotrophic energy production (Robertson and Groffman, 2007). Heterotrophic nitrifiers have been shown to produce NO from organic N and inorganic substrates (e.g. Papen et al., 1989).

Altogether, a large number of heterotrophic bacteria (e.g. *Paracoccus*, *Alcaligenes*, *Thiosphaera*, *Pseudomonas* spp., described by Kuenen and Robertson (1994), Moir et al. (1996), Daum et al. (1998), Nishio et al. (1998)) and fungi (e.g. *Ascomycota* and *Basidiomycota* (Shoun et al., 1992, 2012; Prendergast-Miller et al., 2011), and *Glomeromycota* groups (Cousins et al., 2003; Porras-Alfaro et al., 2011; Bates et al., 2012)) can nitrify.

In soil solution with sufficient oxygen supply, nitrification is controlled predominantly by the availability of NH_4^+ (Robertson, 1989; Ludwig et al., 2001) or easy decomposable organic N (e.g. amines and amides), if fungal nitrification prevails (Conrad, 2002). Many studies support the idea that in a wide range of soils nitrification is the dominating process for soil NO

production as an intermediate in the oxidation of NH_2OH to NO_2^- (Hooper and Terry, 1979; Firestone and Davidson, 1989; Bollmann et al., 1999; Dunfield and Knowles, 1999; Gasche and Papen, 1999; Godde and Conrad, 2000; Venterea and Rolston, 2000; Ludwig et al., 2001; Garrido et al., 2002; Cheng et al., 2004; Wan et al., 2009; Wu et al., 2010; Ju et al., 2011; Mei et al., 2011; Cui et al., 2012; Luo et al., 2012 and others). Rates of nitric oxide formation during nitrification were estimated as 0.1-10% of gross NH_4^+ oxidation (Ludwig et al., 2001 and reference therein), but Garrido et al. (2002) reported a tighter range of 0.6-2.5%. It is also well known that some, but not all, AOB and AOA in both natural and agricultural soils are very sensitive to high substrate concentrations and that nitrification can be inhibited by substrate concentrations in the range of 1.0–5.0 mM NH_4^+ or NH_3 (Anthonisen et al., 1976; Stark and Firestone, 1996; Shi and Norton, 2000; Carrera et al., 2004; Koper et al., 2010; Norton and Stark, 2011). AOB, less sensitive to NH_3 compared to AOA (Prosser and Nicol, 2012), prefer to colonize areas with high soil NH_4^+ or NH_3 concentrations (Hayden et al., 2010; Ollivier et al., 2011).

As for all biological processes, temperature is an important parameter determining the rate of nitrification (Machefert et al., 2002; Robertson and Groffman, 2007) with specific optima depending on the microbial community active in different environments (Singh et al., 1993; Stark, 1996; Stark and Firestone, 1996; Norton and Stark, 2011). In general temperature optima for AOB of temperate climate zone soils are around 22-30 °C (Koops et al., 1991; Singh et al., 1993; Stark, 1996; Stark and Firestone, 1996; Norton and Stark, 2011), however, for tropical soils optima can be

close to 35 °C (Myers, 1975). In spite of these high temperature optima, reasonable rates of nitrifier activity were reported also at low soil temperatures, such as 2–10 °C (Cookson et al., 2002; Avrahami et al., 2003; Avrahami and Conrad, 2005), and were even observed in frozen soil together with detectable NO emission rates (Freppaz et al., 2007). The temperature effect on nitrification has been described by many process models. For example, Stark (1996) tested 5 different models and argued that the best fit model, the generalized Poisson density function (Parton et al., 1987), successfully describes the temperature response of nitrification activity over a temperature range of 5-50 °C. But he also stated that the Arrhenius equation (Laudelout, 1978) can still be used, providing adequate simulation over a more narrow temperature range of 5-28 °C (Fig. 1).

INSERT Fig. 1 HERE

The increase in NO emission rates in response to temperature is site specific (Saad and Conrad, 1993; Martin et al., 1998; Gasche and Papen, 1999; Ludwig et al., 2001; Schindlbacher et al., 2004; Kitzler et al., 2006; Laville et al., 2009; Yao et al., 2010). However, over the temperature range 0 to 35 °C the average NO response shows a Q_{10} of $\approx 2-4$ (Williams and Fehsenfeld, 1991; Martin et al., 1998; Gasche and Papen, 1999; Yu et al., 2008, 2010; Laville et al., 2009; Yao et al., 2010).

Optimum conditions for nitrification are normally met at a water filled pore space (WFPS) of 30-60% (Firestone and Davidson, 1989; Bouwman., 1998; Davidson et al., 2000). Following the conceptual Hole-In-the-Pipe

(HIP) model of Firestone and Davidson (1989), soil moisture content seems to be the most general and robust driver for determining the proportions of soil N gases emitted from different ecosystems, with NO dominating soil N gas emissions at WFPS <30-60%, and N₂O and N₂ dominating soil N gas emissions at WFPS >60-65%.

A soil with near neutral pH of 6.5-7.0 (Killham, 1990; Machefert et al., 2002) generally appears to favor nitrification by AOB and also mesophilic archaea (Jung et al., 2014; Stieglmeier et al., 2014a). The pH optimum is much lower (ca. 4.5) for acidophilic AOA (Nicol et al., 2008; Lehtovirta-Morley et al., 2011). Nitrification rates were found to be strongly ($p < 0.05$) correlated with NO production during incubation experiments for a range of acidic, neutral and alkaline soils (Garrido et al., 2002; Cheng et al., 2004). Highest nitrification rates as well as NO emissions were observed for neutral to alkaline soils (Cheng et al., 2004). For example, nitrification is thought to be the main process for NO production in cropland on calcareous soils (Wan et al., 2009; Ju et al., 2011; Mei et al., 2011; Cui et al., 2012) and in acid forest soils receiving high rates of atmospheric N (Gasche and Papen, 1999; Wu et al., 2010; Luo et al., 2012). In other studies (Nagele and Conrad, 1990; Yamulki et al., 1997; Ste-Marie and Pare, 1999) increasing pH stimulated nitrification rates and N₂O and NO release under aerobic conditions. Prevailing NO production was also shown in aerobic soils by Garrido et al. (2002). In aerobic and anaerobic incubation experiments with five soil types plus or minus the addition of 10 Pa of the nitrification inhibitor acetylene (C₂H₂) to the headspace the authors showed that NO was likely to be produced exclusively from nitrification. Zhu et al. (2013)

suggested that at high O₂ concentration (21%) nitrification seems to be the main responsible process for NO formation from NH₃.

3.2.1.1. AOB vs. AOA: distribution and contribution to nitrification


In terrestrial ecosystems where the total soil N concentration is greater than 0.7%, nitrification is a highly significant and important process (Ollivier et al., 2011). Based on data of alpine glacier forefields in Austria (Nicol et al., 2005; Deiglmayr et al., 2006; Kandeler et al., 2006; Hämmerli et al., 2007) and Switzerland (Duc et al., 2009; Lazzaro et al., 2009; Brankatschk et al., 2011) it was summarized by Ollivier et al. (2011) that nitrification activity was predominantly driven by AOA, despite of its lower abundance compared to AOB. Apparently, archaea were more active compared to bacteria under extreme conditions, such as ammonium-poor environments (Di et al., 2009), low pH (Nicol et al., 2008; Lehtovirta-Morley et al., 2011) and temperature stress (Schleper et al., 2005; Valentine, 2007). A surprisingly large abundance of AOA was also demonstrated by Su et al. (2010) in soils from moderate climatic zones (arable land (Cambisol), Southern Germany), where AOB were exhausted by freeze-thaw cycles, whilst archaeal communities thrived. Thus, AOA may be important players for ammonia oxidation processes, and may contribute substantially to NO production during freeze-thaw events.

So far there is only little evidence that AOA are involved in soil NO production or that AOA do express the HAO enzyme. For example, Vajrjala et al. (2013) demonstrated by a combined physiological and stable isotope

tracer analyses that NH_2OH is an intermediate product of NH_3 oxidation to NO_2^- in the archaeon *Nitrosopumilus maritimus*. The authors proposed that an archaeal AMO homolog is responsible for NH_2OH formation, while the oxidation of NH_2OH to NO_2^- is likely performed by an archaea unique enzyme system. This enzyme system may be connected to soluble periplasmic multicopper oxidases (MCO) and membrane-anchored copper-binding proteins described by Walker et al. (2010). The latter authors also found *nirK* genes in archaea, though its role remained unclarified (Walker et al., 2010 and references therein; Jung et al., 2014; Park et al., 2014). Thus, in analogy to AOB, NO production by AOA may be linked to NH_2OH oxidation to NO_2^- or AOA produced NH_2OH may be used as substrate by other microorganisms to produce NO. Another NO production pathway for AOA may be the formation of nitroxyl hydride (HNO) during NH_3 oxidation (Schleper and Nicol, 2010; Walker et al., 2010), with HNO being converted to NO by copper-complexes/copper-containing proteins (Hughes, 1999). A significant importance of NO in the AOA energy metabolism, earlier postulated by Walker et al. (2010) and Schleper and Nicol (2010), has been recently confirmed experimentally (Yan et al., 2012; Shen et al., 2013). Apparently, AOA can form N_2O by direct oxidation of NH_3 rather than from NH_2OH (Vajrала et al., 2013), while Stieglmeier et al. (2014b) described N_2O formation as a hybrid of NO_2^- reduction and NH_3 oxidation.

3.2.2. Denitrification

Denitrification is the stepwise reduction of nitrate to nitrite, nitric oxide, nitrous oxide and dinitrogen gas (Equation (6)), catalyzed by the enzymes nitrate reductase (membrane-bound (NAR) or periplasmic (NAP)), nitrite reductase (NIR), nitric oxide reductase (NOR), and nitrous oxide reductase (N₂OR) (Payne, 1973, 1981; Knowles, 1982; Stouthamer, 1988; Revsbech and Sørensen, 1990; Zumft, 1992, 1997).



The absence of NO emission during denitrification may be explained by the “diffusion limitation” hypothesis (Firestone and Davidson, 1989; Skiba et al., 1997). This hypothesis suggests that at low O₂ concentrations, i.e. conditions which favour denitrification such as waterlogging, the NO produced is unlikely to escape from the soil to the atmosphere due to limited gas diffusion. Thus, the NO is trapped and is available as denitrification substrate for further reduction to N₂O and/or N₂. This has recently been experimentally confirmed in river sediments using ¹⁵NO stable isotopes (Schreiber et al., 2014). However, under such conditions plant NO production and emission may be an important source of atmospheric NO (see below sections 3.2.9 and 3.2.10).

Controlling factors for denitrification are soil moisture content, soil temperature, N-NO₃⁻ and easily decomposable C availability, soil properties affecting soil aeration and microbial activity (e.g. texture and organic matter content), and agricultural management (Stouthamer, 1988; Revsbech and Sørensen, 1990; Zumft, 1997; Bouwman et al., 2002; Skiba, 2008; Rees et

al., 2013). High rates of denitrification tend to be observed in N fertilized soils and highly irrigated loam soils when mineral N as well as C is not limiting (Barton et al., 1999; Groffman et al., 2009). Based on numerous published studies with agricultural (grassland and cropland) and forest soils, Barton et al. (1999) concluded that denitrification rates tended to be higher in agricultural soils (mean rate $13 \text{ kg N ha}^{-1} \text{ a}^{-1}$) than in natural forest soils (e.g. mean rate $1.9 \text{ kg N ha}^{-1} \text{ a}^{-1}$). However, these estimates are mainly based on the acetylene blockage technique with results being highly questionable if used under aerobic conditions (Bollmann and Conrad, 1997; Butterbach-Bahl et al., 2013).

Soil moisture content and soil temperature are key drivers of denitrification and their alterations can commonly explain up to 95% of the variation of the N_2O emission (Butterbach-Bahl et al., 2013). In addition, freeze-thaw events can trigger pulses of soil N_2O emissions and can contribute significantly to the annual N_2O emission rate in regions experiencing several weeks of subzero winter temperatures (Mørkved et al., 2006; Sharma et al., 2006; Wagner-Riddle et al., 2008; Kim et al., 2012; Luo et al., 2012). Freeze-thaw induced N_2O emissions are due to a complex mix of soil physical and microbial processes that require anaerobic conditions and a surplus of easily degradable substrates (De Bruijn et al., 2009). Little is known if freeze-thaw periods also significantly stimulate soil NO emissions. The multi-year data set on soil NO emissions from an acid forest soil in the South of Germany reported by Gasche and Papen (1999) and Luo et al. (2012) does not indicate that freeze-thaw periods trigger high NO emissions, though at the same site high pulse emissions of N_2O were

observed in approximately 1 out of 3 years (Luo et al., 2012). However, it has been recently confirmed that NO emissions during the cold seasons (16 of October – 15 of April periods) contribute ca. 29% to the annual NO budget based on 16 years of measurement data in a forest stand (Höglwald) in South Germany (Medinets et al., unpublished data). The microbial processes involved have not been identified; however, we assume that denitrification plays an important role, since high denitrifier activity has been demonstrated during freeze-thaw events (Mørkved et al., 2006; Sharma et al., 2006; Wagner-Riddle et al., 2008; Kim et al., 2012; Luo et al., 2012). There is a need for more continuous NO flux measurements during cold periods and winter/spring transition periods, in order to improve our estimates of annual flux rates.

Soil pH is another important factor determining denitrification rates. Bakken et al. (2012) showed that the ratio of $\text{N}_2\text{O}/(\text{N}_2+\text{N}_2\text{O})$ is negatively correlated with soil pH over the pH range 5-8, which is typical for agricultural soils. The authors concluded that low pH interferes with the synthesis of the N_2O reductase enzyme, most likely by affecting the enzyme assembly in the periplasm. Thus, liming can be an efficient way to reduce N_2O (Bakken et al., 2012) and also NO emissions (Gasche and Papen (1999). Comparing limed and non-limed areas in the Höglwald Forest, Gasche and Papen (1999) concluded that an increase in NO consumption rather than a decrease in NO production was driving the decrease in soil surface NO emissions.

3.2.2.1. Heterotrophic (classical) denitrification

567

568 Most denitrifiers are facultative aerobes (including bacteria (e.g.
 569 *Alcaligenes faecalis*, *Pseudomonas stutzeri*, *Paracoccus denitrificans*),
 570 fungi (e.g. *Fusarium oxysporum*, *Cylindrocarpon tonkinense*) and archaea
 571 (e.g. *Methanosaeta concilii*, *Pyrobaculum aerophilum*)), but in case of O₂
 572 depletion they can switch to anaerobic respiration using NO₃⁻ as electron
 573 acceptor (Payne, 1981; Knowles, 1982; Stouthamer, 1988; Revsbech and
 574 Sørensen, 1990; Zumft, 1992, 1997; Kobayashi et al., 1996; Park et al.,
 575 1997; Cabello et al., 2004; Hayatsu et al., 2008; Shoun et al., 2012).
 576 Although large denitrification rates are linked to low O₂ concentrations,
 577 aerobic denitrification has been demonstrated for some bacteria (Lloyd,
 578 1993). For example, Bateman and Baggs (2005) used isotopic tracer to
 579 identify aerobic denitrification in dry soil (20% WFPS).


580 It is well known that NO and N₂O can be produced in soils
 581 simultaneously, and the emission ratio of N-NO/N-N₂O is conventionally
 582 used to assess the dominance of microbial production pathways for NO and
 583 N₂O. At a ratio >1 nitrification is supposed to be the main process, while at
 584 a ratio <1 denitrification is generally assumed to dominate N trace gas
 585 production (Davidson, 1991; FAO and IFA, 2001; Parton et al., 2001;
 586 Garrido et al., 2002; Akiyama and Tsuruta, 2003; Cheng et al., 2004;
 587 Nakajima et al., 2005; del Prado et al., 2006). Contradictory to this
 588 suggestion, Wang et al. (2011) observed during gas-flow-soil-core
 589 incubation experiments of soils enriched with NO₃⁻ and excess glucose
 590 (ratio of C:N = 6) and maintained under anaerobic condition that
 591 denitrification was the main process of NO production even though the N-

NO/N-N₂O ratio was above 1. Similarly in a previous laboratory study (Anderson and Levine, 1986), the emission ratio of N-NO/N-N₂O was 3 for a pure denitrifier culture of *A. faecalis* under micro-aerobic conditions. These results suggest that at high soil NO₃⁻ concentrations and micro-aerobic or anaerobic conditions, NO production is exclusively associated with denitrification (Ludwig et al., 2001; Russow et al., 2009; Wang et al., 2011). Bergaust et al. (2012) observed that NOR-deficient strains of denitrifying bacteria could grow by denitrification under conditions that allow NO to escape and/or be consumed by other organisms, thus avoiding NO toxicity. These findings indicate that the role of denitrification as source of atmospheric NO should be revisited.

3.2.2.2. Nitrifier denitrification

Nitrifier denitrification (Equation (7)) is a process in which NO₂⁻ is reduced to gaseous NO, N₂O and N₂ by AOB with NH₄⁺ as an electron donor under O₂ limitation (Poth and Focht, 1985; Poth, 1986; Wrage et al., 2001). Basically the same enzymes (NIR, NOR, N₂OR) involved in the stepwise denitrification reduction cascade from nitrate to nitrous oxide or di-nitrogen are also activated during nitrifier denitrification. Ammonia oxidizing bacteria are responsible for this process and were found to denitrify under a wide range of environmental conditions from arctic to tropical climatic zones (Kool et al., 2009a, 2009b, 2010; Szukics et al., 2010; Baggs, 2011; Banerjee et al., 2011; Toyoda et al., 2011; Wertz et al., 2012; Vanitchung et al., 2013). This process is important to avoid

accumulation of toxic levels of NO_2^- (Stein and Arp, 1998; Beaumont et al., 2004, 2005; Baggs, 2011).



NO

Nitrifier denitrification is a significant source of NO emitted from soils. NO is an intermediate of NO_2^- reduction by nitrifiers (Remde and Conrad, 1990; Wrage et al., 2001) with N_2O production being often the final step of nitrifier denitrification (Poth and Focht, 1985). However, further reduction to N_2 may also be possible (Poth, 1986), although a NOR homolog has so far not been identified in AOB. WFPS (Garrido et al., 2002) and pH (Nagele and Conrad, 1990; Yamulki et al., 1997; Ste-Marie and Pare, 1999; Cheng et al., 2004) can affect NO and N_2O emission rates under aerobic conditions. Soil core incubation experiments using a range of agricultural soil types collected in France showed that under aerobic condition around 0.6-2.5% of the NH_4^+ applied was emitted as N-NO, while 0.06-1% was emitted as N- N_2O (Garrido et al., 2002). Recently, Zhu et al. (2013) during laboratory experiments on loam, sandy loam and clay loam soils (sampled in California, USA) found under controlled condition (temperature, O_2 concentration, N-application) that at $\text{O}_2 > 0\%$ (0.5-21%) most of the released NO (72-97%) was produced by the NH_3 oxidation pathways (nitrifier denitrification, nitrification-coupled denitrification and nitrification). Moreover NO production increased while the O_2 concentration declined. This demonstrates that nitrifier denitrification and/or partially coupled

nitrification-denitrification processes (Wrage et al., 2001; Zhu et al., 2013) could have been responsible for the observed NO emission. Nitrifier denitrification may contribute significantly to losses of NH_4^+ as NO and N_2O emission from soils (Zumft, 1997; Zhu et al., 2013), however, a contribution of nitrification (at least up to the formation of NO_2^- or directly via NH_2OH aerobically) cannot be excluded (Zhu et al., 2013).

3.2.3. The contribution of nitrification and denitrification to NO production

Both, the nitrifier and denitrifier microbial communities can play significant roles in NO production in the soil of terrestrial ecosystems under a wide range of oxygen concentrations. This was recently confirmed by Russow et al. (2009), who demonstrated significant increases of NO emission rates with declining O_2 partial pressure during soil laboratory incubation experiments. They carried out three separate experiments using the tracers $^{15}\text{N-NH}_4^+$, $^{15}\text{N-NO}_3^-$ or $^{15}\text{N-NO}_2^-$ in a soil reactor with a continuously well-mixed headspace (Russow et al., 2009) under a range of O_2 concentrations (Table 1).

INSERT Table 1 HERE


The results clearly showed that NO_2^- was the main precursor of NO under any oxygen condition, but that the source of NO_2^- was different. Under aerobic conditions ($\text{O}_2 = 20$ vol. %) NO_2^- formed by nitrification from

ammonium contributed 70% of the emitted NO and 10% of the emitted NO came from NO_2^- which was formed from the reduction of nitrate by denitrification (Table 1). However, it is likely that in the described experiment nitrifier denitrification contributed to the aerobic NO production. In contrast, under anaerobic condition 87% of the emitted NO was generated by denitrification of nitrate. However, the emission rate under anaerobic (denitrification prevailing) condition was ca. 4-fold higher than under aerobic (nitrification prevailing) conditions (Table 1). Russow et al. (2009) also reported that the fate of NO_2^- freshly added to the soil was different from endogenous NO_2^- , i.e. NO_2^- generated by nitrification and denitrification in the soil. Apparently, exogenous or freshly added NO_2^- undergoes rapid microbial as well as chemical decomposition (Van Cleemput and Baert, 1976; Van Cleemput and Samater, 1996; Venterea and Rolston, 2000; Islam et al., 2008).

Russow et al. (2009) demonstrated very clearly that NO was the exclusive precursor of N_2O under anaerobic condition, i.e. NO produced by denitrification was also consumed by denitrification. This confirms the “diffusion limitation” hypothesis (Firestone and Davidson, 1989; Skiba et al., 1997), which assumes that diffusion limitation in soils with a WFPS >>60% increases the likeliness that NO produced under anaerobic condition *in situ* is further reduced to N_2O (and N_2) by the denitrifying microbial community.

3.2.4. Codenitrification

692 Codenitrification is a metabolic process, co-occurring during
 693 conventional denitrification, where NO_2^- or NO is reduced by other
 694 nucleophilic N compounds (e.g., amines (R-NH_2), NH_2OH , NH_4^+ , azide (N_3^-
 695), hydrazine (N_2H_4) and salicylhydroxamic acid) to form N_2O and/or N_2
 696 (Shoun et al., 1992; Tanimoto et al., 1992; Spott et al., 2011) (Equation (8)).



700 It is assumed that the codenitrification pathway is based on biotically
 701 mediated N-nitrosation via enzyme (E) bound NO complexes (e.g., E-NO,
 702 E- NO^- and E- NO^+) (Stamler et al., 1992; Kumon et al., 2002; Spott et al.,
 703 2011 and references therein). Both NO_2^- and NO are considered as nitroso
 704 donors for nitrosating agents (e.g., E-NO, E- NO^- and E- NO^+) and the
 705 reaction is catalyzed by *cdI* NIR (Averill, 1996; Kim and Hollocher, 1984;
 706 Weeg-Aerssens et al., 1988). Thus, NO_2^- as well as NO can be directly
 707 involved in the biological formation of hybrid N-N gas, and under certain
 708 conditions the reaction between NO_2^- and NO can be reversible (Su et al.,
 709 2004; Spott et al., 2011 and references therein). This is in-line with the
 710 statement by Averill (1996) that NIR and NOR enzymes of many
 711 denitrifiers are likely to be strongly coupled and may function as multi-
 712 enzyme complexes and, therefore, are likely to play a key role as biotic
 713 catalysts of the codenitrification process.

714 Evidence for codenitrification has been found in archaea (order
 715 *Sulfolobales*) (Immoos et al., 2004), bacteria (orders *Actinomycetales*,
 716 *Burkholderiales*, *Enterobacteriales*, *Pseudomonadales*, *Rhizobiales* and

Rhodobacterales) (e.g., Garber and Hollocher 1982a,b; Goretski and Hollocher, 1991; Ye et al., 1991; Okada et al., 2005) and fungi (order *Hypocreales*) (e.g., Shoun et al., 1992; Tanimoto et al., 1992; Usuda et al., 1995, Sameshima-Saito et al., 2004; Su et al., 2004). Codenitrification seems to be a widely distributed process across terrestrial as well as aquatic ecosystems. But only a few studies provide direct evidence of codenitrification in natural environments, for example in grassland (Laughlin and Stevens, 2002) and agricultural soils (Spott and Stange, 2011; Long et al., 2013).

Controlling factors for codenitrification appear to be closely related to those for denitrification. Accordingly, oxygen availability, pH and availability of respirable organic carbon substrates are the main controllers of codenitrification (Spott et al., 2011), and as for denitrification, may occur under micro-aerobic conditions (Kumon et al., 2002; Okada et al., 2005).

Assuming that most denitrifiers are heterotrophic microorganisms, Spott et al. (2011) have suggested that codenitrification as well as denitrification are related to the availability of respirable organic carbon substances. Short-term experiments showed that decreasing availability of organic carbon compounds (e.g., succinate) diminish denitrification rates, but enhance the codenitrification/denitrification ratio of N_2 produced (Weeg-Aerssens et al., 1998).

In studies where NH_2OH (as naturally occurring nucleophilic compound) was added to denitrifier cultures (Garber and Hollocher, 1982b; Kim and Hollocher, 1984; Weeg-Aerssens et al., 1987, 1988; Goretski and Hollocher,

1991) or soil (Spott and Strange, 2011) 98% of the N₂O produced was formed by codenitrification.


The importance of codenitrification as a key process of N₂O and N₂ production has also been shown under natural conditions. Laughlin and Stevens (2002) showed that up to 92% of released N₂ in grassland soils was produced by codenitrification.

In addition, increasing NO production by denitrification has been observed in the presence of codenitrification (e.g., Garber and Hollocher, 1982a,b, Goretski and Hollocher, 1991). Goretsky and Hollocher (1991) have pointed out that azide (as a nucleophilic compound) partially inhibited NOR activity, thus resulting in NO accumulation. It is quite possible that others nucleophilic compounds could act analogically on NOR enzymes. In addition, it may be also attributed to a sort of abortive reaction of denitrification (Spott and Strange, 2011) as well as may indicate the underconsumption of NO₂⁻ and NO by a microbial N-nitrosation (i.e. codenitrification).

3.2.5. Dissimilatory nitrate reduction to ammonium

Nitrate ammonification or dissimilatory nitrate reduction to ammonium (DNRA) is a fermentative process, using NO₃⁻ as electron acceptor during its conversion via NO₂⁻ to NH₄⁺ (Cole and Brown, 1980; Cole, 1990):





767

768 Two types of DNRA have been determined, acting in different
 769 subcellular compartments. 1) Periplasmic, energy-conserving (respiratory)
 770 nitrate reduction to ammonium, which catalyzes the electron transport from
 771 formate or H_2 to NO_2^- (using NAP-NRF (nitrite reduction to formate
 772 dehydrogenase or hydrogenase enzymes) was described in *Escherichia coli*,
 773 *Desulfovibrio*, and *Wolinella* spp. (Simon, 2002; Simon et al., 2003; Cabello
 774 et al., 2012). 2) Cytoplasmic dissimilatory $\text{NO}_3^-/\text{NO}_2^-$ reduction to NH_4^+ ,
 775 which functions as both electron sink and detoxification of NO_2^- formed in
 776 NO_3^- respiration in the cytoplasm (using NAR-NIR enzymes). Both
 777 processes can result in NO as well as N_2O production. These processes have
 778 been reported for *E. coli* and *Klebsiella* spp. (Moreno-Vivián et al., 1999;
 779 Cabello et al., 2012), but may also occur in other microorganisms.

780 DNRA can be performed by different groups of bacteria, including
 781 obligate anaerobes (e.g. *Clostridium* spp.), facultative anaerobes (e.g.
 782 *Enterobacter* spp.) and aerobes (e.g. *Bacillus* spp.) (Tiedje, 1988). Very
 783 reduced and carbon rich environments (C/N ratio >4) favour DNRA (Buresh
 784 and Patrick, 1978; Tiedje et al., 1982; Tiedje, 1988; Fazzolari et al., 1998).
 785 Positive correlations of DNRA rates with soil pH, C/ NO_3^- ratio, bulk soil
 786 density, sand content and NO_2^- concentration were reported by Schmidt et
 787 al. (2011) for temperate arable soils.

788 The DNRA pathway was reported to be responsible for up to >99% of
 789 the NO_3^- consumption in forest soils (Bengtsson and Bergwall, 2000; Silver
 790 et al., 2001, 2005; Pett-Ridge et al., 2006; Huygens et al., 2007; Rütting et

al., 2008; Templer et al., 2008), and for up to 21% of NO_3^- consumption in rice paddies (Chen et al., 1995a, b; Yin et al., 2002). DNRA was attributed to NO_3^- consumption in calcareous agricultural soils following glucose addition (Wan et al., 2009), and in temperate arable soils, depending on the presence of low weight C sources (Schmidt et al., 2011). Based on correlation and regression analyses, Rütting et al. (2011) concluded that highest gross DNRA rates can be expected in soils with high organic matter content in humid temperate regions in soil with lower soil moisture.

Since NO_2^- was suggested as an intermediate during the reduction of NO_3^- to NH_4^+ (Cole, 1990; Baggs, 2011) in both periplasm and cytoplasm, evidence is increasing that N_2O is produced during DNRA (Stevens et al., 1998; Baggs, 2011; Rütting et al., 2011). Therefore, it may be assumed that NO (e.g. as an intermediate for N_2O) is produced during reduction of NO_2^- in the cytoplasmic or/and the periplasmatic space. Thus, DNRA may be considered as an additional source not only for N_2O , but also for NO in soils. However, the role of DNRA, as a source for soil NO, remains to be investigated.


3.2.6. Anaerobic ammonium oxidation

Anaerobic ammonium oxidation (anammox) is a biological process where NH_4^+ serves as electron donor and NO_2^- as electron acceptor to form N_2 (van de Graaf et al., 1990, 1995; Strous et al., 1996; Kuypers et al., 2003; Kuenen, 2008). The anammox pathway is restricted to some slow-growing, strictly anoxic, and lithotropic bacteria belonging to the order

816 *Planctomycetales* of the phylum *Planctomycetes* (Kartal et al., 2011, 2013).
 817 Up to date 10 species of five genera (*Candidatus Brocadia* (Strous et al.,
 818 1999a; Kartal et al., 2008; Oshiki et al., 2011), *Candidatus Kuenenia*
 819 (Strous et al., 2006), *Candidatus Scalindua* (Schmid et al., 2003; Woebken
 820 et al., 2008; van de Vossenberg et al., 2013), *Candidatus Anammoxoglobus*
 821 (Kartal et al., 2007b) and *Candidatus Jettenia* (Quan et al., 2008; Hu et al.,
 822 2011)) have been described. Representatives of four from five genera,
 823 except *Candidatus Anammoxoglobus*, have been identified in terrestrial
 824 ecosystems (Humbert et al., 2010; Long et al., 2013; Wang and Gu, 2013).

825 Anammox bacteria were first discovered in probes from wastewater
 826 treatment bioreactors (van de Graaf et al., 1995, 1996; Mulder et al., 1995;
 827 Jetten et al., 1997; Strous et al., 1997), but since then have been found in
 828 various ecosystems such as marine oxygen-limited zones and sediments
 829 (Rysgaard et al. 2004; Dalsgaard et al. 2005; Kuypers et al. 2005; Lam et al.
 830 2007; van de Vossenberg et al. 2008; Hong et al. 2011), marine surface
 831 sediments (Hietanen and Kuparinen, 2008; Rich et al., 2008), sea ice
 832 (Rysgaard et al., 2008), estuaries (Trimmer et al., 2003; Dale et al. 2009),
 833 freshwater ecosystem (Schubert et al. 2006; Rich et al. 2008), oil reservoirs
 834 (Li et al. 2010a), marshlands (Koop-Jakobsen and Giblin 2009; Li et al.,
 835 2011a), wetlands (Jetten et al. 2003; Zhu et al. 2010; Humbert et al., 2012),
 836 permafrost soils (Philipot et al., 2007; Humbert et al., 2010), peat soils (Hu
 837 et al., 2011), rice paddy soils (Zhu et a., 2011; Wang and Gu, 2013),
 838 grassland soils (Humbert et al., 2010), agricultural soils (Long et al., 2013),
 839 and the rhizosphere (Humbert et al., 2010)

The anammox process occurs in a special intracytoplasmic compartment (organelle), the anammoxosome, which is surrounded by ladderane lipids (Lindsay et al., 2001; van Niftrik et al., 2004; Kuypers et al., 2003; Kartal et al., 2011). The reaction pathway is likely structured in three distinctive steps (Strous et al., 2006) (Equation (10)): During the first stage NO_2^- is reduced to NO by cytochrome *cdI* NIR. Subsequently, the reaction between NH_4^+ and NO to hydrazine (N_2H_4) is catalyzed by a hydrazine synthase (HZS). Finally N_2H_4 is enzymatically dehydrogenized by a hydrazine dehydrogenase (DHD) resulting in N_2 production. Meanwhile a part of NO_2^- is oxidized for carbon fixation with NO_3^- formation.



Kartal et al. (2010b; 2011) showed that N_2H_4 and NO are obligatory intermediates of anammox, that anammox bacteria are tolerant to extremely high concentrations of NO (3500-5000 ppm), and that the reduction of NO is exclusively linked to the catabolic activity of the anammox pathway (Kartal et al., 2010).

Data describing controlling factors of the anammox process are scarce. Strictly anoxic condition and substrate availability (Kartal et al., 2013) under stable environmental conditions are assumed to favour anammox bacteria in natural ecosystems (Dalsgaard et al., 2003; Humbert et al., 2010). Anammox bacteria can grow at very low substrate concentrations, but require NO_2^- as well as NH_4^+ . Interestingly, NO_2^- serves as both the electron acceptor for the ammonium oxidation and the ultimate electron donor in the

reaction with bicarbonate (HCO_3^-) for biomass formation and NO_3^- production as a by-product (Strous et al., 1998; Kartal et al., 2013). Substrate consumption for anammox, including that for carbon fixation, are 1.27 moles of NO_2^- (including conversion of 1 mole via NO for NH_4^+ oxidation and 0.27 moles for carbon fixation) and 1 mole of NH_4^+ per 0.066 mole of fixed carbon (Strous et al., 1998; Kartal et al., 2013). Therefore, for the fixation of one mole of carbon into biomass 15 catabolic cycles of ammonium oxidation, resulting in significant N_2 production, are needed, which explains the slow growth rate of the bacteria (Kartal et al., 2013). N_2O production has not been observed so far, despite targeted experiments using a range of NO concentrations (Kartal et al., 2010). Anammox bacteria can also grow heterotrophically thereby converting organic compounds, e.g. formate, acetate, propionate, methanol, mono- and dimethylamine into biomass C (Strous et al., 2006; Kartal et al., 2007a,b, 2008, 2013) or even to CO_2 (Kartal et al., 2007a,b, 2008, 2013).

Dalsgaard and Thamdrup (2002) reported that the temperature optimum for NH_4^+ oxidation by anammox bacteria isolated from marine sediments was ca. 15 °C, though it may vary from 6 °C (Dalsgaard and Thamdrup, 2002) to temperatures >50°C (Jaeschke et al., 2009; Byrne et al., 2009; Li et al., 2010).


High NH_3 concentrations as found at high pH values may inhibit anammox (Aktan et al., 2012; Yang et al., 2014), while accumulation of heavy metals (e.g., As, Cd and Pb) in sediments affects the diversity of anammox bacteria (Li et al., 2011a; Yang et al., 2014). Generally, the

diversity of anammox bacteria is higher in terrestrial systems as compared to marine systems (Humbert et al., 2010). Also increased soil or sediment aeration is negatively affecting anammox activity (Long et al., 2013) while reported effects of increasing N availability remains controversial: Koop-Jakobsen and Giblin (2009) did not find statistically significant differences between fertilized and unfertilized marsh lands while Hu et al. (2011) found that in NO_2^- and NH_4^+ amended peat soils the abundance of *Ca. Jettenia asiatica* increased.

However, it remains unknown if anamox bacteria are significant sources of NO or possibly even sinks in terrestrial ecosystems.

3.2.7. Nitrite-dependent anaerobic oxidation of methane

Nitrite-dependent anaerobic oxidation of methane (N-AOM) is an “intra-aerobic” pathway of methane (CH_4) oxidation to CO_2 by O_2 . However, in this reaction the O_2 is produced by NO_2^- reduction via NO dismutation to O_2 and N_2 (Equation (11)) (Ettwig et al., 2010).



The process itself requires a set of enzymes: methane monooxygenase (MMO), methanol dehydrogenase (MDH), formate dehydrogenase (FDH) and nitrite or nitric oxide reductase, which has been found in slow-growing Gram-negative bacteria *Candidatus 'Methyloirabilis oxyfera'* belonging to the phylum NC10 (Ettwig et al., 2010). *M. oxyfera* has been enriched from

freshwater sediments (Raghoebarsing et al., 2006; Ettwig et al., 2008, 2009), and its complete genome has recently been published (Ettwig et al., 2010). Ettwig et al. (2010) have speculated that NOR may be involved in NO detoxification. Exogenous NO as well as NO_2^- has been demonstrated to be rapidly reduced to N_2 and O_2 , thus stimulating CH_4 oxidation.

Very recently Harron et al. (2013) described an anaerobic, methane-oxidizing and nitrate-reducing archaeon *Candidatus 'Methanoperedens nitroreducens'*, which was enriched from a mixture of freshwater sediments and anaerobic wastewater sludge. This archaeon has been demonstrated to oxidize CH_4 to CO_2 while reducing NO_3^- to NO_2^- . Moreover *M. nitroreducens* was able to oxidize CH_4 in the presence of NH_4^+ through a syntrophic relationship with the anaerobic ammonium oxidizing bacteria *Kuenenia* spp. (Harron et al., 2013). Anammox bacteria have been shown to utilize NO_2^- , reduced by *M. nitroreducens* for NH_4^+ oxidation, thereby producing NO_3^- as byproduct (Harron et al., 2013).

Occurrence of N-AOM has been widely reported for freshwater sediments (Deutzmann and Schink, 2011; Kampman et al., 2012; Kojima et al., 2012; Shen et al., 2014a), estuarine sediments (Shen et al., 2014b), wastewater sludge (Luesken et al., 2011a), peat lands (Zhu et al., 2012), wetlands (Hu et al., 2014) and rice paddy soils (Wang et al., 2012; Shen et al., 2013; Hu et al., 2014).

Generally, oxic/anoxic interfaces with high CH_4 and $\text{NO}_3^-/\text{NO}_2^-$ concentrations are favourable for the N-AOM process (Oremland, 2010; Shen et al., 2012). For example, highest N-AOM activity has been found at

a depths of 50-60 cm for wetlands (Hu et al., 2014), 80-85 cm for peatlands (Zhu et al., 2012) and 90-100 cm for paddy soils (Hu et al., 2014).

Temperature optimum for ‘intra-aerobic’ CH₄ oxidation has been detected to be 25-30 °C for bacteria (Ettwig et al., 2010) and a bit widely 22-35 °C for archaea (Harron et al., 2013). N-AOM microorganisms are mesophilic to pH with optimum of 7-8 (Raghoebarsingetal, 2006; Ettwig et al., 2010), although are still active at more acidic (5.9) pH (Zhu et al., 2012).

However, to date there is no evidence that N-AOM contributes to NO production in soils, though NO is an obligatory intermediate.

3.2.8. Unspecific enzymo-oxidative mechanisms related to soil NO contents

The similarity of biochemical processes in different groups of living organisms is not surprising. Therefore, we briefly outline the seven known pathways of NO production in plants (Table 2), as described by Gupta et al. (2011) and compare it with soil microbial processes of NO production were appropriate. In this context, different pathways of NO biosynthesis could be classified either as reductive or as oxidative (Table 2).

INSERT Table 2 HERE

The ‘hydroxylamine-mediated NO production’ in plants may be due to the same or a very similar biochemical mechanism as soil NO production during nitrification. In plants, hydroxylamine reacts directly with superoxide

(O₂⁻) to form NO under aerobic conditions (Vetrovsky et al., 1996), whereas this conversion in nitrifiers (*Nitrosomonas* spp.) is catalyzed by the enzyme hydroxylamine oxidase (Lees, 1952; Hooper and Terry, 1979; Hooper et al., 1997). *In vitro* experiments adding hydroxylamine (NH₂OH) to plant cells confirmed that NH₂OH is indeed converted to NO and NO₂⁻ (Rümer et al., 2009a, 2009b; Gupta et al., 2011).

The enzyme superoxide dismutase (SOD) (Beyer et al., 1991) was considered to be essential for the conversion of NH₂OH to NO and NO₂⁻ in plant cells and cell-free laboratory experiments (Rümer et al., 2009a, b). In cell-free systems NO emissions increased up to 10-fold in air and 25-fold in a N₂ environment in the presence of SOD and hydroxylamine compared to controls where only hydroxylamine was added. As both, substrate (NH₂OH) and by-products (NO and NO₂⁻) are able to penetrate membranes (Rümer et al., 2009a) and extracellular SOD (EC-SOD) can originate from excretion by bacteria (Tullius et al., 2001; Takahashi et al., 2003) as well as plant cells (Alscher et al., 2002), oxidation of both endogenous and exogenous hydroxylamine may take place inside or outside plant cells (Rümer et al., 2009a). Murphy and Sies (1991) reported that SOD can facilitate the reversible conversion of nitroxyl anion (NO⁻) to NO *in vitro*. The actual mechanism of the SOD-catalyzed reaction of NO and NO₂⁻ production from hydroxylamine is still unclear and its presence in the soil so far has not been demonstrated. It is noteworthy, that soil NH₂OH concentrations (e.g., 0.3-34.8 µg N kg⁻¹ dry forest soil) can be comparable with those of NO₂⁻ (Liu et al., 2014).

We assume that similar enzymatic environments as those described above, can be found in soils with high microbial activity and high nutrient concentrations, especially in the rhizosphere, when nutrients and enzymes are released into the soil, for example after rewetting/thawing of dry/frozen soils. Thus, theoretically, an unspecific enzymo-oxidative mechanisms could trigger NO and NO₂⁻ production in soils.


SOD is widely produced by most organisms (Beyer et al., 1991; Scandalios, 1997; Tullius et al., 2001; Alscher et al., 2002; Takahashi et al., 2003). In the soil, SOD is a rather thermo- and chemo-stable protein (Hunter et al., 2002; Khanna-Chopra and Sabarinath, 2004) that may originate from the active microbial community (Tullius et al., 2001; Takahashi et al., 2003), or recently decaying organisms. Considering these processes and mechanisms, we hypothesize that not only nitrifying (AOB and AOA) microbes are responsible for soil NO production, but that also other microbes via the release of extracellular SOD (directly) or SOD (after cell damage) contribute to soil NO production. More research is required to investigate activating factors for SOD in bacteria, since up to now only data for plant (Bowler et al., 1994; Scandalios, 1997; Babithaa et al., 2002; Baranenko, 2006) and animal cells (Yamakura and Kawasaki, 2010; Miller, 2012) are available.

3.2.9. Nitric oxide synthase

Nitric oxide synthase (NOS) is a common ubiquitous enzyme, which is responsible for NO synthesis in cells of bacteria as well as higher

organisms, including mammals. NOS is present in protists, such as myxomycetes (Messner et al., 2009) and eukaryotic single cells (Fritz-Laylin et al., 2009). Active NOS enzymes are ubiquitously present in invertebrates, such as echinoderms, coelenterates, nematodes, annelids, insects, crustaceans and molluscs (Jacklet, 1997 and reference therein). In mammals, many cell types such as endothelial cells, neurons, myocytes, smooth muscle cells, and activated mune cells (e.g. leucocytes and macrophages) produce NO by both enzymatic and non-enzymatic pathways (Zweier et al., 1995; Velayutham and Zweier, 2013 and reference therein). Enzymatic NO synthesis by NOS appears to be much more important than non-enzymatic production (Zhou and Zhu, 2009; Chen et al., 2010; Forstermann and Sessa, 2012).

NOS-derived NO synthesis proceeds in a two step oxidation of the amino acid precursor L-arginine (L-Arg) via N-hydroxy-L-arginine to L-citrulline (L-Cit) (Griffith and Stuehr, 1995) in the presence of 5,6,7,8-tetrahydrobiopterin (BH₄), reduced nicotinamide-adenine-dinucleotide phosphate (NAD(P)H), molecular oxygen (O₂) and Ca²⁺/calmodulin (CaM) (Zhou and Zhu, 2009; Chen et al., 2010; Forstermann and Sessa, 2012):



1033

It is known that three phyla of Gram-positive bacteria (*Firmicutes*, *Actinobacteria*, and *Deinococcus-thermus*), at least one phylum of archaea (*Euryarchaeota*) and one representative of the *Proteobacteria* phylum of Gram-negative bacteria (*Sorangium cellulosum*) possess NOS-like enzymes

that are highly homologous to the oxygenase domain of eukaryotic NOS (Stuehr, 1999; Gusarov et al., 2008; Sudhamsu and Crane, 2009 and reference therein; Crane et al., 2010 and reference therein). These specific proteins were found in pathogenic as well as in non-pathogenic soil bacteria (Gusarov et al., 2008, 2009). Indeed, a high level of functional and structural similarity between bacterial NOS (bNOS) and eukaryotic NOS was reported (Pant et al., 2002; Pant and Crane, 2006; Salard et al., 2006; Gusarov et al., 2008; Sudhamsu and Crane, 2009). Bacterial and archaeal NOS were thought to be unable to produce NO *in vivo* because of a lacking reductase domain (Adak et al., 2002) and only more recent studies have provided evidence of bNOS mediated bacterial NO production thereby using various nonspecific cellular reductases as their redox partners (Johnson et al., 2008; Gusarov et al., 2008; Shatalin et al., 2008). In the mentioned works it is proposed that NO, escaping from the cellular lumen, is readily oxidized in the culture medium under aerobic conditions forming NO₂⁻ and NO₃⁻. Shatalin et al. (2008) and Schreiber et al. (2011) have demonstrated directly that NO was produced by *B. anthracis* and *B. subtilis*, using an NO sensitive dye. Furthermore, it has been shown in plant-pathogenic *Streptomyces* spp. that bNOS-derived NO production considerably exceeds the requirement of phytotoxin thaxtomin A nitration. Johnson et al. (2008) confirmed that surplus NO was produced by bNOS, and was detected in the gas phase above the culture medium by chemiluminescence. It is also known that other, NOS-independent mechanisms of L-Arg conversion to L-Cit in the urea cycle catalysed by arginine deiminase (Yamasaki and Sakihama, 2000, equation 12), arginase

or ornithine carbamoyl transferase (Jansson and Lindblad, 1998; Viator et al., 2008) are present in bacteria (Sudhamsu and Crane, 2009).

In mammals, three isoforms of NOS originating from separate genes have been described, i.e. endothelial NOS (eNOS), neuronal NOS (nNOS) and inducible NOS (iNOS). The first two types of NOS are constitutively expressed in the cells and are called cNOS; iNOS is typically expressed under infectious and inflammatory conditions at dramatically higher rates compared to cNOS (Wu, 1995; Siervo et al., 2011). Furthermore, NO_3^- and NO_2^- can also be reduced to NO via other enzymatic (NAP/NIR) and non-enzymatic reactions (e.g. via deoxygenated haemoglobin in acidic environment, via neuroglobin, by xanthine oxydo-reductase) in mammalian cells (Burmester and Hankeln, 2004; Gladwin and Kim-Shapiro, 2008; Jansson et al., 2008; Li et al., 2009). It is likely that eukaryotes have acquired the NOS enzyme from bacteria, which possess the most ancient primitive NOS type (Gusarov et al., 2008), by horizontal gene transfer, as supported by recent phylogenetic tree analysis (Sudhamsu and Crane, 2009).

In plant cells, a gene with significant homology to that encoding animal NOS has not been detected (Moreau et al., 2010; Gupta et al., 2011), and NOS-derived NO production has not been confirmed as an enzymatic pathway of Arg-derived NO production in plants (Zemojtel et al., 2006; Gas et al., 2009; Moreau et al., 2010). However, several studies showed evidence for an NOS-like enzymatic reaction in plants that is involved in various processes, based on a correlation between the supply with L-Arg and its analogs with NO production (Mackerness et al., 2001; Lum et al., 2002).

There are at least 6 other pathways of NO production in plant cells, mentioned in Table 2, but not described here, because in our opinion they most probably are not relevant for unspecific enzymo-oxidative pathways in soil.

3.2.10. Biotic consumption of NO in the soil

Soils are a net source of NO (IPCC, 2007), but also a sink for atmospheric NO (IPCC, 2007; Slemr and Seiler, 1984, 1991; Ludwig and Meixner, 1994; Ludwig et al., 2001; Laville et al., 2009) or can be re-deposited as NO or/and NO₂ onto plant surfaces (Wesely and Hicks, 2000; Butterbach-Bahl et al., 2004; Horii et al., 2004; Seok et al., 2013; Shen et al., 2013). Plants can use atmospheric NO and NO₂ as additional nitrogen source (Neubert et al., 1993; Geßler et al. 2000; Butterbach-Bahl et al., 2004; Teklemariam and Sparks, 2006;). Plant uptake of atmospheric NO and NO₂ is a diffusive process through the stomata and flux rates depend on the compensation points of NO and NO₂ and their atmospheric gas mixing ratios. The atmospheric gas mixing ratios can vary significantly between ecosystems (Conrad, 1996; Geßler et al., 2000; Ludwig et al., 2001), and enhanced mixing ratios can stimulate the growth of chemolithoautotrophic nitrite oxidizers colonizing the phyllosphere (Geßler et al., 2002; Papen et al., 2002).

NO production during denitrification and nitrifier denitrification, is much larger than the NO emitted (Firestone and Davidson, 1989; Skiba et al, 1997), because a significant proportion of NO produced by denitrification is

immediately consumed by denitrification for energy production (Zumft and Cardenas, 1979) and simultaneous detoxification (Zumft, 1997). Thus, the net NO emission rate from denitrification processes is typically very small.

Some heterotrophic bacteria can oxidize rather than reduce NO via aerobic co-oxidation reactions (Baumgärtner et al., 1996; Koschorreck et al. 1996; Rudolph et al. 1996; Koschorreck and Conrad 1997; Dunfield and Knowles 1997, 1998, 1999; Conrad, 1999). Increased NO consumption was demonstrated after manure or compost application (Dunfield and Knowles 1998). The magnitude of NO consumption in soils remains uncertain, but concentrations in the soil atmosphere can be significant. E.g., in temperate forest soils NO concentrations varied in a range of 60-180 ppbv at 0 to 10 cm soil depth (Dong, Simon and Rennenberg, unpublished data).

It is well known that NO is an important free diffusive signalling molecule in higher organisms with many direct and indirect functions, such as transcriptional gene regulation, post-translational protein modification, cytoprotection, cytotoxicity, pathogenesis, memory modulation and learning, or vasodilation (vascular smooth muscle relaxation) (for detailed information see section 2 below). For these specific purposes, NO is produced by the NOS enzyme or/and other enzymatic reactions, but also exogenous NO is consumed (Gusarov et al., 2013). The contribution of exogenous NO in intracellular signalling processes has rarely been studied and, therefore, is poorly understood. NO is also consumed for cell detoxification mainly via forming reactive N species (RNS), such as the NO radical (NO^*), nitroxyl (NO^-), S-nitrosothiols (RSNOs), NO-soluble guanylyl cyclase (NO-sGC), and dinitrosyl-iron complexes (DNICs). Not

all NOS-derived NO is stored and converted to RNS and surplus will be emitted (Johnson et al., 2008); unfortunately, quantitative data are not available.

Many bacteria (including those not engaged in the N-cycle) are able to detoxify NO by a range of enzymes, such as NO dioxygenase (flavohemoglobin, *Hmp*), flavodiiron NO reductase (flavorubredoxin, *norVW*) and periplasmic cytochrome C nitrite reductase (*NrfA*), under both oxic and anoxic conditions (Poole et al., 2005; Koul et al., 2014; Mühlig et al., 2014). Under aerobic conditions *Hmp* catalizes the oxidation of NO to NO_3^- (Crawford and Goldberg, 1998; Gardner et al., 1998; Hausladen et al., 2001;); and *Hmp* was shown to protect *Salmonella typhimurium* against the growth inhibitory affect of NO (Mills et al. (2008). Meanwhile under anoxic conditions *Hmp* and *NorVW* facilitated the reduction of NO to N_2O (Kim et al. 1999; Gardner et al. 2002; Mills et al., 2005). The enzyme NrfA can catalize the five-electron-reduction of NO to $\text{NH}_3/\text{NH}_4^+$ undr anaerobic conditions (Pooock et al., 2002; van Wonderen et al., 2008) and other proteins possessing an ability to mediate NO detoxification have been found across bacteria, e.g., truncated globin (HbN) in *Mycobacterium bovis* (Ouellet et al., 2002), vitreoscilla globin (Vgb) in *Vitreoscilla* spp. (Frey et al., 2002), cytochrome *c'* (CycP) in *Rhodobacter capsulatus* (Cross et al., 2001) and single-domain globin (Cgb) in *Campilobacter coli* and *C. jejuni* (Elvers et al., 2004).

For eucaryota the rate of NO consumption by cells is directly dependent on, and proportional to, the oxygen concentration. According to Thomas et al. (2001, 2008) this directly points to an important regulatory

relationship between NO signaling and tissue oxygen concentration. Increased oxygen levels will increase NO consumption, and in reverse NO regulates oxygen consumption via inhibition of mitochondrial respiration. This important interdependent relationship between NO and O₂ provides a direct feedback mechanism to regulate their respective concentrations (Thomas et al., 2008). There are indications that such a mechanism may also regulate NO concentration in soil air, though simultaneous measurements of the dynamics of NO and O₂ concentrations in soil air are still needed for further judgement.

Quantification of the contribution of different NO consumption processes has so far not been achieved. However, Koschorreck and Conrad (1997) have measured a pseudo-first-order uptake rate constant (k) of NO consumption in soil samples from four different ecosystems (primary forest, tree seedling plantation, flooded savanna, soil after tree burning). They reported that under aerobic conditions the consumption rate was low and varied between 12 and 28 cm³ h⁻¹ g⁻¹, while at anaerobic condition the consumption rate was 1-2 orders of magnitude higher (227-3861 cm³ h⁻¹ g⁻¹ dw). Further studies are needed to fill this large knowledge gap.

3.3. Interrelation between main abiotic and biotic processes of NO transformations in soils

Based on recently published literature, we have created a conceptual diagram of all known and theoretical microbial, chemical and enzymatic processes where NO is an obligatory player (Fig. 2). It is likely that NO₂⁻, a

precursor of NO, is the central intermediate connecting all microbial processes and processes associated with chemodenitrification.

INSERT Fig. 2 HERE

As shown in Fig. 2, all processes are interrelated, interacting, and can operate in parallel and/or partially stepwise, utilizing intermediates or products, which were formed during other processes. The unique integrity of interconnections between all components of the system *in situ*, presents the greatest challenge for research, in particular under field conditions.

Unraveling these interactions requires controlled laboratory experiments applying state-of-the-art methods such as multi-isotope tracing (e.g. Kool et al., 2009a, b) together with combined gene expression and functional analyses (e.g. Bru et al., 2010) of microbial mono-cultures and mixtures (e.g. Rümer et al., 2009a, b; Russow et al., 2009).

Nitrification and denitrification are considered to be the main soil microbial processes leading to NO production. *In situ* and *in vivo* laboratory studies have suggested that nitrification rates can be estimated from initial and final substrate concentrations, assuming that oxidation of NH_4^+ via NH_2OH to NO_2^- and NO_3^- is prerogative for aerobic nitrification. However, we cannot ignore that part of the NH_2OH formed from NH_4^+ is decomposed chemically or by non-specific enzyme-oxidative mechanisms. Thus, we cannot answer the following simple questions due to a lack of knowledge:

1212 - What is the relative contribution of oxidative (nitrification) and reductive
 1213 (denitrification, codenitrification, DNRA, anammox, N-AOM) processes to
 1214 NO_2^- production in soils; and can nitrifiers also utilize NO_2^- formed by other
 1215 microbial processes?

1216 - What is the exact fate of NO_2^- in soils, i.e. to what extent is NO_2^- further
 1217 oxidized to NO_3^- or reduced to NO, N_2O , N_2 or even NH_3 ?

1218 - What are the dynamics of N oxidizing and reducing processes in soils,
 1219 since current lack of adequate measuring techniques limits the identification
 1220 of individual processes in bulk soil?

1221 - What are the gross NO production and consumption rates and what is
 1222 the contribution of different processes to this consumption?

1223

1224 *In situ* studies have enabled us to estimate with reasonable confidence
 1225 rates of production and consumption of by- or end-products of
 1226 nitrification/denitrification pathways under certain environmental condition.
 1227 However, we can only speculate about the processes involved. In other
 1228 words, we are studying ‘symptoms’ (substances), but not ‘diseases’
 1229 (processes). The future challenge is to characterize and quantify these
 1230 processes with new experimental approaches to better understand drivers
 1231 and processes leading to NO emissions from soil.

1232

1233 **4. Physiological functions of NO in different groups of organisms**

1234

1235 Generally NO-related signalling functions are attributed to various
 1236 reactive N species (RNS), which are derivatives of NO, e.g. NO radical

(NO \cdot), nitroxyl (NO $^-$), nitrosonium (NO $^+$), peroxyxynitrite (ONOO $^-$), S-nitrosothiols (RSNOs), NO-soluble guanylyl cyclase (NO-sGC), dinitrosyl-iron complexes (DNICs), N $_2$ O $_5$, etc.


4.1. Functions of NO in bacteria

In Gram-positive bacteria (e.g. *B. subtilis*, *B. anthracis*), endogenous NO produced by bNOS as well as exogenous NO mainly possess the function of rapid protection against oxidative stress. Direct protection is achieved through catalase activation and transient inhibition of the rate of enzymatic reduction of free cysteine. This sulphur amino acid is involved in the reduction of Fe $^{3+}$ to Fe $^{2+}$, thus suppressing Fe $^{2+}$ -mediated formation of hydroxyl radicals (a Fenton reaction) (Gusarov and Nudler, 2005; Shatalin et al., 2008). In addition, it was demonstrated (Gusarov et al., 2009) that the enzyme bNOS protects bacteria (e.g. *B. subtilis*, *Staphylococcus aureus*) against a wide spectrum of antibiotics by endogenous NO production, either directly by nitrosation (acridines) or indirectly by NO-mediated suppression of oxidative stress (pyocyanin, cephalosporins, lactams). Corker and Poole (2003) showed that anaerobic NO accumulation in *E.coli* grown in the presence of NO $_3^-$ but absence of *Hmp* inactivated the anaerobic regulator *Fnr* (fumarate and nitrate reductase). *Fnr* controls periplasmic cytochrome *c* nitrite reductase (NrfA), Nir and Nar, and thereby blocks further NO production from NO $_3^-$ via NO $_2^-$. In addition, Mühlig et al. (2014) proposed that in *S. typhimurium* NO can initiate detoxification via inactivation of *Fnr* and/or NO-responsive regulator (NsrR) derepressing *Hmp* expression as

well as via activation of an anaerobic nitric oxide reductase transcription regulator (NorR) derepressing NorV expression.

bNOS-dependent NO production is involved in the synthesis of a nitrated phytotoxin thaxtomin A and thereby plays a major role in the pathogenesis of *Streptomyces* spp., (Johnson et al., 2008). Endogenous NO produced by NOS indirectly protects *Deinococcus radiodurans* against ultraviolet radiation (Patel et al., 2009).

In bacteria, where NOS is expressed (e.g. *S. aureus*, *B. subtilis*, *B. anthracis*), flavohemoglobins (flavoHbs) are co-expressed and in the presence of O₂ may convert bNOS-derived NO to NO₃⁻ with electron transfer from NAD(P)H to the ferric heme iron ligand via FAD (Bang et al., 2006; Ilari and Boffi, 2008; Nobre et al., 2008):



Hence NOS-produced NO may be consumed by bacteria in a balanced way, although evidence for and the rate of NO consumption in a reaction with endogenous flavoHbs have to be elucidated in further studies (Rafferty, 2011).

Moreover, it was observed that also SOD A expression in *B. subtilis* is significantly increased by bNOS activity. From this observation it was speculated that NO can act as a transcriptional regulator, however, a

mechanism of this regulation has not been revealed (details in Rafferty, 2011).

Recently, Schreiber et al. (2011) showed that biofilm dispersal of *B. subtilis* appears to be affected by NOS activity. The authors suggested that NO is involved in the fine-tuning decision between adaptation to anoxic conditions (in the biofilm) or dispersal from the biofilm. A role of NO for biofilm dispersion was reported for the pathogens *Pseudomonas aeruginosa* (Barraud et al., 2006, 2009a) and *S. aureus* (Schlag et al., 2007), the myxomycete *Candida albicans*, as well as in mixed-species biofilms (Barraud et al., 2009b). In contrast, in many Gram-negative bacteria, where NO is mainly synthesized as a by-product by NAR/NAP during denitrification, NO can play a signalling function to enhance biofilm formation. For instance, NO not only induces biofilm formation, but also up-regulates the genes involved in NIR and NAP synthesis and oxidative stress tolerance in *Neisseria gonorrhoeae* (Falsetta et al., 2011). Moreover, NO triggers the transcription of a gene obligatory for attachment and initial biofilm formation in a number of nitrifying bacteria (e.g. *Nitrosomonas europaea*, *Nitrosolobus multiformis* and *Nitrospira briensis*) (Schmidt et al., 2004).

Increasing the NO concentration in the medium induced the formation of biofilms by the Gram-negative rhizobacteria *Azospirillum brasilense*, whilst a gradual decrease of NO in the medium appears to mobilize cell motility (Arruebarrena Di Palma et al., 2013). The authors showed that both endogenously produced and exogenously added NO (e.g. GSNO as NO-donor) caused the same response. Apparently, NO-mediated effects on

bacterial biofilm formation or dispersal are species-specific phenomena, depending on N availability (e.g. at the soil microsite or the host environment).

4.2. Functions of NO in protists

NOS activity in myxomycetes (e.g. *Physarum polycephalum*) is induced under nutrient limitation and is involved in sporulation, but the mechanisms responsible so far have not been described (Messner et al., 2009).

Recently, a NOS enzyme without a reductase domain, but resembling bacterial NOS, was found in the eukaryotic unicellular alga *Naegleria gruberi* (Fritz-Laylin et al., 2009). Characterization of the enzyme and its function has not been reported.

4.3. Functions of NO in animals

Invertebrates and vertebrates (i.e. worms, beetles,, rodents, moles,, ruminants) influence the physical and chemical composition of soil, by burrowing, compaction and deposition of nutrients (i.e. faeces, urine and saliva), thereby indirectly influence NO production and consumption processes.

4.3.1. Invertebrates

In invertebrates (e.g. roundworms) NO can prolong life and mediate stress resistance to heat (Gusarov et al., 2013), Cd²⁺ toxicity (Cui et al., 2007) and the response to pathogenic bacteria (e.g. *P. aeruginosa*) (Troemel et al., 2006).

In addition, in invertebrates (e.g. echinoderms, coelenterates, nematodes, annelids, insects, crustaceans and molluscs) NO is of ubiquitous importance as an orthograde transmitter and a co-transmitter in signalling cascades as well as a modulator of conventional transmitter release (Jacklet 1997). These signalling functions of NOS-derived NO include neuronal sensory, including chemosensory (Gelperin, 1994; Jacklet and Gruhn, 1994; Elphick et al., 1995), as well as signalling in learning processes (Robertson et al., 1995; Kendrick et al., 1997; Müller, 1997) and development (Davis and Murphey, 1994; Kuzin et al., 1996; Froggett and Leise, 1997; Jacklet, 1997; Meleshkevitch et al., 1997). In the bug *Rhodnius* NO mediated vasodilation (Nussenzveig et al., 1995); the exact mechanism is not clear, but may resemble that identified in mammals (Jacklet, 1997). Comprehensive studies carried out (Susswein and Chiel, 2012) on the sea slug *Aplysia* elucidated that NO plays a major role in neuron mediated control of food finding and food consumption. Thus, NO is associated with the neural function of the swallow-rejecting mechanism, i.e. the rejection and reposition of mechanically resistant food, and the formation of memories of food inedibility (learning function), when food could not be swallowed successfully (Susswein and Chiel, 2012).

NO produced by bacteria (e.g. *B. subtilis*, *E. coli* with a NOS plasmid), previously eaten by the roundworm *Caenorhabditis elegans* (lacking its own NOS), diffuses into the worm's intestine tissues and triggers a cascade of signalling reactions causing a specific transcriptional response that promotes thermotolerance and prolongs life (Gusarov et al., 2013). The anti-aging effects of bacterial NO, were demonstrated by adding exogenous NO to the growth medium of the worm (Gusarov et al., 2013). The authors suggested that similar mechanisms may be relevant in higher organisms, one example may be the beneficial effect of 'normal' gastrointestinal microbiota. Such gastrointestinal microbes, predominately Gram-positive lactic acid bacteria (e.g. *Lactobacillus*, *Streptococcus*, *Lactococcus* spp.) possess NOS (Masson et al., 2011 and references therein) and NOS-derived NO may be used by the host. Thus, bacterial NO may diffuse into gastrointestinal tract cells and increase the level of available NO, which together with endogenous produced NO by the host may be involved in vasodilation, vasoprotection, cytoprotection, neuroprotection, etc. (Lundberg et al., 1994, 2008; Velayutham and Zweier, 2013 and references therein). However, this suggestion requires further investigations.

4.3.2. Mammals (including humans)

It has been clearly demonstrated that in mammals NO is involved in the regulation of synaptic signalling events, blood pressure, gut peristalsis, vasodilation, penile erection, developing retinal tissue at the level of gene transcription, mRNA translation and post-translational modifications of

proteins (Forstermann and Sessa, 2012; Socodato et al., 2013). Zhou and Zhu (2009) indicated that NO is also engaged in modulating memory, learning and neurogenesis. The functions of NO in mammals include a whole set of both positive and negative effects listed in Table 3.

INSERT Table 3 HERE

The presumably healthy human population of Earth (7.22 billion in March, 2014 according to Worldometers, 2014) annually exhales approximately 92 Gg N-NO a⁻¹ (estimated using Antczak et al. (2011), Davies and Moores (2003) and Levitzky (2003) data), which is equally to 1% of total soil emission (IPCC, 2007). Undoubtedly, this value is an underestimation, as people suffering from inflammatory diseases or physiological problems exhale higher rates of NO (Kharitonov et al., 1996; Fuchs et al., 2012). Exhaled NO has been proposed as an inflammatory disease marker for humans, since iNOS can be triggered to a greater degree by inflammatory cytokines, endotoxines and viral infections (Asano et al., 1994; Hunt et al., 2000; Antczak et al., 2011). We can speculate with confidence that NO is also exhaled by other mammals, including those living in the soil. Hence, the total exhaled NO rate of mammals is likely to be much higher than the estimate for the human population.

4.4. Functions of NO in plants

1409 In plants NO is a ubiquitous endogenous key mediator of numerous
1410 physiological and developmental processes (Guo et al., 2003; Lamattina et
1411 al., 2003; Wendehenne et al., 2004; Delledonne, 2005; Besson-Bard et al.,
1412 2008; Neill et al., 2008). In the aboveground parts of the plant, it is, for
1413 example, involved in flowering, seed germination and floral development;
1414 in belowground parts in root organogenesis, lateral root development, and
1415 formation of root hairs and adventitious roots (see review by Mur et al.,
1416 2012 and references therein). NO also plays a role in plant-microbe
1417 interaction including host defense, pathogen virulence and symbiotic
1418 interaction (Mur et al., 2012). In addition, it fulfills functions in stomatal
1419 regulation (García-Mata and Lamattina, 2001; Desikan et al., 2002; Neill et
1420 al., 2002), root nitrogen uptake and metabolism (Simon et al., 2009; 2013)
1421 and adaptive responses to abiotic stress (Neill et al., 2003; 2008; Besson-
1422 Bard et al., 2008; Mur et al., 2012). Abiotic stress reactions with proven
1423 participation of NO signaling include drought (García-Mata and Lamattina,
1424 2001; Desikan et al., 2002; Neill et al., 2002; Freschi et al. 2010), salinity
1425 (Zhang et al., 2004, 2006; Liu et al., 2007; Shi et al., 2007; Zhao et al.,
1426 2007; David et al., 2010; Chen et al., 2013), heat (Leshem et al., 1998;
1427 Gould et al., 2003), cold (Zhao et al. 2009) and flooding (Dean and Harper,
1428 1986; Guo et al., 2003; Zhang et al., 2006; Ferreira et al., 2010; Gupta and
1429 Kaiser, 2010; Gupta et al., 2012). All these environmental factors cause
1430 oxidative stress in plants; it is therefore suggested that NO stimulates
1431 antioxidative defense mechanisms during periods of elevated production
1432 and abundance of reactive oxygen species (ROS) (Neill et al., 2008).

NO production by plants is of particular significance upon nitrate reduction in roots under hypoxia (Dean and Harper, 1986; Dordas et al., 2003, 2004; Igamberdiev et al., 2004; Igamberdiev and Hill, 2009; Gupta and Kaiser, 2010; Gupta et al., 2012). NO formation was determined in these studies directly in the tissue affected by hypoxia stress. Recently, NO emissions were measured from the leaves of trees, where only the root system was flooded (Copolovici and Niinemets, 2010). Because NO emissions were highest in flooding sensitive and lowest in flooding tolerant species, NO emissions were suggested to be a marker of flooding tolerance. In addition, a regulatory function of NO in stomatal conductance of flooded plants was postulated (Copolovici and Niinemets, 2010). The significance of NO produced in plant roots upon hypoxia for other soil biota has so far not been elucidated. In addition, the contribution of plant derived NO for NO emissions from the soil and from aboveground parts of plants into the atmosphere has so far not been quantified.

In plants, NO is involved in protein modification as posttranslational regulator of enzymes both directly and indirectly via its derivatives (RNS). S-nitrosylation of cysteine, nitrosylation of transition metals and tyrosine nitration appear to be the main NO-associated protein modifications. S-nitrosylation is involved in gene regulation, modulates phytohormone signalling and can control programmed cell death (PCD) in opposing ways (promote or inactivate) (Hara et al., 2005; Melotto et al., 2006; Belenghi, 2007; Forman et al., 2008; Tada et al., 2008). NO regulation of gene expression via S-nitrosylation has been widely reported (Grün et al., 2006 and reference therein). However, the regulatory mechanisms involved in this

regulation are still unclear (Grün et al., 2006; Leitner et al., 2009). NO appears to modulate the response of phytohormones, involved in pathogen-induced stomatal movements via S-nitrosylation of K⁺ outward channels (Sokolovski et al., 2004; Melotto et al., 2006). An opposite function of NO-mediated S-nitrosylation in apoptosis is connected with cytosolic glyceraldehyde 3-phosphate dehydrogenase (GAPDH) inactivation. The role of metal nitrosylation in plants has not been revealed yet, but it seems that cytochrome P450s could be a target (Leitner et al., 2009). NO can easily neutralize harmful O₂⁻ to form peroxynitrite (ONOO⁻); ONOO⁻ can further react with tyrosine residues by nitration, thereby enhancing tyrosine residue containing proteins' susceptibility to proteolysis (Grune et al., 1998; Souza et al., 2000). Tyrosine nitration is associated with disease resistance response (Sailto et al., 2006; Romero-Puertas et al., 2007; Cecconi et al., 2009), plant resistance to abiotic and biotic stresses, but is also important for normal growth, fertility and reproduction of plants (Rusterucci et al., 2007; Lee et al., 2008; Leitner et al., 2009).

The following pathways of NO scavenging have been considered in plant cells. NO can be transformed to nitrate by non-symbiotic haemoglobins under hypoxic stress (Perazolli et al., 2004), providing cells with NO₃⁻, an important nutrient which acts as a signal for plant growth and regulates of genes expression (Crawford and Glass, 1998 and reference therein; Stitt et al., 2002 and reference therein). NO can easily react with glutathione (GSH) to form S-nitrosylated glutathione (GSNO). Further, GSNO can be used as a NO storage pool and/or act a transnitrosylation agent, or can be reduced by S-nitrosogluthathione reductase (GSNOR), producing oxidised glutathione

(GSSG) and NH_3 . Great significance is attributed to the reaction of NO with superoxide to form ONOO^- , which can be detoxified by peroxiredoxins with nitrite production or react with tyrosine residues. Resistance during biotic and abiotic stresses appears to be associated with NO-mediated GSNO formation and transport in systemic stress signalling, as well as tyrosine nitration (Saito et al., 2006; Corpas et al., 2008).

4.4.1. Microbial NO and plant pathogenesis

Plant-pathogenic *Streptomyces* spp. produce endogenous NO catalysed by the bNOS enzyme at the host-pathogen interface, and is induced by cellobiose, a disaccharide product of cellulose degradation (Johnson et al., 2008). In fact, bNOS-derived NO is used for nitration of thaxtomin A, a dipeptide phytotoxin, which inhibits cellulose biosynthesis (Johnson et al., 2008; Fry and Loria, 2002; Scheible et al., 2003). Since NO can easily diffuse through biological membranes and is also well known as a defence and signalling molecule in plants, the NO produced by *Streptomyces* spp. in response to the degradation of the host plant cell wall is likely to penetrate into plant tissues, thereby affecting the plant signalling systems (Johnson et al., 2008).

4.4.2. Soil microbial NO and plant root processes

NO plays a significant role in legume-rhizobium symbiosis, since both plant and bacteria are involved in production and metabolism of NO

(Meilhoc et al., 2011). NOS-like activity was observed in free living rhizobia under anaerobic condition (Pii et al., 2007) as well as during the symbiosis establishment phase (Meilhoc et al., 2011). In mature N₂-fixing nodules denitrification and the plant NR/mitochondrial electron transport chain (ETC) system seem to be basic NO sources under micro-oxic condition (Sanchez et al., 2010; Horchani et al., 2011). Signalling functions of NO are attributed to the expression of genes involved in nodule organogenesis, C- and N-metabolism, redox response, and cell division (Cooper, 2004; Frendo et al., 2005; Pii et al., 2007). It also has been shown that functional nodules of *Glycine max* (Meakin et al., 2007), and *Medicago truncatula* (Horchani et al., 2011) increased their NO production under oxygen limiting condition.

In greenhouse experiments it was shown that the rhizosphere NO concentration modulated uptake of N compounds by tree roots (Simon et al., 2009, 2013). It is therefore assumed that soil microbial NO is sensed by roots and acts as a signal determining the competitive strength of roots in the acquisition of N sources from the soil. At the ecosystem level, this signalling function of NO appears highly important, particularly in low N soils, since plant root and bacteria compete for the same inorganic and organic N sources (Stoelken et al., 2010). It is currently unknown if N acquisition by mycorrhizal fungi is also subject to bacterial NO mediated modulation. It is also unclear if this signalling process is based on root surface interactions or requires NO influx into the root. Since NO action is thought to take place at the level of posttranslational protein modification (Leitner et al., 2009), it is feasible that NO of microbial origin acts on the

outer surface of the plasmalemma on transmembrane proteins responsible of N transport processes. If microbial NO would pass the plasmalemma, it would directly interact with plant responses to abiotic stress such as salinity, high temperature, high light intensity and anoxia. These environmental factors are all subject to signalling by posttranslational modifications mediated by NO internally produced by plants (Leitner et al., 2009). Therefore, it appears that a clear separation of external NO of bacterial origin and internally produced NO is highly desirable for the interaction of plants with its ever changing environment. Still NO influx into the roots is likely to take place, since other trace gases of soil microbial origin such as CH₄ and N₂O, are subject to root influx, plant mediated transport, and release from the shoot into the atmosphere (Schütz et al., 1991; Butterbach-Bahl et al., 1997; Machacova et al., 2013). The contribution of this pathway to the release of soil microbial NO into the atmosphere is currently unknown. It also remains to be analysed if some of the NO produced inside plant cells is emitted into the atmosphere.

5. Conclusions

New approaches and techniques, e.g. stable isotope labelling, inhibitor application, gas-flow-soil-core and chamber methods, “omics” technologies, have improved existing understanding and have discovered new mechanisms of N transformation leading to NO production. It is likely that archaea are important players involved in processes related to ammonia

oxidation especially in NH_4^+ -poor and/or acid environments. It has clearly been demonstrated that:

(a) nitrite is the main precursor for NO under both oxic and anoxic condition, but sources for NO_2^- can be linked either to oxidative or reductive microbial N transformation pathways;

(b) ammonium is the dominant (70%) source of NO under aerobic condition, which confirms previous reports that nitrification is the prevailing process responsible for soil NO production;

(c) nitrate is a dominant (87%) source of NO under anoxic condition, which elucidates the significant role of denitrification in NO production;

(d) nitric oxide is a free (and non-enzyme-bound) precursor for N_2O under anaerobic conditions, thereby confirming the “diffusion limitation” hypothesis.

Our literature review suggests that NO/ N_2O emission ratios are possibly not good predictors of the NO production pathway (nitrification or denitrification). There is some evidence that periplasmic and cytoplasmic DNRA may produce NO, but the significance at ecosystem level needs to be studied. Codenitrification process has been shown to mediate NO production by denitrification. Significance of NO for the anaerobic processes anammox and N-AOM has been elucidated as well as the potential importance of NO loss/leakage; the latter urgent needs for further investigations.

We have described a theoretically feasible unspecific enzymo-oxidative mechanism of NO production in soils, which suggests that not only nitrifying and denitrifying microbes produce NO, but that also extracellular enzymes from a wide range of microorganisms could influence NO production.

NO is a signalling molecule due to its ability to diffuse freely across biological membranes, hence it can directly or indirectly (via RNS) modulate the activities of cellular and extracellular proteins in various groups of organisms, implementing significant physiological functions.

NOS seems to be a ubiquitous trans-species enzyme (although its presence in plants has not been confirmed yet), which is responsible for NO synthesis in various organisms. However, role of NO production via NOS in ecosystem functioning is unknown.

In bacteria NO production is associated with a defence function in early stages of infection. At the same time NO produced by the host organism is part of its protective system against pathogens. Furthermore bNOS-derived NO from non-pathogenic and opportunistic bacteria can diffuse to host cells and can be used by a host for a wide range of physiological purposes, i.e. cause beneficial effect on inter-organismic level.

A new role of soil microbial NO in determining the competition between microbial and plant use of soil nitrogen resources has been recently suggested, but still requires validation at the field and identification on the mechanisms involved. In addition, the role of plants in mediating the exchange of microbial NO into the atmosphere requires further investigations.

A wide range of prokaryotes and eukaryotes are able to produce NO by multiple pathways for its own purposes, since each cell needs a sufficient amount of NO for its normal physiological functioning. However it is unknown to what extent cells rely on NO produced by exogenous processes. Detailed studies of the cellular NO demand in physiological processes will provide a closer understanding of NO exchange at the cellular and the organismic level.

Many NO consumption pathways have been described, both abiotic (e.g., nitrosation and possible reaction with SOM in soil; reactions in soil-atmosphere surface) and biotic processes (e.g., denitrification, codenitrification, anammox, N-AOM, detoxification, for physiological purposes).

Detailed investigations are needed to clarify molecular mechanisms of NO production and consumption, its controlling factors, and the significance of NO as a regulator of microbial, animal and plant processes in order to gain a better understanding of soil NO emissions to the atmosphere.

Acknowledgments

Authors gratefully acknowledge support from the projects "Effects of Climate Change on Air Pollution Impacts and Response Strategies for European Ecosystems" (ÉCLAIRE), funded under the EC 7th Framework Programme (Grant Agreement No. 282910), EU COST Action ES0804 - Advancing the integrated monitoring of trace gas exchange Between Biosphere and Atmosphere (ABBA) and the Transnational Access to Research Infrastructures activity in EC 7th Framework Programme under

the Experimentation in Ecosystem Research (ExpeER, Grant Agreement No. 262060) project for conducting the research. We would like to express our gratitude to anonymous reviewers and the editor, Prof. Dr. Dave Coleman for their constructive comments and suggestions.

References

Adak, S., Aulak, K.S., Stuehr, D.J., 2002. Direct evidence for nitric oxide production by a nitric-oxide synthase-like protein from *Bacillus subtilis*. *Journal of Biological Chemistry* 277, 16167-16171.

Akiyama, H., Morimoto, S., Hayatsu, M., Hayakawa, A., Sudo, S., Yagi, K., 2013. Nitrification, ammonia-oxidizing communities, and N₂O and CH₄ fluxes in an imperfectly drained agricultural field fertilized with coated urea with and without dicyandiamide. *Biology and Fertility of Soils* 49, 213-223.

Akiyama, H., Tsuruta, H., 2003. Nitrous oxide, nitric oxide, and nitrogen dioxide fluxes from soils after manure and urea application. *Journal of Environmental Quality* 32, 423-431.

Aktan, C.K., Yapsakli, K., Mertoglu, B., 2012. Inhibitory effects of free ammonia on Anammox bacteria. *Biodegradation* 23, 751-762.

Alscher, R.G., Erturk, N., Heath, L.S., 2002. Role of superoxide dismutases (SODs) in controlling oxidative stress in plants. *Journal of Experimental Botany* 53, 1331-1341.

Anderson, I.C.; Levine, J.S., 1986. Relative rates of nitric oxide and nitrous oxide production by nitrifiers, denitrifiers, and nitrate respirers. *Applied and Environmental Microbiology* 51, 938-945.

- 1656 Antczak, A., Ciebiada, M., Kharitonov, S.A., Gorski, P., Barnes, P.J.,
1657 2012. Inflammatory markers: exhaled nitric oxide and carbon monoxide
1658 during the ovarian cycle. *Inflammation* 35, 554-559.
- 1659 Anthonisen, A.C., Loehr, R.C., Prakasam, T.B.S., Srinath, E.G., 1976.
1660 Inhibition of nitrification by ammonia and nitrous acid. *Journal of Water*
1661 *Pollution Control Federation*, 835-852.
- 1662 Arah, J.R.M., 1997. Apportioning nitrous oxide fluxes between
1663 nitrification and denitrification using gas-phase mass spectrometry. *Soil*
1664 *Biology and Biochemistry* 29, 1295-1299.
- 1665 Arruebarrena Di Palma, A., Pereyra, C.M., Moreno Ramirez, L.,
1666 Vázquez, X., María, L., Baca, B. E., Pereyra M.A., Lamattina, L., Creus,
1667 C.M., 2013. Denitrification-derived nitric oxide modulates biofilm
1668 formation in *Azospirillum brasilense*. *FEMS Microbiology Letters* 338, 77-
1669 85.
- 1670 Arp, D.J., Sayavedra-Soto, L.A., Hommes, N.G., 2002. Molecular
1671 biology and biochemistry of ammonia oxidation by *Nitrosomonas europaea*.
1672 *Archives of microbiology* 178, 250-255.
- 1673 Asano, K., Chee, C.B., Gaston, B., Lilly, C.M., Gerard, C., Drazen, J.M.,
1674 Stamler, J.S., 1994. Constitutive and inducible nitric oxide synthase gene
1675 expression, regulation, and activity in human lung epithelial cells.
1676 *Proceedings of the National Academy of Sciences* 91, 10089-93.
- 1677 Averill, B.A., 1996. Dissimilatory nitrite and nitric oxide reductases.
1678 *Chemical Reviews* 96, 2951-2964.

- 1679 Avrahami, S., Conrad, R., 2005. Cold-temperate climate: A factor for
1680 selection of ammonia oxidizers in upland soil? Canadian Journal of
1681 Microbiology 51, 709-714.
- 1682 Avrahami, S., Liesack, W., Conrad, R., 2003. Effects of temperature and
1683 fertilizer on activity and community structure of soil ammonia oxidizers.
1684 Environmental Microbiology 5, 691-705.
- 1685 Babitha, M.P., Bhat, S.G., Prakash, H.S., Shetty, H.S., 2002. Differential
1686 induction of superoxide dismutase in downy mildew-resistant
1687 and-susceptible genotypes of pearl millet. Plant Pathology 51, 480-486.
- 1688 Baggs, E.M., 2011. Soil microbial sources of nitrous oxide: recent
1689 advances in knowledge, emerging challenges and future direction. Current
1690 Opinion in Environmental Sustainability 3, 321-327.
- 1691 Bakken, L.R., Bergaust, L., Liu, B., Frostegård, Å., 2012. Regulation of
1692 denitrification at the cellular level: a clue to the understanding of N₂O
1693 emissions from soils. Philosophical Transactions of the Royal Society B:
1694 Biological Sciences 367, 1226-1234.
- 1695 Banerjee, S., Si, B.C., Siciliano, S.D., 2011. Evidence of high microbial
1696 abundance and spatial dependency in three arctic soil ecosystems. Soil
1697 Science Society of America Journal 75, 2227-2232.
- 1698 Bang, I.S., Liu, L., Vazquez-Torres, A., Crouch, M.L., Stamler, J.S.,
1699 Fang, F.C., 2006. Maintenance of nitric oxide and redox homeostasis by the
1700 *Salmonella* flavohemoglobin hmp. Journal of Biological Chemistry 281,
1701 28039-28047.
- 1702 Baranenko, V.V., 2006. Superoxide dismutase in The Plant Cells.
1703 Cytology 48, 465-474 (in Russian).

- 1704 Barraud, N., Hassett, D.J., Hwang, S.H., Rice S.A., Kjelleberg, S., Webb,
 1705 J.S., 2006. Involvement of nitric oxide in biofilm dispersal of *Pseudomonas*
 1706 *aeruginosa*. Journal of Bacteriology 188, 7344-7353.
- 1707 Barraud, N., Schleheck, D., Klebensberger, J., Webb, J.S., Hassett, D.J.,
 1708 Rice, S.A., Kjelleberg, S., 2009a. Nitric oxide signaling in *Pseudomonas*
 1709 *aeruginosa* biofilms mediates phosphodiesterase activity, decreased cyclic
 1710 di-GMP levels, and enhanced dispersal. Journal of Bacteriology 191, 7333-
 1711 7342.
- 1712 Barraud, N., Storey, M.V., Moore, Z.P., Webb, J.S., Rice, S.A.,
 1713 Kjelleberg, S., 2009b. Nitric oxide-mediated dispersal in single- and multi-
 1714 species biofilms of clinically and industrially relevant microorganisms.
 1715 Microbial Biotechnology 2, 370-378.
- 1716 Barton, L., McLay, C.D.A., Schipper, L.A., Smith, C.T., 1999. Annual
 1717 denitrification rates in agricultural and forest soils: a review. Australian
 1718 Journal of Soil Research 37, 1073-1093.
- 1719 Bateman, E.J., Baggs, E.M, 2005. Contributions of nitrification and
 1720 denitrification to N₂O emissions from soils at different water filled pore
 1721 space. Biology and Fertility of Soils 41, 379-388.
- 1722 Bates, S.T., Nash III, T.H., Garcia-Pichel, F., 2012. Patterns of diversity
 1723 for fungal assemblages of biological soil crusts from the southwestern
 1724 United States. Mycologia 104, 353-361.
- 1725 Baumgärtner, M., Koschorreck, M., Conrad, R., 1996. Oxidative
 1726 consumption of nitric oxide by heterotrophic bacteria in soil. FEMS
 1727 Microbiology and Ecology 19, 165-170.

- 1728 Beaumont, H.J., Lens, S.I., Reijnders, W.N., Westerhoff, H.V., Van
1729 Spanning, R.J., 2004. Expression of nitrite reductase in *Nitrosomonas*
1730 *europaea* involves NsrR, a novel nitrite-sensitive transcription repressor.
1731 Molecular microbiology 54, 148-158.
- 1732 Beaumont, H.J., Lens, S.I., Westerhoff, H.V., Van Spanning, R.J., 2005.
1733 Novel nirK cluster genes in *Nitrosomonas europaea* are required for NirK-
1734 dependent tolerance to nitrite. Journal of bacteriology 187, 6849-6851.
- 1735 Belenghi, B., Romero-Puertas, M.C., Vercammen, D., Brackener, A.,
1736 Inzé, D., Delledonne, M., Van Breusegem, F., 2007. Metacaspase activity of
1737 *Arabidopsis thaliana* is regulated by S-nitrosylation of a critical cysteine
1738 residue. Journal of Biological Chemistry 282:1352-1358.
- 1739 Bengtsson, G., Bergwall, C., 2000. Fate of ¹⁵N labelled nitrate and
1740 ammonium in a fertilized forest soil. Soil Biology and Biochemistry 32,
1741 545-557.
- 1742 Bergaust, L., van Spanning, R.J., Frostegård, Å., Bakken, L.R., 2012.
1743 Expression of nitrous oxide reductase in *Paracoccus denitrificans* is
1744 regulated by oxygen and nitric oxide through *FnrP* and *NNR*. Microbiology
1745 158, 826-834.
- 1746 Berthrong, S.T., Schadt, C.W., Pineiro, G., Jackson, R.B., 2009.
1747 Afforestation alters the composition of functional genes in soil and
1748 biogeochemical processes in South American grasslands. Applied and
1749 Environmental Microbiology 75, 6240-6248.
- 1750 Besson-Bard, A., Pugin, A., Wendehenne, D., 2008. New insights into
1751 nitric oxide signaling in plants. Annual Review of Plant Biology 59, 21-39.

- 1752 Beyer, W., Imlay, J., Fridovich, I., 1991. Superoxide dismutases.
1753 Progress in Nucleic Acid Research and Molecular Biology 40, 221-253.
- 1754 Boje, K.M., Arora, P.K., 1992. Microglial-produced nitric oxide and
1755 reactive nitrogen oxides mediate neuronal cell death. Brain Research 587,
1756 250-256.
- 1757 Bolli, R., Li, Q.H., Tang, X.L., Guo, Y., Xuan, Y.T., Rokosh, G., Dawn,
1758 B., 2007. The late phase of preconditioning and its natural clinical
1759 application - gene therapy. Heart Failure Reviews 12, 189-199.
- 1760 Bollmann, A., Conrad, R., 1997. Acetylene blockage technique leads to
1761 underestimation of denitrification rates in oxic soils due to scavenging of
1762 intermediate nitric oxide. Soil Biology and Biochemistry 29, 1067–1077.
- 1763 Bollmann, A., Koschorreck, M., Meuser, K., Conrad, A., 1999.
1764 Comparison of two different methods to measure nitric oxide turnover in
1765 soils. Biology and Fertility of Soils 29, 104–110.
- 1766 Bouwman, A.F., Boumans, L.J.M., Batjes, N.H., 2002. Modeling global
1767 annual N₂O and NO emissions from fertilized fields. Global
1768 Biogeochemical Cycles 16, 1080.
- 1769 Bouwman, A., 1998. Nitrogen oxides and tropical agriculture. Nature
1770 392, 886-887.
- 1771 Bowler, C., Van Camp, W., Van Montagu, M., Inzé, D., Asada, K., 1994.
1772 Superoxide dismutase in plants. Critical Reviews in Plant Sciences 13, 199-
1773 218.
- 1774 Brankatschk, R., Töwe, S., Kleineidam, K., Schlöter, M., Zeyer, J., 2010.
1775 Abundances and potential activities of nitrogen cycling microbial

- 1776 communities along a chronosequence of a glacier forefield. ISME Journal 5,
1777 1025-1037.
- 1778 Bremner, J.M., Fúhr, F., 1966. Tracer studies of the reaction of soil
1779 organic matter with nitrite. In: The use of isotopes in soil organic matter
1780 study. Pergamon Press, New York, pp. 337–346.
- 1781 Broadbent, F.E., Clark, F., 1965. Denitrification. In: Bartholomew, W.V.,
1782 Clark, F. (Eds.), Soil Nitrogen, American Society of Agronomy, Madison,
1783 pp. 344-359.
- 1784 Brochier-Armanet, C., Boussau, B., Gribaldo, S., Forterre, P., 2008.
1785 Mesophilic crenarchaeota: Proposal for a third archaeal phylum, the
1786 *Thaumarchaeota*. Nature Reviews Microbiology 6, 245-252.
- 1787 Brown, G.C., 2010. Nitric oxide and neuronal death. Nitric Oxide 23,
1788 153-165.
- 1789 Brown, G.C., Neher, J.J., 2010. Inflammatory neurodegeneration and
1790 mechanisms of microglial killing of neurons. Molecular Neurobiology 41,
1791 242-247.
- 1792 Bru, D., Ramette, A., Saby, N.P., Dequiedt, S., Ranjard, L., Jolivet, C.,
1793 Arrouays, D., Philippot, L., 2010. Determinants of the distribution of
1794 nitrogen-cycling microbial communities at the landscape scale. ISME
1795 Journal 5, 532-542.
- 1796 Buresh, R.J., Patrick, W.H., 1978. Nitrate reduction to ammonium in
1797 anaerobic soil. Soil Science Society of America Journal 42, 913-918.
- 1798 Burmester, T., Hankeln, T., 2004. Neuroglobin: a respiratory protein of
1799 the nervous system. Physiology 19, 110-113.

- 1800 Butterbach-Bahl, K., Baggs, E.M., Dannenmann, M., Kiese, R.,
 1801 Zechmeister-Boltenstern, S., 2013. Nitrous oxide emissions from soils: how
 1802 well do we understand the processes and their controls? Philosophical
 1803 Transactions of the Royal Society B: Biological Sciences 368.
- 1804 Butterbach-Bahl, K., Gundersen, P., Ambus, P., Augustin, J., Beier, C.,
 1805 Boeckx, P., Dannenmann, M., Sanchez Gimeno, B., Ibrom, A., Kiese, R.,
 1806 Kitzler, B., Rees, R.M., Smith, K.A., Stevens, C., Vesala, T., Zechmeister-
 1807 Boltenstern, S., 2011. Nitrogen processes in terrestrial ecosystems. In:
 1808 Sutton, M.A., Howard, C.M., Erisman, J.W., Billen, G., Bleeker, A.,
 1809 Glennfeld, P., van Grinsven, H., Grizzetti, B. (Eds.), The European Nitrogen
 1810 Assessment , Cambridge University Press, Cambridge, pp. 99-125.
- 1811 Butterbach-Bahl, K., Kahl, M., Mykhayliv, L., Werner, C., Kiese, R., Li,
 1812 C., 2009. A European-wide inventory of soil NO emissions using the
 1813 biogeochemical models DNDC/Forest-DNDC. Atmospheric Environment
 1814 43, 1392-1402.
- 1815 Butterbach-Bahl, K., Kock, M., Willibald, G., Hewett, B., Buhagiar, S.,
 1816 Papen, H., Kiese, R., 2004. Temporal variations of fluxes of NO, NO₂, N₂O,
 1817 CO₂ and CH₄ in a tropical rain forest ecosystem. Global Biogeochemical
 1818 Cycles 18.
- 1819 Butterbach-Bahl, K., Papen, H., Rennenberg, H., 1997. Impact of gas
 1820 transport through rice cultivars on methane emission from rice paddy fields.
 1821 Plant, Cell and Environment 20, 1175-1183.
- 1822 Byrne, N., Strous, M., Crepeau, V., Kartal, B., Birrien, J.L., Schmid, M.,
 1823 Lesongeur, F., Schouten, S., Jaeschke, A., Jetten, M., Prieur, D., Godfroy,

- 1824 A., 2009. Presence and activity of anaerobic ammoniumoxidizing bacteria
1825 at deep-sea hydrothermal vents. *ISME Journal* 3, 117–123.
- 1826 Cabello, P., Roldán, M.D., Castillo, F., Moreno-Vivián, C., 2012. The
1827 Nitrogen Cycle. In: *PIV Biogeochemical Cycles and their consequences*. In:
1828 Schmidt, T.M, Schaechter, M., (Eds.), *Topics in Ecological and*
1829 *Environmental Microbiology*, Elsevier Academic Press, Oxford, pp. 483-
1830 707.
- 1831 Cabello, P., Roldán, M.D., Moreno-Vivián, C., 2004. Nitrate reduction
1832 and the nitrogen cycle in archaea. *Microbiology* 150, 3527-3546.
- 1833 Carrera, J., Jubany, I., Carvallo, L., Chamy, R., Lafuente, J. 2004.
1834 Kinetic models for nitrification inhibition by ammonium and nitrite in a
1835 suspended and an immobilised biomass systems. *Process Biochemistry* 39,
1836 1159-1165.
- 1837 Cecconi, D., Orzetti, S., Vandelle, E., Rinalducci, S., Zolla, L.,
1838 Delledonne, M., 2009. Protein nitration during defence response in
1839 *Arabidopsis thaliana*. *Electrophoresis* 30, 2460-2468.
- 1840 Chalk, P.M., Smith, C.J., 1983. Chemodenitrification. In: Freney, J.R.,
1841 Simpson, J.R. (Eds.), *Gaseous Loss of Nitrogen from Plant-Soil Systems*,
1842 Springer Science + Business Media, Dordrecht, pp. 65-89.
- 1843 Chameides, W.L., Kasibhata, P.S., Yieger, J., Levy, H., 1994. The
1844 growth of continentalscale metro-agro-plexes, regional ozone pollution, and
1845 world food production. *Science* 264, 74–77.
- 1846 Chao, C.C., Hu, S., Molitor, T.W., Shaskan, E.G., Peterson, P. K., 1992.
1847 Activated microglia mediate neuronal cell injury via a nitric oxide
1848 mechanism. *The Journal of Immunology* 149, 2736–2741.

- 1849 Chen, C.A., Wang, T.Y., Varadharaj, S., Reyes, L.A., Hemann, C.,
 1850 Talukder, M.A., Chen, Y.R., Druhan, L.J., Zweier, J.L., 2010. S-
 1851 glutathionylation uncouples eNOS and regulates its cellular and vascular
 1852 function. *Nature* 468, 1115–1118.
- 1853 Chen, D.L., Chalk, P.M., Freney, J.R., Smith, C.J., Luo, Q.X., 1995a.
 1854 Estimation of nitrification rates in flooded soils. *Microbial Ecology* 30, 269-
 1855 284.
- 1856 Chen, D.L., Chalk, P.M., Freney, J.R., 1995b. Distribution of reduced
 1857 products of ^{15}N labelled nitrate in anaerobic soils. *Soil Biology and*
 1858 *Biochemistry* 27, 1539-1545.
- 1859 Chen, J., Xiong, D-Y., Wang, W-H., Hu, W-J., Simon, M., Xiao, Q.,
 1860 Chen, J., Liu, T-W., Liu, X., Zheng, H-L., 2013. Nitric oxide mediates root
 1861 K^+/Na^+ balance in a Mangrove plant, *Kandelia obovata*, by enhancing the
 1862 expression of AKT1-type K^+ channel and Na^+/H^+ antiporter under high
 1863 salinity. *PLoS ONE* 8, e71543. doi:10.1371/journal.pone.0071543.
- 1864 Cheng, W., Tsuruta, H., Chen, G., Yagi, K., 2004. N_2O and NO
 1865 production in various Chinese agricultural soils by nitrification. *Soil*
 1866 *Biology and Biochemistry* 36, 953–963.
- 1867 Clark, F.E., 1962. Losses of nitrogen accompanying nitrification.
 1868 Transactions of the International Society of Soil Science, Communication
 1869 IV and V, 173–176.
- 1870 Cole, J.A., 1990. Physiology, biochemistry and genetics of nitrate
 1871 dissimilation to ammonia. In: Revsbech, N. P., Sørensen, J. (Eds.),
 1872 Denitrification in soil and sediment. Plenum Press, New York, pp. 57-76.

- 1873 Cole, J.A., Brown, C.M., 1980. Nitrite reduction to ammonia by
1874 fermentative bacteria: a short circuit in the biological nitrogen cycle. FEMS
1875 Microbiology Letters 7, 65-72.
- 1876 Colloff, M.J., Wakelin, S.A., Gomez, D., Rogers, S.L., 2008. Detection
1877 of nitrogen cycle genes in soils for measuring the effects of changes in land
1878 use and management. Soil Biology and Biochemistry 40, 1637–1645.
- 1879 Conrad, R., 1996. Metabolism of nitric oxide in soil and soil
1880 microorganisms and regulation of flux into the atmosphere. In: Murrell,
1881 J.C., Kelly, D.P. (Eds.), Microbiology of Atmospheric Trace Gases, NATO
1882 ASI Series, vol. 109, Springer, Berlin, pp. 167–203.
- 1883 Conrad, R., 2002. Microbiological and biochemical background of
1884 production and consumption of NO and N₂O in soil. In: Gasche, R., Papen,
1885 H., Rennenberg, H. (Eds.), Trace Gas Exchange in Forest Ecosystems,
1886 Kluwer Academic Publishers, Dordrecht, pp. 3-33.
- 1887 Cookson, W.R., Cornforth, I.S., Rowarth, J.S., 2002. Winter soil
1888 temperature (2-15 degrees C) effects on nitrogen transformations in clover
1889 green manure amended or unamended soils; a laboratory and field study.
1890 Soil Biology and Biochemistry 34, 1401-1415.
- 1891 Cooper, J.E., 2004. Multiple responses of rhizobia to flavonoids during
1892 legume root infection. Advances in Botanical Research 41, 1-62.
- 1893 Copolovici, L., Niinemets, Ü., 2010. Flooding induced emissions of
1894 volatile signalling compounds in three tree species with differing
1895 waterlogging tolerance. Plant, Cell and Environment 33, 1582-1594.

- 1896 Corker, H., Poole, R. K., 2003. Nitric oxide formation by *Escherichia*
 1897 *coli* dependence on nitrite reductase, the no-sensing regulator Fnr, and
 1898 flavohemoglobin Hmp. *Journal of Biological Chemistry* 278, 31584-31592.
- 1899 Corpas, F.J., Chaki, M., Fernandez-Ocana, A., Valderrama, R., Palma,
 1900 J.M., Carreras, A., Begara-Morales, J.C., Airaki, M., del Rio, L.A., Barroso,
 1901 J.B., 2008. Metabolism of reactive nitrogen species in pea plants under
 1902 abiotic stress conditions. *Plant and Cell Physiology* 49, 1711-1722.
- 1903 Costa, E., Pérez, J., Kreft, J.U., 2006. Why is metabolic labour divided in
 1904 nitrification? *TRENDS in Microbiology* 14, 213-219.
- 1905 Cousins, J., Hope, D., Gries, C., Stutz, J., 2003. Preliminary assessment
 1906 of arbuscular mycorrhizal fungal diversity and community structure in an
 1907 urban ecosystem. *Mycorrhiza* 13, 319-326.
- 1908 Crane, B.R., Sudhamsu, J., Patel, B.A., 2010. Bacterial nitric oxide
 1909 synthases. *Annual Review of Biochemistry* 79, 445-470.
- 1910 Crawford, N.M., Glass, A.D., 1998. Molecular and physiological aspects
 1911 of nitrate uptake in plants. *Trends in Plant Science* 3, 389-395.
- 1912 Crawford, M.J., Goldberg, D.E., 1998. Role for the *Salmonella*
 1913 flavohemoglobin in protection from nitric oxide. *Journal of Biological*
 1914 *Chemistry* 273, 12543-12547.
- 1915 Cross, R., Lloyd, D., Poole, R.K., Moir, J.W., 2001. Enzymatic removal
 1916 of nitric oxide catalyzed by cytochrome c' in *Rhodobacter capsulatus*.
 1917 *Journal of Bacteriology* 183, 3050-3054.
- 1918 Cui, F., Yan, G., Zhou, Z., Zheng, X., Deng, J., 2012. Annual emissions
 1919 of nitrous oxide and nitric oxide from a wheatmaize cropping system on a

- 1920 silt loam calcareous soil in the North China Plain. Soil Biology and
1921 Biochemistry 48, 10-19.
- 1922 Cui, Y., McBride, S.J., Boyd, W.A., Alper, S., Freedman, J.H., 2007.
1923 Toxicogenomic analysis of *Caenorhabditis elegans* reveals novel genes and
1924 pathways involved in the resistance to cadmium toxicity. Genome Biology
1925 8, R122.
- 1926 Dale, O.R., Tobias, C.R., Song, B., 2009. Biogeographical distribution of
1927 diverse anaerobic ammonium oxidizing (anammox) bacteria in Cape Fear
1928 River estuary. Environmental Microbiology 11, 1194–1207.
- 1929 Dalsgaard, T., Canfield, D.E., Petersen, J., Thamdrup, B., Acuna-
1930 Gonzalez, J., 2003. N₂ production by the anammox reaction in the anoxic
1931 water column of Golfo Dulce, Costa Rica. Nature 422, 606–608.
- 1932 Dalsgaard, T., Donald, B.T., Canfield, E., 2005. Anaerobic ammonium
1933 oxidation (anammox) in the marine environment. Research in Microbiology
1934 156, 457–464.
- 1935 Dalsgaard, T., Thamdrup, B., 2002. Factors controlling anaerobic
1936 ammonium oxidation with nitrite in marine sediments. Applied and
1937 environmental microbiology 68, 3802-3808.
- 1938 Daum, M., Zimmer, W., Papen, H., Kloos, K., Nawrath, K., Bothe, H.,
1939 1998. Physiological and molecular biological characterization of ammonia
1940 oxidation of the heterotrophic nitrifier *Pseudomonas putida*. Current
1941 Microbiology 37, 281-288.
- 1942 Davies, A. S., Moores, C., 2003. The Respiratory System. Churchill
1943 Livingstone, London, 197 pp.

- 1944 David, A., Yadav, S., Bhatla, S.C., 2010. Sodium chloride stress induces
 1945 nitric oxide accumulation in root tips and oil body surface accompanying
 1946 slower oleosin degradation in sunflower seedlings. *Physiologia Plantarum*
 1947 140, 342-354.
- 1948 Davidson, E.A., 1991. Fluxes of nitrous oxide and nitric oxide from
 1949 terrestrial ecosystems. In: Roger, J.E., Whitman, W.B. (Eds.), *Microbial*
 1950 *Production and Consumption of Greenhouse Gases: Methane, Nitrogen*
 1951 *Oxide, and Halo-methane*, American Society of Microbiology, Washington,
 1952 DC, pp. 219-235.
- 1953 Davidson, E.A., Keller, M., Erickson, H.E., Verchot, L.V., Veldkamp, E.,
 1954 2000 Testing a Conceptual Model of Soil Emissions of Nitrous and Nitric
 1955 Oxides. *BioScience*, 50, 667-680.
- 1956 Davidson, E.A, Kingeree, W., 1997. A global inventory of nitric oxide
 1957 emissions from soils. *Nutrient Cycling in Agroecosystems* 48, 37-50.
- 1958 Davis, G., Murphey, R., 1994. Long-term regulation of short term
 1959 transmitter release properties: retrograde signaling and synaptic
 1960 development. *TINS* 17, 9-13.
- 1961 De Bruijn, A.M.G., Butterbach-Bahl, K., Blagodatsky, S., Grote, R.,
 1962 2009. Model evaluation of different mechanisms driving freeze-thaw N₂O
 1963 emissions. *Agriculture, Ecosystems and Environment* 133, 196-207.
- 1964 Dean, J.V., Harper, J.E., 1986. Nitric oxide and nitrous oxide production
 1965 by soybean and winged bean during the in vivo nitrate reductase assay.
 1966 *Plant Physiology* 82, 718-723.
- 1967 Deiglmayr, K., Philippot, L., Tscherko, D., Kandeler, E., 2006. Microbial
 1968 succession of nitrate-reducing bacteria in the rhizosphere of *Poa alpina*

- 1969 across a glacier foreland in the Central Alps. *Environmental Microbiology*
 1970 8, 1600-1612.
- 1971 Delledonne, M., 2005. NO news is good news for plants. *Current*
 1972 *Opinion in Plant Biology* 8, 390-396.
- 1973 Delmas, R., Serça, D., Jambert, C., 1997. Global inventory of NO_x
 1974 sources. *Nutrient Cycling in Agroecosystems* 48, 51-60.
- 1975 del Prado, A., Merino, P., Estavillo, J.M., Pinto, M., Gonzalez-Murua,
 1976 C., 2006. N₂O and NO emissions from different N sources and under a
 1977 range of soil water contents. *Nutrient Cycling in Agroecosystems* 74, 229-
 1978 243.
- 1979 Derbyshire, E.R., Marletta, M.A., 2009. Biochemistry of soluble
 1980 guanylate cyclase. In: Schmidt, H.H.H.W., Hofmann, F., Stasch, J-P. (Eds.),
 1981 cGMP: Generators, Effectors and Therapeutic Implications, Springer Berlin
 1982 Heidelberg, Berlin, pp. 17-31.
- 1983 Desikan, R., Griffiths, R., Hancock, J., Neill, S., 2002. A new role for an
 1984 old enzyme: nitrate reductase-mediated nitric oxide generation is required
 1985 for abscisic acid-induced stomatal closure in *Arabidopsis thaliana*.
 1986 *Proceedings of the National Academy of Sciences USA* 99, 16314-16318.
- 1987 Di, H.J., Cameron, K.C., Shen, J.P., Winefield, C.S., O'Callaghan, M.,
 1988 Bowatte, S., He, J.Z., 2009. Nitrification driven by bacteria and not archaea
 1989 in nitrogen-rich grassland soils. *Nature Geoscience* 2, 621-624.
- 1990 Dimmeler, S., Zeiher, A.M. 1997. Nitric oxide and apoptosis: Another
 1991 paradigm for the double-edged role of nitric oxide. *Nitric Oxide* 1, 275-281.
- 1992 Dordas, C., Hasinoff, B.B., Igamberdiev, A.U., Manac'h, N., Rivoal, J.,
 1993 Hill, R.D., 2003. Expression of a stress-induced hemoglobin affects NO

- 1994 levels produced by alfalfa root cultures under hypoxic stress, *The Plant*
 1995 *Journal* 35, 763-770.
- 1996 Dordas, C., Hasinoff, B.B., Rivoal, J., Hill, R.D., 2004. Class-1
 1997 hemoglobins, nitrate and NO levels in anoxic maize cell-suspension
 1998 cultures, *Planta* 219, 66-72.
- 1999 Duc, L., Noll, M., Meier, B.E., Bürgmann, H., Zeyer, J., 2009. High
 2000 diversity of diazotrophs in the forefield of a receding alpine glacier.
 2001 *Microbial Ecology* 57, 179-190.
- 2002 Dunfield, P.F., Knowles, R., 1997. Biological oxidation of nitric oxide in
 2003 a humisol. *Biology and Fertility of Soils* 24, 294-300.
- 2004 Dunfield, P.F., Knowles, R., 1998. Organic matter, heterotrophic activity,
 2005 and NO consumption in soils. *Global Change Biology* 4, 199-207.
- 2006 Dunfield, P.F., Knowles, R., 1999. Nitrogen monoxide production and
 2007 consumption in an organic soil. *Biology and Fertility of Soils* 30, 153-159.
- 2008 El Azhar, S., Vandenabeele, J., Verstraete, W., 1986. Nitrification and
 2009 organic nitrogen formation in soils. *Plant and soil* 94, 383-399.
- 2010 El Azhar, S., Verhe, R., Proot, M., Sandra, P., Verstraete, W., 1986.
 2011 Binding of nitrite-N on polyphenols during nitrification. *Plant and Soil* 94,
 2012 369-382.
- 2013 Elphick, M., Kemenes, G., Staras, K., O'Shea, M., 1995. Behavioral role
 2014 for nitric oxide in chemosensory activation of feeding in a mollusc. *The*
 2015 *Journal of Neuroscience* 15, 7653-7664.
- 2016 Elvers, K.T., Wu, G., Gilberthorpe, N.J., Poole, R.K., Park, S.F., 2004.
 2017 Role of an inducible single-domain hemoglobin in mediating resistance to

2018 nitric oxide and nitrosative stress in *Campylobacter jejuni* and
 2019 *Campylobacter coli*. Journal of Bacteriology 186, 5332-5341.

2020 Erguder, T.H., Boon, N., Wittebolle, L., Marzorati, M., Verstraete, W.,
 2021 2009. Environmental factors shaping the ecological niches of ammonia-
 2022 oxidizing archaea. FEMS Microbiology Reviews 33, 855-869.

2023 Erusalimsky, J. D., Moncada, S., 2007. Nitric oxide and mitochondrial
 2024 signaling: From physiology to pathophysiology. Arteriosclerosis,
 2025 Thrombosis and Vascular Biology 27, 2524-2531.

2026 Ettwig, K.F., Butler, M.K., Le Paslier, D., Pelletier, E., Mangenot, S.,
 2027 Kuypers, M.M., Schreiber, F., Dutilh, B.E., Zedelius, J., de Beer, D.,
 2028 Gloerich, J., Wessels, H.J.C.T., van Alen, T., Luesken, F., Wu, M.L., van de
 2029 Pas-Schoonen, K.T., Op den Camp, H.J.M., Janssen-Megens, E.M.,
 2030 Francoijs, K-J., Stunnenberg, H., Weissenbach, J., Jetten, M.S.M., Strous,
 2031 M., 2010. Nitrite-driven anaerobic methane oxidation by oxygenic bacteria.
 2032 Nature 464, 543-548.

2033 Ettwig, K.F., Shima, S., van de Pas-Schoonen, K.T., Kahnt, J., Medema,
 2034 M. H., Op den Camp, H.J.M., Jetten, M.S.M., Strous, M., 2008.
 2035 Denitrifying bacteria anaerobically oxidize methane in the absence of
 2036 Archaea. Environmental Microbiology 10, 3164–3173.

2037 Ettwig, K.F., van Alen, T., van de Pas-Schoonen, K.T., Jetten, M.S.M.,
 2038 Strous, M., 2009. Enrichment and molecular detection of denitrifying
 2039 methanotrophic bacteria of the NC10 phylum. Applied and Environmental
 2040 Microbiology 75, 3656–3662.

2041 Falsetta, M.L., Steichen, C.T., McEwan, A.G., Cho, C., Ketterer, M.,
 2042 Shao, J., Hunt, J., Jennings, M.P., Apicella, M.A. 2011. The composition

2043 and metabolic phenotype of *Neisseria gonorrhoeae* biofilms. *Frontiers in*
 2044 *Microbiology* 2, 75.

2045 FAO, IFA, 2001. Global estimates of gaseous emissions of NH₃, NO and
 2046 N₂O from agricultural land, Food and Agriculture Organization of the
 2047 United Nations, International Fertilizer Industry Association, Rome, 106 pp.

2048 Fazzolari, E., Nicolardeot, B., Germon, J.C., 1998. Simultaneous effects
 2049 of increasing levels of glucose and oxygen partial pressure on denitrification
 2050 and dissimilatory nitrate reduction to ammonium in repacked soil cores.
 2051 *European Journal of Soil Biology* 34, 47-52.

2052 Ferreira, L.C., Cataneo, A.C., Remaeh, L.M.R., Corniani, N., de Fatima
 2053 Fumis, T., de Souza, Y.A., Scavroni, J., Soares B.J.A., 2010. Nitric oxide
 2054 reduces oxidative stress generated by lactofen in soybean plants. *Pesticide*
 2055 *Biochemistry and Physiology* 97, 47-54.

2056 Firestone, M.K., Davidson, E.A., 1989. Microbiological basis of NO and
 2057 N₂O production and consumption in soil. In: Andreae, M.O., Schimel, D.S.
 2058 (Eds.), *Exchange of Trace Gases between Terrestrial Ecosystems and the*
 2059 *Atmosphere*. John Wiley, New York, pp. 7-21.

2060 Finlayson-Pitts, B.J., Pitts, J.N. Jr., 1986. *Atmospheric Chemistry:*
 2061 *Fundamentals and Experimental Techniques*. John Wiley, New York, 1098
 2062 pp.

2063 Ford, P.C., Lorkovic, I.M., 2002. Mechanistic aspects of the reactions of
 2064 nitric oxide with transition-metal complexes. *Chemical Reviews* 102, 993-
 2065 1018.

2066 Forman, H.J., Fukuto, J.M., Miller, T., Zhang, H., Rinna, A., Levy, S.,
 2067 2008. The chemistry of cell signaling by reactive oxygen and nitrogen

- 2068 species and 4-hydroxynonenal. Archives of Biochemistry and Biophysics
 2069 477, 183-195.
- 2070 Forstermann, U., Sessa, W.C., 2012. Nitric oxide synthases: Regulation
 2071 and function. European Heart Journal 33, 829-837.
- 2072 Fowler, D., Pilegaard, K., Sutton, M.A., Ambus, P., Raivonen, M.,
 2073 Duyzer, J., Simpson, D., Fagerli, H., Fuzzi, S., Schjoerring, J.K., Granier,
 2074 C., Neftel, A., Isaksen, I.S.A., Laj, P., Maione, M., Monks, P.S., Burkhardt,
 2075 J., Daemmgen, U., Neiryneck, J., Personne, E., Wichink-Kruit, R.,
 2076 Butterbach-Bahl, K., Flechard, C., Tuovinen, J.P., Coyle, M., Gerosa, G.,
 2077 Loubet, B., Altimir, N., Gruenhage, L., Ammann, C., Cieslik, S., Paoletti,
 2078 E., Mikkelsen, T.N., Ro-Poulsen, H., Cellier, P., Cape, J.N., Horvath, L.,
 2079 Loreto, F., Niinemets, U., Palmer, P.I., Rinne, J., Misztal, P., Nemitz, E.,
 2080 Nilsson, D., Pryor, S., Gallagher, M.W., Vesala, T., Skiba, U., Bruggemann,
 2081 N., Zechmeister-Boltenstern, S., Williams, J., O'Dowd, C., Facchini, M.C.,
 2082 de Leeuw, G., Flossman, A., Chaumerliac, N., Erisman, J.W., 2009.
 2083 Atmospheric composition change: Ecosystems - Atmosphere interactions.
 2084 Atmospheric Environment 43, 5193-5267.
- 2085 Frendo, P., Harrison, J., Norman, C., Hernandez Jimenez, M.J., Van de
 2086 Sype, G., Gilabert, A., Puppo, A., 2005. Glutathione and homoglutathione
 2087 play a critical role in the nodulation process of *Medicago truncatula*.
 2088 Molecular Plant-Microbe Interactions 18, 254-259.
- 2089 Freppaz, M., Williams, B.L., Edwards, A.C., Scalenghe, R., Zanini, E.,
 2090 2007. Labile nitrogen, carbon, and phosphorus pools and nitrogen
 2091 mineralization and immobilization rates at low temperatures in seasonally
 2092 snow-covered soils. Biology and Fertility of Soils 43, 519-529.

- 2093 Freschi, L., Rodrigues, M.A., Domingues, D.S., Purgatto, E., Van Sluys,
 2094 M.A., Magalhaes, J.R., Kaiser, W.M., Mercier, H., 2010. Nitric oxide
 2095 mediates the hormonal control of Crassulacean acid metabolism expression
 2096 in young pineapple plants. *Plant Physiology* 152, 1971-1985.
- 2097 Frey, A.D., Farres, J., Bollinger, C.J.T., Kallio, P.T., 2002. Bacterial
 2098 hemoglobins and flavohemoglobins for alleviation of nitrosative stress in
 2099 *Escherichia coli*. *Applied and Environmental Microbiology* 68, 4835–4840.
- 2100 Fry, B.A., Loria, R., 2002. Thaxtomin A: evidence for a The Plant Cell
 2101 wall target. *Physiological and Molecular Plant Pathology* 60, 1-8.
- 2102 Fritz-Laylin, L.K., Prochnik, S.E., Ginger, M.L., Dacks, J.B., Carpenter,
 2103 M.L., Field, M.C., Kuo, A., Paredez, A., Chapman, J., Pham, J., Shu, S.,
 2104 Neupane, R., Cipriano, M., Mancuso, J., Tu, H.; Salamov, A., Lindquist, E.,
 2105 Shapiro, H., Lucas, S., Grigoriev, I.V., Cande, W.Z., Fulton, C., Rokhsar,
 2106 D.S., Dawson, S.C., 2010. The genome of *Naegleria gruberi* illuminates
 2107 early eukaryotic versatility. *Cell* 140, 631-642.
- 2108 Froggett, S., Leise, E., 1997. Endogenous nitric oxide inhibits
 2109 metamorphosis in a larval mollusc. *Society for Neuroscience - Abstract* 23,
 2110 1234.
- 2111 Fuchs, O., Genuneit, J., Latzin, P., Büchele, G., Horak, E., Loss, G.,
 2112 Sozanska, B., Weber, J., Boznanski, A., Heederik, D., Braun-Fahrlander, C.,
 2113 Frey, U., von Mutius, E., 2012. Farming environments and childhood atopy,
 2114 wheeze, lung function, and exhaled nitric oxide. *J. Allergy Clin. Immun.*
 2115 130, 382-388.

- 2116 Garber, E.A.E., Hollocher, T.C., 1982a. Positional isotopic equivalence
 2117 of nitrogen in N₂O produced by the denitrifying bacterium *Pseudomonas*
 2118 *stutzeri*. The Journal of Biological Chemistry 257, 4705-4708.
- 2119 Garber, E.A.E., Hollocher, T.C., 1982b. ¹⁵N, ¹⁸O tracer studies on the
 2120 activation of nitrite by denitrifying bacteria. Nitrite/water-oxygen exchange
 2121 and nitrosation reactions as indicators of electrophilic catalysis. Journal of
 2122 Biological Chemistry 257, 8091-8097.
- 2123 García-Mata, C., Lamattina, L., 2001. Nitric oxide induces stomatal
 2124 closure and enhances the adaptive plant responses against drought stress.
 2125 Plant Physiology 126, 1196-1204.
- 2126 Gardner, P.R., Gardner, A. M., Martin, L.A., Salzman, A.L., 1998. Nitric
 2127 oxide dioxygenase: an enzymic function for flavohemoglobin. Proceedings
 2128 of National Academy of Science USA 95, 10378–10383.
- 2129 Gardner, A.M., Helmick, R.A., Gardner, P.R., 2002. Flavorubredoxin, an
 2130 inducible catalyst for nitric oxide reduction and detoxification in
 2131 *Escherichia coli*. Journal of Biological Chemistry 277, 8172-8177.
- 2132 Garrido, F., Hénault, C., Gaillard, H., Pérez, S., Germon, J.C., 2002. N₂O
 2133 and NO emissions by agricultural soils with low hydraulic potentials. Soil
 2134 Biology and Biochemistry 34, 559-575.
- 2135 Gas, E., Flores-Pérez, Ú., Sauret-Güeto, S., Rodríguez-Concepción, M.,
 2136 2009. Hunting for plant nitric oxide synthase provides new evidence of a
 2137 central role for plastids in nitric oxide metabolism. The Plant Cell 21, 18-23
- 2138 Gasche, R., Papen, H., 1999. A 3-year continuous record of nitrogen
 2139 trace gas fluxes from untreated and limed soil of a N-saturated spruce and

- 2140 beech forest ecosystem in Germany: 2. NO and NO₂ fluxes, Journal of
 2141 Geophysical Research 104, 18505-18520.
- 2142 Gelperin, A., 1994. Nitric oxide mediates network oscillations of
 2143 olfactory interneurons in a terrestrial mollusc. Nature 369, 61-63.
- 2144 Geßler, A., Rienks, M., Rennenberg, H., 2000. NH₃ and NO₂ fluxes
 2145 between beech trees and the atmosphere – correlation with climatic and
 2146 physiological parameters. New Phytologist 147, 539-560.
- 2147 Geßler, A., Rienks, M., Rennenberg, H., 2002. Stomatal uptake and
 2148 cuticular adsorption contribute to dry deposition of NH₃ and NO₂ to needles
 2149 of adult spruce (*Picea abies*) trees. New Phytologist 156, 179-194.
- 2150 Gilberthorpe, N.J., Poole, R.K., 2008. Nitric Oxide Homeostasis in
 2151 *Salmonella typhimurium* roles of respiratory nitrate reductase and
 2152 flavohemoglobin. Journal of Biological Chemistry 283, 11146-11154.
- 2153 Gladwin, M.T., Kim-Shapiro, D.B., 2008. The functional nitrite
 2154 reductase activity of the heme-globins. Blood 112, 2636-2647.
- 2155 Godde, M., Conrad, R., 2000. Influence of soil properties on the turnover
 2156 of nitric oxide and nitrous oxide by nitrification and denitrification at
 2157 constant temperature and moisture. Biology and Fertility of Soils 32, 120-
 2158 128.
- 2159 Goretski, J., Hollocher, T.C., 1991. Catalysis of nitrosyl transfer by
 2160 denitrifying bacteria is facilitated by nitric oxide. Biochemical and
 2161 Biophysical Research Communications 175, 901-905.
- 2162 Gotoh, T., Mori, M., 2006. Nitric oxide and endoplasmic reticulum
 2163 stress. Arteriosclerosis, Thrombosis and Vascular Biology 26, 1439-1446.

- 2164 Gould, K., Lamotte, O., Klinguer, A., Pugin, A., Wendehenne, D., 2003.
 2165 Nitric oxide production in tobacco leaf cells: a generalized stress response?
 2166 Plant, Cell and Environment 26, 1851-1862.
- 2167 Granfeldt, A., Lefer, D.J., Vinten-Johansen, J., 2009. Protective
 2168 ischaemia in patients: Preconditioning and postconditioning. Cardiovascular
 2169 Research 83, 234-246.
- 2170 Griffith, O.W., Stuehr, D.J., 1995. Nitric oxides synthases – properties
 2171 and catalytic mechanism. Annual Review of Physiology 57, 707-736.
- 2172 Groffman, P.M., Butterbach-Bahl, K., Fulweiler, R.W., Gold, A.J.,
 2173 Morse, J.L., Stander, E.K., Tague, C., Tonitto, C., Vidon, P., 2009.
 2174 Incorporating spatially and temporally explicit phenomena (hotspots and hot
 2175 moments) in denitrification models. Biogeochemistry 93, 49-77.
- 2176 Grün, S., Lindermayr, C., Sell, S., Durner, J., 2006. Nitric oxide and gene
 2177 regulation in plants. Journal of Experimental Botany 57, 507-516.
- 2178 Grune, T., Blasig, I.E., Sitte, N., Roloff, B., Haseloff, R., Davies, K.J.A.,
 2179 1998. Peroxynitrite increases the degradation of aconitase and other cellular
 2180 proteins by proteasome. Journal of Biological Chemistry 273, 10857-10862.
- 2181 Guo, F-Q., Okamoto, M., Crawford, N.M., 2003. Identification of a plant
 2182 nitric oxide synthase gene involved in hormonal signaling. Science 302,
 2183 100-103.
- 2184 Gupta, K.J., Fernie, A.R., Kaiser, W.M., van Dongen, J.T., 2011. On the
 2185 origins of nitric oxide. Trends in Plant Science 16, 160-168.
- 2186 Gupta, K.J., Kaiser, W.M., 2010. Production and scavenging of nitric
 2187 oxide by barley root mitochondria. Plant and Cell Physiology 51, 576-584.

- 2188 Gupta, K.J., Shah, J.K., Brotman, Y., Jahnke, K., Willmitzer, L., Kaiser,
2189 W.M., Bauwe, H., Igamberdiev, A.U., 2012. Inhibition of aconitase by nitric
2190 oxide leads to induction of the alternative oxidase and to a shift of
2191 metabolism towards biosynthesis of amino acids. *Journal of Experimental*
2192 *Botany* 63, 1773-1784.
- 2193 Gusarov, I., Nudler, E., 2005. NO-mediated cytoprotection: instant
2194 adaptation to oxidative stress in bacteria. *Proceedings of the National*
2195 *Academy of Sciences USA* 102, 13855-13860.
- 2196 Gusarov, I., Gautier, L., Smolentseva, O., Shamovsky, I., Eremina, S.,
2197 Mironov, A., Nudler, E. 2013. Bacterial nitric oxide extends the lifespan of
2198 *C. elegans*. *Cell* 152, 818-830.
- 2199 Gusarov, I., Shatalin, K., Starodubtseva, M., Nudler, E. 2009.
2200 Endogenous nitric oxide protects bacteria against a wide spectrum of
2201 antibiotics. *Science* 325, 1380-1384.
- 2202 Gusarov, I., Starodubtseva, M., Wang, Z.Q., McQuade, L., Lippard, S.J.,
2203 Stuehr, D.J., Nudler, E., 2008. Bacterial nitric-oxide synthases operate
2204 without a dedicated redox partner. *Journal of Biological Chemistry* 283,
2205 13140–13147.
- 2206 Hämmerli, A., Waldhuber, S., Miniaci, C., Zeyer, J., Bunge, M., 2007.
2207 Local expansion and selection of soil bacteria in a glacier forefield.
2208 *European Journal of Soil Science* 58, 1437-1445.
- 2209 Hanson, P. J., Lindberg, S. E., 1991. Dry deposition of reactive nitrogen
2210 compounds: a review of leaf, canopy and non-foliar measurements.
2211 *Atmospheric Environment* 25, 1615-1634.

- 2212 Hara, M.R., Agrawal, N., Kim, S.F., Cascio, M.B., Fujimuro, M., Ozeki,
 2213 Y., Takahashi, M., Cheah, J.H., Tankou, S.K., Hester, L.D., Ferris, C.D.,
 2214 Hayward, S.D., Snyder, S.H., Sawa, A., 2005. Snitrosylated GAPDH
 2215 initiates apoptotic cell death by nuclear translocation following Siah1
 2216 binding. *Nature Cell Biology* 7, 665-674.
- 2217 Haroon, M.F., Hu, S., Shi, Y., Imelfort, M., Keller, J., Hugenholtz, P.,
 2218 Yuan, Z., Tyson, G.W., 2013. Anaerobic oxidation of methane coupled to
 2219 nitrate reduction in a novel archaeal lineage. *Nature* 500, 567-570.
- 2220 Hausladen, A., Gow, A., Stamler, J.S., 2001. Flavohemoglobin
 2221 denitrosylase catalyzes the reaction of a nitroxyl equivalent with molecular
 2222 oxygen. *Proceedings of National Academy of Science USA* 98, 10108-
 2223 10112.
- 2224 Hayatsu, M., Tago, K., Saito, M., 2008. Various players in the nitrogen
 2225 cycle: diversity and functions of the microorganisms involved in
 2226 nitrification and denitrification. *Soil Science and Plant Nutrition* 54, 33-45.
- 2227 Hayden, H.L., Drake, J., Imhof, M., Oxley, A.P.A., Norng, S., Mele,
 2228 P.M., 2010. The abundance of nitrogen cycle genes *amoA* and *nifH* depends
 2229 on land-uses and soil types in South-Eastern Australia. *Soil Biology and*
 2230 *Biochemistry* 42, 1774-1783.
- 2231 Hertel, O., Reis, S., Skjoth, C.A., Bleeker, A., Harrison, R., Cape, J.N.,
 2232 Fowler, D., Skiba, U., Simpson, D., Jickells, T., Baker, A., Kulmala, M.,
 2233 Gyldenkaerne, S., Sorensen, L.L., Erisman, J.W., 2011. Nitrogen processes
 2234 in the atmosphere. In: Sutton, M.A., Howard, C.M., Erisman, J.W., Billen,
 2235 G., Bleeker, A., Glennfeld, P., van Grinsven, H., Grizzetti, B. (Eds.), *The*

- 2236 European Nitrogen Assessment , Cambridge University Press, Cambridge,
2237 pp. 177-207.
- 2238 Hietanen, S., Kuparinen, J., 2008. Seasonal and shortterm variation in
2239 denitrification and anammox at a coastal station on the Gulf of Finland,
2240 Baltic Sea. *Hydrobiologia* 596, 67–77.
- 2241 Hong, Y., Yin, B., Zheng, T.L., 2011. Diversity and abundance of
2242 anammox bacterial community in the deep-ocean surface sediment from
2243 equatorial Pacific. *Applied Microbiology and Biotechnology* 89, 1233–
2244 1241.
- 2245 Hooper, A.B., Terry, K.R., 1979. Hydroxylamine oxidoreductase of
2246 nitrosomonase: production of nitric oxide from hydroxylamine. *Biochimica
2247 et Biophysica Acta* 571, 12–20.
- 2248 Hooper, A.B., Vannelli, T., Bergmann, D.J., Arciero, D.M., 1997.
2249 Enzymology of the oxidation of ammonia to nitrite by bacteria. *Antonie van
2250 Leeuwenhoek* 71, 59–67.
- 2251 Horchani, F., Prevot, M., Boscari, A., Evangelisti, E., Meilhoc, E.,
2252 Bruand, C., Raymond, P., Boncompagni, E., Aschi-Smiti, S., Puppo, A.,
2253 Brouquisse, R., 2011. Both plant and bacterial nitrate reductases contribute
2254 to nitric oxide production in *Medicago truncatula* nitrogen-fixing nodules.
2255 *Plant Physiology* 155, 1023-1036.
- 2256 Horii, C.V., Munger, J.W., Wofsy, S.C., Zahniser, M., Nelson, D.,
2257 McManus, J.B., 2004. Fluxes of nitrogen oxides over a temperate deciduous
2258 forest. *Journal of Geophysical Research-Atmospheres* 109, D08305.
- 2259 Hu, B.L., Rush, D., van der Biezen, E., Zheng, P., van Mullekom, M.,
2260 Schouten, S., Damsté, J.S.S, Smolders, A.J., Jetten, M.S.M., Kartal, B.,

- 2261 2011. New anaerobic, ammonium-oxidizing community enriched from peat
 2262 soil. *Applied and Environmental Microbiology* 77, 966–971.
- 2263 Hu, B.L., Shen, L.D., Lian, X., Zhu, Q., Liu, S., Huang, Q., He, Z.-F.,
 2264 Geng, S., Cheng, D.-K., Lou, L.-P., Xu, X.-Y., Zheng, P., He, Y.-F., 2014.
 2265 Evidence for nitrite-dependent anaerobic methane oxidation as a previously
 2266 overlooked microbial methane sink in wetlands. *PNAS* 111, 4495-4500.
- 2267 Hughes, M.N., 1999. Relationships between nitric oxide, nitroxyl ion,
 2268 nitrosonium cation and peroxyxynitrite. *Biochimica et Biophysica Acta*
 2269 (BBA)-Bioenergetics 1411, 263-272.
- 2270 Humbert, S., Tarnawski, S., Fromin, N., Mallet, M.P., Aragno, M., Zopfi,
 2271 J., 2010. Molecular detection of anammox bacteria in terrestrial ecosystems:
 2272 distribution and diversity. *ISME Journal* 4, 450-454.
- 2273 Humbert, S., Zopfi, J., Tarnawski, S.E., 2012. Abundance of anammox
 2274 bacteria in different wetland soils. *Environmental Microbiology Reports* 4,
 2275 484-490.
- 2276 Hunt, J.F., Fang, K., Malik, R., Snyder, A., Malhorta, N., Platts-Mills,
 2277 T.A.E., 2000. Endogenous airway acidification: Implications for asthma
 2278 pathophysiology. *American Journal of Respiratory and Critical Care*
 2279 *Medicine* 161, 694-699.
- 2280 Hunter, T., Bannister, J.V., Hunter, G.J., 2002. Thermostability of
 2281 manganese- and iron-superoxide dismutases from *Escherichia coli* is
 2282 determined by the characteristic position of a glutamine residue. *European*
 2283 *Journal of Biochemistry* 269, 5137-5148.
- 2284 Huygens, D., Rütting, T., Boeckx, P., Van Cleemput, O., Godoy, R.,
 2285 Müller, C., 2007. Soil nitrogen conservation mechanisms in a pristine south

- 2286 Chilean Nothofagus forest ecosystem. *Soil Biology and Biochemistry* 39,
2287 2448-2458.
- 2288 Ientile, R., Picciurro, V., Pedale, S., Nucci, C., Malecka, B., Nisticò, G.,
2289 Macaione, S., 1996. Nitric oxide enhances amino acid release from
2290 immature chick embryo retina. *Neuroscience Letters* 219, 79-82.
- 2291 Igamberdiev, A.U., Hill, R.D., 2009. Plant mitochondrial function during
2292 anaerobiosis. *Annals of Botany* 103, 259-268.
- 2293 Igamberdiev, A.U., Seregelyes, C., Manac, N., Hill, R.D., 2004. NADH-
2294 dependent metabolism of nitric oxide in alfalfa root cultures expressing
2295 barley hemoglobin. *Planta* 219, 95-102.
- 2296 Ignarro, L.J. (Ed.), 2009. *Nitric Oxide: Biology and Pathology*, second
2297 ed. Elsevier Academic Press, London, 845 pp.
- 2298 Ilari, A., Boffi, A., 2008. Structural studies on flavohemoglobins.
2299 *Methods in Enzymology* 436, 187-202.
- 2300 Immoos, C.E., Chou, J., Bayachou, M., Blair, E., Greaves, J., Farmer,
2301 P.J., 2004. Electrocatalytic reductions of nitrite, nitric oxide, and nitrous
2302 oxide by thermophilic cytochrome P450CYP119 in film-modified
2303 electrodes and an analytical comparison of its catalytic activities with
2304 myoglobin. *Journal of the American Chemical Society* 126, 4934-4942.
- 2305 IPCC, 2007. Agriculture, forestry and other land use. In: Eggleston, H.S.,
2306 Buendia, L., Miwa, K., Ngara, T., Tanabe, K. (Eds.), 2006 IPCC guidelines
2307 for national greenhouse gas inventories, IGES, Hayama.
- 2308 Islam, A., Chen, D., White, R.E., Weatherley, A.J., 2008. Chemical
2309 decomposition and fixation of nitrite in acidic pasture soils and implications

2310 for measurement of nitrification. *Soil Biology and Biochemistry* 40, 262-
2311 265.

2312 Jacklet, J.W., 1997. Nitric oxide signaling in invertebrates. *Invertebrate*
2313 *Neuroscience* 3, 1-14.

2314 Jacklet, J., Gruhn, M., 1994. Co-localization of NADPHdiaphorase and
2315 myomodulin in synaptic glomeruli of *Aplysia*. *NeuroReport* 5, 1841-1844.

2316 Jaeschke, A., den Camp, H.J.M.O., Harhangi, H., Klimiuk, A., Hopmans,
2317 E.C., Jetten, M.S.M., Schouten, S., Damste, J.S.S., 2009. 16S rRNA gene
2318 and lipid biomarker evidence for anaerobic ammonium-oxidizing bacteria
2319 (anammox) in California and Nevada hot springs. *FEMS Microbiology and*
2320 *Ecology* 67, 343–350.

2321 Jansson, E., Lindblad, P., 1998. Cloning and molecular characterization
2322 of a presumptive *argF*, a structural gene encoding ornithine carbamoyl
2323 transferase (OCT), in the cyanobacterium *Nostoc* sp. PCC 73102.
2324 *Physiologia Plantarum* 103, 347-353.

2325 Jansson, E.Å., Huang, L., Malkey, R., Govoni, M., Nihlén, C., Olsson,
2326 A., Stensdotter, M., Petersson, J., Holm, L., Weitzberg, E., Lundberg, J.O.,
2327 2008. A mammalian functional nitrate reductase that regulates nitrite and
2328 nitric oxide homeostasis. *Nature Chemical Biology* 4, 411-417.

2329 Jetten, M.S.M., de Bruijn, P., Kuenen, J.G., 1997. Hydroxylamine
2330 metabolism in *Pseudomonas* PB16: involvement of a novel hydroxylamine
2331 oxidoreductase. *Antonie Van Leeuwenhoek* 71, 69–74.

2332 Jetten, M.S.M., Sliekers, O., Kuypers, M., Dalsgaard, T., van Niftrik, L.,
2333 Cirpus, I., van de Pas-Schoonen, K., Lavik, G., Thamdrup, B., Le Paslier,
2334 D., Op den Camp, H.J.M., Hulth, S., Nielsen, L.P., Abma, W., Third, K.,

- 2335 Engström, P., Kuenen, J.G., Jorgensen, B.B., Canfield, D.E., Damsté, J.S.S.,
 2336 Revsbech, N.P., Fuerst, J., Weissenbach, J., Wagner, M., Schmidt, I.,
 2337 Schmid, M., Strous, M., 2003. Anaerobic ammonium oxidation by marine
 2338 and freshwater planctomycete-like bacteria. *Applied Microbiology and*
 2339 *Biotechnology* 63, 107–114.
- 2340 Johnson, E.G., Sparks, J.P., Dzikovski, B., Crane, B.R., Gibson, D.M.,
 2341 Loria, R., 2008. Plant-pathogenic *Streptomyces* species produce nitric oxide
 2342 synthase-derived nitric oxide in response to host signals. *Chemistry and*
 2343 *Biology* 15, 43-50.
- 2344 Ju, X.T., Lu, X., Gao, Z.L., Chen, X.P., Su, F., Kogge, M., Römheld, V.,
 2345 Christie, P., Zhang, F., 2011. Processes and factors controlling N₂O
 2346 production in an intensively managed low carbon calcareous soil under sub-
 2347 humid monsoon conditions. *Environmental Pollution* 159, 1007-1016.
- 2348 Jung, M.Y., Park, S.J., Kim, S.J., Kim, J.G., Damsté, J.S.S., Jeon, C.O.,
 2349 Rhee, S.K., 2014. A mesophilic, autotrophic, ammonia-oxidizing archaeon
 2350 of *Thaumarchaeal* group I. 1a cultivated from a deep oligotrophic soil
 2351 horizon. *AEM* 80, 3645-3655.
- 2352 Kampman, C., Hendrickx, T.L.G., Luesken, F.A., van Alen, T.A., Op
 2353 den Camp, H.J.M., Jetten, M.S.M., Zeeman, G., Buisman, C.J.N., Temmink,
 2354 H., 2012. Enrichment of denitrifying methanotrophic bacteria for
 2355 application after direct low temperature anaerobic sewage treatment. *Journal*
 2356 *of Hazardous Materials* 227, 164–171.
- 2357 Kandeler, E., Deiglmayr, K., Tscherko, D., Bru, D., Philippot, L., 2006.
 2358 Abundance of *narG*, *nirS*, *nirK*, and *nosZ* genes of denitrifying bacteria

2359 during primary successions of a glacier foreland. Applied and
 2360 Environmental Microbiology 72, 5957-5962.

2361 Kartal, B., Almeida, N.M., Maalcke, W.J., Camp, H.J., Jetten, M.S.,
 2362 Keltjens, J.T., 2013. How to make a living from anaerobic ammonium
 2363 oxidation. FEMS Microbiology Reviews 37, 428-461.

2364 Kartal, B., Kuypers, M.M.M., Lavik, G., Schalk, J., Op den Camp,
 2365 H.J.M., Jetton, M.S.M., Strous, M., 2007a. Anammox bacteria disguised as
 2366 denitrifiers: nitrate reduction to dinitrogen gas via nitrite and ammonium.
 2367 Environmental Microbiology 9, 635–642.

2368 Kartal, B., Maalcke, W.J., de Almeida, N.M., Cirpus, I., Gloerich, J.,
 2369 Geerts, W., Op den Camp¹, H.J.M., Harhangi, H.R., Janssen-Megens, E.M.,
 2370 Francoijs, K.-J., Stunnenberg, H.G., Keltjens, J.T., Jetton, M.S.M., Strous,
 2371 M., 2011. Molecular mechanism of anaerobic ammonium oxidation. Nature
 2372 479, 127-130.

2373 Kartal, B., Rattray, J., van Niftrik, L.A., van de Vossenberg, J., Schmid,
 2374 M. C., Webb, R.I., Schouten, S., Fuerst, J.A., Damsté, J.S.S, Jetten, M.S.M.,
 2375 Strous, M., 2007b. *Candidatus "Anammoxoglobus propionicus"* a new
 2376 propionate oxidizing species of anaerobic ammonium oxidizing bacteria.
 2377 Systematic and Applied Microbiology 30: 39–49.

2378 Kartal, B., Tan, N.C., Van de Biezen, E., Kampschreur, M.J., Van
 2379 Loosdrecht, M.C., Jetten, M.S., 2010. Effect of nitric oxide on anammox
 2380 bacteria. Applied and Environmental Microbiology 76, 6304-6306.

2381 Kartal, B., van Niftrik, L., Rattray, J., van de Vossenberg, J.L., Schmid,
 2382 M.C., Damsté, J.S.S, Jetton, M.S.M., Strous, M., 2008. *Candidatus*

- 2383 *'Brocadia fulgida'*: an autofluorescent anaerobic ammonium oxidizing
 2384 bacterium. FEMS Microbiology and Ecology 63, 46–55.
- 2385 Kendrick, K., Guevara-Guzzman, R., Zorritla, J., Hinton, M., Broad, K.,
 2386 Mimmiack, M., Ohkura, S., 1997. Formation of olfactory memories
 2387 mediated by nitric oxide. Nature 388, 670- 674.
- 2388 Kesik, M., Ambus, P., Baritz, R., Brüggemann, N., Butterbach-Bahl, K.,
 2389 Damm, M., Duyzer, J., Horváth, L., Kiese, R., Kitzler, B., Leip, A., Li, C.,
 2390 Pihlatie, M., Pilegaard, K., Seufert, G., Simpson, D., Skiba, U., Smiatek, G.,
 2391 Vesala, T., Zechmeister-Boltenstern, S. 2005. Inventories of N₂O and NO
 2392 emissions from European forest soils. Biogeosciences 2, 353-375.
- 2393 Kesik, M., Blagodatsky, S., Papen, H., Butterbach-Bahl, K., 2006. Effect
 2394 of pH, temperature and substrate on N₂O, NO and CO₂ production by
 2395 *Alcaligenes faecalis* p. Journal of Applied Microbiology 101, 655 -667.
- 2396 Khanna-Chopra, R., Sabarinath, S., 2004. Heat-stable chloroplastic
 2397 Cu/Zn superoxide dismutase in *Chenopodium murale*. Biochemical and
 2398 Biophysical Research Communications 320, 1187-1192.
- 2399 Kharitonov, S.A., Chung, K.F., Evans, D., O'Connor, B.J., Barnes, P.J.,
 2400 1996. Increased exhaled nitric oxide in asthma is mainly derived from lower
 2401 respiratory tract. American Journal of Respiratory and Critical Care
 2402 Medicine 153, 1773-1780.
- 2403 Killham, K., 1990. Nitrification in coniferous forest soils. Plant and Soil
 2404 128, 31–44.
- 2405 Kim, C.H., Hollocher, T.C., 1984. Catalysis of nitrosyl transfer-reactions
 2406 by a dissimilatory nitrite reductase (cytochrome-cd1). Journal of Biological
 2407 Chemistry 259, 2092-2099.

- 2408 Kim, D.G., Vargas, R., Bond-Lamberty, B., Turetsky, M.R., 2012.
2409 Effects of soil rewetting and thawing on soil gas fluxes: a review of current
2410 literature and suggestions for future research. *Biogeosciences* 9, 2459–2483.
- 2411 Kim, E.K., Kochi, J.K., 1991. Charge-transfer structures of aromatic
2412 electron donor-acceptor complexes leading to electron transfer with the
2413 electrophilic nitrosonium cation. *Journal of the American Chemical Society*
2414 113, 4962-4974.
- 2415 Kim, S.O., Oori, Y., Lloyd, D., Hughes, M.N., Poole, R.K., 1999. Anoxic
2416 function for the *Escherichia coli* flavohaemoglobin (Hmp): reversible
2417 binding of nitric oxide and reduction to nitrous oxide. *FEBS Letters* 445,
2418 389-394.
- 2419 Kitzler, B., Zechmeister-Boltenstern, S., Holtermann, C., Skiba, U.,
2420 Butterbach-Bahl, K., 2006. Controls over N₂O, NO_x and CO₂ fluxes in a
2421 calcareous mountain forest soil. *Biogeosciences* 3, 383-395.
- 2422 Knowles, R., 1982. Denitrification. *Microbiological Reviews* 46, 43-70.
- 2423 Kobayashi, M., Matsuo, Y., Takimoto, A., Suzuki, S., Maruo, F., Shoun,
2424 H., 1996. Denitrification, a novel type of respiratory metabolism in fungal
2425 mitochondrion. *Journal of Biological Chemistry* 271, 16263–16267.
- 2426 Könneke, M., Bernhard, A.E., de la Torre, J.R., Walker, C.B.,
2427 Waterbury, J.B., Stahl, D.A., 2005. Isolation of an autotrophic ammonia-
2428 oxidizing marine archaeon. *Nature* 437, 543-546.
- 2429 Kool, D.M., Müller, C., Wrage, N., Oenema, O., Van Groenigen, J.W.,
2430 2009a. Oxygen exchange between nitrogen oxides and H₂O can occur
2431 during nitrifier pathways. *Soil Biology and Biochemistry* 41, 1632-1641.

- 2432 Kool, D.M., Wrage, N., Oenema, O., Harris, D., Van Groenigen, J.W.,
 2433 2009b. The ^{18}O signature of biogenic nitrous oxide is determined by O
 2434 exchange with water. *Rapid Communications in Mass Spectrometry* 23,
 2435 104-108.
- 2436 Kool, D.M., Wrage, N., Zechmeister-Boltenstern, S., Pfeffer, M., Brus,
 2437 D., Oenema, O., Van Groenigen, J.W., 2010. Nitrifier denitrification can be
 2438 a source of N_2O from soil: a revised approach to the dual-isotope labelling
 2439 method. *European Journal of Soil Science* 61, 759-772.
- 2440 Koop-Jakobsen, K., Giblin, A., 2009. Anammox in tidal marsh
 2441 sediments: The role of salinity, nitrogen loading, and marsh vegetation.
 2442 *Estuaries and Coasts* 32, 238–245.
- 2443 Koops, H.P., Bottcher, B., Moller, U.C., Pommerening-Roser, A., Stehr,
 2444 G., 1991. Classification of eight new species of ammonia-oxidizing
 2445 bacteria: *Nitrosomonas communis* sp nov., *Nitrosomonas ureae* sp nov.,
 2446 *Nitrosomonas aestuarii* sp nov., *Nitrosomonas marina* sp nov.,
 2447 *Nitrosomonas nitrosa* sp nov., *Nitrosomonas oligotropha* sp nov.,
 2448 *Nitrosomonas halophila* sp nov. *Journal of General Microbiology* 137,
 2449 1689-1699.
- 2450 Koper, T.E., Habteselassie, M.Y., Stark, J.M., Norton, J.M., 2010.
 2451 Nitrification exhibits Haldane kinetics in an agricultural soil treated with
 2452 ammonium sulfate or dairy waste compost. *FEMS Microbiology Ecology*
 2453 74, 316-322.
- 2454 Koschorreck, M., Conrad, R., 1997. Kinetics of nitric oxide consumption
 2455 in tropical soils under oxic and anoxic conditions. *Biology and Fertility of*
 2456 *Soils* 25, 82-88.

- 2457 Koschorreck, M., Moore, E., Conrad, R., 1996. Oxidation of nitric oxide
2458 by a new heterotrophic *Pseudomonas* sp. Archives of Microbiology 166, 23-
2459 31.
- 2460 Koul, V., Adholeya, A., Kochar, M., 2014. Sphere of influence of indole
2461 acetic acid and nitric oxide in bacteria. Journal of Basic Microbiology 54, 1-
2462 11.
- 2463 Kroncke, K.D., Fehsel, K., Kolb-Bachofen, V., 1997. Nitric oxide:
2464 Cytotoxicity versus cytoprotection – how, why, when, and where? Nitric
2465 Oxide 1, 107-120.
- 2466 Kuenen, J.G., 2008. Anammox bacteria: from discovery to application.
2467 Nature Reviews Microbiology 6, 320–326.
- 2468 Kuenen, J.G., Robertson, L.A., 1994. Combined nitrification-
2469 denitrification processes. FEMS Microbiology Reviews 15, 109-117.
- 2470 Kumon, Y., Sasaki, Y., Kato, I., Takaya, N., Shoun, H., Beppu, T., 2002.
2471 Codenitrification and denitrification are dual metabolic pathways through
2472 which dinitrogen evolves from nitrate in *Streptomyces antibioticus*. Journal
2473 of Bacteriology 184, 2963-2968.
- 2474 Kuypers, M.M., Lavik, G., Woebken, D., Schmid, M., Fuchs, B.M.,
2475 Amann, R., Jorgensen, B.B., Jetton, M.S., 2005. Massive nitrogen loss from
2476 the Benguela upwelling system through anaerobic ammonium oxidation.
2477 Proceedings of National Academy of Science USA 102, 6478–6483.
- 2478 Kuypers, M.M.M., Sliekers, A.O., Lavik, G., Schmid, M.C., Jørgensen,
2479 B.B., Kuenen, J.G., Damsté, J.S.S., Strous, M., Jetton, M.S.M., 2003.
2480 Anaerobic ammonium oxidation by anammox bacteria in the Black Sea.
2481 Nature 422, 608–611.

- 2482 Kuzin, B., Roberts, I., Peunova, N. Enikolopov, G., 1996. Nitric oxide
2483 regulates cell proliferation during *Drosophila* development. Cell 87, 639-
2484 649.
- 2485 Lam, P., Jensen, M.M., Lavik, G., McGinnis, D.F., Muller, B., Schubert,
2486 C.J., Amann, R., Thamdrup, B., Kuypers, M.M., 2007. Linking crenarchaeal
2487 and bacterial nitrification to anammox in the Black Sea. Proceedings of
2488 National Academy of Science USA 104, 7104–7109.
- 2489 Lamattina, L., García-Mata, C., Graziano, M., Pagnussat, G., 2003. Nitric
2490 oxide: the versatility of an extensive signal molecule. Annual Review of
2491 Plant Biology 54, 109-136.
- 2492 Lange, M., Enkhbaatar, P., Nakano, Y., Traber, D.L., 2009. Role of nitric
2493 oxide in shock: The large animal perspective. Frontiers in Bioscience 14,
2494 1979-1989.
- 2495 Laudelout, H., 1978. Mathematical modeling of biological ammonium
2496 oxidation. In: Schlessinger, D. (Ed.), Microbiology, American Society of
2497 Microbiology, Washington DC, 384-386.
- 2498 Laughlin, R.J., Stevens, R.J., 2002. Evidence for fungal dominance of
2499 denitrification and codenitrification in a grassland soil. Soil Science Society
2500 of America Journal 66, 1540-1548.
- 2501 Laville, P., Flura, D., Gabrielle, B., Loubet, B., Fanucci, O., Rolland, M.-
2502 N., Cellier, P., 2009. Characterisation of soil emissions of nitric oxide at
2503 field and laboratory scale using high resolution method. Atmospheric
2504 Environment 43, 2648-2658.
- 2505 Laville, P., Lehuger, S., Loubet, B., Chaumartin, F., Cellier, P., 2011.
2506 Effect of management, climate and soil conditions on N₂O and NO

- 2507 emissions from an arable crop rotation using high temporal resolution
2508 measurements. *Agricultural and Forest Meteorology* 151, 228–240.
- 2509 Lazzaro, A., Abegg, C., Zeyer, J., 2009. Bacterial community structure of
2510 glacier forefields on siliceous and calcareous bedrock. *European Journal of*
2511 *Soil Science* 60, 860-870.
- 2512 Lee, J., Chen, L., West, A.H., Richter-Addo, G.B., 2002. Interactions of
2513 organic nitroso compounds with metals. *Chemical Reviews* 102, 1019-1066.
- 2514 Lee, J.H., Yang, E.S., Park, J.W., 2003. Inactivation of NADP⁺-
2515 dependent isocitrate dehydrogenase by peroxynitrite. Implications for
2516 cytotoxicity and alcohol-induced liver injury. *Journal of Biological*
2517 *Chemistry* 278, 51360-51371.
- 2518 Lee, K.Y., Kuchynka, D.J., Kochi, J.K., 1990. Redox equilibria of the
2519 nitrosonium cation and of its nonbonded complexes. *Inorganic Chemistry*
2520 29, 4196-4204.
- 2521 Lee, U., Wie, C., Fernandez, B.O., Feelisch, M., Vierling, E., 2008.
2522 Modulation of nitrosative stress by S-nitrosoglutathione reductase is critical
2523 for thermotolerance and plant growth in *Arabidopsis*. *The Plant Cell* 20,
2524 786-802.
- 2525 Lees, H., 1952. Hydroxylamine as an intermediate in nitrification. *Nature*
2526 169, 156-157.
- 2527 Lehtovirta-Morley, L.E., Stoecker, K., Vilcinskas, A., Prosser, J.I., Nicol,
2528 G.W., 2011. Cultivation of an obligate acidophilic ammonia oxidizer from a
2529 nitrifying acid soil. *PNAS*, 108, 15892-15897.

- 2530 Leininger, S., Urich, T., Schloter, M., Schwark, L., Qi, J., Nicol, G.W.,
 2531 Prosser, J.I., Schuster, S.C., Schleper, C., 2006. Archaea predominate
 2532 among ammonia-oxidizing prokaryotes in soils. *Nature* 442, 806-809.
- 2533 Leitner, M., Vandelle, E., Gaupels, F., Bellin, D., Delledonne, M., 2009.
 2534 NO signals in the haze: nitric oxide signalling in plant defence. *Current*
 2535 *Opinion in Plant Biology* 12, 451-458.
- 2536 Leshem, Ya.Y., Wills, R.B.H., Ku, V.V.-V., 1998. Evidence for the
 2537 function of the free radical gas - nitric oxide (NO^{*}) - as an endogenous
 2538 maturation and senescence regulating factor in higher plants. *Plant*
 2539 *Physiology and Biochemistry* 36, 825-833.
- 2540 Levitzky, M.G., 2003. *Pulmonary Physiology*. McGraw Hill
 2541 Professional, New York, 278 pp.
- 2542 Li, H., Forstermann, U., 2000. Nitric oxide in the pathogenesis of
 2543 vascular disease. *The Journal of Pathology* 190, 244-254.
- 2544 Li, H., Kundu, T.K., Zweier, J.L., 2009. Characterization of the
 2545 magnitude and mechanism of aldehyde oxidase-mediated nitric oxide
 2546 production from nitrite. *Journal of Biological Chemistry* 284, 33850–33858.
- 2547 Li, H., Chen, S., Mu, B.Z., Gu, J.-D., 2010. Molecular detection of
 2548 anaerobic ammonium-oxidizing (anammox) bacteria in high-temperature
 2549 petroleum reservoirs. *Microbiology and Ecology* 60, 771–783.
- 2550 Li, M, Cao, H., Hong, Y., Gu, J.-D., 2011a. Seasonal dynamics of
 2551 anammox bacteria in estuarial sediment of the Mai Po Nature Reserve
 2552 revealed by analyzing the 16S rRNA and hydrazine oxidoreductase (hzo)
 2553 genes. *Microbes and Environments* 26, 5–22.

- 2554 Li, X., Jankovic, J., Le, W., 2011b. Iron chelation and neuroprotection in
2555 neurodegenerative diseases. *Journal of Neural Transmission* 118, 473-477.
- 2556 Lindsay, M.R., Webb, R.I., Strous, M., Jetton, M.S.M., Butler, M.K.,
2557 Forde, R.J., Fuerst, J.A., 2001. Cell compartmentalisation in
2558 planctomycetes: novel types of structural organisation for the bacterial cell.
2559 *Archives of Microbiology* 175, 413–429.
- 2560 Liu, S., Vereecken, H., Brüggemann, N., 2014. A highly sensitive
2561 method for the determination of hydroxylamine in soils. *Geoderma* 232,
2562 117-122.
- 2563 Liu, Y., Wu, R., Wan, Q., Xie, G., Bi, Y., 2007. Glucose-6-phosphate
2564 dehydrogenase plays a pivotal role in nitric oxide-involved defense against
2565 oxidative stress under salt stress in red kidney bean roots. *Plant and Cell*
2566 *Physiology* 48, 511-522.
- 2567 Lloyd, D., 1993. Aerobic denitrification in soils and sediments: From
2568 fallacies to facts. *Trends in Ecology and Evolution* 8, 352-356.
- 2569 Long, A., Heitman, J., Tobias, C., Philips, R., Song, B., 2013. Co-
2570 occurring anammox, denitrification, and codenitrification in agricultural
2571 soils. *Applied and Environmental Microbiology* 79, 168-176.
- 2572 Luesken, F.A., van Alen, T.A., van der Biezen, E., Frijters, C., Toonen,
2573 G., Kampman, C., Hendrickx, T.L.G., Zeeman, G., Temmink, H., Strous,
2574 M., Op den Camp, H.J.M., Jetten, M.S.M., 2011. Diversity and enrichment
2575 of nitrite-dependent anaerobic methane oxidizing bacteria from waste water
2576 sludge. *Applied Microbiology and Biotechnology* 92, 845–854.
- 2577 Ludwig, J., Meixner, F.X., 1994. Surface exchange of nitric oxide (NO)
2578 over three European ecosystems. In: Angeletti, G., Restelli, G., (Eds.),

- 2579 Proceedings of the Sixth European Symposium on the Physico-Chemical
 2580 Behaviour of Atmospheric Pollutants, Commission of the European
 2581 Communities, Luxembourg, pp. 587-593.
- 2582 Ludwig, J., Meixner, F.X., Vogel, B., Förstner, J., 2001. Soil-air
 2583 exchange of nitric oxide: an overview of processes, environmental factors,
 2584 and modeling studies. *Biogeochemistry* 52, 225 -257.
- 2585 Lum, H.K., Butt, Y.K.C., Lo, S.C.L., 2002. Hydrogen peroxide induces a
 2586 rapid production of nitric oxide in mung bean (*Phaseolus aureus*). *Nitric*
 2587 *Oxide* 6, 205-213.
- 2588 Lundberg, J.O., Weitzberg, E., Gladwin, M.T., 2008. The nitrate–nitrite–
 2589 nitric oxide pathway in physiology and therapeutics. *Nature Reviews Drug*
 2590 *Discovery* 7, 156-167.
- 2591 Lundberg, J.O., Weitzberg, E.J., Lundberg, M., Alving, K., 1994.
 2592 Intragastric nitric oxide production in humans: Measurements in expelled
 2593 air. *Gut* 35, 1543-1546.
- 2594 Luo, G.J., Brüggemann N., Wolf, B., Gasche, R., Grote, R., Butterbach-
 2595 Bahl, K., 2012. Decadal variability of soil CO₂, NO, N₂O, and CH₄ fluxes at
 2596 the Höglwald Forest, Germany. *Biogeosciences* 9, 1741-1763.
- 2597 Machacova, K., Papen, H., Kreuzwieser, J., Rennenberg, H., 2013.
 2598 Inundation strongly stimulates methane and nitrous oxide emissions from
 2599 stems of the upland tree *Fagus sylvatica*. *Plant and Soil* 364, 287-301.
- 2600 Machefert, S.E., Dise, N.B., Goulding, K.W.T., Whitehead, P.G., 2002.
 2601 Nitrous oxide emission from a range of land uses across Europe. *Hydrology*
 2602 *and Earth System Sciences* 6, 325-337.

- 2603 Mackerness, S.A.H., John, C.F., Jordan, B., Thomas, B., 2001. Early
2604 signalling components in ultraviolet- B responses: distinct roles for different
2605 reactive oxygen species and nitric oxide. *FEBS Letters* 489, 237-242.
- 2606 MacMicking, J., Xie, Q.W., Nathan, C., 1997. Nitric oxide and
2607 macrophage function. *Annual Review of Immunology* 15, 323-350.
- 2608 Maggesissi, R., Gardino, P., Guimarães-Souza, E., Paes-de- Carvalho, R.,
2609 Silva, R., Calaza, K., 2009. Modulation of GABA release by nitric oxide in
2610 the chick retina: different effects of nitric oxide depending on the cell
2611 population. *Vision Research* 49, 2494-2502.
- 2612 Martens-Habbena, W., Berube, P.M., Urakawa, H., de la Torre, J.R.,
2613 Stahl, D.A., 2009. Ammonia oxidation kinetics determine niche separation
2614 of nitrifying Archaea and Bacteria. *Nature* 461, 976-979.
- 2615 Martin, E., Berka, V., Sharina, I., Tsai, A.L., 2012. Mechanism of
2616 binding of NO to soluble guanylyl cyclase: implication for the second NO
2617 binding to the heme proximal site. *Biochemistry* 51, 2737-2746.
- 2618 Martin, R.E., Scholes, M.C., Mosier, A.R., Ojima, D.S., Holland, E.A.,
2619 Parton, W.J., 1998. Controls on annual emissions of nitric oxide from soils
2620 of the Colorado shortgrass steppe. *Global Biogeochemical Cycles* 12, 81-91.
- 2621 McCleverty, J.A., 2004. Chemistry of nitric oxide relevant to biology.
2622 *Chemical Reviews* 104, 403-418.
- 2623 Meakin, G.E., Bueno, E., Jepson, B., Bedmar, E.J., Richardson, D.J.,
2624 Delgado, M.J., 2007. The contribution of bacteroidal nitrate and nitrite
2625 reduction to the formation of nitrosylhaemoglobin complexes in soybean
2626 root nodules. *Microbiology* 153, 411-419.

- 2627 Meleshkevitch, E., Budko, D., Norby, S., Moroz, L., Hadfield, M., 1997.
 2628 Nitric oxide dependent modulation of the metamorphosis in mollusc
 2629 *Peeteilla sibogae* (gastropoda, nudibranchia). Society for Neuroscience -
 2630 Abstract 23, 1233.
- 2631 Melotto, M., Underwood, W., Koczan, J., Nomura, K., He, S.Y., 2006.
 2632 Plant stomata function in innate immunity against bacterial invasion. Cell
 2633 126, 969-980.
- 2634 Mei, B., Zheng, X., Xie, B., Dong, H., Yao, Z., Liu, C., Zhou, Z., Wang,
 2635 R., Deng, J., Zhu, J., 2011. Characteristics of multiple-year nitrous oxide
 2636 emissions from conventional vegetable fields in southeastern China. Journal
 2637 of Geophysical Research 116, 012011.
- 2638 Meilhoc, E., Boscari, A., Bruand, C., Puppo, A., Brouquisse, R., 2011.
 2639 Nitric oxide in legume–rhizobium symbiosis. Plant Science 181, 573-581.
- 2640 Messner, S., Leitner, S., Bommassar, C., Golderer, G., Grobner, P.,
 2641 Werner, E. R., Werner-Felmayer, G., 2009. Physarum nitric oxide
 2642 synthases: genomic structures and enzymology of recombinant proteins.
 2643 Biochemical Journal 418, 691-700.
- 2644 Mikkelsen, R.B., Wardman, P., 2003. Biological chemistry of reactive
 2645 oxygen and nitrogen and radiation-induced signal transduction mechanisms.
 2646 Oncogene 22, 5734-5754.
- 2647 Miller, A.F., 2012. Superoxide dismutases: ancient enzymes and new
 2648 insights. FEBS Letters 586, 585-595.
- 2649 Mills, P.C., Richardson, D.J., Hinton, J.C.D., Spiro, S., 2005.
 2650 Detoxification of nitric oxide by the flavorubredoxin of *Salmonella enterica*
 2651 serovar *Typhimurium*. Biochemical Society Transactions 33, 198-199.

- 2652 Mills, P.C., Rowley, G., Spiro, S., Hinton, J.C.D., Richardson, D.J.,
2653 2008. A combination of cytochrome c nitrite reductase (NrfA) and
2654 flavorubredoxin (NorV) protects *Salmonella enterica* serovar *Typhimurium*
2655 against killing by NO in anoxic environments. *Microbiology* 154, 1218-
2656 1228.
- 2657 Moir, J.W.B., Crossman, L.C., Spiro, S., Richardson, D.J., 1996. The
2658 purification of ammonia mono-oxygenase from *Paracoccus denitrificans*.
2659 *FEBS Letters* 387, 71-74.
- 2660 Moreau, M., Lindermayr, C., Durner, J., Klessig, D.F., 2010. NO
2661 synthesis and signaling in plants – where do we stand? *Physiologia*
2662 *Plantarum* 138, 372-383.
- 2663 Moreno-Vivián, C., Cabello, P., Martínez-Luque, M., Blasco, R.,
2664 Castillo, F., 1999. Prokaryotic nitrate reduction: molecular properties and
2665 functional distinction among bacterial nitrate reductases. *Journal of*
2666 *Bacteriology* 181, 6573-6584.
- 2667 Mørkved, P.T., Dörsch, P., Henriksen, T.M., Bakken, L.R., 2006. N₂O
2668 emissions and product ratios of nitrification and denitrification as affected
2669 by freezing and thawing. *Soil Biology and Biochemistry* 38, 3411-3420.
- 2670 Mühlig, A., Kabisch, J., Pichner, R., Scherer, S., Müller-Herbst, S., 2014.
2671 Contribution of the NO-detoxifying enzymes HmpA, NorV and NrfA to
2672 nitrosative stress protection of *Salmonella Typhimurium* in raw sausages.
2673 *Food Microbiology* 42, 26-33.
- 2674 Mulder, A., van de Graaf, A.A., Robertson, L.A., Kuenen, J.G., 1995.
2675 Anaerobic ammonium oxidation discovered in a denitrifying fluidized bed
2676 reactor. *FEMS Microbiology and Ecology* 16, 177–184.

- 2677 Müller, U., 1997. The nitric oxide system in insects. Progress in
2678 Neurobiology 51, 363-381.
- 2679 Mur, L.A., Sivakumaran, A., Mandon, J., Cristescu, S.M., Harren, F.J.,
2680 Hebelstrup, K.H., 2012. Haemoglobin modulates salicylate and
2681 jasmonate/ethylene-mediated resistance mechanisms against pathogens.
2682 Journal of Experimental Botany 63, 4375-4387.
- 2683 Murphy, M.E., Sies, H., 1991. Reversible conversion of nitroxyl anion to
2684 nitric oxide by superoxide dismutase. Proceedings of the National Academy
2685 of Sciences USA 88, 10860-10864.
- 2686 Myers, R.J.K., 1975. Temperature effects on ammonification and
2687 nitrification in a tropical soil. Soil Biology and Biochemistry 7, 83-86.
- 2688 Nagele, W., Conrad, R., 1990. Influence of pH on the release of NO and
2689 N₂O from fertilized and unfertilized soil. Biology and Fertility of Soils 10,
2690 139-144.
- 2691 Nakajima, Y., Ishizuka, S., Tsuruta, H., Iswandi, A., Murdiyarso, D.,
2692 2005. Microbial processes responsible for nitrous oxide production from
2693 acid soils in different land-use patterns in Pasirmayang, central Sumatra,
2694 Indonesia. Nutrient Cycling in Agroecosystems 71, 33-42.
- 2695 Natalis, P., Collin, J.E., Delwiche, J., Caprace, G., Hubin, M.J., 1979.
2696 Ionization energy values for the transitions $\text{NO}^+, X^1\Sigma^+(v'=0-34) \leftarrow \text{NO}, X^2$
2697 $\Pi(v''=0)$ and molecular constants of nitric oxide ground ionic state,
2698 determined by Ne(I) (73.6 nm) photoelectron spectroscopy. Journal of
2699 Electron Spectroscopy and Related Phenomena 17, 421-423.

- 2700 Nathan, C.F., Hibbs, J.B. Jr., 1991. Role of nitric oxide synthesis in
2701 macrophage antimicrobial activity. *Current Opinion in Immunology* 3, 65-
2702 70.
- 2703 Neill, S., Barros, R., Bright, J., Desikan, R., Hancock, J., Harrison, J.,
2704 Morris, P., Ribeiro, D., Wilson, I., 2008. Nitric oxide, stomatal closure, and
2705 abiotic stress. *Journal of Experimental Botany* 59, 165-176.
- 2706 Neill, S.J., Desikan, R., Clarke, A., Hancock, J.T., 2002. Nitric oxide is a
2707 novel component of abscisic acid signaling in stomatal guard cells. *Plant*
2708 *Physiology* 128, 13-16.
- 2709 Neill, S.J., Desikan, R., Hancock, J.T., 2003. Nitric oxide signalling in
2710 plants. *New Phytologist* 159, 11-35.
- 2711 Neubert, A., Kley, D., Wildt, J., Segschneider, H.J., Förstel, H., 1993.
2712 Uptake of NO, NO₂ and O₃ by sunflower (*Helianthus annuus* L.) and
2713 tobacco plants (*Nicotiana tabacum* L.): dependence on stomatal
2714 conductivity. *Atmospheric Environment* 27, 2137-2145.
- 2715 Nicol, G.W., Leininger, S., Schleper, C., Prosser, J.I., 2008. The
2716 influence of soil pH on the diversity, abundance and transcriptional activity
2717 of ammonia oxidizing archaea and bacteria. *Environmental Microbiology*
2718 10, 2966-2978.
- 2719 Nicol, G.W., Tscherko, D., Embley, T.M., Prosser, J.I., 2005. Primary
2720 succession of soil *Crenarchaeota* across a receding glacier foreland.
2721 *Environmental Microbiology* 7, 337-347.
- 2722 Nishio, T., Yoshikura, T., Mishima, H., Inouye, Z., Itoh, H., 1998.
2723 Conditions for nitrification and denitrification by an immobilized

- 2724 heterotrophic bacterium *Alcaligenes faecalis* OKK17. Journal of
 2725 Fermentation and Bioengineering 86, 351-356.
- 2726 Nobre, L.S.; GonÁalves, V.L.; Saraiva, L.M.; Robert, K.P., 2008.
 2727 Flavohemoglobin of *Staphylococcus aureus*. Methods in Enzymology 436,
 2728 203-216.
- 2729 Norton, J.M., Stark, J.M., 2011. Regulation and measurement of
 2730 nitrification in terrestrial systems. In: Klotz, M.G. (Ed.), Methods in
 2731 Enzymology, Vol. 486, Elsevier Academic Press, Burlington, pp. 343-368.
- 2732 Nussenzveig, R., Bentley, D, Ribeiro, J., 1995. Nitric oxide loading of
 2733 the salivary nitric oxide -carrying hemoprotein (nitrophorins) in the blood-
 2734 sucking bug *Rhodnius prolixus*. Journal of Experimental Biology 198, 1093-
 2735 1098.
- 2736 Okada, N., Nomura, N., Nakajima-Kambe, T., Uchiyama, H., 2005.
 2737 Characterization of the aerobic denitrification in *Mesorhizobium* sp. Strain
 2738 NH-14 in comparison with that in related rhizobia. Microbes and
 2739 Environments 20, 208-215.
- 2740 Ollivier, J., Towe, S., Bannert, A., Hai, B., Kastl, E.-M., Meyer, A., Su,
 2741 M.X., Kleineidam, K., Schlöter, M., 2011. Nitrogen turnover in soil and
 2742 global change. FEMS Microbiology Ecology 78, 3-16.
- 2743 Oremland, R.S., 2010. NO connection with methane. Nature 464, 500–
 2744 501.
- 2745 Oshiki, M., Shimokawa, M., Fujii, N., Satoh, H., Okabe, S., 2011.
 2746 Physiological characteristics of the anaerobic ammoniumoxidizing
 2747 bacterium *Candidatus 'Brocadia sinica'*. Microbiology 157, 1706–1713.

- 2748 Ouellet, H., Ouellet, Y., Richard, C., LaBarre, M., Wittenberg, B.,
2749 Wittenberg, J., Guertin, M., 2002. Truncated hemoglobin HbN protects
2750 *Mycobacterium bovis* from nitric oxide. *Proceedings of National Academy*
2751 *of Science USA* 99, 5902-5907.
- 2752 Oyadomari, S., Araki, E., Mori, M., 2002. Endoplasmic reticulum stress-
2753 mediated apoptosis in pancreatic beta-cells. *Apoptosis* 7, 335-345.
- 2754 Oyadomari, S., Takeda, K., Takiguchi, M., Gotoh, T., Matsumoto, M.,
2755 Wada, I., Akira, S., Araki, E., Mori, M., 2001. Nitric oxide-induced
2756 apoptosis in pancreatic beta cells is mediated by the endoplasmic reticulum
2757 stress pathway. *Proceedings of the National Academy of Sciences USA* 98,
2758 10845-10850.
- 2759 Pant, K., Crane, B.R., 2006. Nitrosyl-heme structures of *Bacillus subtilis*
2760 nitric oxide synthase have implications for understanding substrate
2761 oxidation. *Biochemistry* 45, 2537-2544.
- 2762 Pant, K., Bilwes, A.M., Adak, S., Stuehr, D.J., Crane, B.R., 2002.
2763 Structure of a nitric oxide synthase heme protein from *Bacillus subtilis*.
2764 *Biochemistry* 41, 11071-11079.
- 2765 Papen, H., Geßler, A., Zumbusch, E., Rennenberg, H. 2002.
2766 Chemolithoautotrophic nitrifiers in the phyllosphere of a spruce ecosystem
2767 receiving high atmospheric nitrogen input. *Current Microbiology* 44, 56-60.
- 2768 Papen, H, Von Berg, R., Hinkel, I., Thoene, B., Rennenberg, H., 1989.
2769 Heterotrophic nitrification by *Alcaligenes faecalis*: NO₂⁻, NO₃⁻, N₂O, and
2770 NO production in exponentially growing cultures. *Applied and*
2771 *Environmental Microbiology* 55, 2068-2072.

- 2772 Park, S.J., Ghai, R., Martín-Cuadrado, A.B., Rodríguez-Valera, F.,
 2773 Chung, W.H., Kwon, K., Lee, J-H., Madsen, E.L., Rhee, S.K., 2014.
 2774 Genomes of two new ammonia-oxidizing archaea enriched from deep marine
 2775 sediments. PloS One 9, 96449.
- 2776 Park, S.Y., Shimizu, H., Adachi, S.I., Nakagawa, A., Tanaka, I.,
 2777 Nakahara, K., Shoun, H., Obayashi, E., Nakamura, H., Iizuka, T., Shiro, Y.,
 2778 1997. Crystal structure of nitric oxide reductase from denitrifying fungus
 2779 *Fusarium oxysporum*. Nature Structural and Molecular Biology 4, 827-832.
- 2780 Parton, W.J., Holland, E.A., Del Grosso, S.J., Hartman, M.D., Martin,
 2781 R.E., Mosier, A.R., Ojima, D.S., Schimel, D.S., 2001. Generalized model
 2782 for NO_x and N₂O emissions from soils. Journal of Geophysical Research
 2783 106, 17403-17419.
- 2784 Parton, W.J., Schimel, D.S., Cole, C.V., Ojima, D.S., 1987. Analysis of
 2785 factors controlling soil organic matter levels in great plains Grasslands. Soil
 2786 Science Society of America Journal 51, 1173-1179.
- 2787 Patel, B.A., Moreau, M., Widom, J., Chen, H., Yin, L., Hua, Y., Crane,
 2788 B.R., 2009. Endogenous nitric oxide regulates the recovery of the radiation-
 2789 resistant bacterium *Deinococcus radiodurans* from exposure to UV light.
 2790 Proceedings of the National Academy of Sciences USA 106, 18183-18188.
- 2791 Payne, W.J., 1973. Reduction of nitrogenous oxides by microorganisms.
 2792 Bacteriological Reviews 37, 409-452.
- 2793 Payne, W.J., 1981. Denitrification. John Wiley & Sons, New York, 214
 2794 pp.
- 2795 Perazolli, M., Dominici, P., Romero-Puertas, M.C., Zago, E., Zeier, J.,
 2796 Sonoda, M., Lamb, C., Delledonne, M., 2005. Arabidopsis nonsymbiotic

- 2797 hemoglobin AHb1 modulates nitric oxide bioactivity. *The Plant Cell* 16,
2798 2785-2794.
- 2799 Pett-Ridge, J., Silver, W.L., Firestone, M.K., 2006. Redox fluctuations
2800 frame microbial community impacts on N-cycling rates in a humid tropical
2801 forest soil. *Biogeochemistry* 81, 95-110.
- 2802 Philipot, L., Hallin, S., Schloter, M., 2007. Ecology of denitrifying
2803 prokaryotes in agricultural soil. *Advances in Agronomy* 96, 249 –305.
- 2804 Pilegaard, K., 2013. Processes regulating nitric oxide emissions from
2805 soils. *Philosophical Transactions of the Royal Society B: Biological*
2806 *Sciences* 368.
- 2807 Pii, Y., Crimi, M., Cremonese, G., Spena, A., Pandolfini, T., 2007. Auxin
2808 and nitric oxide control indeterminate nodule formation. *BMC Plant*
2809 *Biology* 7, 21.
- 2810 Poock, S.R., Leach, E.R., Moir, J.W., Cole, J.A., Richardson, D.J., 2002.
2811 Respiratory detoxification of nitric oxide by the cytochrome c nitrite
2812 reductase of *Escherichia coli*. *Journal of Biological Chemistry* 277, 23664-
2813 23669.
- 2814 Poole, R.K., 2005. Nitric oxide and nitrosative stress tolerance in
2815 bacteria. *Biochemical Society Transactions* 33, 176-180.
- 2816 Porras-Alfaro, A., Herrera, J., Natvig, D.O., Lipinski, K., Sinsabaugh,
2817 R.L., 2011. Diversity and distribution of soil fungal communities in a
2818 semiarid grassland. *Mycologia* 103, 10-21.
- 2819 Porter, L.K., 1969. Gaseous products produced by anaerobic reaction of
2820 sodium nitrite with oxime compounds and oximes synthesized from organic
2821 matter. *Soil Science Society of America Journal* 33, 696-702.

- 2822 Portugal, C.C., da Encarnação, T.G., Socodato, R., Moreira, S.R.,
 2823 Brudzewsky, D., Ambrósio, A. F., Paes-de-Carvalho, R., 2012. Nitric oxide
 2824 modulates sodium vitamin C transporter 2 (SVCT-2) protein expression via
 2825 protein kinase G (PKG) and nuclear factor-kappaB (NF-kB). *Journal of*
 2826 *Biological Chemistry* 287, 3860-3872.
- 2827 Poth, M., 1986. Dinitrogen production from nitrite by a *Nitrosomonas*
 2828 isolate. *Applied and Environmental Microbiology* 52, 957-959.
- 2829 Poth, M., Focht, D.D., 1985. ¹⁵N kinetic analysis of N₂O production by
 2830 *Nitrosomonas europaea*: an examination of nitrifier denitrification. *Applied*
 2831 *and Environmental Microbiology* 49, 1134-1141.
- 2832 Prendergast-Miller, M.T., Baggs, E.M., Johnson, D., 2011. Nitrous oxide
 2833 production by the ectomycorrhizal fungi *Paxillus involutus* and *Tylospora*
 2834 *fibrillosa*. *FEMS Microbiology Letters* 316, 31-35.
- 2835 Primblecombe, P., 1996. Air, composition and chemistry, second ed.
 2836 Cambridge University Press, Cambridge, 250 pp.
- 2837 Prosser, J.I., 1989. Autotrophic nitrification in bacteria. *Advances in*
 2838 *Microbial Physiology* 30, 125-181.
- 2839 Prosser, J.I., Nicol, G.W., 2008. Relative contributions of archaea and
 2840 bacteria to aerobic ammonia oxidation in the environment. *Environmental*
 2841 *Microbiology* 10, 2931-2941.
- 2842 Prosser, J.I., Nicol, G.W., 2012. Archaeal and bacterial ammonia
 2843 oxidisers in soil: the quest for niche specialisation and differentiation.
 2844 *Trends in Microbiology* 20, 523-531.
- 2845 Quan, Z.X., Rhee, S.K., Zuo, J.E., Yang, Y., Bae, J.W., Park, J.R., Lee,
 2846 S.T., Park, Y.H., 2008. Diversity of ammonium-oxidizing bacteria in a

2847 granular sludge anaerobic ammoniumoxidizing (anammox) reactor.
2848 Environmental Microbiology 10, 3130–3139.

2849 Rafferty, S., 2011. Nitric oxide synthases of bacteria and other
2850 unicellular organisms. Open Nitric Oxide Journal 3, 25-32.

2851 Raghoebarsing, A.A., Pol, A., van de Pas-Schoonen, K.T., Smolders, A.J.
2852 P., Ettwig, K.F., Rijpstra, W.I.C., Schouten, S., Damsté, J.S.S., Op den
2853 Camp, H.J.M., Jetten, M.S.M., Strous, M., 2006. A microbial consortium
2854 couples anaerobic methane oxidation to denitrification. Nature 440, 918–
2855 921.

2856 Rahmanto, Y.S., Kalinowski, D.S., Lane, D.J., Lok, H.C., Richardson,
2857 V., Richardson, D.R., 2012. Nitrogen monoxide (NO) storage and transport
2858 by dinitrosyl-dithiol-iron complexes: long-lived NO that is trafficked by
2859 interacting proteins. Journal of Biological Chemistry 287, 6960-6968.

2860 Rees, R.M., Augustin, J., Alberti, G., Ball, B.C., Boeckx, P., Cantarel,
2861 A., Castaldi, S., Chirinda, N., Chojnicki, B., Giebels, M., Gordon, H.,
2862 Grosz, B., Horvath, L., Juszczak, R., Klemetsson, A.K., Klemetsson, L.,
2863 Medinets, S., Machon, A., Mapanda, F., Nyamangara, J., Olesen, J., Reay,
2864 D., Sanchez, L., Sanz Cobena, A., Smith, K.A., Sowerby, A., Sommer, M.,
2865 Soussana, J.F., Stenberg, M., Topp, C.F.E., van Cleemput, O., Vallejo, A.,
2866 Watson, C.A., Wuta, M., 2013. Nitrous oxide emissions from European
2867 agriculture - an analysis of variability and drivers of emissions from field
2868 experiments. Biogeosciences, 10, 2671-2682.

2869 Remde, A., Conrad, R., 1990. Production of nitric oxide in *Nitrosomonas*
2870 *europaea* by reduction of nitrite. Archives of Microbiology 154, 187–191.

- 2871 Revsbech, N.P., Sørensen, J. (Eds.), 1990. Denitrification in soil and
2872 sediment. Plenum Press, New York, 349 pp.
- 2873 Rich, J.J., Dale, O.R., Song, B., Ward, B.B., 2008. Anaerobic ammonium
2874 oxidation (Anammox) in Chesapeake Bay sediments. *Microbial Ecology* 55,
2875 311–320.
- 2876 Richardson, D.R., Tran, E.H., Ponka, P., 1995. The potential of iron
2877 chelators of the pyridoxal isonicotinoyl hydrazone class as effective
2878 antiproliferative agents. *Blood* 86, 4295-4306.
- 2879 Ridnour, L.A., Thomas, D.D., Mancardi, D., Espey, M.G., Miranda,
2880 K.M., Paolocci, N., Feelisch, M., Fukuto, J., Wink, D.A., 2004. The
2881 chemistry of nitrosative stress induced by nitric oxide and reactive nitrogen
2882 oxide species. Putting perspective on stressful biological situations.
2883 *Biological Chemistry* 385, 1-10.
- 2884 Robertson, G.P., 1989. Nitrification and denitrification in humid tropical
2885 ecosystems: Potential controls on nitrogen retention. In: Procter, J. (Ed.),
2886 Mineral nutrients in tropical forest and savanna ecosystems, Blackwell
2887 Scientific, Boston, pp. 55-69.
- 2888 Robertson, G.P., Groffman, P.M., 2007. Nitrogen transformations. In:
2889 Paul, E.A. (Ed.), *Soil microbiology, ecology, and biochemistry*, third ed.,
2890 Springer, New York, pp. 341-364.
- 2891 Robertson, J., Bonaventura, J., Kohm, A., 1995. Nitric oxide synthase
2892 inhibition blocks octopus touch learning without producing sensory or
2893 motor dysfunction. *Proceedings of the Royal Society of London B:*
2894 *Biological Sciences* 261, 167-172.

- 2895 Romero-Puertas, M.C., Laxa, M., Matté, A., Zaninotto, F., Finkemeier,
 2896 I., Jones, A.M.E., Perazolli, M., Vandelle, E., Dietz, K.-J., Delledonne, M.,
 2897 2007. S-nitrosylation of peroxiredoxin II E promotes peroxynitrite-mediated
 2898 tyrosine nitration. *The Plant Cell* 19, 4120-4130.
- 2899 Rudolph, J., Koschorreck, M., Conrad, R., 1996. Oxidative and reductive
 2900 microbial consumption of nitric oxide in a heathland soil. *Soil Biology and*
 2901 *Biochemistry* 28, 1389-1396.
- 2902 Ruggiero, C.E., Carrier, S.M., Antholine, W.E., Whittaker, J.W., Cramer,
 2903 C.J., Tolman, W.B., 1993. Synthesis and structural and spectroscopic
 2904 characterization of mononuclear copper nitrosyl complexes: models for
 2905 nitric oxide adducts of copper proteins and copper-exchanged zeolites.
 2906 *Journal of the American Chemical Society* 115, 11285-11298.
- 2907 Rümer, S., Gupta, K.J., Kaiser, W.M., 2009a. The Plant Cells oxidize
 2908 hydroxylamines to NO. *Journal of Experimental Botany* 60, 2065-2072.
- 2909 Rümer, S., Gupta, K.J., Kaiser, W.M., 2009b. Oxidation of
 2910 hydroxylamines to NO by The Plant Cells. *Plant Signaling and Behavior* 4,
 2911 853-855.
- 2912 Russow, R., Stange, C.F., Neue, H.U., 2009. Role of nitrite and nitric
 2913 oxide in the processes of nitrification and denitrification in soil: Results
 2914 from ¹⁵N tracer experiments. *Soil Biology and Biochemistry* 41, 785-795.
- 2915 Rusterucci, C., Espunya, M.C., Díaz, M., Chabannes, M., Martínez,
 2916 M.C., 2007. S-nitrosoglutathione reductase affords protection against
 2917 pathogens in *Arabidopsis*, both locally and systemically. *Plant Physiology*
 2918 143, 1282-1292.

- 2919 Rütting, T., Boeckx, P., Müller, C., Klemetsson, L., 2011. Assessment
2920 of the importance of dissimilatory nitrate reduction to ammonium for the
2921 terrestrial nitrogen cycle. *Biogeosciences* 8, 1779-1791.
- 2922 Rütting, T., Huygens, D., Muller, C., Cleemput, O., Godoy, R., Boeckx,
2923 P., 2008. Functional role of DNRA and nitrite reduction in a pristine south
2924 Chilean Nothofagus forest. *Biogeochemistry* 90, 243-258.
- 2925 Rysgaard, S., Glud, R.N., Risgaard-Petersen, N., Dalsgaard, T., 2004.
2926 Denitrification and anammox activity in Arctic marine sediments.
2927 *Limnology and Oceanography* 49, 1493–1502.
- 2928 Rysgaard, S., Glud, R.N., Sejr, M.K., Blicher, M.E., Stahl, H.J., 2008.
2929 Denitrification activity and oxygen dynamics in Arctic sea ice. *Polar*
2930 *Biology* 31, 527–537.
- 2931 Saad, O.A., Conrad, R., 1993. Temperature dependence of nitrification,
2932 denitrification, and turnover of nitric oxide in different soils. *Biology and*
2933 *Fertility of Soils* 15, 21-27.
- 2934 Saito, S., Yamamoto-Katou, A., Yoshioka, H., Doke, N., Kawakita, K.,
2935 2006. Peroxynitrite generation and tyrosine nitration in defense responses in
2936 tobacco BY-2 cells. *Plant and Cell Physiology* 47, 689-697.
- 2937 Salard, I., Mercey, E., Rekka, E., Boucher, J.L., Nioche, P., Mikula, I.,
2938 Martasek, P., Raman, C.S., Mansuy, D., 2006. Analogies and surprising
2939 differences between recombinant nitric oxide synthase-like proteins from
2940 *Staphylococcus aureus* and *Bacillus anthracis* in their interactions with L-
2941 arginine analogs and iron ligands. *Journal of Inorganic Biochemistry* 100,
2942 2024-2033.

- 2943 Sameshima-Saito, R., Chiba, K., Minamisawa, K., 2004. New method of
 2944 denitrification analysis of *Bradyrhizobium* field isolates by gas
 2945 chromatographic determination of N-15-labeled N₂. Applied and
 2946 Environmental Microbiology 70, 2886-2891.
- 2947 Sanchez, C., Gates, A.J., Meakin, G.E., Uchiumi, T., Girard, L.,
 2948 Richardson, D.J., Bedmar, E.J., Delgado, M.J., 2010. Production of nitric
 2949 oxide and nitrosylhemoglobin complexes in soybean nodules in response
 2950 to flooding. Molecular Plant-Microbe Interactions 23, 702-711.
- 2951 Scandalios, J.G., 1997. Molecular genetics of superoxide dismutase in
 2952 plants. In: Scandalios, J.G. (Ed.), Oxidative stress and the molecular biology
 2953 of antioxidant defences, Cold Spring Harbor Lab. Press, New York, pp.
 2954 527-568.
- 2955 Schauss, K., Focks, A., Leininger, S., Kotzerke, A., Heuer, H.,
 2956 Thiele-Bruhn, S., Sharma, S., Wilke, B.-M., Matthies, M., Smalla, K.,
 2957 Munch, J.C., Amelung, W., Kaupenjohann, M., Schloter, M., Schleper, C.,
 2958 2009. Dynamics and functional relevance of ammonia-oxidizing archaea in
 2959 two agricultural soils. Environmental Microbiology 11, 446-456.
- 2960 Scheible, W.R., Fry, B., Kochevenko, A., Schindelasch, D., Zimmerli, L.,
 2961 Somerville, S., Loria, R., Somerville, C.R., 2003. An Arabidopsis mutant
 2962 resistant to thaxtomin A, a cellulose synthesis inhibitor from *Streptomyces*
 2963 species. The Plant Cell 15, 1781-1794.
- 2964 Schindlbacher, A., Zechmeister-Boltenstern, S., Butterbach-Bahl, K.,
 2965 2004. Effects of soil moisture and temperature on NO, NO₂, and N₂O
 2966 emissions from European forest soils. Journal of Geophysical Research-
 2967 Atmos. 109, D17302.

- 2968 Schlag, S., Ners, C., Birkenstock, T.A., Altenberend, F., Gotz, F., 2007.
 2969 Inhibition of staphylococcal biofilm formation by nitrite. *Journal of*
 2970 *Bacteriology* 189, 7911-7919.
- 2971 Schleper, C., Jurgens, G., Jonuscheit, M., 2005. Genomic studies of
 2972 uncultivated archaea. *Nature Reviews Microbiology* 3, 479-488.
- 2973 Schleper, C., Nicol, G.W., 2010. Ammonia-oxidising archaea–
 2974 physiology, ecology and evolution. *Advances in Microbial Physiology* 57,
 2975 1–41.
- 2976 Schmid, M., Walsh, K., Webb, R., Rijpstra, W.I.C., van de Pas-
 2977 Schoonen, K., Verbruggen, M.J., Hill, T., Moffett, B., Fuerst, J., Schouten,
 2978 S., Damsté, J.S.S., Harris, J., Shaw, P., Jetten, M., Strous, M., 2003.
 2979 *Candidatus “Scalindua brodae”*, sp. nov., *Candidatus “Scalindua*
 2980 *wagneri”*, sp. nov., two new species of anaerobic ammonium oxidizing
 2981 bacteria. *Systematic and Applied Microbiology* 26, 529–538.
- 2982 Schmidt, C.S., Richardson, D.J., Baggs, E.M., 2011. Constraining the
 2983 conditions conducive to dissimilatory nitrate reduction to ammonium in
 2984 temperate arable soils. *Journal of Inorganic Biochemistry* 43, 1607-1611.
- 2985 Schmidt, I., Steenbakkers, P.J.M., op den Camp, H.J.M., Schmidt, K.,
 2986 Jetten, M.S.M., 2004. Physiologic and proteomic evidence for a role of
 2987 nitric oxide in biofilm formation by *Nitrosomonas europaea* and other
 2988 ammonia oxidizers. *Journal of Bacteriology* 186, 2781-2788.
- 2989 Schreiber, F., Beutler, M., Enning, D., Lamprecht-Grandio, M., Zafra,
 2990 O., González-Pastor, J.E., de Beer, D., 2011. The role of nitric-oxide-
 2991 synthase-derived nitric oxide in multicellular traits of *Bacillus subtilis* 3610:
 2992 biofilm formation, swarming, and dispersal. *BMC Microbiology* 11, 111.

- 2993 Schreiber, F., Stief, P., Kuypers, M.M., de Beer, D., 2014. Nitric oxide
2994 turnover in permeable river sediment. *Limnology and Oceanography* 59,
2995 1310-1320.
- 2996 Schreiber, F., Wunderlin, P., Udert, K.M., Wells, G.F., 2012. Nitric oxide
2997 and nitrous oxide turnover in natural and engineered microbial
2998 communities: biological pathways, chemical reactions, and novel
2999 technologies. *Frontiers in microbiology* 3, 372.
- 3000 Schubert, C.J., Durisch-Kaiser, E., Wehrli, B., Thamdrup, B., Lam, P.,
3001 Kuypers, M.M., 2006. Anaerobic ammonium oxidation in a tropical
3002 freshwater system (Lake Tanganyika). *Environmental Microbiology* 8,
3003 1857–1863.
- 3004 Schütz, H., Schröder, P., Rennenberg, H., 1991. Role of plants in
3005 regulating the methane flux to the atmosphere. In: Sharkey, T.D, Holland,
3006 E.A., Mooney, H.A. (Eds.), *Trace Gas Emissions by Plants*, Academic
3007 Press, San Diego, pp. 29-63.
- 3008 Seok, B., Helmig, D., Ganzeveld, L., Williams, M.W., Vogel, C.S., 2013.
3009 Dynamics of nitrogen oxides and ozone above and within a mixed
3010 hardwood forest in northern Michigan. *Atmospheric Chemistry and Physics*
3011 13, 7301-7320.
- 3012 Sharma, S., Szele, Z., Schilling, R., Munch, J.C., Schlöter, M., 2006.
3013 Influence of freeze-thaw stress on the structure and function of microbial
3014 communities and denitrifying populations in soil. *Applied and*
3015 *Environmental Microbiology* 72, 2148-2154.
- 3016 Shatalin, K., Gusarov, I., Avetissova, E., Shatalina, Y., McQuade, L.E.,
3017 Lippard, S.J., Nudler, E., 2008. *Bacillus anthracis*-derived nitric oxide is

- 3018 essential for pathogen virulence and survival in macrophages. Proceedings
3019 of the National Academy of Sciences USA 105, 1009-1013.
- 3020 Shen, J., Li, Y., Liu, X., Luo, X., Tang, H., Zhang, Y., Wu, J., 2013.
3021 Atmospheric dry and wet nitrogen deposition on three contrasting land use
3022 types of an agricultural catchment in subtropical central China. Atmospheric
3023 Environment 67, 415-424.
- 3024 Shen, L.-D., He, Z.-F., Zhu, Q., Chen, D.-Q., Lou, L.-P., Xu, X.-Y.,
3025 Zheng, P., Hu, B.-L., 2012. Microbiology, ecology, and application of the
3026 nitrite-dependent anaerobic methane oxidation process. Frontiers in
3027 Microbiology 3, 269.
- 3028 Shen, T., Stieglmeier, M., Dai, J., Urich, T., Schleper, C., 2013.
3029 Responses of the terrestrial ammonia-oxidizing archaeon *Ca.*
3030 *Nitrososphaera viennensis* and the ammonia-oxidizing bacterium
3031 *Nitrospira multiformis* to nitrification inhibitors. FEMS Microbiology
3032 Letters 344, 121–129.
- 3033 Shi, W., Norton, J.M., 2000. Effect of long-term, biennial, fall-applied
3034 anhydrous ammonia and nitrapyrin on soil nitrification. Soil Science Society
3035 of America Journal 64, 228-234.
- 3036 Shi, Q., Ding, F., Wang, X., Wei, M., 2007. Exogenous nitric oxide
3037 protect cucumber roots against oxidative stress induced by salt stress. Plant
3038 Physiology and Biochemistry 45, 542-550.
- 3039 Shoun, H., Kim, D.H., Uchiyama, H., Sugiyama, J., 1992. Denitrification
3040 by fungi. FEMS Microbiology Letters 94, 277-281.
- 3041 Shoun, H., Fushinobu, S., Jiang, L., Kim, S.W., Wakagi, T., 2012.
3042 Fungal denitrification and nitric oxide reductase cytochrome P450nor.

- 3043 Philosophical Transactions of the Royal Society B: Biological Sciences 367,
3044 1186-1194.
- 3045 Shoun, H., Kim, D.H., Uchiyama, H., Sugiyama, J., 1992. Denitrification
3046 by fungi. FEMS Microbiology Letters 94, 277-281.
- 3047 Siervo, M., Stephan, B.C., Feelisch, M., Bluck, L.J., 2011. Measurement
3048 of in vivo nitric oxide synthesis in humans using stable isotopic methods: A
3049 systematic review. Free Radical Biology and Medicine 51, 795-804.
- 3050 Silver, W.L., Herman, D.J., Firestone, M.K., 2001. Dissimilatory nitrate
3051 reduction to ammonium in upland tropical forest soils. Ecology 8, 2410-
3052 2416.
- 3053 Silver, W.L., Thompson, A.W., Reich, A., Ewel, J.J., Firestone, M.K.,
3054 2005. Nitrogen cycling in tropical plantation forests: potential controls on
3055 nitrogen retention. Ecological Applications 15, 1604-1614.
- 3056 Simon, J., 2002. Enzymology and bioenergetics of respiratory nitrite
3057 ammonification. FEMS Microbiology Reviews 26, 285-309.
- 3058 Simon, J., Dong, F., Buegger, F., Rennenberg, H., 2013. Rhizospheric
3059 NO affects N uptake and metabolism in Scots pine (*Pinus sylvestris* L.)
3060 seedlings depending on soil N availability and N sources. Plant, Cell and
3061 Environment 36, 1019-1026.
- 3062 Simon, J., Sanger, M., Schuster, S.C., Gross, R., 2003. Electron transport
3063 to periplasmic nitrate reductase (NapA) of *Wolinella succinogenes* is
3064 independent of a NapC protein. Molecular Microbiology 49, 69-79.
- 3065 Simon, J., Stoelken, G., Rienks, M., Rennenberg, H., 2009. Rhizospheric
3066 NO interacts with the acquisition of reduced nitrogen sources by the roots of
3067 European beech (*Fagus sylvatica*). FEBS Letters 583, 2907-2910.

- 3068 Singh, B., Singh, B., Singh, Y., 1993. Potential and kinetics of
3069 nitrification in soils from semiarid regions of northwestern India. *Arid Land*
3070 *Research and Rehabilitation* 7, 39-50.
- 3071 Skiba, U., 2008. Denitrification. In: Jørgensen, S.E., Fath, B.D. (Eds.),
3072 *Ecological Processes*, vol. 2 of *Encyclopedia of Ecology*, Elsevier, Oxford,
3073 pp. 866-871.
- 3074 Skiba, U., Fowler, D., Smith, K.A., 1997. Nitric oxide emissions from
3075 agricultural soils in temperate and tropical climates: Sources, control and
3076 mitigation options. *Nutrient Cycling in Agroecosystems* 48, 139–153.
- 3077 Slemr, F., Seiler, W., 1984. Field measurement of NO and NO₂
3078 emissions from fertilized and unfertilized soils. *Journal of Atmospheric*
3079 *Chemistry* 2, 1-24.
- 3080 Slemr, F., Seiler, W., 1991. Field study of environmental variables
3081 controlling the NO emission from soil and the NO compensation point.
3082 *Journal of Geophysical Research* 96, 13017-13031.
- 3083 Smith, C.J., Chalk, P.M., 1980. Fixation and loss of nitrogen during
3084 transformations of nitrite in soils. *Soil Science Society of America Journal*
3085 44, 288-291.
- 3086 Smith, N.A., Smith, P., 1992. Nitrate reduction and N-nitrosation by
3087 *Obesumbacterium proteus*. *Letters in Applied Microbiology* 14, 61-64.
- 3088 Socodato, R., Portugal, C.C., Encarnação, T.G., Domith, I., Cossenza,
3089 M., Paes-de-Carvalho, R., 2013. Studying nitric oxide in the developing
3090 retina: neuromodulatory functions and signaling mechanisms. *Open Nitric*
3091 *Oxide Journal* 4, 1-15.

- 3092 Sokolovski, S., Blatt, M.R., 2004. Nitric oxide block of outward-
3093 rectifying K^+ channels indicates direct control by protein nitrosylation in
3094 guard cells. *Plant Physiology* 136, 4275-4284.
- 3095 Souza, J.M., Choi, I., Chen, Q., Weisse, M., Daikhin, E., Yudkoff, M.,
3096 Spang, A., Hatzenpichler, R., Brochier-Armanet, C., Rattei, T., Tischler, P.,
3097 Spieck, E., Streit, W., Stahl, D.A., Wagner, M., Schleper, C., 2010. Distinct
3098 gene set in two different lineages of ammonia-oxidizing archaea supports
3099 the phylum *Thaumarchaeota*. *Trends in Microbiology* 18, 331-340.
- 3100 Sparks, J.P., 2009. Ecological ramifications of the direct foliar uptake of
3101 nitrogen. *Oecologia* 159, 1-13.
- 3102 Spott, O., Russow, R., Stange, C.F., 2011. Formation of hybrid N_2O and
3103 hybrid N_2 due to codenitrification: First review of a barely considered
3104 process of microbially mediated N-nitrosation. *Soil Biology and*
3105 *Biochemistry* 43, 1995-2011.
- 3106 Spott, O., Stange, C.F., 2011. Formation of hybrid N_2O in a suspended
3107 soil due to co-denitrification of NH_2OH . *Journal of Plant Nutrition and Soil*
3108 *Science* 174, 554-567.
- 3109 Staffelbach, T., Neftel, A., Horowitz, L.W., 1997. Photochemical oxidant
3110 formation over southern Switzerland: 2 model results. *Journal of*
3111 *Geophysical Research* 102, 23363-23373.
- 3112 Stamler, J.S., Singel, D.J., Loscalzo, J., 1992. Biochemistry of nitric-
3113 oxide and its redox-activated forms. *Science* 258, 1898-1902.
- 3114 Stanbury, D.M., 1989. Reduction potentials involving inorganic free
3115 radicals in aqueous solution. *Advances in Inorganic Chemistry* 33, 69-138.

- 3116 Stark, J.M., 1996. Modeling the temperature response of nitrification.
3117 Biogeochemistry 35, 433-445.
- 3118 Stark, J.M., Firestone, M.K., 1996. Kinetic characteristics of ammonium-
3119 oxidizer communities in a California oak woodland-annual grassland. Soil
3120 Biology and Biochemistry 28, 1307-1317.
- 3121 Ste-Marie, C., Pare, D., 1999. Soil, pH and N availability effects on net
3122 nitrification in the forest floors of a range of boreal forest stands. Soil
3123 Biology and Biochemistry 31, 1579-1589.
- 3124 Stein, L.Y., Arp, D.J., 1998. Loss of ammonia monooxygenase activity in
3125 *Nitrosomonas europaea* upon exposure to nitrite. Applied and
3126 Environmental Microbiology 64, 4098-4102.
- 3127 Stevens, R.J., Laughlin, R.J., Malone, J.P., 1998. Soil pH affects the
3128 processes reducing nitrate to nitrous oxide and di-nitrogen. Soil Biology and
3129 Biochemistry 30, 1119-1126.
- 3130 Stevenson, F.J., Harrison, R.M., Wetselaar, R., Leeper, R.A., 1970.
3131 Nitrosation of soil organic matter: III. Nature of gases produced by reaction
3132 of nitrite with lignins, humic substances, and phenolic constituents under
3133 neutral and slightly acidic conditions. Soil Science Society of America
3134 Journal 34, 430-435.
- 3135 Stieglmeier, M., Klingl, A., Alves, R.J., Simon, K.M.R., Melcher, M.,
3136 Leisch, N., Schleper, C., 2014a. *Nitrososphaera viennensis* sp. nov., an
3137 aerobic and mesophilic ammonia-oxidizing archaeon from soil and member
3138 of the archaeal phylum *Thaumarchaeota*. International Journal of
3139 Systematic and Evolutionary Microbiology, ijs.0.063172-0.

- 3140 Stieglmeier, M., Mooshammer, M., Kitzler, B., Wanek, W., Zechmeister-
 3141 Boltenstern, S., Richter, A., Schleper, C., 2014b. Aerobic nitrous oxide
 3142 production through N-nitrosating hybrid formation in ammonia-oxidizing
 3143 archaea. ISME Journal 8, 1135-1146.
- 3144 Stitt, M., Müller, C., Matt, P., Gibon, Y., Carillo, P., Morcuende, R.,
 3145 Krapp, A., 2002. Steps towards an integrated view of nitrogen metabolism.
 3146 Journal of Experimental Botany 53, 959-970.
- 3147 Stoelken, G., Simon, J., Ehrling, B., Rennenberg, H., 2010. The presence
 3148 of amino acids affects inorganic N uptake in non-mycorrhizal seedlings of
 3149 European beech (*Fagus sylvatica* L.). Tree Physiology 30, 1118-1128.
- 3150 Stouthamer, A.H., 1988. Dissimilatory reduction of oxidized nitrogen
 3151 compounds. In: Zehnder, A.J.B. (Ed.), Biology of anaerobic
 3152 microorganisms, John Wiley & Sons, New York, pp. 245-303.
- 3153 Strous, M., Fuerst, J.A., Kramer, E.H., Logemann, S., Muyzer, G., van de
 3154 Pas-Schoonen, K.T., Webb, R., Kuenen, J.G., Jetton, M.S.M., 1999.
 3155 Missing lithotroph identified as new planctomycete. Nature 400, 446-449.
- 3156 Strous, M., Heijnen, J.J., Kuenen, J.G., Jetton, M.S.M., 1998. The
 3157 sequencing batch reactor as a powerful tool for the study of slowly growing
 3158 anaerobic ammonium-oxidizing microorganisms. Applied Microbiology and
 3159 Biotechnology 50, 589-596.
- 3160 Strous, M., Pelletier, E., Mangenot, S., Rattei, T., Lehner, A., Taylor,
 3161 M.W., Horn, M., Daims, H., Bartol-Mavel, D., Wincker, P., Barbe, V.,
 3162 Fonknechten, N., Vallenet, D., Segurens, B., Schenowitz-Truong, C.,
 3163 Medigue, C., Collingro, A., Snel, B., Dutilh, B.E., Op den Camp, H.J.M.,
 3164 van der Drift, C., Cirpus, I., van de Pas- Schoonen, K.T., Harhangi, H.R.,

- 3165 van Niftrik, L., Schmid, M., Keltjens, J., van de Vossenberg, J., Kartal, B.,
 3166 Meier, H., Frishman, D., Huynen, M.A., Mewes, H.W., Weissenbach, J.,
 3167 Jetten, M.S.M., Wagner, M., Le Paslier, D., 2006. Deciphering the evolution
 3168 and metabolism of an anammox bacterium from a community genome.
 3169 *Nature* 440, 790-794.
- 3170 Strous, M., van Gerven, E., Kuenen, J.G., Jetton, M.S.M, 1997. Effects
 3171 of aerobic and microaerobic conditions on anaerobic ammoniumoxidizing
 3172 (anammox) sludge. *Applied and Environmental Microbiology* 63, 2446–
 3173 2448.
- 3174 Stuehr, D.J., 1999. Mammalian nitric oxide synthases. *Biochimica et*
 3175 *Biophysica Acta* 1411, 217-230.
- 3176 Su, F., Takaya, N., Shoun, H., 2004. Nitrous oxide-forming
 3177 codenitrification catalyzed by cytochrome P450_{nor}. *Bioscience*
 3178 *Biotechnology and Biochemistry* 68, 473-475.
- 3179 Su, H., Cheng, Y., Oswald, R., Behrendt, T., Trebs, I., Meixner, F. X.,
 3180 Andreae M.O., Cheng P., Zhang Y., Pöschl, U., 2011. Soil nitrite as a
 3181 source of atmospheric HONO and OH radicals. *Science* 333, 1616-1618.
- 3182 Su, M.X., Kleineidam, K., Schlöter, M., 2010. Influence of different litter
 3183 quality on the abundance of genes involved in nitrification and
 3184 denitrification after freezing and thawing of an arable soil. *Biology and*
 3185 *Fertility of Soils* 46, 537-541.
- 3186 Subbarao, G.V., Ito, O., Sahrawat, K.L., Berry, W.L., Nakahara, K.,
 3187 Ishikawa, Watanabe T., Suenaga K., Rondon M., Rao, I.M., 2006. Scope
 3188 and strategies for regulation of nitrification in agricultural systems -

- 3189 challenges and opportunities. *Critical Reviews in Plant Sciences* 25, 303-
3190 335.
- 3191 Sudhamsu, J., Crane, B.R., 2009. Bacterial nitric oxide synthases: what
3192 are they good for? *Trends in Microbiology* 17, 212-218.
- 3193 Susswein, A.J., Chiel, H.J., 2012. Nitric oxide as a regulator of behavior:
3194 New ideas from *Aplysia* feeding. *Progress in Neurobiology* 97, 304-317.
- 3195 Szukics, U., Abell, G. C., Hödl, V., Mitter, B., Sessitsch, A., Hackl, E.,
3196 Zechmeister-Boltenstern, S., 2010. Nitrifiers and denitrifiers respond rapidly
3197 to changed moisture and increasing temperature in a pristine forest soil.
3198 *FEMS Microbiology Ecology* 72, 395-406.
- 3199 Tada, Y., Spoel, S.H., Pajerowska-Mukhtar, K., Mou, Z., Song, J., Wang,
3200 C., Zuo, J., Dong, X., 2008. Plant immunity requires conformational charges
3201 of NPR1 via S-nitrosylation and thioredoxins. *Science* 321, 952-956.
- 3202 Takahashi, Y., Katoh, S., Shikura, N., Tomoda, H., Omura, S., 2003.
3203 Superoxide dismutase produced by soil bacteria increases bacterial colony
3204 growth from soil samples. *The Journal of General and Applied*
3205 *Microbiology* 49, 263-266.
- 3206 Talukder, M.A., Yang, F., Shimokawa, H., Zweier, J.L., 2010. eNOS is
3207 required for acute in vivo ischemic preconditioning of the heart: Effects of
3208 ischemic duration and sex. *American Journal of Physiology-Heart and*
3209 *Circulatory Physiology* 299, H437-445.
- 3210 Tanimoto, T., Hatano, K., Kim, D.H., Uchiyama, H., Shoun, H., 1992.
3211 Co-denitrification by the denitrifying system of the fungus *Fusarium*
3212 *oxysporum*. *FEMS Microbiology Letters* 93, 177-180.

- 3213 Teillet-Billy, D., Fiquet-Fayard, F., 1977. The NO-3 Σ -and 1 Δ
 3214 resonances: theoretical analysis of electron scattering data. Journal of
 3215 Physics B: Atomic and Molecular Physics 10, L111.
- 3216 Teklemariam, T.A., Sparks, J.P., 2006. Leaf fluxes of NO and NO₂ in
 3217 four herbaceous plant species: The role of ascorbic acid. Atmospheric
 3218 Environment 40, 2235-2244.
- 3219 Templer, P.H., Silver, W.L., Pett-Ridge, J., DeAngelis, K.M., Firestone,
 3220 M.K., 2008. Plant and microbial controls on nitrogen retention and loss in a
 3221 tropical humid forest. Ecology 89, 3030-3040.
- 3222 Tiedje, J.M., 1988. Ecology of denitrification and dissimilatory reduction
 3223 of nitrate to ammonium. In: Zinder, A.J.B. (Ed.), Biology of Anaerobic
 3224 Microorganisms, John Wiley and Sons, New York, pp.179-244.
- 3225 Tiedje, J.M., Sexstone, A.M., Myrold, D.D., Robinson, J.A., 1982.
 3226 Denitrification: ecological niches, competition and survival. Antonie van
 3227 Leeuwenhoek 48, 569-583.
- 3228 Thomas, D.D., Liu, X., Kantrow, S.P., Lancaster, J.R., 2001. The
 3229 biological lifetime of nitric oxide: implications for the the perivascular
 3230 dynamics of NO and O₂. Proceedings of National Academy of Science 98,
 3231 355-360.
- 3232 Thomas, D.D., Ridnour, L.A., Isenberg, J.S., Flores-Santana, W.,
 3233 Switzer, C.H., Donzelli, S., Hussain, P., Vecoli, C., Paolozzi, N., Ambs, S.,
 3234 Colton, C.A., Harris, C.S., Roberts, D.D., Wink, D.A., 2008. The chemical
 3235 biology of nitric oxide: implications in cellular signaling. Free Radical
 3236 Biology and Medicine 45, 18-31.

- 3237 Thorn, K.A., Mikita, M.A., 2000. Nitrite fixation by humic substances:
 3238 nitrogen-15 nuclear magnetic resonance evidence for potential intermediates
 3239 in chemodenitrification. *Soil Science Society of America Journal* 64, 568-
 3240 582.
- 3241 Tourna, M., Freitag, T.E., Nicol, G.W., Prosser, J.I., 2008. Growth,
 3242 activity and temperature responses of ammonia-oxidizing archaea and
 3243 bacteria in soil microcosms. *Environmental Microbiology* 10, 1357-1364.
- 3244 Toyoda, S., Yano, M., Nishimura, S.I., Akiyama, H., Hayakawa, A.,
 3245 Koba, K., Sudo, S., Yagi, K., Makabe, A., Tobari, Y., Ogawa, N.O.,
 3246 Ohkouchi, N., Yamada, K., Yoshida, N., 2011. Characterization and
 3247 production and consumption processes of N₂O emitted from temperate
 3248 agricultural soils determined via isotopomer ratio analysis. *Global*
 3249 *Biogeochemical Cycles* 25, GB2008.
- 3250 Trimmer, M., Nicholls, J.C., Deflandre, B., 2003. Anaerobic ammonium
 3251 oxidation measured in sediments along the Thames estuary, United
 3252 Kingdom. *Applied and Environmental Microbiology* 69, 6447–6454.
- 3253 Troemel, E.R., Chu, S.W., Reinke, V., Lee, S.S., Ausubel, F.M., Kim,
 3254 D.H., 2006. p38 MAPK regulates expression of immune response genes and
 3255 contributes to longevity in *C. elegans*. *PLoS Genet.* 2, e183.
- 3256 Tullius, M.V., Harth, G., Horwitz, M.A., 2001. High extracellular levels
 3257 of *Mycobacterium tuberculosis* glutamine synthetase and superoxide
 3258 dismutase in actively growing cultures are due to high expression and
 3259 extracellular stability rather than to a protein-specific export mechanism.
 3260 *Infection and Immunity* 69, 6348-6363.

- 3261 Usuda, K., Toritsuka, N., Matsuo, Y., Kim, D.H., Shoun, H., 1995.
 3262 Denitrification by the fungus *Cylindrocarpon tonkinense* - anaerobic cell-
 3263 growth and 2 Isozyme forms of cytochrome P-450Nor. Applied and
 3264 Environmental Microbiology 61, 883-889.
- 3265 Vajrala, N., Martens-Habbena, W., Sayavedra-Soto, L.A., Schauer, A.,
 3266 Bottomley, P.J., Stahl, D.A., Arp, D.J., 2013. Hydroxylamine as an
 3267 intermediate in ammonia oxidation by globally abundant marine archaea.
 3268 PNAS 110, 1006–1011.
- 3269 Valentine, D.L., 2007. Adaptations to energy stress dictate the ecology
 3270 and evolution of the archaea. Nature Reviews Microbiology 5, 316-323.
- 3271 Van Cleemput, O., Baert, L., 1976. Theoretical considerations on nitrite
 3272 self-decomposition reactions in soils. Soil Science Society of America
 3273 Journal 40, 322-324.
- 3274 Van Cleemput, O., Baert, L., 1984. Nitrite a key compound in N loss
 3275 processes under acid condition? Plant and Soil 76, 233-241.
- 3276 Van Cleemput, O., Samater, A.H., 1996. Nitrite in soils: accumulation
 3277 and the role in the formation of gaseous N compounds. Fertilizer Research
 3278 45, 81-89.
- 3279 van de Graaf, A., de Bruijn P., Robertson, L.A., Jetton, M.S.M., Kuenen
 3280 J.G., 1996. Autotrophic growth of anaerobic ammonium-oxidizing micro-
 3281 organisms in a fluidized bed reactor. Microbiology 142, 2187–2196.
- 3282 van de Graaf, A., Mulder, A., de Bruijn, P., Jetten, M.S.M., Robertson,
 3283 L.A., Kuenen, J.G., 1995. Anaerobic oxidation of ammonium is a
 3284 biologically mediated process. Applied and Environmental Microbiology
 3285 61, 1246–1251.

- 3286 van de Graaf, A., Mulder, A., Slikhuis, H., Robertson, L.A., Kuenen,
3287 J.G., 1990. Anoxic ammonium oxidation. In: C. Christiansen, L. Munck, J.
3288 Villadsen (eds.), Proceedings of the Fifth European Congress on
3289 Biotechnology, vol. 1. Munksgaard, Copenhagen, Denmark, p. 388–391.
- 3290 van de Vossenberg, J., Rattray, J.E., Geerts, W., Kartal, B., van Niftrik,
3291 L., van Donselaar, E.G., Damsté J.S.S., Strous, M., Jetton, M.S.M., 2008.
3292 Enrichment and characterization of marine anammox bacteria associated
3293 with global nitrogen gas production. Environmental Microbiology 10,
3294 3120–3129.
- 3295 van de Vossenberg, J., Woebken, D., Maalcke, W.J., Wessels, H.J.C.T.,
3296 Dutilh, B.E., Kartal, B., Janssen-Megens, E.M., Roeselers, G., Yan, J.,
3297 Speth, D., Gloerich, J., Geerts, W., van der Biezen, E., Pluk, W., Francoijs,
3298 K.-J., Russ, L., Lam, P., Malfatti, S.A., Tringe, S.G., Haaijer, S.C.M., Op
3299 den Camp, H.J.M., Stunnenberg, H.G., Amann, R., Kuypers, M.M.M.,
3300 Jetton, M.S.M., 2013. The metagenome of the marine anammox bacterium
3301 '*Candidatus Scalindua profunda*' illustrates the versatility of this globally
3302 important nitrogen cycle bacterium. Environmental Microbiology 15, 1275-
3303 1289.
- 3304 Van Niftrik, L.A., Fuerst, J.A., Damsté, J.S.S., Kuenen, J.G., Jetton,
3305 M.S.M., Strous, M., 2004. The anammoxosome: an intracytoplasmic
3306 compartment in anammox bacteria. FEMS Microbiology Letters 233, 7–13.
- 3307 van Wonderen, J.H., Burlat, B., Richardson, D.J., Cheesman, M.R., Butt,
3308 J.N., 2008. The nitric oxide reductase activity of cytochrome c nitrite
3309 reductase from *Escherichia coli*. Journal of Biological Chemistry 283, 9587-
3310 9594.

- 3311 Vanitchung, S., Conrad, R., Harvey, N.W., Chidthaisong, A., 2011.
 3312 Fluxes and production pathways of nitrous oxide in different types of
 3313 tropical forest soils in Thailand. *Soil Science and Plant Nutrition* 57, 650-
 3314 658.
- 3315 Velayutham, M., Zweier, J.L., 2013. Nitric oxide signaling in biology.
 3316 *Me*
 3317 *ssenger*, 2, 1-18.
- 3318 Venterea, R.T., Rolston, D.E., 2000. Mechanism and kinetics of nitrite
 3319 and nitrous oxide production during nitrification in agricultural soil. *Global*
 3320 *Change Biology* 6, 303-316.
- 3321 Venterea, R.T., Rolston, D.E., Cardon, Z.G., 2005. Effects of soil
 3322 moisture, physical, and chemical characteristics on abiotic nitric oxide
 3323 production. *Nutrient Cycling in Agroecosystems* 72, 27-40.
- 3324 Vetrovsky, P., Stoclet, J.-C., Entlicher, G., 1996. Possible mechanism of
 3325 nitric oxide production from NG-hydroxy-L-arginine or hydroxylamine by
 3326 superoxide ion. *The International Journal of Biochemistry and Cell Biology*
 3327 28, 1311-1318.
- 3328 Viator, R.J., Rest, R.F., Hildebrandt, E., McGee, D.J., 2008.
 3329 Characterization of *Bacillus anthracis* arginase: effects of pH, temperature,
 3330 and cell viability on metal preference. *BMC Biochemistry* 9, 15.
- 3331 Wagner-Riddle, C., Hu, Q., Van Bochove, E., Jayasundara, S., 2008.
 3332 Linking nitrous oxide flux during spring thaw to nitrate denitrification in the
 3333 soil profile. *Soil Science Society of America Journal* 72, 908-916.
- 3334 Walker, C.B., De La Torre, J.R., Klotz, M.G., Urakawa, H., Pinel, N.,
 3335 Arp, D.J., Brochier-Armanet, C., Chain, P.S.G., Chan, P.P., Gollabgir, A.,

- 3336 Hemp, J., Hügler, M., Karr, E.A., Könneke, M., Shin, M., Lawton, T.J.,
 3337 Lowe, T., Martens-Habbena, W.L., Sayavedra-Soto, A., Lang, D., Sievert,
 3338 S.M., Rosenzweig, A.C., Manning, G., Stahl, D.A., 2010. *Nitrosopumilus*
 3339 *maritimus* genome reveals unique mechanisms for nitrification and
 3340 autotrophy in globally distributed marine crenarchaea. PNAS 107, 8818-
 3341 8823.
- 3342 Wan, Y., Ju, X., Ingwersen, J., Schwarz, U., Stange, C.F., Zhang, F.,
 3343 Streck, T., 2009. Gross nitrogen transformations and related nitrous oxide
 3344 emissions in an intensively used calcareous soil. Soil Science Society of
 3345 America Journal 73, 102-112.
- 3346 Wang, J., Gu, J.D., 2013. Dominance of *Candidatus Scalindua* species in
 3347 anammox community revealed in soils with different duration of rice paddy
 3348 cultivation in Northeast China. Applied Microbiology and Biotechnology
 3349 97, 1785-1798.
- 3350 Wang, P., Zweier, J.L., 1996. Measurement of nitric oxide and
 3351 peroxynitrite generation in the postischemic heart. Evidence for
 3352 peroxynitrite-mediated reperfusion injury. Journal of Biological Chemistry
 3353 271, 29223-29230.
- 3354 Wang, R., Willibald, G., Feng, Q., Zheng, X., Liao, T., Brüggemann, N.,
 3355 Butterbach-Bahl, K., 2011. Measurement of N₂, N₂O, NO, and CO₂
 3356 emissions from soil with the gas-flow-soil-core technique. Environmental
 3357 Science and Technology 45, 6066-6072.
- 3358 Weeg-Aerssens, E., Tiedje, J.M., Averill, B.A., 1987. The mechanism of
 3359 microbial denitrification. Journal of the American Chemical Society 109,
 3360 7214-7215.

- 3361 Weeg-Aerssens, E., Tiedje, J.M., Averill, B.A., 1988. Evidence from
3362 isotope labeling studies for a sequential mechanism for dissimilatory nitrite
3363 reduction. *Journal of the American Chemical Society* 110, 6851-6856.
- 3364 Wei, X.Q., Charles, I.G., Smith, A., Ure, J., Feng, G.J., Huang, F.P., Xu,
3365 D., Muller, W., Moncada, S., Liew, F.Y., 1995. Altered immune responses
3366 in mice lacking inducible nitric oxide synthase. *Nature* 375, 408-411.
- 3367 Wendehenne, D., Durner, J., Klessig, D.F., 2004. Nitric oxide: a new
3368 player in plant signalling and defence responses. *Current Opinion in Plant*
3369 *Biology* 7, 449-455.
- 3370 Wertz, S., Goyer, C., Zebarth, B.J., Burton, D.L., Tatti, E., Chantigny,
3371 M.H., Filion, M., 2013. Effects of temperatures near the freezing point on
3372 N₂O emissions, denitrification and on the abundance and structure of
3373 nitrifying and denitrifying soil communities. *FEMS Microbiology Ecology*
3374 83, 242-254.
- 3375 Wesely, M.L., Hicks, B.B., 2000. A review of the current status of
3376 knowledge on dry deposition. *Atmospheric Environment* 34, 2261-2282.
- 3377 West, M.B., Rokosh, G., Obal, D., Velayutham, M., Xuan, Y.T., Hill,
3378 B.G., Keith, R.J., Schrader, J., Guo, Y., Conklin, D.J., Prabhu, S. D.,
3379 Zweier, J.L., Bolli, R., Bhatnagar, A., 2008. Cardiac myocytespecific
3380 expression of inducible nitric oxide synthase protects
3381 againstischemia/reperfusion injury by preventing mitochondrial
3382 permeability transition. *Circulation* 118, 1970-1978.
- 3383 Williams, D.L.H., 2004. Nitrosation reactions and the chemistry of nitric
3384 oxide. Elsevier, Amsterdam, 280 p.

- 3385 Williams, E.J., Fehsenfeld, F.C., 1991. Measurement of soil nitrogen
3386 oxide emissions at three North American ecosystems. Journal of
3387 Geophysical Research - Atmospheres 96, 1033-1042.
- 3388 Woebken, D., Lam, P., Kuypers, M.M.M., Naqvi, S.W., Kartal, B.,
3389 Strous, M., Jetton, M.S.M., Fuchs, B.M., Amann, R., 2008. A
3390 microdiversity study of anammox bacteria reveals a novel *Candidatus*
3391 *Scalindua* phylotype in marine oxygen minimum zones. Environmental
3392 Microbiology 10, 3106–3119.
- 3393 Wong, J.M., Billiar, T.R., 1995. Regulation and function of inducible
3394 nitric oxide synthase during sepsis and acute inflammation. Advances in
3395 Pharmacology 34, 155-170.
- 3396 Wong, M.W., Nobes, R.H., Bouma, W.J., Radom, L., 1989. Isoelectronic
3397 analogs of molecular nitrogen: Tightly bound multiply charged species. The
3398 Journal of Chemical Physics 91, 2971-2979.
- 3399 Worldometers, 2014. Real time world statistics,
3400 <http://www.worldometers.info> (accepted 2014.05.14).
- 3401 Wrage, N., Velthof, G.L., Beusichem van, M.L., Oenema, O., 2001. Role
3402 of nitrifier denitrification in the production of nitrous oxide. Soil Biology
3403 and Biochemistry 33, 1723-1732.
- 3404 Wu, K.K., 1995. Inducible cyclooxygenase and nitric oxide synthase.
3405 Advances in Pharmacology 33, 179-207.
- 3406 Wu, X., Brüggemann, N., Gasche, R., Shen, Z.Y., Wolf, B., Butterbach-
3407 Bahl, K., 2010. Environmental controls over soil-atmosphere exchange of
3408 N₂O, NO and CO₂ in a temperate Norway spruce forest. Global
3409 Biogeochemical Cycles 24, GB2012.

- 3410 Wullstein, L., Gilmour, C., 1966. Nonenzymatic formation of nitrogen
3411 gas. *Nature* 210, 1150-1151.
- 3412 Yamakura, F., Kawasaki, H., 2010. Post-translational modifications of
3413 superoxide dismutase. *Acta (BBA)-Proteins and Proteomics* 1804, 318-325.
- 3414 Yamasaki, H., Sakihama, Y., 2000. Simultaneous production of nitric
3415 oxide and peroxynitrite by plant nitrate reductase: in vitro evidence for the
3416 NR-dependent formation of active nitrogen species. *FEBS Letters* 468, 89-
3417 92.
- 3418 Yamulki, S., Harrison, R.M., Goulding, K.W.T., Webster, C.P., 1997.
3419 N₂O, NO and NO₂ fluxes from a grassland: effect of soil pH. *Soil Biology*
3420 *and Biochemistry* 29, 1199-1208.
- 3421 Yan, J., Haaijer, S.C., Op den Camp, H.J., van Niftrik, L., Stahl, D.A.,
3422 Konneke, Darci Rush, M., Sinninghe Damsté, J.S., Hu, Y.Y., Jetten,
3423 M.S.M., 2012. Mimicking the oxygen minimum zones: stimulating
3424 interaction of aerobic archaeal and anaerobic bacterial ammonia oxidizers in
3425 a laboratory-scale model system. *Environmental Microbiology* 14, 3146–
3426 3158.
- 3427 Yang, A., Zhang, X., Agogué, H., Dupuy, C., Gong, J., 2014. Contrasting
3428 spatiotemporal patterns and environmental drivers of diversity and
3429 community structure of ammonia oxidizers, denitrifiers, and anammox
3430 bacteria in sediments of estuarine tidal flats. *Annals of Microbiology*, 1-12.
- 3431 Yao, Z., Wu, X., Wolf, B., Dannenmann, M., Butterbach-Bahl, K.,
3432 Brüggemann, N., Chen, W., Zheng, X., 2010. Soil-atmosphere exchange
3433 potential of NO and N₂O in different land use types of Inner Mongolia as

- 3434 affected by soil temperature, soil moisture, freeze-thaw, and drying-wetting
 3435 events. *Journal of Geophysical Research - Atmospheres* 115, D17116.
- 3436 Ye, R.W., Torosuares, I., Tiedje, J.M., Averill, B.A., 1991. (H₂O)-O-18
 3437 isotope exchange studies on the mechanism of reduction of nitric-oxide and
 3438 nitrite to nitrous oxide by denitrifying bacteria e evidence for an
 3439 electrophilic nitrosyl during reduction of nitric oxide. *Journal of Biological*
 3440 *Chemistry* 266, 12848-12851.
- 3441 Yin, S.X., Chen, D., Chen, L.M., Edis, R., 2002. Dissimilatory nitrate
 3442 reduction to ammonium and responsible microorganisms in two Chinese
 3443 and Australian paddy soils. *Soil Biology and Biochemistry* 34, 1131-1137.
- 3444 Yu, J.B., Meixner F.X., Sun W.D., Liang Z.W., Chen Y., Mamtimin B.,
 3445 Wang G.P., Sun, Z.G., 2008. Biogenic nitric oxide emission from saline
 3446 sodic soils in a semiarid region, northeastern China: A laboratory study,
 3447 *Journal of Geophysical Research* 113, G04005.
- 3448 Yu, J.B., Meixner, F. X., Sun, W.D., Mamtimin, B., Xia, C., Xie, W.,
 3449 2010. Biogenic nitric oxide emission of mountain soils sampled from
 3450 different vertical landscape zones in the Changbai Mountains, Northeastern
 3451 China. *Environmental Science and Technology* 44, 4122-4128.
- 3452 Zemojtel, T., Fröhlich, A., Palmieri, M.C., Kolanczyk, M., Mikula, I.,
 3453 Wyrwicz, L.S., Wanker, E.E., Mundlos, S., Vingron, M., Martasek, P.,
 3454 Durner, J., 2006. Plant nitric oxide synthase: a never-ending story? *Trends*
 3455 *in Plant Science* 11, 524-525.
- 3456 Zhang, Y.-Y., Liu, J., Liu, Y.-L., 2004. Nitric oxide alleviates growth
 3457 inhibition of maize seedlings under NaCl stress. *Journal of Plant Physiology*
 3458 *and Molecular Biology* 30, 455.

- 3459 Zhang, Y., Wang, L., Liu, Y., Zhang, Q., Wie, Q., Zhang, W., 2006.
3460 Nitric oxide enhances salt tolerance in maize seedlings through increasing
3461 activities of proton-pump and Na^+/H^+ antiport in the tonoplast. *Planta* 224,
3462 545-555.
- 3463 Zhao, M.-G., Chen, L., Zhang, L.-L., Zhang, W.-H., 2009. Nitric
3464 reductase-dependent nitric oxide production is involved in cold acclimation
3465 and freezing tolerance in *Arabidopsis*. *Plant Physiology* 151, 755-767.
- 3466 Zhao, M.-G., Tian, Q.-Y., Zhang, W.-H., 2007. Nitric oxide synthase-
3467 dependent nitric oxide production is associated with salt tolerance in
3468 *Arabidopsis*. *Plant Physiology* 144, 206-217.
- 3469 Zhou, L., Zhu, D.Y., 2009. Neuronal nitric oxide synthase: Structure,
3470 subcellular localization, regulation, and clinical implications. *Nitric Oxide*
3471 20, 223-230.
- 3472 Zhu, X., Burger, M., Doane, T.A., Horwath, W.R., 2013. Ammonia
3473 oxidation pathways and nitrifier denitrification are significant sources of
3474 N_2O and NO under low oxygen availability. *Proceedings of the National*
3475 *Academy of Sciences USA* 110, 6328-6333.
- 3476 Zhu, B., van Dijk, G., Fritz, C., Smolders, A.J., Pol, A., Jetten, M.S.,
3477 Ettwig, K.F., 2012. Anaerobic oxidization of methane in a minerotrophic
3478 peatland: enrichment of nitrite-dependent methane-oxidizing bacteria.
3479 *Applied and Environmental Microbiology* 78, 8657-8665.
- 3480 Zhu, G., Jetton, M.S., Kusch, P., Ettwig, K.F., Yin, C., 2010. Potential
3481 roles of anaerobic ammonium and methane oxidation in the nitrogen cycle
3482 of wetland ecosystems. *Applied Microbiology and Biotechnology* 86, 1043–
3483 1055.

- 3484 Zhu, G., Wang, S., Wang, Y., Wang, C., Risgaard-Petersen, N., Jetton,
3485 M.S., Yin, C., 2011. Anaerobic ammonia oxidation in a fertilized paddy
3486 soil. *ISME Journal* 5, 1902–1912.
- 3487 Zumft, W.G., 1992. The denitrifying prokaryotes. In: Balows, A., Trüper,
3488 H.G., Dworkin, M., Harder, W., Schleifer, K.-H. (Eds.), *The prokaryotes. A*
3489 *handbook on the biology of bacteria: ecophysiology, isolation,*
3490 *identification, applications, second ed., vol. 1.* Springer-Verlag, New York,
3491 pp. 554–582.
- 3492 Zumft, W.G., 1997. Cell biology and molecular basis of denitrifi cation.
3493 *Micobiology and Molecular Biology Reviews* 61, 533-616.
- 3494 Zumft, W.G., Cardenas, J., 1979. The inorganic biochemistry of nitrogen.
3495 *Bioenergetic processes. Naturwissenschaften* 66, 81-88.
- 3496 Zweier, J.L., Talukder, M.A., 2006. The role of oxidants and free radicals
3497 in reperfusion injury. *Cardiovascular Research* 70, 181-190.
- 3498 Zweier, J.L., Wang, P., Samouilov, A., Kuppusamy, P., 1995. Enzyme-
3499 independent formation of nitric oxide in biological tissues. *Nature medicine*
3500 1, 804-809.
- 3501
- 3502
- 3503
- 3504
- 3505
- 3506
- 3507
- 3508

Figure captions and Tables

Fig. 1. Example of models simulating the temperature effect on nitrification rates (adopted from Stark, 1996). Curves were reconstructed using coefficients for temperature response functions, taken from Stark (1996) [Table 1, p. 440] for open grassy interspaces with the temperature optimum of 35.9 °C.





Fig. 2. Schematic diagram of NO transformations mediated by microbial, enzymatic and chemical processes in soils.

DNRA (dissimilatory nitrate reduction to ammonium); anammox (anaerobic ammonium oxidation); N-AOM (nitrite-dependent anaerobic oxidation of methane); RNS (reactive N species); enzymes: AMO (ammonia monooxygenase); HAO (hydroxylamine oxidoreductase); NAR (membrane-bound nitrate reductase); NAP (periplasmic nitrate reductase); NirK (copper-containing nitrite reductase); NirS (cytochrome *cd1* nitrite reductase); NirB (cytoplasmic nitrite reductase); Nrf (cytochrome *c* nitrite reductase); NrfA (periplasmic cytochrome *c* nitrite reductase); NXR (nitrite oxidoreductase); cNor (nitric oxide reductase that accepts electrons from c-type cytochromes); qNor (nitric oxide reductase that accepts electrons from quinols); NorVW (flavorubredoxin), Hmp, (flavo-hemoglobins); N₂OR (nitrous oxide reductase); HZS complex (hydrazine synthase enzyme complex); HDH (hydrazine dehydrogenase); "NOD" (undefined hypothetical nitric oxide dismutase); NOS (nitric oxide synthase); SOD (super oxide dismutase); XOD (xanthine oxidase); E-NO complex (enzyme (E) bound NO complexes, e.g. E-NO, E-NO⁻ and E-NO⁺, which mediates biotical N-nitrosation).



3544 Table 1. Emission rates and sources of nitric oxide under a range of oxygen
 3545 concentrations (from Russow et al., 2009).

O ₂ content (vol. %)	Emission ($\mu\text{g N kg}^{-1} \text{ h}^{-1}$)	NO formation from ^a		
		NH ₄ ⁺ (%)	NO ₃ ⁻ (%)	NH ₄ ⁺ + NO ₃ ⁻ (%)
20.0	0.92±0.35	70	10	80
2.0	1.16±0.24	26	53	79
0.3	1.90±0.88	1.7	81	83
0	3.71±1.40	0	87	87

3546 ^astandard error of the mean (1 σ), n=6

3547

3548

3549 Table 2. Pathways of NO production in plants (Gupta et al., 2011 and
 3550 reference therein).

Reductive pathways	Oxidative pathways
Nitrate reductase (NR)	Nitric oxide synthase (NOS) -like activity
Plasma membrane-bound nitrite: NO reductase (NiNOR)	Arginine-dependent, polyamine-mediated NO Production
Mitochondrial nitrite reduction	Hydroxylamine-mediated NO production
Xanthine oxidoreductase in plant peroxisomes	

3551

3552

3553 Table 3. Positive and negative effects of NO and its derivatives in
 3554 mammals.

Effect/function	Agent	Location	Reference
Positive effect			
Vasodilation (vascular smooth muscle relaxation)	Formation of NO-sGC or (NO) ₂ -sGC complexes, with releasing of His-105 triggers various cellular signalling pathways (e.g. cGMP formation with further cGK, PDE and iongated channels regulation)	Endothelium eNOS-derived NO could immediately diffuse across the cell membrane to smooth muscle cells	Li and Forstermann, 2000; Derbyshire and Marletta, 2009; Martin et al., 2012
Neurotransmission			
Vasoprotection via inhibiting platelet aggregation			
Stimulating smooth muscle proliferation			
Protection against atherogenesis on its early stages, preventing leukocyte adhesion to the vascular endothelium			
Protective function via cytotoxic effect on intracellular bacteria, cancer cells and tumor tissues	NO-mediated	Activated macrophages	Nathan and Hibbs, 1991; Wei et al., 1995; MacMicking et al., 1997; Forstermann and Sessa, 2012; Rahmanto et al., 2012;
Cardioprotection (e.g. against ischemic and reperfusion injury)	NO-mediated	Cardiocytes	Bolli et al., 2007; West et al., 2008; Granfeldt et al., 2009; Talukder et al., 2010
Antitumor activity	NO-mediated via reduced glutathione (GSH)	multidrug resistance protein (MRP) 1 channel in various cells	Richardson et al., 1995; Li et al., 2011b;
Neuroprotection			
Regulating release of several neuromodulators in the developing retina (e.g. glutamate, gamma-aminobutyric acid (GABA), glutamine, ascorbate)	NO as an atypical retinal messenger	Retina	Ientile et al., 1996; Maggesissi et al., 2009; Portugal et al., 2012
Negative effect			
Cytotoxicity (e.g. reaction with proteins and nucleic acids), leading to apoptosis and cell death	Overproduction of NO	Various types of cells	Boje and Arora, 1992; Dimmeler and Zeiher, 1997; Kroncke et al., 1997; Gotoh and

			Mori, 2006; Erusalimsky and Moncada, 2007; Forstermann and Sessa, 2012;
Attenuation of energy production by inhibiting mitochondrial respiration and glycolysis	Overproduction of NO	Mitochondria and cytoplasm	Erusalimsky and Moncada, 2007; Brown, 2010;
Neurodegenerative disorders and cerebral infarction	Overproduction of NO by activated macrophages or microglia cells	Neurons	Chao et al., 1992; Kroncke et al., 1997; Ignarro, 2009;
Septic shock due to vasodilation and hypotension	Overproduction of NO	Vascular system	Wong and Billiar, 1995; Lange et al., 2009;
Pathogenesis of Type I diabetes due to NO induced islet cell death	Overproduction of NO	Endocrine system	Oyadomari et al., 2002
Apoptosis due to eliminating Ca^{2+} from endoplasmic reticulum	Overproduction of NO	Pancreatic β - cells	Oyadomari et al., 2001, 2002
Damaging DNA, proteins and lipids	NO-mediated oxidative reaction products (e.g. ONOO^-)	Various types of cells	Lee et al., 2003; Mikkelsen and Wardman, 2003; Ridnour et al., 2004
Brain pathology		Neurons	Brown and Neher, 2010
Myocardial injury		Cardiocytes	Wang and Zweier, 1996; Zweier and Talukder, 2006