

Article (refereed) - postprint

Granroth-Wilding, H.M.V.; Daunt, F.; Cunningham, E.J.A.; Burthe, S.J. 2017.
Between-individual variation in nematode burden among juveniles in a wild host.

This version available <http://nora.nerc.ac.uk/515928/>

NERC has developed NORA to enable users to access research outputs wholly or partially funded by NERC. Copyright and other rights for material on this site are retained by the rights owners. Users should read the terms and conditions of use of this material at <http://nora.nerc.ac.uk/policies.html#access>

This is the author's final manuscript version of the journal article, incorporating any revisions agreed during the peer review process.

This article has been published in a revised form in *Parasitology* (2017), 144(2): 248-258 <http://dx.doi.org/10.1017/S0031182016001700>. This version is free to view and download for private research and study only. Not for re-distribution, re-sale or use in derivative works. © Cambridge University Press 2016.

There may be differences between this version and the publisher's version. You are advised to consult the publisher's version if you wish to cite from this article.

The definitive version is available at <http://www.cambridge.org>

Contact CEH NORA team at
noraceh@ceh.ac.uk

1 **Between-individual variation in nematode burden among juveniles in a wild host**

2

3 Granroth-Wilding H.M.V.^{1 + #}, Daunt F.², Cunningham E.J.A.^{1*} & Burthe S.J.^{2*}

4

5 1. Wellcome Centre for Infection, Immunity and Evolution/Institute of Evolutionary Biology,
6 University of Edinburgh, Ashworth Laboratories, Charlotte Auerbach Road, Edinburgh EH9 3FL,
7 UK

8 2. NERC Centre for Ecology & Hydrology, Bush Estate, Penicuik EH26 0QB, UK

9 ⁺ Corresponding author: Hanna Granroth-Wilding; hanna@granroth-wilding.co.uk

10 [#] Current address: Department of Biosciences, University of Helsinki, Viikinkaari 1, P.O. Box 65
11 00014 Helsinki, Finland

12 ^{*}Equal author contributions

13

14

15 **Running title:** Variation in nematode burdens of juvenile birds

16

17

18 ABSTRACT

19 Parasite infection in young animals can affect host traits related to demographic processes such as
20 survival and reproduction, and is therefore crucial to population viability. However, variation in
21 infection among juvenile hosts is poorly understood. Experimental studies have indicated that
22 effects of parasitism can vary with host sex, hatching order and hatch date, yet it remains unclear
23 whether this is linked to differences in parasite burdens. We quantified gastrointestinal nematode
24 burdens of wild juvenile European shags (*Phalacrocorax aristotelis*) using two *in situ* measures
25 (endoscopy of live birds and necropsy of birds that died naturally) and one non-invasive proxy
26 measure (faecal egg counts). *In situ* methods revealed that almost all chicks were infected (98%),
27 that infections established at an early age, and that older chicks hosted more worms, but faecal egg
28 counts underestimated prevalence. We found no strong evidence that burdens differed with host sex,
29 rank or hatch date. Heavier chicks had higher burdens, demonstrating that the relationship between
30 burdens and their costs is not straightforward. *In situ* measures of infection are therefore a valuable
31 tool in building our understanding of the role that parasites play in the dynamics of structured
32 natural populations.

33

34 **Keywords:** Parasite burden, endoscope, dissection, *Contracaecum*, anisakid, seabird,
35 macroparasite, prevalence, FEC, demographic trait, growth, host-parasite interaction

36 KEY FINDINGS

37

- 38 • We quantified nematode burdens of seabird chicks using necropsy, endoscopy and FECs
- 39 • *In situ* techniques showed 98% prevalence, early establishment and higher burdens with age
- 40 • Faecal egg counts, a proxy measure, underestimated prevalence
- 41 • Chicks with higher burdens weighed more, contrary to expectations if infection is costly
- 42 • Endoscopy of juveniles enables monitoring of wild hosts' infections across their lifetime

43 INTRODUCTION

44

45 The costs that parasite infection can impose on their hosts can influence key demographic traits,
46 such as reproductive success and survival, which are crucial to the growth rate and hence viability
47 of populations (Albon et al., 2002; Newey et al., 2005; Redpath et al., 2006, Tompkins 2011).
48 However, parasitism is unlikely to affect all individuals in a population in the same way. Firstly,
49 individuals may host different burdens as a result of differences in exposure to parasites,
50 susceptibility to infection and resistance to its impacts. This contributes to parasite abundance
51 typically showing a skewed distribution among hosts, which is particularly well documented in
52 macroparasites (Randolph et al., 1999; Shaw and Dobson, 1995). Secondly, once parasitized, the
53 relationship between parasite load and host fitness may vary between individuals due to differences
54 in tolerance for a given parasite load. Siblings, for example, may vary in the level of maternal
55 antibodies they receive (Pihlaja et al., 2006), males may be affected more than females due to
56 immunosuppressive effects of testosterone (Mougeot et al., 2009), and the relative benefits of
57 allocating resources between fighting infection and reproduction may vary with age (Adamo et al.,
58 2001). These factors may lead to different types of host responding differently to infection, with
59 consequences for key host traits related to fitness such as weight gain during critical periods of
60 growth. Understanding how parasite burdens and their impacts on hosts vary between different
61 classes of individual may therefore be crucially important for understanding the impacts of parasites
62 on heterogeneous host populations.

63 A key process for population viability is the level of offspring recruitment to the breeding
64 population. Understanding how parasitism impacts on the juvenile subset of the population is
65 therefore important for modelling population growth. Infection in early life can alter juveniles'
66 developmental trajectories (Fitze et al., 2004; Romano et al., 2011), with potentially life-long fitness
67 consequences that may further influence demographic processes such as reproduction and survival
68 long after recruitment (Lindström, 1999; Metcalfe and Monaghan, 2001; Monaghan, 2008).
69 However, despite the importance of early-life infection, between-individual patterns of parasitism
70 and the development of infections in juvenile hosts have not been widely investigated in the wild.
71 Although young hosts have been shown to exhibit systematic between-individual differences in
72 their response to experimental infection or anti-parasite treatments according to characteristics such
73 as hatching order (Granroth-Wilding et al., 2014), sex (Romano et al., 2011) or timing of breeding
74 (Reed et al., 2008), it remains unclear whether variation in response is associated with differences in

75 burden or differences in tolerance. Thus, quantifying individual parasite burdens across the juvenile
76 component of the population, where individuals' responses to infection are also known by
77 measuring key fitness-related traits, is central to accurately predicting parasite impacts at the
78 population level.

79 Quantifying parasite burdens is logistically challenging in the wild, particularly for
80 endoparasites that often make up the majority of a host's parasite biomass (Hoberg, 2005). Necropsy
81 allows direct counts of parasites in the host and is widely considered to give the most reliable
82 measure. However, this destructive method prevents longitudinal studies, which are crucial for
83 detecting associated fitness consequences such as recruitment probability and future reproductive
84 success (Fitze et al., 2004). In juvenile hosts, such sublethal impacts typical of macroparasites have
85 the potential to affect key demographic parameters over a range of timescales; avoiding destructive
86 sampling is therefore particularly important to understand the full fitness consequences of infection
87 in young hosts. Moreover, necropsy may not be viable for hosts of conservation importance. Faecal
88 egg counts (FECs) are a common non-destructive and non-invasive proxy measure of endoparasite
89 burden (e.g. Bowman and Georgi, 2009; Craig *et al.* 2006; Seivwright *et al.* 2004), but may not
90 always reflect true parasite burden due to variable rates of helminth egg production (Shaw and
91 Moss, 1989; Tompkins and Hudson, 1999), poor sensitivity at low burdens (Levecke *et al.* 2009),
92 and not representing larval helminths that do not produce eggs but can nonetheless be costly to the
93 hosts (Fagerholm and Overstreet, 2008). Recent work in wild adult seabirds has pioneered
94 endoscopy as an additional, direct and reliable method to obtain an index of gastrointestinal
95 nematode burdens in live individuals (Burthe et al., 2013). This approach has great potential for
96 quantifying the development of an individual's infection from an early age, but has not previously
97 been applied to juveniles in the wild.

98 Here, we use two *in situ* measures of parasite burden – necropsy and endoscopy – and the
99 proxy measure of FECs to quantify patterns of between-individual variation in the trophically-
100 transmitted gastrointestinal nematode burden of juvenile European shags (*Phalacrocorax*
101 *aristotelis*, henceforth “shag”), a piscivorous seabird. Experimental manipulations of parasite load
102 in adults and chicks has shown that responses to treatment vary with host phenotype: treatment of
103 parents increases male chicks' survival, particularly late in the season, but not female chicks' (Reed
104 et al., 2008); treatment of chicks generally affects the growth rate of last-hatched siblings but not
105 the older brood members (Granroth-Wilding et al., 2014); and the impact of simultaneous treatment
106 of parents and their offspring differs between early- and late-nesting families (Granroth-Wilding et

107 al. 2015). Endoscopy of adults has found males to host more worms than females and late breeders
108 more than early breeders (Burthe et al., 2013), but among juveniles, patterns of variation in parasite
109 abundance and their link with variation in host fitness are not well quantified. It is hence unclear
110 whether these differences in treatment responses between types of juveniles arise from differences
111 in nematode burden or differences in the impact of a similar burden. Moreover, the link between
112 parasite burden and demographically important host traits is unexplored. Our objectives were
113 therefore to: 1. quantify individual parasite burdens of juveniles using two *in situ* methods,
114 endoscopy and necropsy, and compare these to a proxy measure of prevalence based on FECs; 2.
115 identify whether burdens vary with host age, sex, hatching order and hatch date; 3. examine whether
116 natural variation in parasite abundance is associated with a fitness-related trait, host mass.

117

118 METHODS

119

120 *Host-parasite system*

121 This study was carried out in 2012 in the breeding population of shags on the Isle of May National
122 Nature Reserve in south-east Scotland (56°11 N, 2°33 W) that has been the subject of an individual-
123 based long-term demographic study for several decades. Shags are sexually dimorphic, with males
124 growing faster to reach an adult size c. 20% bigger than females (Daunt *et al.* 2001). The modal
125 clutch size in this population is three eggs and these hatch asynchronously, with the second and
126 third siblings (B and C chicks) hatching on average 1 and 2-3 days after the first (A chick). This
127 asynchrony results in a hierarchy of size within the brood, in which youngest siblings generally
128 grow more slowly and have higher mortality but are more plastic in response to changing
129 environmental conditions than their older counterparts (Granroth-Wilding *et al.* 2014; Stokland and
130 Amundsen, 1988). Breeding success declines through the season, with later breeders fledging fewer
131 chicks and producing fewer recruits (Daunt *et al.* 1999; Harris *et al.* 1994).

132 Shags on the Isle of May are infected with the gastrointestinal nematode *Contracaecum*
133 *rudolphii* (Anisakidae: Ascaridoidea; Hartwich 1964), which occur in the GI tract of nestling and
134 adult shags in this population (Reed 2007; Burthe *et al.* 2013; E. Harris, pers. comm.; S. Burthe, J.
135 Chantrey & D. Kowalek, unpublished data). All but one of 146 naturally infected adults endoscoped
136 to date have hosted worms, with wide variation in burdens from 2 to >80 worms (Burthe *et al.*
137 2013; S. Burthe & E. Butterfield, unpublished data). *C. rudolphii* is a widely distributed seabird
138 specialist, now recognised to comprise a complex of morphologically similar species (Anderson,

2000; Fagerholm and Overstreet, 2008; Hoberg, 2005; Moravec, 2009). Nestling shags obtain regurgitate fish directly from their parents' throats and are infected with larval worms in the fish tissue. Direct infection of chicks with adult worms dislodged from the parent's proventriculus could also occur during feeding, but the importance of this transmission route is not well understood (Dubinin, 1949; Fagerholm and Overstreet, 2008; Hoberg, 2005; Huizinga, 1971). Anisakid infection can cause costly pathology at attachment sites such as inflammation, necrosis, haemorrhaging and perforation of the stomach wall (Hoberg, 2005; Kuiken, 1999; McClelland, 2005), which may be compounded by secondary bacterial infections (Fagerholm and Overstreet, 2008), and is expected to activate a costly immune response (Colditz, 2008; Hasselquist and Nilsson, 2012). Moreover, *Contracaecum* is thought to feed on fish ingested by the bird and thus directly competes with the host for resources (Abollo *et al.* 2001; Anderson, 2000; Dubinin, 1949; Huizinga, 1971).

151

152 ***Quantifying nematode burdens***

We quantified the nematode burden of individual shag chicks using two *in situ* techniques, endoscopy of targeted individuals or necropsy (dissections) of a subset of the study population that died naturally during a severe storm. We also conducted faecal egg counts (FECs) on faecal material collected opportunistically from both endoscoped and dissected chicks (all detailed methods below). Not all individuals produced faecal samples, precluding FECs, and no birds were both endoscoped and dissected, as endoscoped chicks were not sacrificed and endoscopy of dead animals is not reliable (S. Burthe, unpublished data).

160

161 ***Endoscopy***

We used a refurbished 9mm diameter medical endoscope (Olympus®, UK) to view the oesophagus and proventriculus of conscious chicks under licence (full details of endoscopy procedure in (Burthe *et al.* 2013). Endoscopy was undertaken by a trained and experienced operator (S. Burthe) while an assistant held the bird still and its bill open. A cloth was placed over the bird's eyes to reduce stress while the endoscope operator inserted the endoscope into the proventriculus. All worms that were visible were counted as the endoscope was slowly withdrawn from the bird. We noted whether the worms were large or small. Visibility was scored on a scale of 1 to 5 (worst to best, as in Burthe *et al.* (2013)) and included in all analyses as poorer visibility could hinder accurate quantification. Endoscopy was carried out when chicks were large enough for the

171 endoscope to be comfortably inserted, around 25 days of age. Throughout the process, there was no
172 evidence of discomfort (e.g. rapid breathing). All endoscoped chicks resumed normal behaviour
173 immediately on being returned to the nest and all fledged successfully. All endoscopy was carried
174 out early in the morning, before parents had returned with the first food load of the day, to avoid
175 views being obstructed by recently ingested food.

176 At endoscopy, chicks were assigned a rank in the brood hierarchy according to size: in broods
177 of three, the heaviest two chicks were designated AB and the lightest C, and in broods of one or
178 two, all chicks were designated AB. Wing length was used as an additional indicator if mass
179 difference was not greater than 20g. Mass at day 25-30 accurately identifies the last-hatched chick
180 in 83% of cases but only distinguishes the first- and second-hatched (A and B) in 47% of cases,
181 whereas A and B are accurately assigned as AB in 89% of cases (data from 27 nests, with three
182 chicks surviving to day 10, in 2010 and 2011 with accurate hatch dates; Granroth-Wilding *et al.*
183 2014). We used chick mass at endoscopy as an indicator of chick performance. At endoscopy age,
184 the majority of growth is completed (Daunt *et al.* 2001), and fledging mass has been shown to
185 correlate with recruitment in a range of species (Magrath, 1991; Schwagmeyer and Mock, 2008).
186 All endoscoped chicks were blood sampled for molecular sexing (Griffiths *et al.* 1996).

187 In total, we endoscoped 45 chicks in 20 nests, of which 18 were undisturbed before
188 endoscopy and 27 were sham-treated controls from a parallel parasite-removal experiment (full
189 details in Supplementary Information; no individuals treated with anti-parasite drugs are included in
190 the results presented here), injected with 0.05ml saline solution at age 10-12 days and subsequently
191 weighed at ages 10, 15 and 25 days. A subset of chicks that remained safely accessible as they got
192 older and more mobile were endoscoped twice (2 untreated chicks and 4 sham-treated).

193

194 *Necropsy*

195 Sacrificing individuals for systematic necropsies was not possible as this would prevent
196 longitudinal investigations of the link between parasite burden and host survival, and moreover
197 removing individuals from this long-term study population is not desirable. However, in 2012, there
198 was an unusually prolonged period of rain and cold weather in the middle of the peak chick-rearing
199 period, lasting over two days. This caused considerable natural juvenile mortality due to
200 waterlogging and chilling of chicks that were still downy (not yet waterproof) but too large to be
201 efficiently sheltered by their parents. Mortality was thus not a direct consequence of overall poor
202 condition nor of parasitism, though both factors may have contributed. Similar weather-related mass

203 mortality events of chicks during the breeding season have only occurred once in the last 15 years,
204 so this was a rare opportunity to obtain a sample of birds for dissection. When the weather
205 improved and it was safe to approach nests, c. 12-36 hours after death, we collected 28 carcasses
206 (median 20 days old, interquartile range 18-26 days; median hatch date 27th May, IQR 21st-29th
207 May; cf. endoscoped chicks, median age 31 days, IQR 28-34 days, median hatch date 17th May,
208 IQR 15th-22nd May). Nine of these were sham-treated controls from the parallel experiment. We also
209 collected 6 further carcasses resulting from other natural mortality, found within a day of death, for
210 necropsy (median age 25 days, IQR 25-29 days; median hatch date 2nd June, IQR 20th May-3rd
211 June). For the 10 dissected chicks that were not of known age, we estimated age from wing length
212 based on the growth rate of chicks from the same year with known hatch dates ($\text{Wing} = 5.81 \times \text{Age}$
213 $- 27.75$; in mixed model accounting for repeated measures within chick, $F_{1,147} = 9795$, $p < 0.001$;
214 without random effects, $r^2 = 0.954$). We assigned ranks to dissected chicks in cases where the whole
215 brood could be assessed either dead or alive, based on the structural measure of wing length: in
216 broods of three, the two chicks with longest wings were assigned AB and the shortest C, and in
217 broods of one or two, all chicks were assigned AB. A sample of blood or tissue was taken from
218 every carcass for molecular sexing (Griffiths *et al.* 1996).

219 Where possible, carcasses were dissected fresh within 6 hours of recovery, or kept at +4°C
220 for up to 24 hours (16 carcasses). If dissections could not be carried out within this time (17
221 carcasses), they were stored at -20°C for up to one week and defrosted before dissection. The
222 proventriculus was removed together with 3cm of oesophagus and small intestine. The removed
223 gastrointestinal portion was then opened out using one medial ventral cut and the stomach contents
224 thoroughly examined, then rinsed with water through a fine mesh. The body cavity was examined
225 for evidence of nematodes migrating away from the proventriculus following host death, and we
226 additionally examined the whole intestine of four individuals; no other visible macroparasites were
227 found (further descriptions in Supplementary Information). All worms were counted, removed and
228 stored in ethanol. To obtain an index of the maturity of the infection in the bird, during which stage
229 *Contracaecum* undergoes substantial growth (Fagerholm and Overstreet, 2008), worms were
230 categorized into size classes based on width ($>0.75\text{mm}$ wide, large; $<0.5\text{mm}$ wide, small; 159 out of
231 1436 worms (11%) in between the categories).

232

233 *Faecal egg counts (FECs)*

234 During endoscopy, we opportunistically collected faecal samples from 19 chicks that defecated

235 during handling, and from 24 dissected chicks, we obtained a faecal sample from the cloaca after
236 carcasses had been frozen at -20°C for long-term storage. All faecal samples were therefore stored
237 at -20°C after collection; previous work in this system has given no evidence that freezing affects
238 egg counts or prevalence (Supplementary Information). FECs were carried out using a flotation
239 technique (Bowman and Georgi, 2009; detailed methods in Supplementary Information). Each
240 sample was suspended in 20 ml saturated salt solution per 1 g of faeces and nematode eggs were
241 counted in 0.45 ml (0.02 g faeces) of the suspension examined under a McMaster slide.

242

243 *Statistical analysis*

244 We first quantified patterns in parasite abundance obtained by each *in situ* parasite measure,
245 endoscopy and dissection. We considered two aspects of nematode infection: total worm burden,
246 indicating overall parasite abundance, and the proportion of worms that were large, which is likely
247 to reflect the duration of the infection. We then tested whether these indices were associated with
248 host age, as expected if chicks are exposed to infective larvae throughout their development, and
249 with phenotypic traits known to affect responses to infection: host sex, rank (AB vs. C) and hatch
250 date (Granroth-Wilding *et al.* 2014; Reed *et al.* 2008, 2012). Lastly, in endoscoped chicks, we
251 examined the association between parasite abundance and chick performance by testing whether
252 chick mass at endoscopy varied with worm count and the proportion of worms that were large.

253 In all analyses of dissected chicks, we excluded two outliers with high statistical leverage:
254 one old chicks with a very high load (a male, 45 days old, hosting 243 worms; range of other chicks
255 8-148 worms) and one very young chick (ca. 2 days old) which was the only dissection that yielded
256 a zero burden. Neither exclusion qualitatively affected any results. Although mortality is generally
257 higher for C chicks in this population (Granroth-Wilding *et al.* 2014), all ranks were equally
258 represented among endoscoped and dissected birds, as were males and females (for ranks across
259 techniques, $\chi^2 = 4.50$, $df = 2$, $p = 0.105$; for sexes, $\chi^2 = 1.32$, $df = 1$, $p = 0.251$). Among endoscoped
260 chicks, we confirmed that visibility score was not related to age, sex, rank or hatch date (all $p > 0.4$
261 in a linear model). Among endoscoped chicks, hatch date (from which age was calculated) was only
262 available for the first-hatched chicks in each nest, so C chicks were assigned an age 2.5 days
263 younger than their AB siblings (median age difference across 42 nests in 2010 and 2011 with
264 accurate hatch date data) to avoid within-brood age differences confounding rank effects. Among
265 dissected chicks, the effects of age and hatch date could only be examined in separate models as the
266 age-specific main mortality event meant that they were closely correlated (in linear model, $r^2 = 0.72$,

267 $p < 0.001$). In these analyses, models containing hatch date instead of age gave almost identical fits
268 ($\Delta AICc \leq 0.1$) and for brevity we present only the age models.

269 All analysis was carried out in R 3.0.2 (R Core Team, 2013) using the packages lme4 (Bates
270 *et al.* 2011) and nlme (Pinheiro *et al.* 2012), using (generalised) linear mixed models (LM Ms or
271 GLMMs). To account for repeated sampling of some individuals and non-independence of siblings
272 within a brood, we fitted chick within nest as nested random factors to the endoscopy data, and nest
273 as a random factor to the dissection data. Total burden was fitted as count data with poisson errors
274 and logistic link function, and proportion of large worms with binomial errors, weighted by the total
275 count, and a logit link function. Effect sizes for the proportion of large worms are presented as the
276 log odds of a worm being large. Mass was modelled in a linear mixed model including log(age) and
277 sex as fixed effects, to account for the non-linear growth curve and sexually dimorphic growth. Due
278 to the low egg prevalence in faeces preventing robust analysis of relationships between FECs and
279 host phenotypic traits or *in situ* worm burdens, we present only descriptive statistics of prevalence
280 (but see Supplementary Information for a preliminary analysis).

281 We used an information theoretic approach to model selection (Burnham and Anderson,
282 2002), identifying important explanatory variables based on the best-fitting model(s) from a
283 candidate set, which is well suited to an exploratory analysis. For each measure of parasite burden,
284 our set of candidate models contained all combinations of the explanatory variables as main effects
285 (age, hatch date, sex and rank, and additionally for endoscopy analyses, visibility) as well as an
286 intercept-only (null) model. The best-fit model was the one that had the lowest AICc (corrected
287 Akaike's Information Criterion, suitable for small sample sizes) in the set, and models with a $\Delta AICc$
288 ≤ 2 from the best fit model were considered an equivalent fit. Model selection based on significance
289 testing gave the same conclusions. All parameters are presented ± 1 standard error, not back-
290 transformed from the log (worm counts) or logit (proportion of large worms) link functions.

291 RESULTS

292 *Quantifying worm burden in situ*

293 The ages of birds available for necropsy ranged from 2 to 45 days and for endoscopy from 25 to 49
294 days. Worm burden measured using necropsy varied from 0 to 243 worms per chick; the youngest
295 and oldest chicks were excluded from further analysis due to their strong leverage, giving an age
296 range of 12–31 days and worm counts of 8 to 148 worms per chick ($n = 31$; mean 36.0 ± 4.9 ;
297 prevalence 100%) (fig. 1). Worm burden measured using endoscopy ranged from 0 to 30 worms
298 per chick (mean worm burden 11.7 ± 1.0 worms; prevalence 98%) (fig. 1). The proportion of large
299 worms ranged from 0 to 35.7% (mean $12.9 \pm 1.9\%$) by necropsy and 0 to 100% (mean $29 \pm 5\%$) by
300 endoscopy.

301 Using necropsy, the youngest chick to host large worms was aged 15 days and the oldest
302 chick without large worms was 18 days. Using endoscopy, large worms were found in chicks from
303 the age of 26 days, (earliest available age 25 days), although chicks with no large worms occurred
304 up to the age of 36 days.

305 Visibility during endoscopy was generally poorer for chicks than for adults endoscoped in
306 parallel studies, mainly due to the presence of semi-digested food. Visibility scores among the
307 chicks in this study ranged from 1 to 4 (mean 2.7 ± 0.1) compared to a range of 3–5 (mean 4.24;
308 $n=17$) for adult shags endoscoped in the same year (S. Burthe, unpublished data).

309

310 *FECs as an indicator of worm burden*

311 We obtained faecal egg counts from 19 endoscoped and 24 dissected chicks from birds aged 25–36
312 days and 12–45 days respectively. Nematode eggs were only found in one third of the 43 samples
313 available (prevalence 37%), despite a prevalence of 99% in individuals sampled using *in situ*
314 measures. Out of the 16 faecal samples that contained worm eggs, only 7 contained more than 1 egg
315 (4 samples with 2 eggs, 2 with 3 eggs and one with 42) and 5 were from chicks in which no large
316 worms were seen (1 necropsy, 4 endoscopies).

317

318

319 *Nematode burden in relation to host traits*

320 In necropsied chicks, aged 12–31 days, worm count was best explained by a model containing
321 only age, with older chicks hosting more worms. A model with age and sex had similar support
322 (table 1, fig. 2). The proportion of worms that were large was best explained by an intercept-only

323 model, with no host traits providing similar explanatory power (table 1, fig. 3).

324 Among endoscoped chicks, aged 25–49 days, total worm burden was best described by a
325 model containing age and visibility (table 1, fig. 2), with older chicks hosting more worms and
326 better visibility resulting in slightly higher worm counts (age effect size 0.10 ± 0.02
327 $\log(\text{worms})/\text{day}$, visibility effect size $0.10 \pm 0.06 \log(\text{worms})$ per score increment). Age was
328 supported in all five top models. Out of three equivalent-fit models, two contained a rank term (in
329 addition to age, C chicks hosted -0.21 ± 0.16 fewer worms than AB chicks). The proportion of large
330 worms was best described by a model containing only age (effect size 0.11 ± 0.04 increase in
331 proportion of large worms/day) (fig. 3), with hatch date and rank each occurring twice in the three
332 equivalent-fit models (in addition to age, effect of hatch date: 0.05 ± 0.00 greater proportion of large
333 worms per day; effect of rank: C chicks 0.83 ± 0.50 greater proportion of large worms) (table 1).

334 A summary of the host traits identified as important to parasite abundance and size
335 distribution by the two measurement techniques is given in table 2. For both necropsy and
336 endoscopy, it is notable that individuals varied considerably in their parasite load, which contributed
337 to many analyses yielding several equivalent-fit models that made it difficult to robustly identify
338 phenotypic traits that influenced parasite load.

339

340 *Effect of infection on host performance*

341 Chick mass at endoscopy was best explained by a model containing main effects of age and worm
342 count (table 3, fig. 4): heavier chicks were older and had higher worm counts (in best-fit model,
343 effect of age $241.4 \pm 43.2 \text{ g}/\log(\text{day})$; effect of worm count, $11.8 \pm 4.8 \text{ g}/\text{worm}$). There was one
344 model of equivalent fit, which contained an additional sex term (males $62.4 \pm 46.6 \text{ g}$ heavier than
345 females).

346

347

348 DISCUSSION

349 The juvenile period is an energetically expensive phase for an individual when the costs associated
350 with parasite infection are likely to have substantial impacts on hosts. Despite this, in comparison to
351 adults, there is very little information for wild juvenile hosts on patterns of parasite prevalence or
352 abundance, particularly internal parasites. Here we have used necropsy and endoscopy,
353 implemented for the first time in juveniles in the wild, to show that infection with gastrointestinal
354 nematodes is near-universal among nestling shags (98% prevalence) and establishes at an early age,

355 and that nematode burden increases with chick age. In contrast, the common proxy measure of
356 FECs suggested a prevalence of only 37%, demonstrating the value of endoscopy as a non-
357 destructive index of *in situ* parasite burden. Previous studies have found chick sex, hatch date and
358 rank to be important in determining responses to anti-parasite treatment (Reed *et al.* 2008, 2012;
359 Granroth-Wilding *et al.* 2014, 2015), yet we found no strong evidence that worm burden varied
360 with any of these host traits. This suggests that differences in response may arise due to variation in
361 tolerance between the subclasses of juvenile as opposed to differences in burden. Further, contrary
362 to predictions, we found that individuals with high worm burdens were heavier than similar-aged
363 individuals with lower burdens.

364

365 *Comparison of techniques for quantifying worm burden*

366 Both necropsy and endoscopy captured the same main pattern of infection in chicks but
367 unfortunately we did not have the opportunity to directly compare counts from the two techniques
368 in the same individuals. None of the birds that suffered natural mortality had been endoscoped,
369 endoscoped chicks could not be sacrificed for necropsy as this would prevent long-term monitoring
370 of infection and its consequences, and endoscopy of carcasses is not feasible as reliable counts are
371 difficult to obtain from the collapsed stomach of a dead bird. Comparisons of necropsied and
372 endoscoped individuals was further constrained by the limited overlap in the ages of chicks used in
373 each technique: endoscopy was carried out on generally older birds and tended to yield lower
374 overall burdens but a higher proportion of large worms than necropsies of generally younger birds.
375 Endoscopy may have yielded lower counts because chicks' stomachs frequently contained residual
376 food that partially obscured the view through the endoscope, a constraint that is more easily avoided
377 when endoscoping adults in this system (Burthe *et al.* 2013). Nonetheless, endoscopy counts from
378 shags have been shown to be repeatable (Burthe *et al.* 2013), and our successful application of this
379 technique to developing hosts thus opens opportunities for monitoring individuals' worm burdens
380 from an early stage in their long lives. Moreover, both *in situ* techniques identified similar
381 prevalences and an increase in burden with chick age, indicating that endoscopy provides a useful
382 index of between-individual variation in worm burdens. This index has already been shown to be
383 valuable for quantifying the effect of anti-parasite treatment in both adults and juveniles, even at
384 low doses (Burthe *et al.* 2013; Supplementary Information, this study).

385 Necropsy, on the other hand, allows complete examination of the gut of the animal at any age
386 and is likely to yield more accurate counts. However, as a destructive sampling technique, necropsy

387 is of limited application because removing individuals from the population is not desirable when
388 investigating longitudinal effects of parasite infection or working with protected natural
389 populations. In such cases, obtaining samples relies on natural mortality that may more strongly
390 affect certain parts of the population, such as those already paying the costs of a high parasite
391 burden. Moreover, necropsy of recovered carcasses may underestimate infection intensity due to
392 post-mortem migration of nematodes away from attachment sites. Given that the endoscope counts,
393 also likely underestimates, captured the same broad patterns of infection as necropsy, we suggest
394 that endoscopy provides an informative non-destructive index, albeit not true counts, of between-
395 host variation in total parasite burden. The repeated measurement of an index of infection intensity
396 across individuals' lives that this enables, while also allowing quantification of its long-term
397 consequences for host fitness, is likely also to be of practical use in other systems.

398 Measuring long-term patterns in individuals' parasite burdens could potentially be made more
399 logistically tractable if a non-invasive proxy for worm burden was available, such as FECs.
400 However, in our system, FECs failed to detect the same levels of infection revealed by *in situ*
401 measures. Although worms were found in 98% of all chicks examined, the majority of faecal
402 samples (63%) did not contain eggs, and faecal egg presence was not related to *in situ* worm burden
403 (Supplementary Information). This may be due in part to chicks hosting a high proportion of worms
404 that were small, likely immature and thus not egg-producing individuals. Variation in this
405 component of the parasite community may nonetheless be important for its impacts on host fitness,
406 as larval worms can still cause severe pathology and thus have non-negligible costs (Fagerholm and
407 Overstreet, 2008; H.-P. Fagerholm, pers. comm.). The limited presence and low counts of nematode
408 eggs in host faeces in this system appears to be a feature of this system, but we cannot rule out that
409 FECs more closely reflecting natural variation in true burdens could be obtained by examining
410 larger amount of faecal material (but see Supplementary Information), which is logistically difficult
411 in the field.

412

413 *Nematode burden in relation to host traits*

414 The positive relationship between worm burden and chick age is consistent with expectations that
415 chicks' infections should intensify throughout the nestling period. This increase suggests that chicks
416 are continuously exposed to either infective larvae in fish and/or adult worms dislodged from the
417 parent's proventriculus during feeding. Continuous exposure among chicks accords with the near-
418 universal prevalence of worms among endoscoped adult shags over 6 study years (S. Burthe,

419 unpublished data), which in turn indicates regular exposure to infected fish (Anderson, 2000;
420 Fagerholm and Overstreet, 2008; Huizinga, 1971). Two further observations can also be interpreted
421 as indicative both of larval worms establishing and growing inside the chick and of ongoing direct
422 transmission of larger worms from the parent's proventriculus: the increase in the proportion of
423 large worms with age in endoscoped chicks, and the presence of nematode eggs in the faeces of a
424 12-day-old chick (lowest estimates for maturation time of larval *C. rudolphii* in the definitive host,
425 c. 1 week; Dubinin, 1949; Huizinga, 1971). Regardless of transmission mechanism, we found
426 established nematode infections in all chicks from early on in their period of rapid growth (from 6-9
427 days; Daunt *et al.* 2001). This supports the potential of parasitism in juvenile shags to influence
428 developmental trajectories and hence long-term performance and fitness in this long-lived species
429 (Lindström 1999; Monaghan 2008).

430 Previous studies have found host sex, timing of breeding and hatching order to be important
431 in shaping individual chicks' responses to anti-parasite treatment (Granroth-Wilding *et al.* 2014,
432 2015; Reed *et al.* 2008), yet here we found little evidence that these traits were strongly associated
433 with worm burden. This contrasts with adult shags, which display variation in burdens related to sex
434 and timing of breeding, traits that also affect responses to treatment (Burthe *et al.* 2013, Reed *et al.*
435 2008). Moreover, in our opportunistic necropsies, selective mortality may have confounded the
436 effects of certain host traits: similarly-aged individuals that died in the storm event had similar
437 hatch dates, for example, yet these two traits may influence infection intensity in different ways (for
438 example, burdens increasing due to continuous exposure with age versus a seasonal increase in
439 exposure to infective larvae) whose effects we were not able to separate.

440

441 *Effect of infection on host performance*

442 Parasitism, by definition, is considered to be costly, yet we found a positive correlation
443 between parasite burden and chick mass, a fitness-related trait that is positively associated with
444 recruitment in many bird species (Schwagmeyer and Mock, 2008). This correlation may arise as
445 chicks fed at a higher rate are likely to have higher levels of exposure to parasites, yet parasite
446 infection in both parents and chicks can also affect how resources are distributed among family
447 members (Granroth-Wilding *et al.* 2014, 2015). Experimental approaches that tease apart the
448 relative effects of exposure, burden and host condition are therefore needed to quantify the effect of
449 parasitism on individual performance. Examining the longer-term association between juvenile
450 worm burden and success in later life should also be a priority for future endoscopy studies in this

451 system, taking advantage of the non-destructive technique to quantify the accumulation of sub-
452 lethal impacts typical of macroparasites. Such a chain of fitness effects is of particular importance
453 where parasite infection can shape hosts' developmental trajectories and life histories (Fitze *et al.*
454 2004; Granroth-Wilding *et al.* 2014; Romano *et al.* 2011).

455

456 *Conclusions*

457 Measuring natural variation between hosts in parasite burdens is an essential link in
458 understanding the role of parasites in regulating natural populations. Here, we have developed
459 endoscopy as a non-destructive method to quantify relative parasite burdens in juveniles and
460 revealed prevalence to be significantly higher than expected from more traditional proxy measures.
461 Our demonstration of widespread infection that is established and increases from as early as 12 days
462 of age highlights the potential importance of nematode infection in shaping the contribution of
463 individual shags to population processes throughout their long life (over 20 years). However, we
464 found no evidence to suggest that parasite burdens differ between subgroups of hosts that have
465 previously been found to respond differently to parasite removal. Variation in tolerance among
466 different parts of the population may therefore play a role in governing variation between hosts in
467 how they are impacted by parasitism. Our findings suggest that endoscopy of live juveniles is an
468 informative index of natural variation in parasite burdens, finding the same patterns of infection
469 across the host population as the more direct but destructive index of necropsy. In addition, our
470 results showed that relationships between parasite burden and fitness-related traits in early life are
471 not straightforward. Hence, in combination with experimental approaches, endoscopy provides a
472 powerful tool to link variation in nematode burden with its impact on host success across a wild
473 animal's life and across subgroups of the population, enabling predictions of how parasitism
474 influences on demographic processes in structured natural populations.

475

476

477 ACKNOWLEDGEMENTS

478 Many thanks to Scottish Natural Heritage for permission to work on the Isle of May National
479 Nature Reserve, to Mark Newell and Katherine Herborn for helping collect carcasses, and to Mike
480 Harris and Sarah Wanless for establishing and maintaining the long-term study on the Isle of May.
481 Thanks also to Jarrod Hadfield for statistical advice. Francesca Walker and Jarod Benton
482 contributed to pilot work on the FEC methods. Endoscopy was carried out under UK Home Office
483 Project Licence nr. PPL 60/4115.

484

485

486 FINANCIAL SUPPORT

487 Development of this methodology was funded by the Centre for Ecology & Hydrology project
488 QUIP (QUantifying the Impact of Parasites on seabirds). HGW was supported by a NERC Doctoral
489 Training Grant and EC by a Royal Society University Research Fellowship.

- Abollo, E., Gestal, C., and Pascual, S.** (2001). Anisakid infection in the European shag *Phalacrocorax aristotelis aristotelis*. *Journal of Helminthology* **75**, 209–214.
- Adamo, S.A., Jensen, M., and Younger, M.** (2001) Changes in lifetime immunocompetence in male and female *Gryllus texensis* (formerly *G. integer*): trade-offs between immunity and reproduction. *Animal Behaviour* **62**, 417–425
- Albon, S.D., Stien, A., Irvine, R.J., Langvatn, R., Ropstad, E., and Halvorsen, O.** (2002). The role of parasites in the dynamics of a reindeer population. *Proceedings of the Royal Society B: Biological Sciences* **269**, 1625–1632.
- Anderson, R.C.** (2000). *Nematode Parasites of Vertebrates: Their Development and Transmission* Wallingford, UK: CABI.
- Bates, D., Maechlar, M., Bolker, B., and Walker, S.** (2015). Fitting linear mixed-effects models using lme4. *Journal of Statistical Software* **67**, 1–48
- Bowman, D.D., and Georgi, J.R.** (2009). *Georgis' Parasitology for Veterinarians* (Elsevier Health Sciences).
- Burnham, K.P., and Anderson, D.R.** (2002). *Model Selection and Multi-Model Inference: A Practical Information-Theoretic Approach* (New York: Springer).
- Burthe, S., Newell, M.A., Goodman, G., Butler, A., Bregnballe, T., Harris, E., Wanless, S., Cunningham, E.J.A., and Daunt, F.** (2013). Endoscopy as a novel method for assessing endoparasite burdens in free-ranging European shags (*Phalacrocorax aristotelis*). *Methods in Ecology and Evolution* **4**, 207–216.
- Colditz, I.G.** (2008). Six costs of immunity to gastrointestinal nematode infections. *Parasite Immunology* **30**, 63–70.
- Craig, B.H., Pilkington, J.G., and Pemberton, J.M.** (2006). Gastrointestinal nematode species burdens and host mortality in a feral sheep population. *Parasitology* **133**, 485–496.
- Daunt, F., Wanless, S., Harris, M.P., and Monaghan, P.** (1999). Experimental evidence that age-specific reproductive success is independent of environmental effects. *Proceedings of the Royal Society B: Biological Sciences* **266**, 1489–1493.
- Daunt, F., Monaghan, P., Wanless, S., Harris, M.P., and Griffiths, R.** (2001). Sons and daughters: age-specific differences in parental rearing capacities. *Functional Ecology* **15**, 211–216.
- Dubin, V.B.** (1949). Experimental study of the life cycles of some parasitic worms of animals in the delta of the Volga river (in Russian). *Paraziticheskoe Sbornik Zool Inst SSSR Parasitol. Dig. Zool. Institute Inst. Sci. USSR* **11**, 126–160.
- Fagerholm, H.P., and Overstreet, R.M.** (2008). Ascaridoid Nematodes: *Contracaecum*, *Porrocaecum*, and *Baylisascaris*. In *Parasitic Diseases of Wild Birds*, C.T. Atkinson, N.J. Thomas, and D.B. Hunter, eds. (New Jersey, USA: Wiley- Blackwell), pp. 413–433.
- Fitze, P.S., Clobert, J., and Richner, H.** (2004). Long-term life-history consequences of ectoparasite-modulated growth and development. *Ecology* **85**, 2018–2026.
- Galaktionov, K.V.** (1996). Impact of seabird helminths on host populations and ecosystems.

Bulletin of the Scandinavian Society for Parasitology **6**, 50–61.

Granroth-Wilding, H.M.V., Burthe, S.J., Lewis, S., Reed, T.E., Herborn, K.A., Newell, M.A., Takahashi, E.A., Daunt, F., and Cunningham, E.J.A. (2014). Parasitism in early life: environmental conditions shape within-brood variation in responses to infection. *Ecology and Evolution* **4**, 3408–3419.

Granroth-Wilding, H.M.V., Burthe, S.J., Lewis, S., Herborn, K.A., Takahashi, E.A., Daunt, F., and Cunningham, E.J.A. (2015). Indirect effects of parasitism: costs of infection to other individuals can be greater than direct costs borne by the host. *Proceedings of the Royal Society B: Biological Sciences* **282**, art. 20150602.

Griffiths, R., Daan, S., and Dijkstra, C. (1996). Sex identification in birds using two CHD genes. *Proceedings of the Royal Society B: Biological Sciences* **263**, 1251–1256.

Harris, M.P., Buckland, S.T., Russell, S.M., and Wanless, S. (1994). Post fledging survival to breeding age of Shags *Phalacrocorax aristotelis* in relation to year, date of fledging and brood size. *Journal of Avian Biology* **25**, 268–274.

Hasselquist, D., and Nilsson, J.-A. (2012). Physiological mechanisms mediating costs of immune responses: what can we learn from studies of birds? *Animal Behaviour* **83**, 1303–1312.

Hoberg, E.P. (2005). Marine birds and their helminth parasites. In *Marine Parasitology*, K. Rohde, ed. (Australia: CSIRO), pp. 414–420.

Huizinga, H.W. (1971). Contraecaeciasis in pelicaniform Birds. *Journal of Wildlife Diseases* **7**, 198–204.

Kuiken, T. (1999). Review of Newcastle Disease in Cormorants. *Waterbirds* **22**, 333–347.

Langston, N., and Hillgarth, N. (1995). Molt varies with parasites in Laysan albatrosses. *Proceedings of the Royal Society B: Biological Sciences* **261**, 239–243.

Levecke, B., De Wilde, N., Vandenhouste, E., and Vercruysse, J. (2009). Field validity and feasibility of four techniques for the detection of *Trichuris* in simians: a model for monitoring drug efficacy in public health? *PLOS Neglected Tropical Diseases* **3**, e366.

Lindström, J. (1999). Early development and fitness in birds and mammals. *Trends in Ecology and Evolution* **14**, 343–348.

Magrath, R. (1991). Nestling Weight and Juvenile Survival in the Blackbird, *Turdus merula*. *Journal of Animal Ecology* **60**, 335–351.

McClelland, G. (2005). Nematoda (roundworms). In *Marine Parasitology*, K. Rohde, ed. (Australia: CSIRO), pp. 104–116.

Metcalf, N.B., and Monaghan, P. (2001). Compensation for a bad start: grow now, pay later? *Trends in Ecology and Evolution* **16**, 254–260.

Monaghan, P. (2008). Review. Early growth conditions, phenotypic development and

environmental change. *Philosophical Transactions of the Royal Society B: Biological Sciences* **363**, 1635–1645.

Moravec, F. (2009). Experimental studies on the development of *Contracaecum rudolphii* (Nematoda: Anisakidae) in copepod and fish paratenic hosts. *Folia Parasitologica* **56**, 185–193.

Mougeot, F., Martínez-Padilla, J., Webster, L.M.I., Blount, J.D., Pérez-Rodríguez, L., and Piertney, S.B. (2009) Honest sexual signalling mediated by parasite and testosterone effects on oxidative balance. *Proceedings of the Royal Society B: Biological Sciences* **276**, 1093–1100.

Newey, S., Shaw, D.J., Kirby, A., Montieth, P., Hudson, P.J., and Thirgood, S.J. (2005). Prevalence, intensity and aggregation of intestinal parasites in mountain hares and their potential impact on population dynamics. *International Journal for Parasitology* **35**, 367–373.

Pihlaja, M., Siitari, H., and Alatalo, R.V. (2006) Maternal antibodies in a wild altricial bird: effects on offspring immunity, growth and survival. *Journal of Animal Ecology* **75**, 1154–1164 .

Pinheiro, J., Bates, D., DebRoy, S., Sarkar, D., and Team, R.D.C. (2012). nlme: Linear and nonlinear mixed effects models. R package version 3.1-126

Randolph, S.E., Miklisova, D., Lysy, J., Rogers, D.J., and Labuda, M. (1999). Incidence from coincidence: patterns of tick infestations on rodents facilitate transmission of tick-borne encephalitis virus. *Parasitology* **118**, 177–186.

R Core Team (2013). *R: A language and environment for statistical computing*. (Vienna, Austria: R Foundation for Statistical Computing).

Redpath, S.M., Mougeot, F., Leckie, F.M., Elston, D.A., and Hudson, P.J. (2006). Testing the role of parasites in driving the cyclic population dynamics of a gamebird. *Ecology Letters* **9**, 410–418.

Reed, T.E., Daunt, F., Hall, M.E., Phillips, R.A., Wanless, S., and Cunningham, E.J.A. (2008). Parasite treatment affects maternal investment in sons. *Science* **321**, 1681–1682.

Reed, T.E., Daunt, F., Kiploks, A.J., Burthe, S.J., Granroth-Wilding, H.M.V., Takahashi, E.A., Newell, M., Wanless, S., and Cunningham, E.J.A. (2012). Impacts of parasites in early life: contrasting effects on juvenile growth for different family members. *PLoS ONE* **7**, e32236.

Romano, A., Rubolini, D., Caprioli, M., Boncoraglio, G., Ambrosini, R., and Saino, N. (2011). Sex-related effects of an immune challenge on growth and begging behavior of barn swallow nestlings. *PLoS ONE* **6**, e22805.

Roulin, A., Brinkhof, M.W.G., Bize, P., Richner, H., Jungi, T.W., Bavoux, C., Boileau, N., and Burneleau, G. (2003). Which chick is tasty to parasites? The importance of host immunology vs. parasite life history. *Journal of Animal Ecology* **72**, 75–81.

Royle, N.J., Smiseth, P.T., and Kölliker, M. (2012). *The Evolution of Parental Care* (Oxford: Oxford University Press).

Schwagmeyer, P.L., and Mock, D.W. (2008). Parental provisioning and offspring fitness: size

matters. *Animal Behaviour* **75**, 291–298.

Seivwright, L.J., Redpath, S.M., Mougeot, F., Watt, L., and Hudson, P.J. (2004). Faecal egg counts provide a reliable measure of *Trichostrongylus tenuis* intensities in free-living red grouse *Lagopus lagopus scoticus*. *Journal of Helminthology* **78**, 69–76.

Shaw, D., and Dobson, A. (1995). Patterns of macroparasite abundance and aggregation in wildlife populations: a quantitative review. *Parasitology* **111**, S111–S133.

Shaw, J.L., and Moss, R. (1989). The role of parasite fecundity and longevity in the success of *Trichostrongylus tenuis* in low density red grouse populations. *Parasitology* **99**, 253–258.

Simon, A., Thomas, D.W., Blondel, J., Lambrechts, M.M., and Perret, P. (2003). Within-brood distribution of ectoparasite attacks on nestling blue tits: a test of the tasty chick hypothesis using inulin as a tracer. *Oikos* **102**, 551–558.

Snow, B. (1960). The breeding biology of the shag *Phalacrocorax aristotelis* on the island of Lundy, Bristol Channel. *Ibis* **102**, 554–575.

Stokland, J.N., and Amundsen, T. (1988). Initial size hierarchy in broods of the shag: relative significance of egg size and hatching asynchrony. *The Auk* **105**, 308–315.

Tompkins, D.M., and Hudson, P.J. (1999). Regulation of nematode fecundity in the ring-necked pheasant (*Phasianus colchicus*): not just density dependence. *Parasitology* **118**, 417–423.

Tompkins, D.M., Dunn, A.M., Smith, M.J., and Telfer, S. (2011) Wildlife disease: from individuals to ecosystems. *Journal of Animal Ecology* **80**, 19–38.

Wiehn, J., and Korpimäki, E. (1998). Resource levels, reproduction and resistance to haematozoan infections. *Proceedings of the Royal Society B: Biological Sciences* **265**, 1197–1201.

Yoder, M., De Ley, I.T., King, I.W., Mundo-Ocampo, M., Mann, J., Blaxter, M., Poiras, L., and De Ley, P. (2006). DESS: a versatile solution for preserving morphology and extractable DNA of nematodes. *Nematology* **8**, 367–376.

Table 1. The top five best-fitting models of worm burden (left columns) and the proportion of worms that were large (right columns) as measured by necropsy (top model set) or endoscopy (bottom model set) in relation to host phenotypic traits. Models in each set are shown in order of decreasing fit with their AICc and Δ AICc relative to the best-fit model. The candidate model set for each variable included all combinations of the following predictor variables: age, hatch date, rank, sex, and for endoscopy also visibility. In the necropsy models, age and hatch date and could not be included in the same models as they were closely correlated. Accordingly, models containing hatch date gave almost identical fits to those instead containing age; to illustrate a broader range of model fits, we show only the age models here.

Model (total worm count)	d.f.	AICc	Δ AICc	Model (proportion of worms large)	d.f.	AICc	Δ AICc
<u>Necropsy</u>							
Age	3	207.7	0.0	(intercept only)	2	117.5	0.0
Age + Sex	4	208.3	0.6	Rank	3	119.9	2.4
Age + Rank	4	210.2	2.5	Sex	3	120.0	2.5
Age + Sex + Rank	5	211.5	3.8	Age	3	120.0	2.6
Sex	3	215.6	7.9	Rank + Sex	4	122.7	5.3
<u>Endoscopy</u>							
Age + Visibility	5	320.2	0.0	Age	4	190.4	0.0
Age	4	320.4	0.2	Age + Rank	5	190.5	0.1
Age + Rank	5	321.1	0.9	Age + Rank + Hatch date	6	190.5	0.1
Age + Visibility + Rank	6	321.4	1.2	Age + Hatch date	5	190.9	0.4
Age + Visibility + Hatch date	6	322.6	2.5	Age + Visibility	5	192.5	2.1

502 Table 2. A summary of patterns of variation in nematode burdens between shag chicks, as quantified
503 using necropsy or endoscopy, in relation to phenotypic host traits. Patterns were investigated in both
504 the total worm burden (top set of variables) and the proportion of worms that were large, indicative
505 of how long the chick had been infected (bottom set of variables). Traits that robustly affected
506 worm measures (occurred in all equivalent-fit models) are indicated with a tick, traits that had some
507 support (occurred in more than one equivalent-fit model) are shown with a tick in brackets, and
508 traits with no robust effects are shown with a cross. Hatch date for dissected chicks is indicated with
509 a dash to show that it could not be tested simultaneously with age, as they were tightly correlated.
510


Explanatory variable	Affects necropsy counts	Affects endoscopy counts
<i>Total burden</i>		
Age	√	√
Hatch date	-	x
Rank	x	(√)
Sex	x	x
<i>Proportion large worms</i>		
Age	x	√
Hatch date	-	(√)
Rank	x	(√)
Sex	x	x

511 Table 3. The top 5 best fit models of mass of endoscoped chicks. The set of candidate models
 512 included all combinations of the following variables: worm count (measured by endoscopy),
 513 log(age), sex and rank.
 514

Model	d.f.	AICc	$\Delta AICc$
log(Age) + Worm count	5	553.8	0.0
log(Age) + Worm count + Sex	6	554.6	0.8
log(Age) + Worm count + Rank	6	556.1	2.2
log(Age) + Sex	5	557.1	3.3
log(Age) + Worm count + Sex + Rank	7	557.3	3.5

515 Figure 1. Histograms showing the spread of worm counts from necropsy (left panel) and endoscopy
516 (right panel). The dissection data does not show two high-leverage individuals excluded from the
517 analysis, a hatchling with no worms and a near-fledgling with 243 worms.

518




519

520


521

522 Figure 2. Total worm burden in relation to chick age for necropsied chicks (left panel) and
 523 endoscoped chicks (right panel). Among endoscoped chicks (which covered an older age range than
 524 necropsied chicks) there was some evidence that rank affected worm count, and to illustrate this, in
 525 the endoscopy panel AB chicks are shown with solid symbols and C chicks with open symbols. The
 526 regression line shown is for the best-fit model, which did not include a rank term. Excluding the
 527 oldest chicks, which did not include any C chicks were found, did not alter the ordering of best-fit
 528 models. Note the difference in scale for worm counts and age ranges between the two measures.
 529 The mean lines show a fitted model without random effects using poisson errors and a log link, with
 530 95% confidence intervals shown by the fine-dotted lines.
 531



532


533 Figure 3. The proportion of worms that were large in relation to chick age for necropsied (left
534 panel) and endoscoped (right panel) chicks. In contrast to the worm count, excluding the oldest
535 chicks here slightly changed the order of the best-fit models to: Age + Hatch date; Age; Age +
536 Hatch date + Rank, Age + Rank. The mean lines show a fitted model without random effects and
537 the fine-dotted lines show its 95% confidence intervals.
538



539

540 Figure 4. The relationship among endoscoped chicks between mass at endoscopy and worm count.
541 The solid line shows the fitted relationship and the dotted line the 95% confidence intervals. To
542 account for other factors affecting mass, mass is shown as the residual from a LMM containing age
543 as the only predictor, following the best-fit model for chick mass.

544



Between-individual variation in nematode burden among juveniles in a wild host

Supplementary Information:

*Using endoscopy to test the efficacy of anti-parasite treatment,
observations from dissections, and patterns in faecal egg counts*

Efficacy of anti-parasite treatment

Introduction

Our study system, the European shag (*Phalacrocorax aristotelis*, henceforth “shag”) and its gastrointestinal nematodes, is increasingly yielding valuable insights into the effects of parasitism on individual fitness-related traits and population-level consequences in wild hosts. Although parasites are known to be important influences on host demography and evolution in wild vertebrates (Hudson *et al.* 2002; Tompkins *et al.* 2011), they are rarely considered as factors in ecological processes in seabirds, a globally threatened group whose members are often used as indicators of the state of their marine environment (Piatt, Sydeman & Wiese 2007; Croxall *et al.* 2012).

Several studies of parasitism in the shag have used anti-helminthic treatment as an experimental approach to investigate the effects of nematode infection (Reed *et al.* 2008, 2012; Burthe *et al.* 2013; Granroth-Wilding *et al.* 2014, 2015). The injectable, broad-spectrum antihelminthic drug, Ivermectin (Panomec®, Merial UK) has thus been shown to affect chick survival and growth, adult condition, and behaviour of both adults and chicks, strongly suggesting that treatment affects worm burden. Ivermectin also impacts on ectoparasites, yet previous evidence from this system suggests that ectoparasites contribute little to the cost of a shag's overall parasite burden (Daunt *et al.* 2001). Indeed, Burthe *et al.* (2013) used endoscopy to show that a high dose of ivermectin significantly reduced or removed worm burdens in the proventriculus of adult shags, with no evidence that infection returned for at least 20 days after treatment. In chicks, faecal egg counts (FECs) have provided an indication that treatment reduces affects worm burden, but direct evidence is lacking in chicks of how treatment at the doses used in previous studies affects worm burden. Demonstrating a real effect of treatment on *in situ* nematode burden is particularly important given that, as we show in the main manuscript, FECs in this low-shedding system may not be sensitive to small-scale variation in worm burdens.

577 Understanding the effect of treatment on parasite load is an important link in understanding
578 how between-individual variation in fitness is linked to infection status in juveniles, given that anti-
579 parasite treatment experiments have suggested that infection in chicks can affect both chick growth
580 and parental condition, with long-lasting effects that may be important in population processes
581 (Granroth-Wilding *et al.* 2014, 2015). Here, we use endoscopy of chicks to quantify the effect of
582 treatment with ivermectin on worm burden in shag chicks, at the dosage used in previous work.

583

584 *Methods*

585 We combined the main endoscopy study of natural variation in parasite burden with an experiment
586 to investigate the efficacy of anti-parasite treatment, following protocol from previous parasite
587 removal experiments in shag chicks (full details in Granroth-Wilding *et al.* 2014). We visited nests
588 of three eggs every two days around predicted hatching to obtain hatching dates. When the oldest
589 chick in a brood was 10–12 days old, if all three chicks were still alive, the whole brood was
590 injected with either 0.05ml ivermectin (Panomec© by Merial, 1% wt/vol) (drug-treated broods) or
591 veterinary saline solution (sham-treated control broods). At treatment, we blood sampled chicks for
592 molecular sexing (Griffiths, Daan & Dijkstra 1996) and assigned a rank in the brood hierarchy to
593 each chick according to size, with the heaviest two assigned AB and the lightest C. We have
594 previously shown that mass at this age correctly identifies the C chick in 90% of broods (Granroth-
595 Wilding *et al.* 2014). Previous work has shown that responses to treatment, and therefore potentially
596 the effect of treatment on worm burden, varied between chicks according to differences in rank, sex
597 and hatch date (Reed *et al.* 2008, 2012; Granroth-Wilding *et al.* 2014, 2015). At or after age 25
598 days, we endoscoped all surviving experimental chicks (66 chicks in 29 nests; detailed endoscopy
599 methods in the main text). We also endoscoped unmanipulated chicks from 6 nests known to have
600 had an initial brood size of three.

601 We examined the efficacy of treatment on worm burden by testing whether it affected the total
602 number of worms and the proportion of worms that were large (an indicator of the maturity of the
603 infection). We also examined the impact of treatment on chick performance by testing whether it
604 affected mass at endoscopy, which was positively associated with natural worm burdens in the main
605 investigation. Unmanipulated and sham-treated chicks were pooled as the control group (see main
606 text). All models included age as a predictor, given that older chicks host more worms and a greater
607 proportion of large worms (see main text) and are heavier. For all three response variables (worm
608 count, proportion large, chick mass), treatment was tested as a main effect and in interactions with

sex, rank or hatch date, factors which have previously been shown to influence the impact of treatment (Granroth-Wilding *et al.* 2014, 2015; Reed *et al.* 2008, 2012). In this directed analysis we used hypothesis-testing to assess the importance of each tested factor, in contrast to the more exploratory AIC-based model selection in the main manuscript. We were unable to robustly test the effect of ivermectin treatment on FECs as only 3 drug-treated chicks yielded faecal samples, but we provide a qualitative discussion of these data. All modelling was conducted in R 3.0.2 (R Core Team 2013) using the packages lme4 (Bates, Maechlar & Bolker 2011) and nlme (Pinheiro *et al.* 2012). Worm count was modelled with poisson errors and a log link, the proportion that were large was modelled as a binomial response (weighted by total count), and mass at endoscopy was modelled as a Gaussian response. All parameters are presented as the mean \pm 1 standard error.

Results & discussion

Ivermectin-treated chicks had lower worm burdens than control chicks (mean burden of ivermectin-treated chicks 8.7 ± 1.3 worms; mean burden of control chicks 11.0 ± 1.1 worms; log-transformed effect size in addition to age $-0.54 \pm 0.26 \log(\text{worms})$, $z = -2.12$, $p = 0.034$) (fig. S1). However, treatment did not affect the proportion of worms that were large (in addition to age, effect of treatment 0.34 ± 0.55 , $z = 0.62$, $p = 0.537$). Sex, age and hatch date did not change the effect of treatment on either worm count or the proportion of worms that were large (all interactions $p > 0.1$). This demonstrates that ivermectin is an effective anti-helminthic in live juveniles in the wild, and indicates that it acts equally on all parts of the worm population. These results support the continued use of ivermectin in long-term experiments into the fitness impacts of parasite infection in the wild, enabling experimental work that is valuable in teasing apart correlative patterns in natural burdens and concurrent variation in host fitness.

Chick mass at endoscopy did not differ between any ivermectin-treated and control chicks (in addition to age, effect of treatment 18.3 ± 61.9 , $t = 0.30$, $p = 0.771$; interactions with sex, rank and hatch date all $p > 0.3$). This is perhaps unexpected given that treatment reduced worm burden and that, among naturally infected chicks, heavier chicks had higher burdens (see main text). However, the lack of an effect of treatment on mass is consistent with between-year variation in the impacts of anti-parasite treatment on shag chicks: breeding conditions in the experimental year were such that we would expect little impact of treatment or variation between individuals (Granroth-Wilding *et al.*, 2014).

641 Although we could not explicitly test the effect of treatment on FECs as a proxy indicator of
642 worm burden, we noted that eggs were detected in the faeces of 16 out of 43 control or
643 unmanipulated chicks (37% prevalence) but in none of the four drug-treated chicks for which we
644 had faecal samples (0% prevalence). This points towards treatment reducing FECs as well as
645 reducing worm burdens measured *in situ*. Although our main study found that egg presence in
646 faeces does not, in this system, provide sufficient resolution to reflect natural variation in worm
647 burdens, it is notable that previous work has shown ivermectin treatment to prevent an increase in
648 FEC with age in shag chicks (Granroth-Wilding *et al.*, 2014). Together, this suggests that FECs may
649 be a useful indicator of artificial differences in worm burden in this system, providing an accessible
650 though crude tool to validate the efficacy of experimental anti-parasite treatment.
651

652 **Observations from dissections**

653

654 As part of the main study, 33 chicks that had died naturally were dissected to obtain an alternative,
655 direct measure of worm burden. Findings concerning between-individual variation in worm burdens
656 are described in the main text; here, we provide a qualitative summary of observations made during
657 dissections concerning the biology of the parasite within the host and pathology of infection.

658

659 All dissected chicks contained food, ranging from a heavily digested paste to almost-whole fish
660 from recent feeds. Worms were found almost exclusively in the proventriculus; some worms were
661 present in the oesophagus of two chicks, but never in the intestine. On no occasion were worms or
662 other visible parasites observed in the body cavity outside the digestive tract. Smaller worms were
663 found predominantly in digested food at the bottom of the stomach, whereas larger worms were
664 found predominantly in or on recently ingested or semi-digested boluses of fish. In most
665 dissections, worms were also found in and under the mucous lining of the stomach. Some
666 attachment points on the stomach wall were characterised by hardened ulcerations, which were all
667 in the upper part of the stomach, more concentrated towards the oesophagus.

668

669 **Patterns in FECs**

670

671 As part of the main study, we collected faecal material from 43 unmanipulated or control-treated
672 chicks to examine how well this proxy measure reflects the more reliable indices of worm burden
673 obtained through endoscopy and necropsy. FECs are commonly used as a non-invasive indicator of
674 variation in worm burden, but their reliability is variable and must be verified in each new system in
675 which they are used. In this paper, our main study revealed that eggs could only be detected at very
676 low levels in faecal material (see main manuscript), and that FECs therefore did not capture the full
677 extent of infection in juveniles, possibly as worms have not yet reached sexual maturity at these
678 early stages of infection. We therefore instead investigate whether the presence/absence of eggs,
679 indicative of an established infection, varies with *in situ* indices of worm burden and with host
680 phenotypic traits that have previously been reported to affect how individual traits are affected by
681 infection.

682

683 *Methods*

684 We opportunistically collected faecal samples from 19 endoscoped chicks that defecated during
685 handling. From 24 dissected chicks, we obtained a faecal sample from the cloaca after carcasses had
686 been frozen at -20°C for long-term storage. All faecal samples were therefore stored at -20°C after
687 collection. Previous work in this system has given no evidence that freezing affects egg counts or
688 prevalence (in 138 faecal samples across 3 years of chicks with natural worm burdens, stored either
689 frozen or at room temperature in the non-distorting preservative DESS (Yoder et al., 2006), in a
690 binomial GLMM including year as a random effect and storage method and age as fixed effects:
691 effect of freezing compared to room-temperature DESS on egg count 0.09 ± 0.8 , $z = -0.11$, $p =$
692 0.910 ; effect on egg presence -1.0 ± 0.8 , $z = -1.31$, $p = 0.191$).

693 FECs were carried out using a flotation technique (Bowman and Georgi, 2009). The sample
694 was fully defrosted and mixed well with 20ml saturated salt solution per 1g of faeces (sample sizes,
695 including a variable proportion of nitrogenous waste, ranged from 0.1 to 1.2 g; mean 0.6 g). The
696 mixture was left for c. 10 minutes to allow organic debris to settle out and the lipid-rich eggs to
697 float up. The upper two-thirds of the water column was then mixed gently using a pipette, and an
698 aliquot taken while raising the pipette slowly through the liquid to ensure sampling of any eggs that
699 had not yet reached the surface. The aliquot was placed in a McMaster slide and the portion under
700 the grid (0.15 ml) was systematically searched for nematode eggs at 40x magnification using a light

701 microscope. Three aliquots were examined from each bird, totalling 0.0225g of faecal material. This
702 is sufficient to detect egg presence in adult birds in this low-shedding system (egg presence/absence
703 in 42 adult shags with natural burdens quantified using a variable number of aliquots: using 4-10
704 aliquots, effect of number of aliquots on egg presence 0.02 ± 0.15 , $z = 0.14$, $p = 0.882$; mean
705 prevalence with 95% confidence intervals across 23 individuals with 10 aliquots $32 \pm 20\%$, across
706 19 individuals with 4-8 aliquots $35 \pm 23\%$; across 43 chicks with 3 aliquots in this study, $37 \pm$
707 15%).

708 Most of our 43 faecal samples contained no eggs and only 7 contained more than 1 egg (9
709 with 1 egg, 4 samples with 2 eggs, 2 with 3 eggs and one with 42). To overcome the statistical
710 challenges presented by such a skewed distribution, FECs were analysed as a binary
711 presence/absence response with binomial errors and a logit link. Fitting egg counts with poisson
712 errors and an observation-level random effect to allow for this overdispersion gave qualitatively
713 similar results. We tested whether the probability of egg presence in FECs varied with total worm
714 burden or the number of large worms (more likely to be mature and thus producing eggs) as
715 quantified using either endoscopy or necropsy. Model selection used AICc (details in main
716 manuscript).

717

718 *Results and discussion*

719 Among models examining the effect of worm burden as measured *in situ* (endoscopy and necropsy
720 combined) on FECs, nematode egg presence in faeces was best explained by a model containing
721 only a measurement technique term (log odds of egg presence in endoscoped chicks 1.17 ± 0.70
722 compared to dissected chicks), although this was of an equivalent fit to an intercept-only model
723 ($\Delta AIC = 0.3$) and a model containing a single large worm count term (log odds of egg
724 presence -0.05 ± 0.06 per large worm). In relation to chick phenotypic traits, egg presence was best
725 explained by chick age, which appeared in all three best-fit models (table S1, fig. S2). There was no
726 strong support for any other chick traits being associated with egg presence in faeces.

727 Despite the lack of evidence for any relationship between the presence of nematode eggs in
728 faeces and the more direct *in situ* indices of infection intensity, this proxy measure did reflect the
729 increase in worm burden with chick age that we found with both necropsy and endoscopy. As worm
730 eggs were more likely to be found in older chicks, FECs may thus have some utility for capturing
731 natural variation (or experimental changes to natural burdens; see above) in established infections
732 across the population. However, the variation in the data resulting from the low prevalences mean

733 that we cannot confidently rule out some zero counts being false negatives, and the results of the
734 FEC analyses should therefore be interpreted with caution. Thus, endoscopy remains a more useful
735 technique for capturing the full extent of infection for any given bird and across the population at
736 any point in its lifetime.

737 **Bibliography**

738

Bates, D., Maechlar, M., and Bolker, B. (2011). *lme4: Linear mixed-effects models using S4 classes*.

Bowman, D.D., and Georgi, J.R. (2009). *Georgis' Parasitology for Veterinarians* (Elsevier Health Sciences).

Burthe, S., Newell, M.A., Goodman, G., Butler, A., Bregnballe, T., Harris, E., Wanless, S., Cunningham, E.J.A., and Daunt, F. (2013). Endoscopy as a novel method for assessing endoparasite burdens in free-ranging European shags (*Phalacrocorax aristotelis*). *Methods in Ecology and Evolution* **4**, 207–216.

Croxall, J.P., Butchart, S.H.M., Lascelles, B., Stattersfield, A.J., Sullivan, B., Symes, A., and Taylor, P. (2012) Seabird conservation status, threats and priority actions: a global assessment. *Bird Conservation International* **22**, 1-34

739 **Daunt, F., Monaghan, P., Wanless, S., and Harris, M.** (2001) Parental age and offspring
740 ectoparasite load in European Shags *Stictocarbo aristotelis*. *Ardea* **89**, 449-455.

741

Granroth-Wilding, H.M.V., Burthe, S.J., Lewis, S., Reed, T.E., Herborn, K.A., Newell, M.A., Takahashi, E.A., Daunt, F., and Cunningham, E.J.A. (2014). Parasitism in early life: environmental conditions shape within-brood variation in responses to infection. *Ecology and Evolution*.

Granroth-Wilding, H.M.V., Burthe, S.J., Lewis, S., Herborn, K.A., Takahashi, E.A., Daunt, F., and Cunningham, E.J.A. (2015). Indirect effects of parasitism: costs of infection to other individuals can be greater than direct costs borne by the host. *Proceedings of the Royal Society B: Biological Sciences* **282**.

Griffiths, R., Daan, S., and Dijkstra, C. (1996). Sex identification in birds using two CHD genes. *Proceedings of the Royal Society B: Biological Sciences* **263**, 1251–1256.

Hudson, P.J., Rizzoli, A., Grenfell, B.T., Heesterbeek, H., and Dobson, A.P. (2002) *The ecology of wildlife diseases*. Oxford: Oxford University Press.

Piatt, J.F., Sydeman, W.J., and Wiese, F. (2007) Introduction: a modern role for seabirds as indicators. *Marine Ecology Progress Series* **352**, 199–204

Pinheiro, J., Bates, D., DebRoy, S., Sarkar, D., and Team, R.D.C. (2012). *Linear and Nonlinear Mixed Effects Models*.

Randolph, S.E., Miklisova, D., Lysy, J., Rogers, D.J., and Labuda, M. (1999). Incidence from coincidence: patterns of tick infestations on rodents facilitate transmission of tick-borne encephalitis virus. *Parasitology* **118**, 177–186.

R Core Team (2013). *R: A language and environment for statistical computing*. (Vienna, Austria: R Foundation for Statistical Computing).

Reed, T.E., Daunt, F., Hall, M.E., Phillips, R.A., Wanless, S., and Cunningham, E.J.A. (2008). Parasite treatment affects maternal investment in sons. *Science* **321**, 1681–1682.

Reed, T.E., Daunt, F., Kiploks, A.J., Burthe, S.J., Granroth-Wilding, H.M.V., Takahashi, E.A., Newell, M., Wanless, S., and Cunningham, E.J.A. (2012). Impacts of parasites in early life: contrasting effects on juvenile growth for different family members. *PLoS ONE* **7**, e32236.

Tompkins, D.M., Dunn, A.M., Smith, M.J., and Telfer, S. (2011) Wildlife disease: from individuals to ecosystems. *Journal of Animal Ecology* **80**, 19-38.


Yoder, M., De Ley, I.T., King, I.W., Mundo-Ocampo, M., Mann, J., Blaxter, M., Poiras, L., and De Ley, P. (2006). DESS: a versatile solution for preserving morphology and extractable DNA of nematodes. *Nematology* **8**, 367–376.

742 Table S1. The top five models of best fit explaining the presence of nematode eggs in shag chick
 743 faeces. The top set of models investigated the relationship of FECs with worm counts as measured
 744 by one of two *in situ* techniques (endoscopy or dissection) on a set of candidate models using the
 745 variables technique, worm count and proportion of large worms. The bottom set investigated
 746 variation in FECs in relation to host traits, and explanatory variables used in building the candidate
 747 model set were age, sex, rank and hatch date. Models are shown with their ΔAICc relative to the
 748 best-fit model, in order of decreasing fit. All models included a random effect of nest.

749

Model terms	d.f.	ΔAICc
<u>Relationship with <i>in situ</i> worm measures</u>		
Technique	3	0.0
(intercept only)	2	0.3
Large worm count	3	1.9
Large worm count + technique	4	2.3
Total worm count + technique	4	2.4
<u>Host traits</u>		
Age	3	0.0
Age + Hatch date	4	1.0
Age + Sex	4	1.7
(intercept only)	2	2.1
Age + Rank	4	2.1

750 Figure S1. Worm counts measured using endoscopy in chicks of varying ages that had been treated
751 with ivermectin (solid symbols and line) or sham-treated not manipulated before endoscopy (hollow
752 symbols, dotted line). The line shows the fitted mixed-effects model.



753

754 Figure S2. The relationship of egg prevalence, quantified using FECs, with chick age. The solid line
755 shows the fitted relationship and the dotted lines its 95% confidence intervals.

756

