

## **C/N ratios and Carbon Isotope Composition of Organic Matter in Estuarine Environments**

**Melanie J. Leng<sup>1</sup> and Jonathan P. Lewis<sup>2</sup>**

The sediments that are preserved in estuarine environments (saltmarsh, riverine estuaries, mangrove habitats, lagoons, isolation basins and fjords) contain organic matter that allows investigation of the provenance of that material. These data can then be used specifically to investigate past sea level/land level changes and changes in freshwater flux. Where microfossils are poorly preserved or absent, C/N and  $\delta^{13}\text{C}$  analyses offer an alternative method to deduce environmental histories, but they are especially useful when used in conjunction with a range of other proxies, and when local modern end-member organic variables can be measured to 'calibrate' the sedimentary C/N and  $\delta^{13}\text{C}$ . There are a wide range of C/N- $\delta^{13}\text{C}$  based carbon studies, here we describe examples of studies in a variety of estuarine environments.

**Keywords:** C/N ratios, carbon isotope ratios, carbon cycle, saltmarsh, riverine estuaries, mangrove habitats, lagoons, isolation basins, fjords

### **1. Introduction**

Here, an estuary is defined as a partially enclosed environment, connected to both freshwater (e.g. a river or stream) and the ocean, where sediment accumulates. Estuaries are effectively a transition zone between the terrestrial and ocean environments, are characterised by environmental gradients (particularly salinity, nutrient and geochemical), and are subjected to rapidly fluctuating conditions due to tidal cycles and periods of high river flow. This makes them extremely important as hosts to diverse habitats and complex ecosystems that require

---

<sup>1</sup>M.J. Leng (corresponding author)  
British Geological Survey  
and University of Nottingham, Nottingham,  
United Kingdom  
email: mjl@bgs.ac.uk

<sup>2</sup>J.P. Lewis  
Department of Geography, Loughborough University  
Loughborough,  
United Kingdom  
email: J.P.Lewis@lboro.ac.uk

Careful monitoring and management, and contain sediments able to record both terrestrial and marine past changes. The temporal longevity of modern estuarine records is often limited to the Holocene as most modern estuaries were formed by ice scouring during the last glacial period (high latitudes) and/or by sea level rise during the deglacial (lower latitudes). Being highly productive and receiving sediment and organic matter from catchments, they often accumulate thick sedimentary sequences providing high-resolution archives of environmental change (e.g. Jennings and Weiner 1996). Many components of estuarine deposits have been used to characterise estuary environmental change (e.g. see chapters 7-18 in this volume) and here we describe the carbon/nitrogen (C/N) ratios and carbon isotope composition ( $\delta^{13}\text{C}$ ) of organic matter. One of the main justifications for investigating C/N and  $\delta^{13}\text{C}$  of organic matter in estuaries relies on the fact that micro- and macrofossils (e.g. diatoms, pollen, foraminifera, plant macrophytes; chapters 9-16, 22, this Volume), which provide information on environmental change, can be spatially restricted to particular environments (e.g. Shennan et al. 2000); and in a dynamic environment it is not uncommon to encounter sediment deposits containing a discontinuous fossil record that may be a function of rapidly changing environments or differential preservation (e.g. Shennan et al. 1999; Wilson et al. 2005a). The C/N and  $\delta^{13}\text{C}$  nature of organic matter within sediments (including estuarine) are generally presented together as they provide complementary palaeoenvironmental information. Evidence of  $\delta^{13}\text{C}$  can enable reconstruction of organic matter type, habitat, in addition to productivity in some circumstances; although the signals can also be impacted by other effects such as post depositional change (Figure 1). C/N data can provide information on the carbon source, in terms of both predominant type of plant and their habitat (i.e. aquatic versus terrestrial). Overall though, the use of C/N and  $\delta^{13}\text{C}$  in estuarine systems has been rather limited to date, particularly in comparison to lakes and the marine environment, where the use of these geochemical proxies in environmental reconstructions are commonplace (e.g. Leng and Marshall 2004; Leng et al. 2005; Maslin and Swann 2005 and references therein).

In this chapter we outline the sources of organic matter in estuarine systems and the general method involved in obtaining C/N and  $\delta^{13}\text{C}$  from bulk sediments. We consider the major controls, the limitations and problems to this approach, and discuss what palaeoenvironmental information can be obtained. The latter is largely represented via specific case studies where C/N and  $\delta^{13}\text{C}$  have been used to either independently reconstruct palaeoenvironmental change or enhance multi-proxy studies. The case studies cover a range of “estuary” types, including: saltmarsh, riverine estuaries, mangroves, lagoons, isolation basins and fjords and glacial embayments.

## 2. Analytical Considerations

The accurate determination of C/N and  $\delta^{13}\text{C}$  composition of organic matter requires the complete removal of all inorganic carbonate from the total carbon within a sample. This is usually achieved through treating the sample with an acid. There are a variety of pre-analysis acid treatment methods that have been used from which it is clear there is no consensus on standard practice (see Brodie et al. 2011a). There are potentially significant non-linear biases on measured C/N and  $\delta^{13}\text{C}$  values (and  $\delta^{15}\text{N}$ ; Brodie et al. 2011b) directly associated with the pre-analysis acid treatment methods, which could undermine the environmental interpretation of the data where variations are relatively small. For example, Brodie et al. (2011a) noted C/N ranges of ~ 6 to 13 and  $\delta^{13}\text{C}$  ranges of  $-27.0\text{‰}$  to  $-28.4\text{‰}$  for a terrestrial plant across different pre-analysis acid treatments (when compared with the raw, untreated value of C/N = 9.5;  $\delta^{13}\text{C} = -27.4 \pm 0.1\text{‰}$ ). These C/N values could suggest organic matter from both an aquatic source and from an admixture of aquatic and terrestrial sources. Brodie et al. (2011c)

recommended that the minutiae of the bulk C/N and  $\delta^{13}\text{C}$ , and of organic matter, are not interpreted, but instead changes in the down-core data sets are used in addition to changes that are significantly greater than an estimate of the inaccuracy and imprecision of the data. This is often considerably greater than the standard instrument precision and may be method dependent. They finally recommended that C/N and  $\delta^{13}\text{C}$  data should only be used as an indication of environmental change (non-quantifiable) and should be supported with other proxy data such as micro/macro fossil data. To date there is no consensus about how to treat samples, but what is clear from the Brodie et al. (2011c) experiments is that pre-treatment methods should be consistent within any study.

The traditional method to determine  $\delta^{13}\text{C}$  of organic material within sediments is by oxidation of the carbonate-free sample off-line, via a vacuum-furnace system, to produce and collect  $\text{CO}_2$  for isotope analysis (Boutton 1991). The  $\text{CO}_2$  is produced by reaction of the organic matter with an oxidant, such as cupric oxide ( $\text{CuO}$ ), in an evacuated sealed tube at 600-800°C. The  $^{13}\text{C}/^{12}\text{C}$  ratio of the  $\text{CO}_2$  is measured with an isotope ratio mass spectrometer (IRMS). Although very accurate, this off-line method is slow and requires large (milligram) sample sizes (Meyers and Teranes 2001). On-line techniques now predominate, using continuous flow or dual inlet mass spectrometry. The on-line techniques require an IRMS coupled to a device that produces  $\text{CO}_2$  from the organic matter, usually an elemental analyser, which can also be used to measure C and N concentrations simultaneously. The on-line measurement of  $^{13}\text{C}/^{12}\text{C}$  allows for much reduced sample size (down to micrograms of C) and quicker/automated analysis. The reproducibility of  $\delta^{13}\text{C}$  data is commonly of the order of  $\pm 0.1\%$ . Carbon and nitrogen weight percentages are used to calculate C/N mass ratios, which can be multiplied by 1.167 (the ratio of atomic weights of carbon and nitrogen) to provide C/N atomic ratios. Both types of values are published, but most often the weight ratios are reported. However, atomic C/N ratios are preferable because they reflect biochemical stoichiometry (Meyers and Teranes 2001).

### 3. Sources of Organic Matter in Estuarine Systems

Estuarine sediments receive both autochthonous (derived *in situ*, such as plants growing on the sediment surface) and allochthonous (transported to the sediment from elsewhere) organic matter. In saltmarsh sediments the dominant autochthonous component is derived from *in situ* vascular plant material. Infrequent tidal inundation allows extensive vegetation cover, and therefore highly organic-rich sediments. In contrast, the area between low and high tide, the tidal flats, are frequently inundated by marine water, thus inhibiting vascular plant vegetation, and consequently the predominant organic matter is allochthonous. Allochthonous organic matter sources include marine particulate organic carbon (POC), mainly from phytoplankton, as well as river-derived terrestrial and fluvial material.

Different sources of organic matter in the estuarine environment have (mostly) different C/N and  $\delta^{13}\text{C}$  compositions. In theory this means the bulk organic C/N and  $\delta^{13}\text{C}$  should reflect the relative amounts of the organic matter from the different sources (e.g. Middelburg et al. 1997), assuming the different sources have relatively uniform compositions (Fry and Sherr 1989) and preservation is similar. In environments that are not regularly flushed by fresh and/or marine water, such as lagoons or isolation basins, autochthonous sources of carbon will dominate the supply to the sediments rather than the balance between autochthonous and allochthonous components (Lamb et al. 2006). The range in C/N and  $\delta^{13}\text{C}$  values for the main sources of organic matter in estuarine environments is given in Figures 2 and 3. However, while the type of plant matter has a significant effect on C/N and  $\delta^{13}\text{C}$ , there are other environmental variables that have been shown to affect specific plant  $\delta^{13}\text{C}$ , such as

drought (Farquhar et al. 1989), salinity (Seemann and Critchley 1985), vapour pressure deficit (Turney 1999) and climate (both temperature and changing isotope composition and concentrations of atmospheric CO<sub>2</sub>; Mackie et al. 2005).

Each of the broad plant types and the undifferentiated organic matter are described in more detail below in terms of their C/N and  $\delta^{13}\text{C}$  compositions.

### 3.1 Terrestrial Plants

Terrestrial plants use atmospheric CO<sub>2</sub>, preferentially taking up  $^{12}\text{C}$ , resulting in organic matter that is depleted in  $^{13}\text{C}$  compared with the atmosphere (current atmospheric CO<sub>2</sub>  $\delta^{13}\text{C}_{\text{PDB}} = -8\text{‰}$ ; Keeling et al. 1995). Terrestrial plants are divided into three main types: C3 plants constitute about 90% of all plants, and typically have  $\delta^{13}\text{C}$  values that range between  $-32\text{‰}$  and  $-21\text{‰}$  (Deines 1980); C4 plants have  $\delta^{13}\text{C}$  values of around  $-13\text{‰}$  (range =  $-17\text{‰}$  to  $-9\text{‰}$ ; Deines 1980; e.g. O'Leary et al. 1992); and CAM photosynthesisers (crassulacean acid metabolism; this includes most succulents), use both the C3 photosynthetic pathway and a C4-like pathway, typically resulting in a wide range of  $\delta^{13}\text{C}$  values ( $-11\text{‰}$  to  $-28\text{‰}$ ; Schleser 1995; Figure 2). Terrestrial plants grow in estuarine environments although the type depends on the elevation above the tidal frame (Gray 1992), but terrestrial plant material also enters estuaries via rivers and streams. Saltmarsh species vary with elevation within the tidal frame as a function of their salinity tolerance, but those plants that have adapted to tolerate very saline conditions are commonly of C4 or CAM type (Rundel et al. 1988). Terrestrial vegetation is predominantly composed of lignin and cellulose, which are nitrogen poor (Tyson 1995) and lead to high C/N (Prahl et al. 1980), often much greater than 10. However, data can be variable, plants from a modern saltmarsh (both terrestrial and saltmarsh) along western Greenland, for example, have variable C/N (10-110), while their fossil equivalents in the saltmarsh sediments have low C/N (ca. 5), although these are also likely to contain some algal material (Leng et al. 2012).

### 3.2 Aquatic Plants

Aquatic plants can utilise both bicarbonate ( $\text{HCO}_3^-$ ) and dissolved CO<sub>2</sub> (Benedict et al. 1980). Dissolved CO<sub>2</sub> has lower  $\delta^{13}\text{C}$  values ( $-8\text{‰}$  if in equilibrium with atmospheric CO<sub>2</sub>) than  $\text{HCO}_3^-$  ( $\sim 0\text{‰}$ ; Keeley and Sandquist 1992), and aquatic plants will preferentially take up dissolved CO<sub>2</sub> (Degens et al. 1968; Falkowski 1991). Because the ratio of CO<sub>2</sub> to  $\text{HCO}_3^-$  is a function of pH, and there is less CO<sub>2</sub> in the marine environment than in freshwater, marine plants have higher  $\delta^{13}\text{C}$  values, and they will consequently utilise predominantly  $\text{HCO}_3^-$  (Lamb et al. 2006). Freshwater algae in C3-dominated environments tend to have lower  $\delta^{13}\text{C}$  values ( $-26\text{‰}$  to  $-30\text{‰}$ ; Schidlowski et al. 1983; Meyers 1994) than marine algae ( $-16\text{‰}$  to  $-23\text{‰}$ ; Haines 1976; Meyers 1994), but algae in C4-vegetation catchments can have relatively high  $\delta^{13}\text{C}$  values ( $> -16\text{‰}$ ) as they also use dissolved organic carbon from decomposition and oxidation of C4 plants (Chivas et al. 2001; Figure 2).

Some studies have shown that bacteria may also have an impact on the  $\delta^{13}\text{C}$  (and C/N) of coastal sediments (e.g. Malamud-Roam and Ingram 2001). Peterson et al. (1980) argued that the  $\delta^{13}\text{C}$  values of living *Spartina alterniflora* ( $-11\text{‰}$  to  $-13\text{‰}$ ) and underlying surface sediments ( $-18\text{‰}$  to  $-24\text{‰}$ ) are different in some saltmarshes of the eastern USA because of the presence of bacteria in the sediment. In another study, cyanobacterial blooms in San Francisco Bay (USA) saltmarsh sediments had low  $\delta^{13}\text{C}$  ( $-26.6\text{‰}$  to  $-28.0\text{‰}$ ), suggesting

that bacteria in large quantities, even in marine-influenced sediments, will cause lower  $\delta^{13}\text{C}$  values (Cloern et al. 2002).

Algae and bacteria have C/N ratios of  $< 10$  (Meyers 1994; Tyson 1995) because nitrogen occurs preferentially in proteins and nucleic acids which are relatively abundant in aquatic plants (Talbot and Johannessen 1992). In particular, bacterial activity can introduce nitrogen to the sediment, and thus decrease C/N ratios (Rice and Hanson 1984). However, aquatic plants and bacteria contain a high level of labile compounds and decompose more rapidly than vascular plants (Valiela et al. 1985), and therefore become less significant to the bulk organic C/N and  $\delta^{13}\text{C}$  (e.g. Andrews et al. 1998).

### 3.3 Particulate and Dissolved Organic Carbon

POC represents one of the main organic carbon pools in the oceans, and is mainly composed of living (phytoplankton, zooplankton and bacteria) and non-living (plant detritus and sewage) material (Lamb et al. 2006). Marine POC typically ranges from  $-21\text{‰}$  to  $-18\text{‰}$  (Peters et al. 1978; Wada et al. 1987; Middelburg and Nieuwenhuize 1998). Fluvial POC is a function of contributions mainly from freshwater phytoplankton ( $\delta^{13}\text{C}_{\text{POC}}$  values  $-30\text{‰}$  to  $-25\text{‰}$ ) and particulate terrestrial organic matter ( $\delta^{13}\text{C}_{\text{POC}}$  values  $-25\text{‰}$  to  $-33\text{‰}$ ; e.g. Salomons and Mook 1981; Barth et al. 1998; Middelburg and Nieuwenhuize 1998). Therefore, mean  $\delta^{13}\text{C}$  values of fluvial POC are nearly always very low. As a result of the difference between marine and fluvial POC,  $\delta^{13}\text{C}_{\text{POC}}$  values commonly increase systematically towards the mouth of estuaries due to different contributions from the two end-members. POC has a large phytoplankton component and therefore C/N values tend to be low at around 5-7 (Meyers 1994; Tyson 1995; Figure 2). Diatoms, a particular type of phytoplankton with a silica skeleton, are common in estuarine environments (see chapter 9) and have particularly low C/N (ca. 5). Dissolved organic carbon (DOC) is principally derived from the decomposition of phytoplankton in the marine environment and from mixtures of terrigenous organic matter and phytoplankton in the fluvial environment (Rashid 1985). The "dissolved" fraction of organic carbon is a classification used to define compounds below  $0.45\text{ }\mu\text{m}$ . Most studies of  $\delta^{13}\text{C}_{\text{DOC}}$  in estuarine waters have shown that the isotope composition reflects the fluvial and/or marine source of the DOC (e.g. Peterson et al. 1994; Van Heemst et al. 2000; Goñi et al. 2003; Otero et al. 2003).

### 3.4 Decomposition of Plants

Organic matter decomposition can potentially have a significant influence on both the C/N and  $\delta^{13}\text{C}$  composition of organic matter (Lamb et al. 2006). Early loss of labile compounds in vascular vegetation can lead to significant changes in  $\delta^{13}\text{C}$  in saltmarsh organic matter, but is usually insufficient to prevent the distinction between organic matter from C3 and C4 plants on marshes (e.g. Malamud-Roam and Ingram 2004). Therefore,  $\delta^{13}\text{C}$  is usually considered to remain a faithful indicator of its precursor. The loss of organic carbon in sediments as decomposition progresses can sometimes complicate the use of C/N ratios due to differential loss of one of the elements and through an increase in inorganic nitrogen relative to organic nitrogen (Sampei and Matsumoto 2001). However, the direction of change in C/N and  $\delta^{13}\text{C}$ , rather than absolute values, is generally more important for detecting changes in the environment. Relative changes are commonly retained in Holocene sediments, suggesting that C/N and  $\delta^{13}\text{C}$  can be employed, even in the most dynamic of coastal environments. To confirm this there is often close agreement between diatom and C/N- $\delta^{13}\text{C}$  records (e.g. Mackie et al. 2005; Wilson et al. 2005b; Zong et al. 2006), despite each responding to

different variables (organic source as opposed to salinity), allowing for greater confidence in palaeoenvironmental interpretations using the geochemistry of organic matter.

#### 4. Estuarine Palaeoenvironments

##### 4.1 Saltmarsh

The saltmarsh environment is essentially the uppermost coastal, intertidal zone between the land and the sea that is flooded regularly by tides. On saltmarshes there are both C3 and C4 plants; C4 plants have evolved to tolerate water stress and high salinity enabling them to survive on brackish/saline marshes. Such C4 grasses include those in the genus *Spartina* which can have very distinct  $\delta^{13}\text{C}$  (Haines 1976; Middleberg et al. 1997; Cloern et al. 2002). In regions where both C3 and C4 plants are native to coastal marshes, the sediment  $\delta^{13}\text{C}$  composition can directly relate to the ratio of C3/C4 plants, and thus is used as a proxy for changes in marine versus freshwater contributions to a saltmarsh (Fry et al. 1977; Ember et al. 1987; Middelburg et al. 1997; Johnson et al. 2007). The most successful application of the variation in C3/C4 plants is in the San Francisco Basin (Chmura and Aharon 1995; Byrne et al. 2001; Malamud-Roam and Ingram 2004). Following the rapid postglacial sea level rise culminating ~6,000  $^{14}\text{C}$  yrs BP, thick sequences of estuarine marsh sediments have accumulated in the bay (Ingram et al. 1996a,b; Byrne et al. 2001). Studies have shown that the  $\delta^{13}\text{C}$  composition of the surface soil organic matter on the marsh reflects the distribution of C3 and C4 plants growing in the local area (Stout et al. 1975; Chmura et al. 1987; Chmura and Aharon 1995; Malamud-Roam and Ingram 2001; Kemp et al. 2010). Therefore, it is assumed that down-core changes in  $\delta^{13}\text{C}$  are a record of marsh C3 and C4 palaeovegetation and associated palaeosalinity (e.g. DeLaune 1986; Byrne et al. 2001; Kemp et al. 2012).  $\delta^{13}\text{C}$  evidence, combined with that from other proxies such as diatoms and pollen, from the north San Francisco Bay over the last 3,000 years were used to distinguish changes in salinity and human impacts (Byrne et al. 2001). Changes in the amount of freshwater discharge flowing into the bay were suggested as the key reason for changes in salinity, and thus vegetation type (C3 and C4) on the marshes, which is both a function of river discharge and sea level. All of these studies assume that the vegetation preserved in the surface organic matter has its origins in the plants growing in the local area. Studies have highlighted issues relating to different size fractions having different provenance as well as contamination from root penetration (Lamb et al. 2006). Lamb et al. (2006) discussed issues relating to the use of simple C3/C4 mixing models. For example, some C4 plants grow on freshwater as well as saline marshes. CAM plants complicate bulk  $\delta^{13}\text{C}$  mixing models as they can have intermediate  $\delta^{13}\text{C}$  values, and where sediments receive a significant proportion of allochthonous material (such as tidal-derived phytoplankton), this could increase the sedimentary  $\delta^{13}\text{C}$  values due to the contribution from marine organic matter. In addition, hypersaline environments can include edaphic algal growth and this can lead to algae forming a significant proportion of the autochthonous carbon input (Zedler 1980; Sullivan and Moncreif 1990).

Many mid- or high-latitude saltmarsh vegetation communities, including many saltmarshes in northern Europe, do not have native C4 plants (e.g. Preston et al. 2002). Consequently, the range in organic matter  $\delta^{13}\text{C}$  values on these saltmarshes is much reduced. In the Mersey Estuary (UK), Wilson et al. (2005b) measured the C/N and  $\delta^{13}\text{C}$  of the two main sources of organic carbon present, the inter-tidal sediments mainly derived from saltmarsh plants, and the suspended estuarine POC. It was shown that ground elevation, in relation to the tidal frame, was a key factor controlling bulk surface inter-tidal sediment C/N and  $\delta^{13}\text{C}$  values. Supra-tidal and high saltmarsh sediments consisted almost entirely of

organic carbon derived from the overlying C3 vegetation, and sub-tidal and tidal-flat sediments comprised mainly tidal-derived POC. The different sources of organic carbon in high saltmarsh and sub-tidal sediments resulted in differing bulk sediment  $\delta^{13}\text{C}$  ( $-27.8\text{‰}$  and  $-22.8\text{‰}$ , respectively) and C/N (11.6 and 9.3) values. This modern calibration study was then applied to a Holocene-aged sediment core taken from the Mersey Estuary. The major findings of the study showed that the interpretation of C/N and  $\delta^{13}\text{C}$  within the Mersey Estuary agreed with other proxies and that inter-tidal and sub-tidal surface sediment values are preserved in the Holocene record (Figure 4). However, Wilson et al. (2005b) noted that the detail of the complementary fossil analysis is often missing in the C/N and  $\delta^{13}\text{C}$  data, but sea level variations can still be clearly identified. This is useful where microfossils are poorly preserved (Wilson et al. 2005b).

## 4.2 Riverine Estuaries

A riverine estuary is a partly enclosed coastal body of water with one or more rivers or streams flowing into it, and with a free connection to the open sea. Many studies have shown a gradient in  $\delta^{13}\text{C}$  (low to high) and C/N (high to low) along the outer reaches of rivers as they approach the sea, from fluvial to marine sediments, especially in the suspended POM C/N and  $\delta^{13}\text{C}$  (e.g. Lamb et al. 2006). This systematic shift in C/N and  $\delta^{13}\text{C}$  is commonly preserved in the sedimentary record (Corbett et al. 2007), which also integrates seasonal fluctuations in C/N and  $\delta^{13}\text{C}$ . Along the Atlantic Coast of the USA, Hunt (1970) measured  $\delta^{13}\text{C}$  from surface sediments in rivers and found that the river mouth sediments had  $\delta^{13}\text{C}$  values of around  $-20\text{‰}$ , whereas several kilometres upstream, sediment  $\delta^{13}\text{C}$  was around  $-26\text{‰}$ . In the Winyah Bay Estuary (USA), Goñi et al. (2003) showed that upstream POM had a C/N of 13 and  $\delta^{13}\text{C}$  of  $-26.3\text{‰}$ , whereas in the lower reaches, C/N was around 6 and  $\delta^{13}\text{C}$   $-22.7\text{‰}$ . Similar patterns in C/N and  $\delta^{13}\text{C}$  from suspended and deposited organic matter have been observed in numerous estuaries across the world (e.g. Sackett and Thompson 1963; Salomons and Mook 1981; Thornton and McMannus 1994; Ruttenberg and Goñi 1997; Andrews et al. 1998; Middelburg and Nieuwenhuize 1998; Zong et al. 2006; Yu et al. 2010). The gradient in C/N and  $\delta^{13}\text{C}$  commonly breaks down in the lower reaches of rivers/estuaries due to efficient sediment mixing and homogenisation (Lamb et al. 2006).

The existence of a gradient in sediment C/N and  $\delta^{13}\text{C}$  along the axis of many poorly mixed riverine estuaries can be used as a palaeoenvironmental tool for investigating changes in sea level and palaeoriver discharge. In the Pearl River Estuary (southern China), analysis of C/N and  $\delta^{13}\text{C}$  in surface sediments showed the gradual transition from fully marine conditions at the margins of the South China Sea to fully freshwater conditions in the inner estuary (Zong et al. 2006; Yu et al. 2010). This relationship was confirmed by diatom species analysis (Zong et al. 2006). Based on this relationship between modern diatom flora, C/N,  $\delta^{13}\text{C}$ , and converting diatom flora data to inferred salinity, Zong et al. (2006) were able to reconstruct the Holocene palaeodischarge history of the Pearl River Estuary from estuarine sediment cores. They showed increasing freshwater flux during the early Holocene reaching a peak at ca. 7,500 – 6000 cal. yrs BP. From 6000 yrs BP, a reduction in freshwater flux was interpreted as being due to a weakening of the summer monsoon (Figure 5).

C/N and  $\delta^{13}\text{C}$  (commonly with  $\delta^{15}\text{N}$ ) data can also be indicative of nutrient enrichment, eutrophication and anoxia (oxygen depletion), especially in riverine estuaries due to increased productivity and progressive alteration of ecosystem structure, and subsequent changes in the predominant source of organic matter supply to the sediment (e.g. Cifuentes et al. 1988; Eadie et al. 1994; Voss et al. 2000; Zimmerman and Canuel 2002; Bratton et al. 2003; Church et al. 2006; Thibodeau et al. 2006; Ellegaard et al. 2006; Rosenbauer et al.

2009). For example, a classic symptom of eutrophication in coastal systems is increased abundance of planktonic algae (Nixon 1995; Anderson et al. 2002), particularly diatoms and dinoflagellates, which has been shown to substantially shift C/N and  $\delta^{13}\text{C}$  in some estuarine systems, largely through dilution of the other sources of organic matter (e.g. Ellegaard et al. 2006; Zimmerman and Canuel 2002). This is particularly well demonstrated in the eutrophic Mariager Fjord in Denmark over the last ca. 100 years, where nutrient loading has resulted in an approximately 4‰ increase in  $\delta^{13}\text{C}$  and a decrease of up to 2 in C/N, concurrent with increased  $\delta^{15}\text{N}$ , diatom-inferred total nitrogen, sedimentary pigment abundances and a shift to a nutrient-rich dinoflagellate flora (Ellegaard et al. 2006).

Chesapeake Bay, the largest riverine estuary in the USA (situated between the states of Maryland and Virginia), is a classic example of a eutrophic estuary. It has received much attention over the past few decades due to increasing nutrient concentrations impacting the natural estuarine ecosystem (Seigler et al. 1985; Cooper and Brush 1991; Cornwell et al. 1996; Bratton et al. 2003; Brush 2009). Numerous multi-proxy palaeoenvironmental studies have been undertaken to reconstruct and monitor the extent, scale, longevity and long-term trends in nutrient concentrations (see Brush 2009 and references therein), and to study ecological responses, including the use of sedimentary C/N and  $\delta^{13}\text{C}$  (Zimmerman and Canuel 2002; Bratton et al. 2003).

Zimmerman and Canuel (2002) used sediment geochemical analyses to reconstruct eutrophic progression over the past five centuries in cores collected from Chesapeake Bay. They found that  $\delta^{13}\text{C}$  (and  $\delta^{15}\text{N}$ ) began to increase in the mid-late 18<sup>th</sup> century, concurrent with significant land clearance (Figure 6). Accompanied by increases in the flux of TOC, and plankton and bacterial biomarker abundances (Figure 6), this pattern generally continues up until the present day, but with increased intensity in the late 1800's and 1900's, reaching an advanced state of eutrophication in the 1950's, likely due to increased rates of land clearance following the introduction of dynamite for blasting of stumps (in the 1880's), "superphosphates", and later mechanisation (i.e. since the 1920's). The increase in  $\delta^{13}\text{C}$  (of 0.5-2‰) is attributed to increasing marine primary productivity in the surface waters, carbon limitation (which uses the lighter carbon more quickly) and/or increasing phytoplankton-derived organic matter supply to the sediments. There was little change in the supply of terrestrial organic matter over this period, suggesting that this is not significantly affecting C/N or  $\delta^{13}\text{C}$ .

Importantly, human impact on the Chesapeake Bay watershed (beginning in the 17<sup>th</sup> century) pre-dates estuarine response, suggesting human impact did not trigger eutrophication and anoxia in the basin. Zimmerman and Canuel (2002) suggested climatic forcing caused the onset of eutrophication due to increased freshwater input (c.f. Stahle et al. 1998; Cronin et al. 2000) bringing more nutrients. Since the mid-late 18<sup>th</sup> century eutrophication occurs, even in normal rainfall years, due to the high levels of nutrients present within the watershed. In a longer term record (ca. 2,700 yrs) from Chesapeake Bay, Bratton et al. (2003) analysed TOC, C/N,  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  and showed that the  $\delta^{13}\text{C}$  signal of the estuary reflects the relative mixing of terrestrial and marine carbon (end-members  $\sim -26\text{‰}$  and  $-20\text{‰}$ , respectively) with a trend towards increasing  $\delta^{13}\text{C}$  values moving in the seaward direction from the head of estuary. In Chesapeake Bay, marine POC is mostly produced by algae, whereas fluvial POC is isotopically lower ( $< -25\text{‰}$ ). However, C/N ratios are more ambiguous possibly due to diagenesis (Bratton et al. 2003). They concluded that  $\delta^{13}\text{C}$  reflects a greater input of terrestrial material, which occurs under wetter conditions, so it was possible to use  $\delta^{13}\text{C}$  as a long-term proxy of wetness/dryness. They described wet-dry cycles identifiable over the last ca. 2,700 years, and correlated them with environmental data from tree rings (Stahle et al.



1998), foraminifera (Cronin et al. 2000) and known historical wet/dry periods (e.g. droughts of the 1950's and 1960's).

### 4.3 Mangrove Habitats

Mangroves (tropical and temperate trees, shrubs and associated vegetation) live at the land-sea interface and their presence-absence are indicators of sea level change (e.g. Woodroffe and Chappell 1985; Woodroffe 1990; Schaeffer-Novelli et al. 2002; McKee et al. 2007). The mangrove biome (or mangle) consists of tidal swamps, common in coastal tropics and subtropics, where fine-grained sediments accumulate, and salinity can range from brackish to hypersaline conditions (e.g. Chapman 1976; Benfer et al. 2007; Krauss et al. 2008). The C/N and  $\delta^{13}\text{C}$  values (and  $\delta^{15}\text{N}$ ) of mangrove peat sediments (usually leaf fragments) are used mainly as a proxy of mangrove physiology and stand structure, i.e. the presence of tall versus dwarf plants at a particular site (Wooller et al. 2003a,b; Monacci et al. 2009). The tall/dwarf isotope differentiation is thought to be a function of nutrient limitation ( $\delta^{15}\text{N}$ ) to the dwarf trees due to the more limited availability of nutrient-rich seawater, whereas differences in  $\delta^{13}\text{C}$  are more likely a function of position/shading of the leaves on the plant. There are obvious difficulties with this interpretation, especially where visible leaf fragments are analysed, as leaf  $\delta^{13}\text{C}$  will also vary up through the plant and be dependent on the amount of shading. However, the use of  $\delta^{13}\text{C}$  (and  $\delta^{15}\text{N}$ ) of mangle leaf fragments through time has been demonstrated over the last 8,000 years from Twin Cays in Belize, where isotope variations are interpreted as indicative of changes between dwarf (mean  $\delta^{13}\text{C} = -24.6\text{‰}$ ,  $\delta^{15}\text{N} = -10\text{‰}$ ), transition and tall mangle (mean  $\delta^{13}\text{C} = -28.3\text{‰}$ ,  $\delta^{15}\text{N} = 0\text{‰}$ ). The interpretation is supported by  $\delta^{18}\text{O}$  (of mangrove leaf fragments) as an indicator of sea level (seawater versus freshwater) (Wooller et al. 2007). While the isotope composition of mangrove plants can infer mangrove ecosystem and sea level variation, it is also obvious that local variation can lead to changes in sedimentation rate, deposition versus erosion, and fresh versus sea water availability. Monacci et al. (2009) showed that there can be significant local responses in a mangrove habitat and that regional environmental change can be manifested very differently between sites.

Mangrove sediments can be used in a similar way to estuarine sediments, allowing for physiology and stand structure variation. Peat cores from the mangrove belt of the Brazilian Amazon region, for example, have been used over the Holocene (with other proxies) to identify incidences of marine influence (Cohen et al. 2012). The authors showed enhanced variations in tidal water salinity during the early and middle Holocene (mangrove habitats), which they took to infer increased aridity (less freshwater flux). The later Holocene showed an increase in fluvial influence that resulted in fragmentation of the mangrove belt as mangrove peat was replaced by sediments containing organic matter of a freshwater origin.

### 4.4 Lagoons

Lagoons are typically shallow water bodies partially separated from the sea by a physical barrier (e.g. sandbanks, shingle bars) and most are highly sensitive to sea level changes. Organic C/N and  $\delta^{13}\text{C}$  analyses have been successfully applied to lagoon sequences to infer freshwater and seawater fluctuations (Müller and Mathesius 1999; Müller and Voss 1999; Yamamuro 2000). Most lagoons receive freshwater via surface inflow and seawater through narrow inlets or overbanking. Lagoons are not continuously flushed by fresh or marine water and, therefore, contain a significant portion of autochthonous organic matter, including plankton and macrophytes, but also can contain some allochthonous marginal terrestrial plant

matter. Lagoons can be strongly influenced by human activities, especially the amount and flux of freshwater, which effects the types of autochthonous organic matter (Jia et al. 2012). Yamamuro (2000) examined a series of surface sediments from two large, connected, brackish lagoons in southwestern Japan using isotopes as well as carbon, nitrogen and phosphorous concentrations: the recent eutrophic sediments have C/N ratios of 8.7 to 11.4; suggesting that they contain a high proportion of plankton (either marine or freshwater) relative to terrestrial plant organic matter. The  $\delta^{13}\text{C}$  values decreased with increasing distance from the seawater inlet into the lagoon, from  $-21.7\text{‰}$  at 2.4 km from the inlet to  $-23.9\text{‰}$  at 32.1 km, suggesting a decrease in marine organic matter which correlates with salinity. Since C/N ratios were unaffected by changes in salinity, there is no correlation between the C/N and  $\delta^{13}\text{C}$  of the organic matter.

Lagoonal sediments can in places contain high amounts of inorganic nitrogen associated with clays. The amount of inorganic nitrogen becomes significant (when compared with %C) in sediments that contain very low concentrations of organic matter. In a study from Nakaumi Lagoon, southwest Japan, Sampei and Matsumoto (2001) investigated C/N ratios over the last ca. 8,000  $^{14}\text{C}$  yrs BP. Their data showed two peaks in bulk C/N of  $> 15$  which could be interpreted as increases in terrestrial input, although this did not agree with the interpretation from other organic proxies. It was thought that high levels of inorganic nitrogen, throughout the core ( $\sim 0.04$  wt%) had the effect of lowering the C/N ratio where %C was less than 1%. This study also highlighted the fact that high levels of phytoplankton can lead to problems when using C/N as a simple two end-member proxy. Sampei and Matsumoto (2001) modelled the relationship between the relative proportions of terrestrial and planktonic organic matter and C/N ratios, and demonstrated that the relationship was non-linear. C/N is therefore not just a function of the balance between the two end-member inputs.

In a Holocene study of coastal lagoons in the southern Baltic Sea region, Müller and Mathesius (1999) and Müller and Voss (1999), compared the C/N and  $\delta^{13}\text{C}$  of modern plants and sediments to investigate sea level variation. The lagoons are currently separated from the Baltic Sea by low barrier islands, which have developed since the mid Holocene. C/N values were generally  $> 12$ , with low  $\delta^{13}\text{C}$  between  $-25\text{‰}$  and  $-30\text{‰}$ , suggesting mainly freshwater-derived organic material (Müller and Voss 1999) with some terrestrial material (Müller and Mathesius 1999), although they also identified intermediary values of autochthonous macrophytes and plankton which complicated the interpretation. From a sediment core, the organic matter in the late Holocene had low C/N values, which was interpreted as less macrophyte and increasingly more phytoplankton input due to eutrophication (Müller and Mathesius 1999). Within the sediment core, the authors also distinguished (based on geochemistry and sedimentology) lacustrine phases ( $\delta^{13}\text{C} = -29.1\text{‰}$ ), sandy shallow lake environments with some aquatic macrophyte input ( $\delta^{13}\text{C} -24.8\text{‰}$ , C/N  $> 12$ ), as well as the more typical lagoonal phases following sand spit formation. Cores from different lagoons showed similar patterns (Müller 2002), although changes were not entirely synchronous suggesting the use of lagoonal sediments to decipher regional, as opposed to local, sea level change is not infallible.

#### 4.5 Isolation Basins

Isolation basins are natural rock depressions that have been separated from the sea due to falls in relative sea level. They are extremely important as they provide long, high-resolution records of sea level change reaching back to the Late Glacial period (e.g. Shennan et al. 1999). Environmental data from isolation basins are extensively used to contribute to glacial-

isostatic adjustment models. In a similar way to lagoons, isolation basins may be connected to the sea intermittently over a period of time depending on their height relative to sea level. The ability of C/N and  $\delta^{13}\text{C}$  to record changes in the source of organic matter in C3 dominated systems has resulted in their use in isolation basin sediments as a proxy for shoreline displacement during the late Quaternary (e.g. Chivas et al. 2001; Westman and Hedenström 2002; Mackie et al. 2005; Mills et al. 2009). As UK isolation basins (common on the west coast of Scotland) are naturally C3-vegetated environments, Holocene sea level studies have traditionally relied on microfossil-based palaeoenvironmental reconstructions (e.g. Shennan et al. 2000; Berglund et al. 2005).

Mackie et al. (2005) measured a range of plants and surface sediment samples from around marine and freshwater contemporary isolation basins in western Scotland to characterise end-member values for C/N and  $\delta^{13}\text{C}$ . They found that the surface sediment C/N and  $\delta^{13}\text{C}$  reflected a mixture of the end-members from the range of modern inputs and thus concluded that, as with other “types” of estuaries, the dominant controlling variable on C/N and  $\delta^{13}\text{C}$  of surface sediment was the source of the organic matter (terrestrial, marine). In isolation basins, C/N and  $\delta^{13}\text{C}$  should therefore be a reliable proxy for past changes in basin salinity and sea level. In a Holocene study of C/N and  $\delta^{13}\text{C}$  profiles from bulk organics from Upper Loch nan Eala (Scotland), Mackie et al. (2005) described a variety of conditions in the basin from fully marine, through to transitional/brackish and freshwater (Figure 7). These palaeoenvironmental changes, inferred from the core C/N and  $\delta^{13}\text{C}$  data, coincide well with the diatom-inferred environments and with previously interpreted changes in sea level (Shennan et al. 2000). However, the relationship breaks down around 6,000 cal yrs BP when diatoms indicate fully marine conditions, whereas C/N and  $\delta^{13}\text{C}$  are more indicative of freshwater. Several possible explanations were discussed including varying amounts of marine plankton or a change in temperature or  $\text{CO}_2$  concentration that might affect the fractionation of  $\delta^{13}\text{C}$ . This discrepancy between the classical diatom salinity reconstruction and geochemical proxies suggests that although the technique appears to work, it is important to use a multi-proxy approach as factors other than organic matter source can influence the interpretation of C/N and  $\delta^{13}\text{C}$ . Philippsen et al. (2013) also demonstrated a strong relationship between diatom-inferred salinity and bulk organic  $\delta^{13}\text{C}$  from a small estuary (Kilen) opening into the Danish Limfjord (a shallow sound subject to changing exposure to, and occasional isolation from, the North Sea and Skagerrak), which was further supported by mollusc, foraminifera and plant macrofossil data (Lewis et al. 2013).

The Gulf of Carpentaria (between Australia and New Guinea) has periodically been cut off from the Indian and Pacific Oceans during the Late Quaternary, forming Lake Carpentaria (Chivas et al. 2001). During periods of low sea level the lake forms an isolation basin with an outlet channel into the Arafura Sea. A multi-proxy study of sediment cores from the lake (going back to 130,000  $^{14}\text{C}$  yrs BP) showed periods of open ocean, estuarine and lacustrine conditions, as well as sub-aerial exposure. C/N and  $\delta^{13}\text{C}$  from 9,700  $^{14}\text{C}$  yrs BP showed that C/N was predominantly higher in the non-marine facies, while  $\delta^{13}\text{C}$  values showed differences between the non-marine (–14‰ to –15‰) and marine (–19‰ to –21‰) phases. The non-marine  $\delta^{13}\text{C}$  values, with high C/N of 10-30 were thought to reflect the abundance of C4 grasses growing around Lake Carpentaria during periods of low sea level (Chivas et al. 2001).

#### 4.6 Fjords and Glacial Embayments

Fjords are deep, submerged, long and narrow inlet valleys formed by glacial activity. Mixing occurs between the freshwater-terrestrial inflow of plant material and marine POM. Due to

the depth of many fjords, tidally derived water can also enter a fjord through deep-water inflows that can reach as far as the fjord head. There are relatively few examples where C/N and  $\delta^{13}\text{C}$  have been applied to fjord sediments. In general though, we expect the distribution between marine and terrestrial-freshwater organic matter to change along the fjord much as it does in a river estuary.  $\delta^{13}\text{C}$  values increase towards the marine environment whereas C/N decreases (St-Onge and Hillaire-Marcel 2001; Sepúlveda et al. 2011). Similar to fjords are polar embayments formed by indentations in the shoreline. These are often ice dammed, creating semi-enclosed bays which contain a lid of freshwater. One such embayment occurs along the Antarctic Peninsula and Alexander Island where  $\delta^{13}\text{C}$  has been used to investigate past phases of collapse in the George VI Ice Shelf (Bentley et al. 2005). Lakes forming behind the ice shelf have freshwater lids but retain a deep marine connection under the ice shelf. Bentley et al. (2005) used both the presence of marine-dominated organisms and bulk organic  $\delta^{13}\text{C}$  to prove the presence/absence of the ice shelves through the Holocene. Specifically, where the organic matter in the sediments had a high  $\delta^{13}\text{C}$  (between  $-23$  and  $-22\text{‰}$ ), there was more marine organic matter, suggesting the absence of ice shelves at that time (and no freshwater lid). When the sediments contained organic matter with low  $\delta^{13}\text{C}$  (around  $-26\text{‰}$ ), then there was likely to be a lid of freshwater and the shelves were more likely to have been present (Figure 8).

## 5. Summary

The sediments that are preserved in estuarine environments (saltmarsh, riverine estuaries, mangrove habitats, lagoons, isolation basins and fjords) contain organic matter that allows investigation of the provenance of that material. These data can then be used specifically to investigate past sea level/land level changes and changes in freshwater flux. Where microfossils are poorly preserved or absent, C/N and  $\delta^{13}\text{C}$  analyses offer an alternative method to deduce environmental histories, especially when used in conjunction with a range of other proxies, and when local modern end-member organic variables can be measured to 'calibrate' the sedimentary C/N and  $\delta^{13}\text{C}$ . There are a wide range of C/N- $\delta^{13}\text{C}$  based carbon budget studies, providing a database of examples of different coastal conditions that can be used for end-member composition for C/N and  $\delta^{13}\text{C}$ .

The most successful studies are probably those that have utilised a known relationship between C3 and C4 plant distribution and salinity in coastal environments, and have then applied this relationship back through the Holocene. In particular, the pioneering work in the San Francisco Basin demonstrated that the relative proportions of C3 and C4 plants growing in an area were preserved in the  $\delta^{13}\text{C}$  composition of the underlying surface sediment, and thus past changes in salinity could be interpreted by observing sedimentary changes in  $\delta^{13}\text{C}$ . Where there are no native C4 plants,  $\delta^{13}\text{C}$  and C/N have also been successfully used as palaeoenvironmental indicators, as bulk inter-tidal sedimentary  $\delta^{13}\text{C}$  and C/N can reflect elevation within the tidal frame. Supra-tidal and high saltmarsh sediments consist almost entirely of organic carbon derived from the overlying C3 vegetation, and sub-tidal and tidal-flat sediments of tidal-derived POM.

In riverine estuaries the use of C/N and  $\delta^{13}\text{C}$  works because of the systematic shift of the provenance of organic matter along the estuary. Several examples can be found all over the world, but in particular along the Pearl River Estuary (China) and Chesapeake Bay (USA), although in the latter case sea level indicators are overprinted by anthropogenic effects in the more recent sediments. The use of C/N and  $\delta^{13}\text{C}$  to provenance organic matter has also been used to varying degrees of success in mangrove habitats and lagoons, although other environmental effects, like tree stands, sediment dispersal and eutrophication, can over print

the primary signal. In isolation basins, C/N and  $\delta^{13}\text{C}$  work particularly well because the change from marine to non-marine conditions can be abrupt, as the isolation is due to a physical barrier. This has been proven particularly well in northwest Scotland. C/N and  $\delta^{13}\text{C}$  in fjords and glacial embayments are used in a similar way to riverine estuaries in that the distribution of freshwater/terrestrial and marine organic matter, as a result of variations in sea level, freshwater flux, and in the case of glacial embayments the presence/absence of ice shelves, may be determined.

## 6. References

- Anderson DM, Gilbert PM, Burkholder JM (2002) Harmful algal blooms and eutrophication: nutrient sources, composition and consequences. *Estuaries* 25:704–726
- Andrews JE, Greenway AM, Dennis PF (1998) Combined carbon isotope and C/N ratios as indicators of source and fate of organic matter in a poorly flushed, tropical estuary: Hunts Bay, Kingston Harbour, Jamaica. *Estuar Coast Shelf Sci* 46:743–756
- Barth JAC, Veizer J, Mayer B (1998) Origin of particulate organic carbon in the upper St. Lawrence: isotopic constraints. *Earth Planet Sc Lett* 162:111–121
- Benedict CR, Wong WWL, Wong JHH (1980). Fractionation of the stable isotopes of inorganic carbon by seagrasses. *Plant Physiol* 65:512–517
- Benfer NP, King BA, Lemckert (2007) Salinity observations in a subtropical estuarine system on the Gold Coast, Australia. *J Coastal Res SI 50* (Proceedings of the 9th International Coastal Symposium, Gold Coast, Australia), pp. 646–651
- Bentley MJ, Hodgson DA, Sugden DE, Roberts SJ, Smith JA, Leng MJ, Bryant C (2005) Early Holocene retreat of the George VI Ice Shelf, Antarctic Peninsula. *Geology* 33:173–176
- Berglund BE, Sandgren P, Barnekow L, Hannon G, Jiang H, Skog G, Yu SY (2005) Early Holocene history of the Baltic Sea, as reflected in coastal sediments in Blekinge, southeastern Sweden. *Quatern Int* 130:111–139
- Boutton, TW (1991) Stable carbon isotope ratios of natural materials: II. Atmospheric, terrestrial, marine, and freshwater environments. In: Coleman DC, Fry B (Eds.), *Carbon Isotopes Techniques*. Academic Press, Inc., San Diego, pp. 173–185
- Bratton JF, Colman SM, Seal II RR, (2003) Eutrophication and carbon sources in Chesapeake Bay over the last 2700 yr: Human impacts in context. *Geochem Cosmochim Ac* 67:3385–3402
- Brodie CR, Casford JSL, Lloyd JM, Leng MJ, Heaton THE, Kendrick CP, Zong YQ (2011c) Evidence for bias in C/N,  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  values of bulk organic matter, and on environmental interpretation, from a lake sedimentary sequence by pre-analysis acid treatment methods. *Quaternary Sci Rev* 30:3076–3087
- Brodie CR, Heaton THE, Leng MJ, Kendrick CP, Casford JSL, Lloyd JM (2011b) Evidence for bias in measured  $\delta^{15}\text{N}$  values of terrestrial and aquatic organic materials due to pre-analysis acid treatment methods. *Rapid Commun Mass Sp* 25: 1089–1099
- Brodie CR, Leng MJ, Casford JSL, Kendrick CP, Lloyd JM, Yongqiang Z, Bird MI (2011a) Evidence for bias in C and N concentrations and  $\delta^{13}\text{C}$  composition of terrestrial and aquatic organic materials due to pre-analysis acid preparation methods. *Chem Geol* 282:67–83
- Brush GS (2009) Historical land use, nitrogen, and coastal eutrophication: a paleoecological perspective. *Estuar Coast* 32:18–28
- Byrne R, Ingram LB, Starratt S, Malamud-Roam F (2001) Carbon-isotope, diatom, and pollen evidence for late Holocene salinity change in a brackish marsh in the San Francisco Estuary. *Quaternary Res* 55:66–76
- Chapman VJ (1976) *Mangrove Vegetation*. J. Cramer, Vaduz, Germany
- Chivas AR, Garcia A, van der Kaars S, Couapel MJJ, Holt S, Reeves JM, Wheeler DJ, Switzer AD, Murray-Wallace CV, Banerjee D, Price DM, Wang SX, Pearson G, Edger NT, Beaufort L, De Deckker P, Lawson E, Cecil CB (2001) Sea-level and environmental changes since the last interglacial in the Gulf of Carpentaria, Australia: an overview. *Quatern Int* 83–85:19–46
- Chmura GL, Aharon P, Socki RA, Abernethy R (1987) An inventory of  $^{13}\text{C}$  abundances in coastal wetlands of Louisiana, USA: vegetation and sediments. *Oecologia* 74:264–271

- Chmura G.L, Aharon P (1995) Stable carbon isotope signatures of sedimentary carbon in coastal wetlands as indicators of salinity regime. *J Coastal Res* 11:124–135
- Church TM, Sommerfield CK, Velinsky DJ, Point D, Benoit C, Amouroux D, Plaa D, Donard OFX (2006) Marsh sediments as records of sedimentation, eutrophication and metal pollution in the urban Delaware Estuary. *Mar Chem* 102:72–95
- Cifuentes LA, Sharp JH, Fogel ML (1988) Stable carbon and nitrogen isotope biogeochemistry in the Delaware Estuary. *Limnol Oceanogr* 33:1102–1115
- Cloern JE, Canuel EA, Harris D, (2002) Stable carbon and nitrogen isotope composition of aquatic and terrestrial plants of the San Francisco Bay estuarine system. *Limnol Oceanogr* 47:713–729
- Cohen MCL, Pessenda LCR, Behling H, Rossetti DDeF, França MC, Guimaraes JTF, Beltrão Smith C (2012) Holocene palaeoenvironmental history of the Amazonian mangrove belt. *Quaternary Sci Rev* 5550–58
- Cooper SR, Brush GS (1991) Long-term history of Chesapeake Bay anoxia. *Science* 254:992–996
- Corbett DR, Vance D, Letrick E, Mallinson D, Culver S (2007) Decadal-scale sediment dynamics and environmental change in the Albermarle estuarine system, North Carolina. *Estuar Coast Shelf Sci* 71:717–729
- Cornwell JC, Conley DJ, Owens M, Stevenson JC (1996) A sediment chronology of the eutrophication of Chesapeake Bay. *Estuaries* 19:488–499
- Cronin T, Willard D, Karlsen A, Ishman S, Verardo S, McGeehin J, Kerhin R, Holmes C, Colman S, Zimmerman A (2000) Climatic variability in the eastern United States over the past millennium from Chesapeake Bay sediments. *Geology* 28:3–6
- Degens ET, Gullard RR, Sackett WM, Hellebust JA (1968) Metabolic fractionation of carbon isotopes in marine plankton. 1. Temperature and respiration experiments. *Deep-Sea Re* 15:1–9
- Deines P (1980) The isotopic composition of reduced organic carbon. In: Fritz P, Fontes JC (Eds.), *Handbook of Environmental Isotope Geochemistry. Volume 1, The Terrestrial Environment*, A. Elsevier, Amsterdam, pp. 329–406
- DeLaune RD (1986) The use of  $\delta^{13}\text{C}$  signature of C3 and C4 plants in determining past depositional environments in rapidly accreting marshes of the Mississippi River deltaic plain, Louisiana, U.S.A. *Chem Geol* 59:315–320
- Eadie BJ, McKee BA, Lansing MB, Robbins JA, Metz S, Trefry JH (1994) Records of nutrient-enhanced coastal ocean productivity in sediments from the Louisiana Continental Shelf. *Estuaries* 17:754–765
- Ellegaard M, Clarke AL, Reuss N, Drew S, Weckström K, Juggins S, Anderson NJ, Conley DJ (2006) Multi-proxy evidence of long-term changes in ecosystem structure in a Danish marine estuary, linked to increased nutrient loading. *Estuar Coast Shelf Sci* 68:567–578
- Ember LM, Williams DF, Morris JT (1987) Processes that influence carbon isotope variations in salt marsh sediments. *Mar Ecol Prog Ser* 36:33–42
- Falkowski PG (1991) Species variability in the fractionation of  $^{13}\text{C}$  and  $^{12}\text{C}$  by marine phytoplankton. *J Plankton Res* 13:21–28
- Farquhar GD, Hubick KT, Condon AG, Richards RA (1989) Carbon isotope fractionation and plant water-use efficiency. In: Rundel PW, Ehleringer JR, Nagy KA (Eds.), *Stable Isotopes in Ecological Research*. Springer-Verlag, New York, pp. 21–40
- Fry B, Scalan RS, Parker PL (1977) Stable carbon isotope evidence for two sources of organic matter in coastal sediments: seagrass and plankton. *Geochim Cosmochim Acta* 41:1875–1877

- Fry B, Sherr EB (1989)  $\delta^{13}\text{C}$  measurements as indicators of carbon flow in marine and freshwater ecosystems. In: Rundel PW, Ehleringer JR, Nagy KA (Eds.), *Stable Isotopes in Ecological Research*. Springer-Verlag, New York, pp. 196–229
- Goñi MA, Teixeira MJ, Perkey DW (2003) Sources and distribution of organic matter in a river-dominated estuary (Winyah Bay, SC, USA). *Estuar Coast Shelf Sci* 57:1023–1048
- Gray AJ (1992) Saltmarsh plant ecology: zonation and succession revisited. In: Allen JRL, Pye K (Eds.), *Saltmarshes: morphodynamics, conservation and engineering significance*. Cambridge University Press, Cambridge, pp. 63–79
- Haines EB (1976) Stable carbon isotope ratios in biota, soils and tidal water of a Georgia salt marsh. *Estuar Coast Mar Sci* 4:609–616
- Hunt JM (1970) The significance of carbon isotope variations in marine sediments. In: Hobson GD, Speers GC (Eds.), *Advances in Organic Geochemistry. Proceedings of the Third International Congress, London, 1966*. Int Ser Monogr Earth Sci 32. Pergamon Press, Oxford, pp. 27–35
- Ingram LB, Ingle JC, Conrad ME (1996a) A 2000-year record of Sacramento-San Joaquin River inflow to San Francisco Bay estuary, California. *Geology* 24:331–334
- Ingram LB, Ingle JC, Conrad ME (1996b) Stable isotope record of late Holocene salinity and river discharge in San Francisco Bay, California. *Earth Planet Sci Lett* 141:237–247
- Jennings AE, Weiner NJ (1996) Environmental change in eastern Greenland during the last 1300 years: evidence from foraminifera and lithofacies in Nansen Fjord, 68°N. *Holocene* 6:79–191
- Jia J, Gao JH, Liu YF, Gao S, Yang Y (2012) Environmental changes in Shamei Lagoon, Hainan Island, China: Interactions between natural processes and human activities. *J Asian Earth Sci* 52:158–168
- Johnson BJ, Moore KA, Lehmann C, Bohlen C, Brown TA (2007) Middle to late Holocene fluctuations of C3 and C4 vegetation in a Northern New England Salt Marsh, Sprague Marsh, Phippsburg Maine. *Org Geochem* 38:394–403
- Keeley JE, Sandquist DR (1992) Carbon: freshwater plants. *Plant Cell Env* 15:1021–1035
- Keeling CD, Whorf TP, Wahlen M, van der Plicht J (1995) Interannual extremes in the rate of rise of atmospheric carbon dioxide since 1980. *Nature* 375:666–670
- Kemp AC, Vane CH, Horton BP, Cuver SJ (2010) Stable carbon isotopes as potential sea-level indicators in salt marshes, North Carolina, USA. *Holocene* 20:623–636
- Kemp AC, Vane CH, Horton BP, Engelhart SE, Nikitina D (2012) Application of stable carbon isotopes for reconstructing salt-marsh floral zones and relative sea level, New Jersey, USA. *J Quaternary Sci* 27:404–414
- Krauss KW, Lovelock CE, McKee KL, López-Hoffman L, Ewe SML, Sousa WP (2008) Environmental drivers in mangrove establishment and early development. A review. *Aquat Bot* 89:105–127
- Lamb AL, Wilson GP, Leng MJ (2006) A review of coastal and palaeoclimate and relative sea-level reconstructions using  $\delta^{13}\text{C}$  and C/N ratios in organic material. *Earth-Science Reviews* 75:29–57
- Leng MJ, Marshall JD, (2004) Palaeoclimate interpretation of stable isotope data from lake sediment archives, *Quaternary Sci Rev* 23:811–831
- Leng MJ, Lamb AL, Heaton THE, Marshall JD, Wolfe BB, Jones MD, Holmes JA, Arrowsmith CA (2005) Isotopes in lake sediments. In: MJ Leng (Ed.) *Isotopes in Palaeoenvironmental Research. Developments in Palaeoenvironmental Research. Volume 10*. Springer, Dordrecht, pp. 147–184
- Leng MJ, Wagner B, Anderson NJ, Bennike O, Woodley E, Kemp SJ (2012) Deglaciation and catchment ontogeny in coastal southwest Greenland: implications for terrestrial and aquatic carbon cycling. *J Quaternary Sci* 27:575–584



Lewis JP, Ryves DB, Rasmussen P, Knudsen KL, Petersen KS, Olsen J, Leng MJ, Kristensen P, McGowan S, Phillipsen B (2013) Environmental change in the Limfjord, Denmark (ca. 7,500–1,500 cal yrs BP): a multiproxy study. *Quaternary Sci Rev* 78:126–140

Mackie EAV, Leng MJ, Lloyd JM, Arrowsmith C (2005) Bulk organic  $\delta^{13}\text{C}$  and C/N ratios as palaeosalinity indicators within a Scottish isolation basin. *J Quaternary Sci* 20:301–408

Malamud-Roam F, Ingram BL (2001) Carbon isotopic compositions of plants and sediments of tide marshes in the San Francisco Estuary. *J Coast Res* 17:17–29

Malamud-Roam F, Ingram BL (2004) Late Holocene  $\delta^{13}\text{C}$  and pollen records of palaeosalinity from tidal marshes in the San Francisco Bay Estuary, California. *Quaternary Res* 62:134–145

Maslin MA, Swann G (2005) Isotopes in Marine Sediments. In: MJ Leng (Ed.) *Isotopes in Palaeoenvironmental Research. Developments in Palaeoenvironmental Research. Volume 10.* Springer, Dordrecht, pp. 227–290

McKee KL, Cahoon DR, Feller I.C (2007) Caribbean mangroves adjust to rising sea level through biotic controls on change in soil elevation. *Global Ecology and Biogeography* 16:545–556

Meyers PA (1994) Preservation of elemental and isotopic source identification of sedimentary organic matter. *Chem Geol* 114:289–302

Meyers PA, Teranes JL (2001) Sedimentary organic matter. In: Last WM, Smol JP (Eds). *Tracking Environmental Change Using Lake Sediments, Volume 2: Physical and geochemical methods.* Springer, Dordrecht, pp. 239–269

Middelburg JJ, Nieuwenhuize J, Lubberts RK, van de Plassche O (1997) Organic carbon isotope systematics of coastal marshes. *Estuar Coast Shelf Sci* 45:681–687

Middelburg JJ, Nieuwenhuize J (1998) Carbon and nitrogen stable isotopes in suspended matter and sediments from the Schelde Estuary. *Mar Chem* 60:217–225

Mills K, Mackay AW, Bradley RS, Finney B (2009) Diatom and stable isotope records of late-Holocene lake ontogeny at Indrepollen, Lofoten, NW Norway: a response to glacio-isostasy and Neoglacial cooling. *Holocene* 19:261–271

Monacci N, Meier-Grünhagen U, Finney BP, Behling H, Wooller MJ (2009) Mangrove ecosystem changes during the Holocene at Spanish Lookout Cay, Belize: Palaeogeography, Palaeoclimatology, Palaeoecology 280:37–46

Müller A (2002) Geochemical expressions of late- and post-glacial land-sea interactions in the southern Baltic Sea. *Boreal Env Res* 7:13–25

Müller A, Mathesius U (1999) The palaeoenvironments of coastal lagoons in the southern Baltic Sea, I. The application of sedimentary  $\text{C}_{\text{org}}/\text{N}$  ratios as source indicators of organic matter. *Palaeogeography, Palaeoclimatology, Palaeoecology* 145:1–16

Müller A, Voss M (1999) The palaeoenvironments of coastal lagoons in the southern Baltic Sea, II.  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  ratios of organic matter – sources and sediments. *Palaeogeography, Palaeoclimatology, Palaeoecology* 145:17–32

Nixon SW (1995) Coastal eutrophication: A definition, social causes, and future concerns. *Ophelia* 41:199–220

O’Leary MH, Madhavan S, Paneth P (1992) Physical and chemical basis of carbon isotope fractionation in plants. *Plant Cell Env* 15:1099–1104

Otero E, Culp R, Noakes JE, Hodson RE (2003) The distribution and  $\delta^{13}\text{C}$  of dissolved organic carbon and its humic fraction in estuaries of southeastern USA. *Estuar Coast Shelf Sci* 56:1187–1194

Peters KE, Sweeney RE, Kaplan IR (1978) Correlation of carbon and nitrogen stable isotopes in sedimentary organic matter. *Limnol Oceanog* 23:598–604

- Peterson BJ, Howarth RW, Lipschutz F, Ashendorf D (1980) Salt marsh detritus: an alternative interpretation of stable carbon isotope ratios and fate of *Spartina alterniflora*. *Oikos* 34:173–177
- Peterson BJ, Fry B, Hullar M, Saupe S, Wright R (1994) The distribution and stable carbon isotope composition of dissolved organic carbon in estuaries. *Estuaries* 17:111–121
- Philippsen B, Olsen J, Lewis JP, Rasmussen P, Ryves DB, Knudsen KL (2013) Mid- to late-Holocene reservoir-age variability and isotope-based palaeoenvironmental reconstruction in the Limfjord, Denmark. *Holocene* 23:1017–1027
- Prahl FG, Bennett JT, Carpenter R (1980) The early diagenesis of aliphatic hydrocarbons and organic matter in sedimentary particulates from Dabob Bay, Washington. *Geochim Cosmochim Acta* 44:1967–1976
- Preston CD, Pearman DA, Dines TD (2002) *New Atlas of the British and Irish Flora*. Oxford University Press, Oxford
- Rashid MA (1985) *Geochemistry of Marine Humic Compounds*. Springer-Verlag, New York
- Rice DL, Hanson RB (1984) A kinetic model for detritus nitrogen: role of the associated bacteria in nitrogen accumulation. *Bull Mar Sci* 35:326–340
- Rosenbauer RJ, Swarzenski PW, Kendall C, Orem WH, Hostettler FD, Rollog ME (2009) A carbon, nitrogen, and sulfur elemental and isotopic study in dated sediment cores from the Louisiana shelf. *Geo-Mar Lett* 29:415–429
- Rundel PW, Ehleringer JR, Nagy KA (1988) *Stable Isotopes in Ecological Research*. Springer-Verlag.
- Ruttenberg KC, Goñi MA (1997) Phosphorus distribution, C:N:P ratios, and  $\delta^{13}\text{C}_{\text{OC}}$  in Arctic, temperate, and tropical coastal sediments: tools for characterizing bulk sedimentary organic matter. *Mar Geol* 139:123–145
- Sackett WM, Thompson RR (1963) Isotopic organic carbon composition of recent continental derived clastic sediments of Eastern Gulf Coast, Gulf of Mexico. *Bulletin of the American Association of Petroleum Geologists* 47:525–531
- Salomons W, Mook WG (1981) Field observations of the isotopic composition of particulate organic carbon in the southern North Sea and adjacent estuaries. *Marine Geology* 41:M11-M20
- Sampei Y, Matsumoto E (2001) C/N ratios in a sediment core from Nakaumi Lagoon, southwest Japan – usefulness as an organic indicator. *Geochem J* 35:189–205
- Schaeffer-Novelli Y, Cintron-Molero G, Soares MLG (2002) Mangroves as indicators of sea level change in the muddy coasts of the world. In Wang Y, Healy T (Eds.), *Muddy Coasts*. Elsevier, pp. 245-262
- Schidlowski M, Hayes JM, Kaplan IR (1983) Isotopic inferences of ancient biochemistries: carbon, sulphur, hydrogen and nitrogen. In: Scholf JW (Ed.), *Earth's Earliest Biosphere, Its Origin and Evolution*. Princeton University Press, Princeton, pp. 149–186
- Schleser GH (1995) Parameters determining carbon isotope ratios in plants. In: Frenzel B, Stauffer B, Weib M (Eds.), *Problems of Stable Isotopes in Tree-Rings, Lake Sediments and Peat Bogs as Climate Evidence for the Holocene*. Fischer Verlag, Stuttgart, pp. 71–96
- Seemann JR, Critchley C (1985) Effects of salt stress on the growth, ion content, stomatal behaviour and photosynthetic capacity of a salt-sensitive species, *Phaseolus vulgaris* L. *Planta* 164:151–162
- Seliger HH, Boggs JA, Biggley WH (1985) Catastrophic anoxia in the Chesapeake Bay in 1984. *Science* 228:70–73
- Sepúlveda J, Pantoja S, Hughen KA (2011) Sources and distribution of organic matter in northern Patagonia fjords, Chile (~44–47°S): A multi-tracer approach for carbon cycling assessment. *Continental Shelf Res* 31:315–329

- Shennan I, Tooley MJ, Green F, Innes J, Kennington K, Lloyd J, Rutherford M (1999) Sea level, climate change and coastal evolution in Morar, northwest Scotland. *Geol Mijnbouw* 77:247–262
- Shennan I, Lambeck K, Horton B, Innes J, Lloyd J, McArthur J, Purcell T, Rutherford M (2000) Late Devensian and Holocene records of relative sea-level changes in northwest Scotland and their implications for glacio-hydro-isostatic modelling. *Quaternary Sci Rev* 19:1103–1135
- Stahle DW, Cleaveland MK, Blanton DB, Therrell MD, Gay DA (1998) The lost colony and Jamestown droughts. *Science* 280:564–567
- St-Onge G, Hillaire-Marcel C (2001) Isotopic constraints of sedimentary inputs and organic carbon burial rates in the Saguenay Fjord, Quebec. *Mar Geol* 176:1–22
- Stout JD, Rafter TA, Troughton JH (1975) The possible significance of isotope ratios in palaeoecology. *B Roy S NZ* 13:279–286
- Sullivan MJ, Moncreiff CA (1990) Edaphic algae are an important component of salt marsh food-webs: evidence from multiple stable isotope analyses. *Mar Ecol Prog Ser* 62:149–159
- Talbot MR, Johannessen T (1992) A high resolution palaeoclimatic record for the last 27,500 years in tropical West Africa from the carbon and nitrogen isotopic composition of lacustrine organic matter. *Earth Planet Sci Lett* 110:23–37
- Thibodeau B, de Vernal A, Mucci A (2006) Recent eutrophication and consequent hypoxia in the bottom waters of the Lower St. Lawrence Estuary: Micropaleontological and geochemical evidence. *Mar Geol* 231:37–50
- Thornton SF, McManus J (1994) Applications of organic carbon and nitrogen stable isotope and C/N ratios as source indicators of organic matter provenance in estuarine systems: Evidence from the Tay Estuary, Scotland. *Estuar Coast Shelf Sci* 38:219–233
- Turney CSM (1999) Lacustrine bulk organic  $\delta^{13}\text{C}$  in the British Isles during the last Glacial-Holocene transition (14-9 ka  $^{14}\text{C}$  BP). *Arctic, Antarctic and Alpine Research* 31:71–81
- Tyson RV (1995) *Sedimentary Organic Matter: Organic Facies and Palynofacies*. Chapman and Hall, London
- Valiela I, Teal JM, Allen SD (1985) Decomposition in salt marsh ecosystems: the phases and major factors affecting disappearance of above-ground organic matter. *J Exp Mar Biol Ecol* 89:29–54
- Van Heemst JDH, Megens L, Hatcher PG, de Leeuw JW (2000) Nature, origin and average age of estuarine ultrafiltered dissolved organic matter as determined by molecular and carbon isotope characterization. *Org Geochem* 31:847–857
- Voss M, Larsen B, Leivuori M, Vallius H (2000) Stable isotope signals of eutrophication in coastal Baltic Sea sediments. *J Mar Syst* 25:287–298
- Wada E, Minagawa M, Mizutani H, Tsuji T, Imaizumi R, Karasawa K (1987) Biogeochemical studies on the transport of organic matter along the Otsuchi River watershed, Japan. *Estuar Coast Shelf Sci* 25:321–336
- Westman P, Hedenström A (2002) Environmental changes during isolation processes from the Litorina Sea as reflected by diatoms and geochemical parameters – a case study. *Holocene* 12:531–540
- Wilson GP, Lamb AL, Leng MJ, Gonzalez S, Huddart D (2005a) Variability of organic  $\delta^{13}\text{C}$  and C/N in the Mersey Estuary, U.K. and its implications for sea-level reconstruction studies. *Estuar Coast Shelf Sci* 64:685–698
- Wilson GP, Lamb AL, Leng MJ, Gonzalez S, Huddart D (2005b)  $\delta^{13}\text{C}$  and C/N as potential coastal palaeoenvironmental indicators in the Mersey Estuary, UK. *Quaternary Sci Rev* 24:2015–2029
- Woodroffe CD, Thom BG, Chappell J (1985) Development of widespread mangrove swamps in mid-Holocene times in northern Australia. *Nature* 317:711–713

- Woodroffe CD (1990) The impacts of sea level rise on Mangrove shorelines. *Prog Phys Geol* 14:483–520
- Wooller M, Smallwood B, Jacobsen M, Fogel M (2003a) Carbon and nitrogen stable isotopic variation in *Laguncularia racemosa* (L.) (white mangrove) from Florida and Belize: Implications for trophic level studies. *Hydrobiologia* 499:13–23
- Wooller M, Smallwood B, Scharler U, Jacobson M, Fogel M (2003b) A taphonomic study of  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  values in *Rhizophora* mangrove leaves for a multi-proxy approach to mangrove palaeoecology. *Org Geochem* 34:1259–1275
- Wooller MJ, Morgan R, Fowell S, Behling H, Fogel M (2007) A multiproxy peat record of Holocene mangrove palaeoecology from Twin Cays, Belize. *Holocene* 17:1129–1139
- Yamamuro M (2000) Chemical tracers of sediment organic matter origins in two coastal lagoons. *J Mar Syst* 26:127–134
- Yu F, Zong Y, Lloyd JM, Huang G, Leng MJ, Kendrick C, Lamb AL, Yim WW-S (2010) Bulk organic  $\delta^{13}\text{C}$  and C/N as indicators for sediment sources in the Pearl River delta and estuary, southern China. *Estuar Coast Shelf Sci* 87:618–630
- Zedler JB (1980) Algal mat productivity: comparisons in a salt marsh. *Estuaries* 3:122–131
- Zimmerman AR, Canuel EA (2002) Sediment geochemical records of eutrophication in the mesohaline Chesapeake Bay. *Limnol Oceanogr* 47:1084–1093
- Zong Y, Lloyd JM, Leng MJ, Yim WW-S, Huang G (2006) Reconstruction of Holocene monsoon history from the Pearl River Estuary, southern China, using diatoms and carbon isotope ratios. *Holocene* 16:251–163

## Figure Captions

**Fig. 1** Controls on the carbon isotope composition of organic matter in estuarine sediments. The main factor is the source of the organic matter, although decompositional processes and even analytical preparation techniques can affect  $\delta^{13}\text{C}$ . The main sources are aquatic and terrestrial. Aquatic plant  $\delta^{13}\text{C}$  is a function of the ratio of  $\text{CO}_2$  to  $\text{HCO}_3^-$ . Marine plants tend to utilise  $\text{HCO}_3^-$  (with higher  $\delta^{13}\text{C}$ ) so have higher  $\delta^{13}\text{C}$  values, whereas freshwater plants utilise  $\text{CO}_2$  (with lower  $\delta^{13}\text{C}$ ) so have lower  $\delta^{13}\text{C}$  values. In the aquatic environment where nutrients can be limiting, increased productivity can lead to higher  $\delta^{13}\text{C}$ . The main difference in the  $\delta^{13}\text{C}$  of terrestrial plants is their photosynthetic pathway; C3 plants have lower  $\delta^{13}\text{C}$  than C4. However, catchment processes can have a major impact on terrestrial plant  $\delta^{13}\text{C}$  including water stress, the amount of sunshine, the development of soils etc. The global carbon cycle, in particular the atmospheric  $\text{CO}_2$  budget affects all plants as does climate and human impacts

**Fig. 2** The range in weight C/N and  $\delta^{13}\text{C}$  values for the main sources of organic matter in the estuarine environment.  $\delta^{13}\text{C}$  is the biggest distinguisher between freshwater and marine as well as C3 and C4 terrestrial plants, while C/N is commonly used to separate aquatic from terrestrial. The fields of data are averages and “end-member” modern values are often measured at specific sites to narrow the ranges. (Modified from Lamb et al. 2006).

**Fig. 3** Schematic summary diagram illustrating the major controls on bulk sediment C/N and  $\delta^{13}\text{C}$  in a C3-dominated catchment. Five of the estuarine-type habitats discussed in the text are included: salt marsh, riverine estuary, mangrove swamp, lagoon and isolation basin. The C/N and  $\delta^{13}\text{C}$  signatures of the major components of estuarine organic matter are included: relative mixing of these components define the C/N and  $\delta^{13}\text{C}$  of estuarine organic matter, though decomposition and/or preparation techniques might also affect values. Fjords generally act in a similar manner to riverine estuaries, whilst C/N and  $\delta^{13}\text{C}$  might fluctuate in glacial embayments due to the presence of ice shelves, which can regulate the relative contribution of marine-sourced organic matter. Symbols: arrows denote inwash. Note,  $\delta^{13}\text{C}$  values are likely to be higher in a C4-dominated catchment due to utilisation of dissolved organic carbon from decomposition and oxidation of C4 plants. The star denotes a hypothetical core location to reconstruct long-term changes in estuarine C/N and  $\delta^{13}\text{C}$

**Fig. 4** Mean early to mid-Holocene and modern  $\delta^{13}\text{C}$  and C/N ratios from the various sub environments along the Mersey Estuary showing the transition from more freshwater-dominated supra-tidal to marine-dominated saltmarsh, channels and tidal flat environments (Wilson et al. 2005b)

**Fig. 5** Reconstruction of the Holocene river palaeodischarge history of the Pearl River Estuary from sediment cores taken in the Pearl River Estuary, southern China (Zong et al. 2006). C/N and  $\delta^{13}\text{C}$  were first calibrated from modern surface sediments which showed a transition from fully marine conditions at the margins of the South China Sea to fully freshwater conditions in the inner estuary. Diatom flora, C/N and  $\delta^{13}\text{C}$  were used as proxies for salinity and showed a significant increase in freshwater flux from 8500 cal yrs BP resulting from the enhanced summer monsoon regime in early Holocene. The strength of freshwater flux peaked between 7500 and 6000 cal yrs BP. In the last 6000 years, freshwater

flux decreased towards present, reflecting a gradual weakening of the summer monsoon (Zong et al. 2006)

**Fig. 6**  $\delta^{13}\text{C}_{\text{org}}$ ,  $\delta^{15}\text{N}_{\text{org}}$ , total organic carbon (TOC) and biomarker evidence for organic matter source (i.e. algal, terrestrial or bacterial) from core M3 (Zimmerman and Canuel, 2002) for the last ~500 years. Increases in algal- (and bacterial-) derived OM, relative to terrestrial matter is clearly shown in the biomarker record. Similar results are evident in the two other cores (Cores RD and RR; Zimmerman and Canuel, 2002)

**Fig. 7** Upper Loch nan Eala sediment core showing the diatom palaeosalinity reconstruction, C/N, and  $\delta^{13}\text{C}$ . The sequence is divided into seven zones based on changes in the  $\delta^{13}\text{C}$  and C/N data. These zones closely correlate with changes in the diatom flora, suggesting a strong salinity control on the type of the organic material entering the basin


**Fig. 8** Multi-proxy analysis of sediments from Moutonnee Lake on the Antarctic Peninsula. Proxies include  $\delta^{13}\text{C}$  of organic matter in bulk sediment, and foraminiferal and diatom concentrations. The bulk organic  $\delta^{13}\text{C}$  as well as the presence of marine- dominated organisms were used to prove the presence/absence of the ice shelves through the Holocene. Specifically, where the organic matter in the sediments has a high  $\delta^{13}\text{C}$  then there is more marine organic matter suggesting the absent of the ice shelves. When the sediments contain organic matter with low  $\delta^{13}\text{C}$  then there was likely to be a lid of freshwater and the shelves are more likely to have been present (from Bentley et al. 2005)








Figure 1



**Figure 2**




Figure 3





Figure 4

**Figure 5**




Figure 6





Figure 7

**Figure 8**