

1 **Accumulation of polybrominated diphenyl ethers and microbiome response**
2 **in the great pond snail *Lymnaea stagnalis* with exposure to nylon (polyamide)**
3 **microplastics**

4
5 Alice A. Horton^{† a,b,c*}, Lindsay K. Newbold^{† b}, Angela M Palacio-Cortés^d, David J. Spurgeon^b, M.
6 Glória Pereira^e, Heather Carter^e, Hyun S. Gweon^{a,f}, Martina G. Vijver^c, Peter M. van Bodegom^c,
7 Mario Antonio Navarro da Silva^d and Elma Lahive^b

8
9 ^a National Oceanography Centre, European Way, Southampton, SO14 SZH, UK.

10 ^b Centre for Ecology and Hydrology, Maclean Building, Benson Lane, Wallingford, Oxfordshire, OX10
11 8BB, UK.

12 ^c Institute of Environmental Sciences, University of Leiden, P.O. Box 9518, 2300 RA Leiden, The
13 Netherlands.

14 ^d Zoology department, Universidade Federal do Paraná, Avenida Coronel Francisco H. dos Santos,
15 Jardim das Americas, Curitiba - PR, 81531-981, Brazil.

16 ^e Centre for Ecology and Hydrology, Library Avenue, Lancaster Environment Centre, Lancaster,
17 Bailrigg, LA1 4AP, UK.

18 ^f School of Biological Sciences, University of Reading, Reading, RG6 6UR, UK.

19

20

21 [†]These authors contributed equally to this work

22 *Corresponding author. Email address: alihort@noc.ac.uk (A. A. Horton)

23 **Keywords**

24 Plastic, organic chemicals, PBDEs, flame retardants, bacteria, microbiology.

25 **Abstract**

26 Microplastics attract widespread attention, including for their potential to transport toxic chemicals in
27 the form of plasticisers and associated hydrophobic organic chemicals, such as polybrominated
28 diphenyl ethers (PBDEs). The aims of this study were to investigate how nylon (polyamide)
29 microplastics may affect PBDE accumulation in snails, and the acute effects of nylon particles and
30 PBDEs on survival, weight change and inherent microbiome diversity and community composition of
31 the pond snail *Lymnaea stagnalis*. Snails were exposed for 96 hours to BDEs-47, 99, 100 and 153 in
32 the presence and absence of 1% w/w nylon microplastics in quartz sand sediment. No mortality was
33 observed over the exposure period. Snails not exposed to microplastics lost significantly more weight
34 compared to those exposed to microplastics. Increasing PBDE concentration in the sediment resulted
35 in an increased PBDE body burden in the snails, however microplastics did not significantly influence
36 total PBDE uptake. Based on individual congeners, uptake of BDE 47 by snails was significantly
37 reduced in the presence of microplastics. The diversity and composition of the snail microbiome was
38 not significantly altered by the presence of PBDEs nor by the microplastics, singly or combined.
39 Significant effects on a few individual operational taxonomic units (OTUs) occurred when comparing
40 the highest PBDE concentration with the control treatment, but in the absence of microplastics only.
41 Overall within these acute experiments, only subtle effects on weight loss and slight microbiome
42 alterations occurred. These results therefore highlight that *L. stagnalis* are resilient to acute exposures
43 to microplastics and PBDEs, and that microplastics are unlikely to influence HOC accumulation or the
44 microbiome of this species over short timescales.

45

46 **1. Introduction**

47 Microplastics are a widely-recognised pollutant. The impacts of microplastics on biota and ecosystems,
48 and their interactions with other environmental pollutants under various environmental conditions, are
49 highly uncertain and existing studies have produced contradictory results (see discussions of expert
50 committee summarised in the report published by SAPEA (2019)). Due to the high affinity of
51 microplastic surfaces for hydrophobic organic chemicals (HOCs), there is potential for particles to sorb
52 HOCs (Hirai et al., 2011; Karapanagioti et al., 2011; Rochman et al., 2013b), which may lead to elevated
53 or reduced bioaccumulation of HOCs by organisms that ingest these microplastics (Bakir et al., 2016;
54 Besseling et al., 2013; Rochman et al., 2013a). However, other studies have not found clear evidence
55 for microplastics altering bioaccumulation or toxicity of HOCs (Ašmonaitė et al., 2018; Beiras and
56 Tato, 2019; Besseling et al., 2017; Horton et al., 2018). The question therefore remains as to whether
57 microplastics will significantly alter the impacts of HOCs on organisms.

58 Within the group of HOCs, polybrominated diphenyl ethers (PBDEs), brominated hydrocarbons
59 commonly used as flame-retardants, are one of the priority pollutant groups. They are found widely
60 throughout the environment (Guan et al., 2007; Hassanin et al., 2004), including in riverine sediments
61 (up to 16088 ng g⁻¹ dry weight total PBDEs in riverbank sediment in China (Luo et al., 2007)). Although
62 the majority of PBDE effects have been reported on mammals (He et al., 2011; Ji et al., 2011), PBDEs
63 have been shown to have a range of toxic effects on aquatic invertebrates including neurotoxicity,
64 genotoxicity and endocrine disrupting properties (Breitholtz and Wollenberger, 2003; Díaz-Jaramillo
65 et al., 2016).

66 As with other persistent organic pollutants, due to their relatively high log Kow, PBDEs sorb to
67 particulate and organic matter within the environment, and to fatty tissues of organisms where they can
68 bioaccumulate (Rahman et al., 2001). Where microplastics and PBDEs occur together, there is the
69 likelihood of interactions. One environmental study found microplastics from the Central Pacific Gyre
70 had surface concentrations of PBDEs up to 9900 ng g⁻¹ (Hirai et al., 2011). Teuten et al. (2009) found
71 lower concentrations of up to 57 ng g⁻¹ from microplastics at the same location, however both these
72 studies suggest the potential for such interactions to influence organism exposure. While some of these

73 measured concentrations likely derive from flame-retardant additives rather than sorbed chemicals, the
74 hydrophobic nature of PBDEs and widespread presence of PBDEs suggests that a proportion is sorbed
75 from the environment (Hirai et al., 2011; Mizukawa et al., 2009). Chua et al. (2014) and Rochman et
76 al. (2013a) have shown that the presence of microplastics within experimental systems can lead to
77 increased body burdens of PBDEs in amphipods and fish, with the type and concentration of
78 microplastics affecting the dynamics of bioaccumulation. Microplastics can also change the way in
79 which different PBDE congeners are accumulated, with higher brominated congeners more likely to be
80 accumulated when microplastics are present (Chua et al., 2014).

81 The gut microbiome is important for nutrition, metabolic function and immunity, with perturbations to
82 the microbial community understood to have implications for organism health and fitness (Licht and
83 Bahl, 2018; Zhu et al., 2018a). A number of studies have been carried out to determine the effects of
84 PBDEs on the gut microbiome of various organisms. Chen et al. (2018) investigated the effects of BDE-
85 71 on the gut microbiome of zebrafish, finding that, in the presence of BDE-71, bacterial diversity was
86 significantly reduced, and bacterial metabolic functioning was altered in a 7-day exposure. Li et al.
87 (2018) showed BDEs-47 and 99 to significantly affect the gut microbial diversity of mice, leading to
88 up- and down-regulation of 45 bacterial OTUs (5-day exposure), while Wang et al. (2018) also found
89 BDE-47 to also lead to a significant reduction in mouse gut microbial diversity and an alteration in the
90 community structure (21-day exposure). Studies have shown that microplastics can similarly alter the
91 gut microbiome of both vertebrates (Jin et al., 2018; Lu et al., 2018) and invertebrates (Zhu et al., 2018a;
92 Zhu et al., 2018b). These studies clearly show that microbiome alterations, expressed as species richness
93 and diversity, are a sensitive endpoint responding to HOC and microplastic exposure, even over short
94 timescales. Therefore, microbiome analysis together with host fitness could provide a fast screening
95 tool for assessing the effects of combined HOCs and microplastics during acute exposures.

96 The aim of this study was to investigate the effects of microplastics and PBDEs, individually and in
97 combination, on the accumulation, physiology and microbiome of the great pond snail *Lymnaea*
98 *stagnalis* (Linnaeus 1758). Molluscs have been shown to bioaccumulate organic chemicals (and metals)
99 as they lack the oxidase systems to metabolise xenobiotic substances (Geyer et al., 1982). These traits

100 make them well suited as a test organism for investigating organic pollutant accumulation (Amorim et
101 al., 2019). Although microplastics and PBDEs have been shown to individually alter the gut
102 microbiome of organisms once ingested, no studies to date have investigated the effects of co-exposure
103 to these pollutants with respect to microbiome responses. We hypothesise that increasing PBDE
104 sediment concentrations will lead to significant changes in the microbiome community (diversity and
105 composition) and that the presence of microplastics will reduce this effect through strong binding of
106 PBDEs, making them less bioavailable to microbiota within the gut. We also hypothesise that the
107 presence of microplastics will reduce PBDE accumulation in the snail.

108

109 **2. Materials and methods**

110 **2.1. Organisms**

111 Adult *Lymnaea stagnalis* were obtained from Blades Biological, UK, and were acclimatised for one
112 week under laboratory conditions prior to the exposure. Cultures were maintained and exposure studies
113 carried out using ISO artificial freshwater as recommended by the OECD for *L. stagnalis* (OECD,
114 2016). An air pump with an air stone was provided for system oxygenation. Stock cultures and
115 exposures were maintained at 20°C with a 16:8 h light:dark cycle. Snails in culture were fed well-
116 washed iceberg lettuce *ad libitum*.

117

118 **2.2. Microplastic particles**

119 Nylon 6 powder (mono-constituent substance, density 1.13 g cm⁻³) was purchased from Goodfellow
120 (Huntingdon, UK). This powder consisted of heterogeneous fragments <50 µm, with a mean size of
121 13-19 µm, measured using a Coulter Counter (Multisizer 3, Beckman, USA) and had been stained with
122 Nile Red dye.

123

124

125 **2.3. PBDEs**

126 Method 527 PBDE Mixture was purchased from LGC Standards (Teddington, UK). This mixture
127 contained BDE- 47, 99, 100, 153 and PBB- 153 (PBB-153 was not considered or measured throughout
128 this study), each at a concentration of 500 $\mu\text{g ml}^{-1}$ in ethyl acetate. With respectively log Kows of 6.81,
129 7.32, 7.24, and 7.9, these BDEs were all highly hydrophobic. These congeners are commonly detected
130 within aquatic organisms and have a high propensity for bioaccumulation (Hirai et al., 2011;
131 Shanmuganathan et al., 2011). A serial dilution was carried out in ethyl acetate in order to provide the
132 ultimate concentrations of each BDE congener in sediment of 3000, 1500, 750, 375, 188 and 94 ng g^{-1} .
133 These concentrations were chosen to reflect concentrations found within freshwater sediments (Luo et
134 al., 2007; Sellström et al., 1998; Yin et al., 2017).

135

136 **2.4. Experimental setup**

137 All experimental treatments (PBDEs and controls) consisted of either sediment without added
138 microplastics, or sediment with added microplastics (1% nylon powder by sediment mass, *i.e.* 10 g kg^{-1}
139 ¹). This is a higher microplastic concentration than has previously been found in freshwater sediments,
140 for example up to 71 mg kg^{-1} found in the Antuã River, Portugal (Rodrigues et al., 2018), and 1 g kg^{-1}
141 found by Klein et al. (2015) along the river Rhine in Germany. This concentration was therefore not
142 chosen to represent environmental relevance, but to enable maximum binding of PBDEs to a large
143 available microplastic surface area, with the intention of assessing how this may influence
144 bioaccumulation in a controlled and heavily contaminated exposure. Each microplastic treatment was
145 prepared by weighing 0.8 g nylon powder and mixing with white quartz sand (SiO_2 , particle size 210-
146 300 μm , Sigma-Aldrich, Poole, UK) to make up to 80 g. All treatments were also run without
147 microplastics, in which case only 80 g quartz sand was weighed into a vessel. For each PBDE treatment,
148 1 ml of each diluted PBDE stock was added to the 80 g quartz sand substrate (with or without
149 microplastics, hereafter referred to as ‘sediment’) and stirred for 2 minutes 30 seconds using a glass
150 rod. This bulk mixture was divided between six replicate 100 ml glass exposure vessels (13 g per

151 vessel). A solvent carrier was used for spiking the PBDEs into the sediment, therefore an ethyl acetate
152 solvent control was also set up (1.25 % ethyl acetate in sediment) by carrying out this procedure with
153 ethyl acetate only. Following dosing, the vessels were left under a fume hood for two days with
154 occasional agitation to ensure complete evaporation of the solvent. Blank controls containing no PBDEs
155 or solvent were also set up both with and without microplastics, but without the need for solvent
156 evaporation. This gave a total of 96 vessels to include blank controls (6 replicates), solvent controls (6
157 replicates) and PBDE treatments (6 concentrations x 6 replicates) without microplastics, and the same
158 treatments with 1% microplastics.

159 To prevent suspension of nylon particles due to water surface tension, a small spray bottle of ISO test
160 water was used to spray eight times onto the surface of the dry sediment. 100 ml of ISO test water was
161 then gently introduced to the vessel and the water surface sprayed another seven times to break the
162 water surface tension and allow any floating nylon particles to sink (15 sprays total). Vessels were left
163 to equilibrate for 48 hours prior to introducing the organisms.

164 One snail was exposed per vessel. Before being added to the test vessels, each snail was rinsed in ISO
165 test water and the shell gently rubbed with a gloved finger to remove any faeces/algae present and patted
166 dry with a tissue. Each snail was weighed and length of shell measured; only snails > 25mm were used
167 in the bioassays at which size all individuals can be expected to be mature (Coeurdassier et al., 2004;
168 Zonneveld and Kooijman, 1989).

169 During exposures, jars were covered with Parafilm® to prevent escape of snails, pierced 10 times to
170 allow for oxygenation. Exposures ran for 96 hours. No food was provided during test exposures in order
171 to run as a simple an exposure as possible, to avoid complex organic interactions which may confound
172 interpretation of the processes occurring. Snails were observed daily to check for mortality. At the end
173 of the exposure, snails were removed from the water, washed in DI water, patted dry with tissue and
174 weighed. Snails were euthanised and preserved: of the six replicate snails for each treatment, three were
175 preserved for microbiome analysis (directly placed into ethanol) and three for tissue PBDE
176 concentration analysis (immediately frozen at -80°C). Snails were not depurated before weighing or
177 preservation as it was decided that analysing organisms with a full gut would give a more natural

178 representation of environmental exposure and associated internal concentration. The overlying water
179 from the exposure vessels was poured away and sediments were dried in a temperature-controlled
180 chamber at 25°C until dry (approx. 2 days). Sediment PBDE concentrations were measured in the dried
181 samples at the end of the experiment.

182

183 **2.5. Chemical analysis**

184 Half of a snail was thawed, removed from the shell and dissected lengthways to obtain a representative
185 sample of the whole body. This tissue was then weighed, ground with sand and dried with anhydrous
186 sodium sulphate. The remaining half snail was refrozen and stored for fluorescence microscopy, to
187 verify microplastic ingestion. Each sample (snail/sediment) was spiked with labelled recovery standards
188 (¹³C BDE 47, ¹³C BDE 126 and ¹³C BDE 153; Cambridge Isotope Laboratories) and soxhlet extracted
189 in dichloromethane (DCM) for 16 h. A small portion of the extract was evaporated to zero volume and
190 the lipid content was determined gravimetrically. The remaining extract was cleaned using automated
191 size exclusion chromatography followed by deactivated (5% deionised water; w/w) alumina column.

192 The clean extract, was then spiked with labelled internal standards (BDE 77 and ¹³C BDE 138;
193 Cambridge Isotope Laboratories) and 100 µl of sample was injected into a GC-MS (Agilent) with
194 programmable temperature vaporization (PTV) inlet. The PTV injector was kept at 55°C for 0.45 min,
195 and heated to 325°C at a rate of 700°C min⁻¹ and kept at 325°C for 5 min. Then the temperature was
196 reduced to 315°C min⁻¹ at a rate of 10°C min⁻¹. The GC-MS had a 25 m HT8 column (0.22 mm internal
197 diameter and 0.25 µm film thickness, SGE Milton Keynes, UK) and the carrier gas was helium (2.0 ml
198 min⁻¹). The temperature programme was: isothermal at 80°C for 2.4 min, 25°C min⁻¹ to 200°C, 5°C
199 min⁻¹ to 315°C and was held at 315°C for 9.8 min. Residues were quantified using internal standard
200 method and also calibration curves of the standard PBDEs (Cambridge Isotope Laboratories) and were
201 recovery corrected. The mean recoveries were: ¹³C BDE 47- 85%, ¹³C BDE 126 – 105% and ¹³C BDE
202 153- 96% and the LOD was 0.109 ng g⁻¹ wet weight.

203

204 **2.6. Ingestion of microplastics**

205 The snail tissue remaining following the chemical analysis was analysed using a fluorescence
206 microscope (Olympus BX41 microscope with an Olympus U-LH100HG 100W mercury lamp using the
207 green filter of the Cy3 (Olympus U-M39004) filter cube, with Olympus analySIS software) to image
208 and qualitatively verify ingestion of microplastics by the snails.

209

210 **2.7. Microbiome analysis**

211 **2.7.1. DNA extraction and sequencing**

212 DNA was extracted from three snails per treatment (whole snail excluding shell) following the protocol
213 described in the SI. Sample DNA required an additional cleaning step through the application of
214 Genomic DNA Clean & Concentrator kit (Zymo research, USA) under the manufacturer's
215 recommended protocol. Resultant DNA was quantified using the nanodrop 8000 UV-Vis
216 spectrophotometer (ThermoFisher scientific, USA).

217 Approximately 40 ng of template DNA was amplified using Q5 high-fidelity DNA polymerase (New
218 England Biolabs, Hitchin, UK) each with a unique dual-index barcode primer combination (Kozich et
219 al., 2013). Individual PCR reactions employed 25 cycles of an initial 30 s, 98°C denaturation step,
220 followed by an annealing phase for 30 s at 53°C, and a final extension step lasting 90 s at 72°C. Primers
221 were based upon the universal primer sequence 341F and 806R (Takahashi et al., 2014). An amplicon
222 library consisting of ~550 bp amplicons spanning the V3-V4 hypervariable regions of encoding for the
223 16S small subunit ribosomal RNA gene (16S rRNA), was sequenced at a concentration of 6 pM with a
224 10% addition of control phiX DNA, on an Illumina MiSeq platform using V3 chemistry (Illumina Inc.,
225 San Diego, CA, USA).

226

227

228

229 **2.7.2. Bioinformatics analysis**

230 Sequenced paired-end reads were joined using VSEARCH (Rognes et al., 2016), quality filtered using
231 FASTX tools (hannonlab.cshl.edu), length filtered with the minimum length of 300 bp, presence of
232 PhiX and adapters were checked and removed with BBTools (jgi.doe.gov/data-and-tools/bbtools/), and
233 chimeras were identified and removed with VSEARCH_UCHIME_REF (Rognes et al., 2016) using
234 Greengenes Release 13_5 (at 97%) (DeSantis et al., 2006). Singletons were removed and the resulting
235 sequences were clustered into operational taxonomic units (OTUs) with VSEARCH_CLUSTER
236 (Rognes et al., 2016) at 97% sequence identity (Tindall et al., 2010). Representative sequences for each
237 OTU were taxonomically assigned by RDP Classifier with the bootstrap threshold of 0.8 or greater
238 (Wang et al., 2007) using the Greengenes Release 13_5 (full) (DeSantis et al., 2006) as the reference.
239 Unless stated otherwise, default parameters were used for the steps listed. The raw sequence data
240 reported in this study have been deposited in the European Nucleotide Archive under study accession
241 number PRJEB27672 (ERP109787).

242

243 **2.8. Statistical analysis**

244 **2.8.1. Chemistry data**

245 Sediment concentration and snail body concentration data were log transformed for normality. As only
246 one sediment concentration was measured per treatment, it was assumed that each of the three snails
247 analysed per treatment was exposed to this measured concentration. To compare the concentrations of
248 PBDEs in sediment and organisms with and without microplastics, only treatments with added PBDEs
249 were included in the analyses of chemical data (i.e. no control treatments) as the control treatments
250 showed very low or non-detected values which could not be log-transformed. Two-way ANOVAs were
251 carried out for each BDE congener, and the total PBDEs, to determine the relationship between snail
252 tissue concentration, the concentration of PBDEs in the sediment and the presence of microplastics (R
253 statistical software).

254

255 **2.8.2. Snail weight data**

256 A two-way ANOVA was conducted considering the effects on snail weight change of PBDE
257 concentration and presence of microplastics as factors, and also their interaction.

258

259 **2.8.3. Microbiome data**

260 After quality filtering, a total of 2626755 sequences remained. One sample was removed from the
261 analysis due to low sequencing efficiency (<6000 sequences). Rarefaction curves were used to ensure
262 the sample depth represented the full community. To account for uneven sequencing depth (inherent in
263 NGS platforms) samples were normalized to lowest sequence depth using the `rarefy_even_depth`
264 function in the R package 'Phyloseq V 1.22.3' (McMurdie and Holmes, 2013). For simplicity, for
265 microbiome analysis with respect to PBDE concentration, nominal PBDE concentrations were used. In
266 order to assess any subtle changes, communities were subdivided into 'core' OTUs (occurring in >50%
267 of samples, at an abundance of >2%) and 'non-core' (all other community members), using the function
268 'prevalence' in the R package 'microbiome' (McMurdie and Holmes, 2013). Analyses were firstly
269 carried out on the whole community and subsequently on the subdivided core and non-core
270 communities.

271 To visualise the relationship between 16 rRNA sequence-based community profiles from different
272 treatments, nonmetric multidimensional scaling (NMDS) was performed using the 'metaMDS'
273 function, based on dissimilarities calculated using the Bray–Curtis index. Additionally, bacterial
274 diversity were assessed using Fishers log series [alpha], as this is largely unaffected by sample sizes >
275 1000 (Magurran, 2004). Differences in bacterial diversity for each PBDE compound and nominal PBDE
276 concentration were tested through the multiple Kruskal-Wallis (H) test, a test which does not assume
277 data normality, using the function 'kruskalmc' in R package 'Pgirmess' version 1.6.9 (Giraudoux et al.,
278 2018). An additional Kruskal-Wallis test was run to determine whether there were differences in
279 microbiome diversity between control and solvent control treatments (Fig. 3). Similarity percentages
280 breakdown procedure (SIMPER) was used to infer the importance of community members within
281 treatments (Clarke, 1993) and again Kruskal-Wallis was used to test significance. Finally, the effect of
282 PBDE concentration, presence of microplastics and their interaction upon community dissimilarity was

283 assessed using the Bray–Curtis index through Permutational Multivariate Analysis of Variance
284 (PERMANOVA, using the ‘ADONIS’ function in R package ‘Vegan’ v2.0-10 (Anderson, 2001;
285 Oksanen et al., 2013)). Taxonomic composition was plotted using the R package ‘ggplot2’ (Wickham,
286 2016). For each treatment, relative abundances per treatment were calculated to account for unequal
287 sampling, taking into account the combined data of the three replicates (Figs. 5, S4 and S5).

288

289 **3. Results**

290 **3.1. Concentration of PBDEs in the presence and absence of microplastics**

291 The control treatments (no PBDEs, with and without microplastics) contained trace concentrations of
292 PBDEs in some instances, although most (overall 73%) were below the detection limit of 0.108 ng g⁻¹.
293 The concentrations of different PBDE congeners in relation to its nominal concentrations varied
294 between 41% and 74% (Table 1). When considering all congeners and concentrations both with and
295 without microplastics, measured sediment concentrations overall were 54% of the nominal
296 concentration. PBDEs were present within sediment at statistically significantly comparable
297 concentrations regardless of the presence or absence of microplastics ($p > 0.05$, ANOVA).

298 The (measured) sediment PBDE concentrations significantly affected PBDE uptake by snails, for all
299 PBDEs independently and combined, higher sediment concentrations resulted in a significantly higher
300 snail body burden ($p < 0.01$, ANOVA, Fig. 1). BDE 47 was the only PBDE congener that showed a
301 significant effect of microplastics on the uptake of PBDEs by snails, with microplastics leading to a
302 significantly lower body burden ($p < 0.01$, ANOVA, Fig. 1). There were no significant interactions
303 between the concentration of PBDEs within the sediment and the presence of microplastics for any of
304 the congeners ($p > 0.05$, two-way ANOVA).

305

306

307 Table 1. Nominal and measured sediment concentrations for each BDE congener, for all PBDE
 308 treatments with and without microplastics. Sediment PBDE concentrations were measured at the end
 309 of the experiment (one replicate per treatment).

		Measured sediment concentration (ng g ⁻¹)				
Nominal concentration		BDE 47	BDE 100	BDE 99	BDE 153	Total BDEs
(ng g ⁻¹ , per BDE)						
	0	0	0	0	0	0
	0 (solvent control)	0	0	0	0	0
	94	34.42	42.67	66.63	88.41	232.12
	188	46.80	67.44	108.19	109.85	332.27
Without	375	61.92	117.09	191.70	215.99	586.70
microplastics	750	233.24	332.20	499.45	456.24	1521.14
	1500	341.34	494.64	805.90	906.48	2548.36
	3000	1776.98	1765.79	2681.38	2290.35	8514.49
	0	0	0	0	0.41	0.41
	0 (solvent control)	0	0	6.93	0.54	7.47
	94	50.77	39.42	68.30	58.05	216.54
	188	56.24	50.58	78.46	81.65	266.92
With	375	170.47	141.11	209.81	197.16	718.55
microplastics	750	326.44	281.85	408.44	367.71	1384.44
	1500	825.13	667.41	978.23	832.01	3302.77
	3000	1449.14	1242.28	1804.11	1593.52	6089.05

310

3.2. Survival and weight change

There was 100% survival throughout the exposure. A significant difference was observed in snail wet weight change between microplastic and non-microplastic treatments, with non-microplastic treatments losing significantly more weight on average (0.11 ± 0.13 g) than microplastic treatments (0.03 ± 0.12

g) (two-way ANOVA, $p < 0.01$, Fig. 2). Concentration of PBDE had no effect on weight change (two-way ANOVA, $p > 0.05$) and there was no interaction between PBDEs and microplastics (two-way ANOVA, $p > 0.05$).

3.3. Ingestion of microplastics

Using fluorescence microscopy, ingestion of microplastics was evident within all snails exposed to microplastics (Fig. S1 D-I). Microplastics could be observed both on the surface of the sample (Fig. S1, G and I) and behind membranes (i.e. within organs, Figs. S1. D, F and H). Based on the way the samples were prepared and analysed, it is not possible to quantitatively analyse ingestion, nor to identify the specific locations where microplastics were found or accumulated. However, ingestion could be qualitatively confirmed.

3.4. Microbiome data

3.4.1. Control treatments

Using a multiple-comparison Kruskal-Wallis test, there were no significant differences in microbial diversity (Fisher's Log alpha) between blank controls and solvent controls, nor between control treatments with and without microplastics (Fig. 3. $p > 0.05$, Kruskal-Wallis). This highlights that there was no effect of the solvent control, or of microplastics alone (in the absence of PBDEs), on snail microbiome structure.

3.4.2. Community composition and diversity

Community diversity assessed by Fisher's log alpha (Fig. 4) showed no significant differences between different PBDE concentrations (all $p < 0.05$, multiple Kruskal-Wallis, H). However, although not significant, it should be noted that diversity does appear to appear to be lower at higher PBDE

concentrations when microplastics are absent, while the diversity of communities in treatments with microplastics appear largely unaffected by PBDE concentration (Fig. 4). This decline in diversity pattern could be related to a loss in richness (count) of non-core OTUs in treatment with high concentrations of PBDE, in the absence of microplastics (Fig. 4 and Fig. S2). Permutational Multivariate Analysis of Variance showed no significant effect on community dissimilarity based on PBDE concentration, presence of microplastics and their interactions ($p > 0.05$ for all comparisons, Fig. S3).

3.4.3. Taxonomic microbiome composition

The greatest number of 16S rRNA gene sequences within the *L. stagnalis* microbiome were found to be from the Gammaproteobacteria, Betaproteobacteria, Alphaproteobacteria, Flavobacteria and Bacilli (Fig. S4) irrespective of treatment. The most dominant order across all treatments are the Enterobacteriales (Fig. 5), and within that order the genus *Klebsiella* (Fig. S5). When comparing individual OTUs in the controls vs the highest PBDE concentration, similarity of percentage (SIMPER) analysis shows that in the absence of microplastics there was a significantly higher relative abundance of OTUs 5512 and 4432 (both identified as belonging to the Enterobacteriaceae) in the highest PBDE concentration treatment (Table S1). There was also a significant reduction in OTU 8733 (identified as belonging to the Flavobacteriaceae), in the highest dose treatment compared to the control (Kruskal-Wallis test $P < 0.05$, $df = 1$). In contrast, no significant differences were observed in individual relative OTU abundance when microplastics were present (Table S1). Some orders are present only in PBDE treatments, notably sulfate-reducing bacteria (Desulfobacteriales and Syntrophobacteriales).

4. Discussion

4.1. The snail microbiome

The individuals used in this test were taken from the field and acclimated for this test, therefore the microbial data acquired here is likely representative of biological variability within wild *L. stagnalis*. Laboratory conditions and the lack of food throughout the exposure would likely have led to changes over time, but this would not have been expected to lead to individual or treatment-specific differences. While the microbiome analysis in this study considered the whole microbiome of the snail, it is expected that the majority of OTUs derive from the gut bacterial community. This is highlighted in the communities across all treatments being dominated by Enterobacterales (Fig. 5), a common order comprising gut bacteria (Hu et al., 2018). For example, *Klebsiella*, the most dominant genus observed here within the family Enterobacteriaceae (Fig. S5), is a polysaccharide degrader linked to the presence of cellulase (Imran et al., 2016). The *L. stagnalis* core microbiome in this study appears to be similar to other freshwater snails and associated habitats including Enterobacterales, Flavobacterales and Bacillales corresponding to lactic acid production (food fermentation) and cellulose degradation (Béguin, 1990; Dar et al., 2017; Hu et al., 2018).

4.2. PBDE accumulation and effects on microbiome

All sediment PBDE concentrations were lower than the nominal concentrations when measured at the end of the exposures. The half-lives of BDEs-47, 99, 100 and 153 are all estimated to be approximately 14,400 hours in sediment (Wania and Dugani, 2003), therefore degradation over the experimental time period is not likely to have been a significant factor leading to the discrepancies between nominal and measured concentrations observed here (estimated 0.3% loss due to degradation over 96 hours based on a half-life of 14,400 hours). Some loss of PBDEs may have occurred as a result of volatilisation during the solvent evaporation step, and some may also have bound to the walls of the glass exposure vessels.

BDE 47 has the lowest log Kow, which would indicate a greater (although still low) partitioning into the water phase than for the other more hydrophobic PBDEs. In a marine study, Mizukawa et al. (2009) found that proportionally, higher brominated BDE congeners (BDE 209) associated most strongly with sediments, while the composition within overlying seawater was dominated by lower brominated congeners (predominantly BDE 47, but also including BDEs 99 and 100). In our study, BDE 47 followed by BDE 99 accumulated most in the snail tissue, with higher internal concentrations compared to the other congeners (Fig. 1). This corresponds with evidence which shows that BDEs 47 and 99 are the most bioavailable PBDE congeners, due to a lower molecular weight and smaller molecules than higher brominated congeners (Liang et al., 2010; Mizukawa et al., 2009; Watanabe and Sakai, 2003; Zhang et al., 2016).

There were a number of sulfate reducing bacteria observed within snails exposed to PBDEs (most notably Desulfobacterales and Syntrophobacterales, Fig. 5), bacteria also recognised to be associated with the debromination of PBDEs (Zhao et al., 2018). These bacteria have not commonly been described in relation to other freshwater snail species (Hu et al., 2018) and were not present within the controls. Burkholderiales, one of the dominant orders found within these snails across all treatments, are also associated with PBDE degradation, especially lower brominated congeners (Robrock et al., 2009).

PBDE concentration had no significant influence on the microbiome, a result which is in contrast to other studies which found that PBDEs affected bacterial community composition and diversity in sediments and within guts, with changes being congener-dependent (Li et al., 2018; Wang et al., 2018; Yen et al., 2009). We therefore reject the starting hypothesis that increasing PBDE sediment concentrations lead to significant structural changes in the microbiome community over an acute timescale. The difference between this and other studies is likely to be because these represent different exposure scenarios (via food or water), plus other studies have usually run for longer time periods, and have generally used much higher PBDE concentrations (e.g. $\mu\text{g g}^{-1}$ concentrations in food)

4.3. Effects of microplastics on snail physiology and microbiome

There was no effect of any exposure condition on survival. Microplastics did subtly affect the wet weight of the snails. In general, the weight of all snails declined throughout the experiment, likely due to the lack of food within the exposure. However, this decline was less pronounced in snails exposed to microplastics (average 0.03 g weight decline in microplastic-exposed snails, compared to average 0.11 g decline in non microplastic-exposed snails, Fig. 4). The reasons for this difference are not clear; most microplastic exposure studies observe a more pronounced weight decline in exposed organisms (Besseling et al., 2013; Zhu et al., 2018a).

The lack of significant influence of microplastics on the microbiome (Fig. 3) is in contrast to other studies on the microbiome response in invertebrates (Zhu et al., 2018a; Zhu et al., 2018b). For example, Zhu et al. (2018b) found a significant increase in the family Bacillaceae within collembolan guts following exposure to microplastics, while our analysis found the order Bacillales to be present in both the microplastic and non-microplastic treatments (Fig. 5). Many gut bacteria are derived from, and influenced by, ingested material, therefore feeding behaviour is likely to have a significant influence on the gut microbiome (Turnbaugh et al., 2009; Zhu et al., 2018b). It was chosen not to feed the snails during the acute exposure, and hence any alterations within the microbiome community could be ascribed solely to the microplastic, PBDEs and their interaction. Despite the lack of significance of microplastics alone, the microbiome analysis suggests that microplastics can subtly influence PBDE impacts on the microbiome. For example, while not significant, there appears to be a tendency for the diversity of non-core bacteria to be lower at higher PBDE concentrations in the absence of microplastics, a trend which is not evident when microplastics were present (Fig. 4, Fig. S2). Microplastics also appear to slightly reduce variability between individuals within the microplastic controls compared to non-microplastic controls i.e. ‘reference’ gut conditions (Figs. 3 and 4). Within natural conditions, a higher microbial diversity between individuals may be beneficial for populations, increasing resilience to perturbation (Heiman and Greenway, 2016; Lozupone et al., 2012).

4.4. Influence of microplastics and PBDE co-exposure on accumulation and microbiome

Microplastics did not influence sediment PBDE concentrations. This result was expected as the microplastics were not removed from the sediment samples before analysis, therefore during analysis, PBDEs were likely to have been extracted from both the sediment and microplastics simultaneously. The concentrations of PBDEs within the sediment significantly affected the amount of PBDEs taken up within the snail, in line with the expected relationship between external exposure concentration and snail body burden.

Given that snails were not depurated before chemical analysis of the whole body, this analysis took into account any chemicals present within the gut content, in addition to those in snail tissues. Microplastics did not influence the uptake of BDEs-99, 100, 153, nor PBDE uptake as a whole. Therefore these PBDEs were equally available regardless of the presence of microplastics and our hypothesis was not supported. This is in contrast to previous studies carried out on microplastic and PBDE interactions, where microplastics have been shown to enhance uptake of PBDEs into fish tissue (Rochman et al., 2013a) and amphipods (Chua et al., 2014). This could be a result of different polymer types used, different exposure conditions (e.g. temperature, pH) and also the susceptibility of the organism to accumulate or metabolise hydrophobic chemicals.

Previous studies have shown that PBDEs can transfer from microplastics into body tissues (Chua et al., 2014; Rochman et al., 2013a). Hence, the concentrations measured here are likely to be a combination of both gut content and tissue concentrations, especially as fluorescence analysis showed that the nylon particles were ingested by snails (Fig. S1). PBDEs entering tissues are unlikely to be taken up only by ingestion of contaminated particles, as the foot of the snail will be exposed to the sediment-based PBDEs by direct contact with the sediment, and to aqueous phase PBDEs through contact with the water phase (Bakir et al., 2016). To allow uptake into tissues, desorption of the chemical from the sediment (or microplastic) surface, whether externally or within the gut, is needed as a prelude to uptake. While it is anticipated that the main route of exposure to PBDEs was via the sediment (either dermally or via ingestion) (Mizukawa et al., 2009), aqueous phase uptake may also be important and the precise nature of exposure may also vary dependent on the behaviour of the BDE congener: BDE 47 was the

only PBDE whose concentration in snails was significantly reduced in the presence of microplastics. BDE 47 is the congener with the lowest log Kow at 6.81, which would be expected to sorb the least strongly to particles (both microplastics and sediment) compared to the other congeners (although it is still highly hydrophobic). This reduced binding affinity could have led to greater BDE 47 partitioning into the water phase in the absence of microplastics, facilitating uptake. The presence of microplastics may have increased the partitioning of BDE 47 to sediment through the addition of a further surface binding phase with a high affinity for HOCs, thus reducing BDE 47 in the more bioavailable water phase, resulting in reduced bioavailability and uptake (Fig. 1).

While microplastics can sorb chemicals, other media (e.g. organic matter, sediment) may also accumulate HOCs and therefore should be also be taken into account when considering pathways for exposure and bioavailability (Bakir et al., 2016; Koelmans et al., 2016). Further, if considering trophic transfer, the interactions with the sediment also indicate the importance of measuring organisms with a full gut, as we did within this study (rather than depurated organisms as is usually the case in chemical bioaccumulation studies), given that PBDEs associated with the gut content may also be bioavailable.

No consistent significant differences were observed in snail microbiome community diversity in response to either the microplastic or PBDE treatments, although a trend for reduced diversity at high PBDE concentrations in the absence of microplastics was suggested, which warrants further investigation. Hence, our hypothesis of chemical effects on the snail microbiome, influenced by microplastics, was not supported over the short exposure timescale used. When investigating the differences in abundance of specific OTUs, significant differences were seen in the abundance of Enterobacteriaceae and Flavobacteriaceae between the control and high PBDE concentration, only when microplastics were absent (Table S1). Enterobacterales can be induced to bloom within the gut under conditions of stress, for example inflammatory responses produced by the gut immune system (Stecher et al., 2012), which may explain their increase in the presence of high PBDE concentrations. Flavobacterales have been associated with polymer degradation (Mergaert and Swings, 1996; Nogales et al., 2011) and have been commonly found associated with marine plastic debris (Bryant et al., 2016; Oberbeckmann et al., 2018) which could explain their decline in the absence of microplastics (combined

with high PBDE concentrations), although it is not possible to link those characteristics directly to this study. The fact that these results were seen only in the absence of microplastics suggests that microplastics may be buffering the effects of PBDEs on the microbiota, although only subtly.

4.5. Long term implications and outlook

Short and long-term exposure are likely to lead to very different microbial community responses, therefore acute exposures can provide information on initial responses to perturbation that would be not be observed during chronic tests (Shade et al., 2012). There is evidence to suggest that microbiomes will respond very quickly to perturbations, for example a study by Yen et al. (2009) found that BDEs 153 and 154 rapidly and irreversibly changed the bacterial community within sediment (within 24 hours). Studies which have found significant changes in organism microbiomes following invertebrate exposure to microplastics usually run for longer timescales, e.g. enchytraeids exposed for seven days (Zhu et al., 2018a) and collembolans exposed for 56 days (Zhu et al., 2018b).

It is recognised that given the simplicity of this exposure, this does not well represent the complexity of an environmental exposure. However, the data here improve our understanding of the interactions between microplastics PBDEs and organisms within a simple controlled exposure, where any effects will be easier to observe and interpret than under complex realistic environmental conditions. The subtle variations in response of the snail microbiome to microplastic exposure, PBDE exposure and co-exposure over a 96-hour exposure indicated that these stressors do affect the structure of the gut community. However, overall response to aspects such as overall diversity were not evident to the same extent as for studies with other species conducted over longer exposure times. These results, therefore, highlight the complexity of responses of organisms to microplastics and organic chemicals, and show the importance of carrying out further studies to understand the interaction between microplastics and HOCs and their influence on organisms in a variety of exposure scenarios and time-scales.

5. Conclusions

Microplastics did not affect survival of the snails. The weight of all snails generally declined throughout the exposure period, however, this decline was less in snails exposed to microplastics. An increased concentration of PBDE in the sediment led to an increased body burden within the snails, however microplastics did not significantly influence this uptake when considering all PBDE congeners overall. BDE 47 was the only congener influenced by the presence of microplastics, leading to a significantly reduced internal concentration in the presence of microplastics. Overall, the diversity and composition of the snail microbiome was not significantly altered by the presence of PBDEs or microplastics, or both combined. However, when considering individual OTUs, significant effects on individual responses were found that can be functionally linked to the exposure of snails to the PBDEs added, a result only observed in the absence of microplastics. This suggests that microplastics influence how PBDEs will impact on specific OTUs. In summary, these results suggest that microplastics and PBDEs have a limited effect both individually and when combined on HOC accumulation and the microbiome of *Lymnaea stagnalis* within an acute exposure. However the subtle effects seen highlight the importance of carrying out further studies to better understand the mechanisms causing the interaction between microplastics and HOCs given that these relationships may become more pronounced over extended time-scales.

Acknowledgements

This study was funded by the Newton Fund, grant number UKNE/N00065X/1 awarded to the Universidade Federal do Paraná, Brazil and the Centre for Ecology and Hydrology, UK, and Conselho Nacional das Fundações Estaduais de Amparo à Pesquisa - Fundação Araucária, grant number 45191.460.30251.29012015. We would like to thank Alex Walton for characterising the nylon particles.

References

- Amorim, J., Abreu, I., Rodrigues, P., Peixoto, D., Pinheiro, C., Saraiva, A., Carvalho, A.P., Guimarães, L., Oliva-Teles, L., 2019. *Lymnaea stagnalis* as a freshwater model invertebrate for ecotoxicological studies. *Science of the Total Environment* 669, 11-28.
- Anderson, M.J., 2001. A new method for non-parametric multivariate analysis of variance. *Austral Ecology* 26, 32-46.
- Ašmonaitė, G., Larsson, K., Undeland, I., Sturve, J., Carney Almroth, B., 2018. Size Matters: Ingestion of Relatively Large Microplastics Contaminated with Environmental Pollutants Posed Little Risk for Fish Health and Fillet Quality. *Environmental Science & Technology* 52, 14381-14391.
- Bakir, A., O'Connor, I.A., Rowland, S.J., Hendriks, A.J., Thompson, R.C., 2016. Relative importance of microplastics as a pathway for the transfer of hydrophobic organic chemicals to marine life. *Environmental Pollution* 219, 56-65.
- Béguin, P., 1990. Molecular Biology of Cellulose Degradation. *Annual Review of Microbiology* 44, 219-248.
- Beiras, R., Tato, T., 2019. Microplastics do not increase toxicity of a hydrophobic organic chemical to marine plankton. *Marine Pollution Bulletin* 138, 58-62.
- Besseling, E., Foekema, E.M., van den Heuvel-Greve, M.J., Koelmans, A.A., 2017. The Effect of Microplastic on the Uptake of Chemicals by the Lugworm *Arenicola marina* (L.) under Environmentally Relevant Exposure Conditions. *Environmental Science & Technology* 51, 8795-8804.
- Besseling, E., Wegner, A., Foekema, E.M., van den Heuvel-Greve, M.J., Koelmans, A.A., 2013. Effects of microplastic on fitness and PCB bioaccumulation by the lugworm *Arenicola marina* (L.). *Environmental Science & Technology* 47, 593-600.
- Breitholtz, M., Wollenberger, L., 2003. Effects of three PBDEs on development, reproduction and population growth rate of the harpacticoid copepod *Nitocra spinipes*. *Aquatic Toxicology* 64, 85-96.
- Bryant, J.A., Clemente, T.M., Viviani, D.A., Fong, A.A., Thomas, K.A., Kemp, P., Karl, D.M., White, A.E., DeLong, E.F., 2016. Diversity and activity of communities inhabiting plastic debris in the North Pacific Gyre. *MSystems* 1, e00024-00016.

Chen, L., Hu, C., Lok-Shun Lai, N., Zhang, W., Hua, J., Lam, P.K.S., Lam, J.C.W., Zhou, B., 2018. Acute exposure to PBDEs at an environmentally realistic concentration causes abrupt changes in the gut microbiota and host health of zebrafish. *Environmental Pollution* 240, 17-26.

Chua, E.M., Shimeta, J., Nugegoda, D., Morrison, P.D., Clarke, B.O., 2014. Assimilation of Polybrominated Diphenyl Ethers from Microplastics by the Marine Amphipod, *Allorchestes Compressa*. *Environmental Science & Technology* 48, 8127-8134.

Clarke, K.R., 1993. Non-parametric multivariate analyses of changes in community structure. *Australian Journal of Ecology* 18, 117-143.

Coeurdassier, M., De Vauflery, A., Scheifler, R., Morhain, E., Badot, P.-M., 2004. Effects of cadmium on the survival of three life-stages of the freshwater pulmonate *Lymnaea stagnalis* (Mollusca: Gastropoda). *Bulletin of Environmental Contamination and Toxicology* 72, 1083-1090.

Dar, M.A., Pawar, K.D., Pandit, R.S., 2017. Gut Microbiome Analysis of Snails: A Biotechnological Approach, Organismal and Molecular Malacology. In: *Organismal and Molecular Malacology*, InTech.

DeSantis, T.Z., Hugenholtz, P., Larsen, N., Rojas, M., Brodie, E.L., Keller, K., Huber, T., Dalevi, D., Hu, P., Andersen, G.L., 2006. Greengenes, a chimera-checked 16S rRNA gene database and workbench compatible with ARB. *Applied and Environmental Microbiology* 72, 5069-5072.

Díaz-Jaramillo, M., Miglioranza, K.S.B., Gonzalez, M., Barón, E., Monserrat, J.M., Eljarrat, E., Barceló, D., 2016. Uptake, metabolism and sub-lethal effects of BDE-47 in two estuarine invertebrates with different trophic positions. *Environmental Pollution* 213, 608-617.

Geyer, H., Sheehan, P., Kotzias, D., Freitag, D., Korte, F., 1982. Prediction of ecotoxicological behaviour of chemicals: relationship between physico-chemical properties and bioaccumulation of organic chemicals in the mussel *Mytilus edulis*. *Chemosphere* 11, 1121-1134.

Giraudoux, P., Antonietti, J.-P., Beale, C., Pleydell, D., Treglia, M., 2018. pgirmess: Spatial Analysis and Data Mining for Field Ecologists. R package version 1.6.9.

Guan, Y.-F., Wang, J.-Z., Ni, H.-G., Luo, X.-J., Mai, B.-X., Zeng, E.Y., 2007. Riverine inputs of polybrominated diphenyl ethers from the Pearl River Delta (China) to the coastal ocean. *Environmental Science & Technology* 41, 6007-6013.

Hassanin, A., Breivik, K., Meijer, S.N., Steinnes, E., Thomas, G.O., Jones, K.C., 2004. PBDEs in European Background Soils: Levels and Factors Controlling Their Distribution. *Environmental Science & Technology* 38, 738-745.

He, P., Wang, A., Niu, Q., Guo, L., Xia, T., Chen, X., 2011. Toxic effect of PBDE-47 on thyroid development, learning, and memory, and the interaction between PBDE-47 and PCB153 that enhances toxicity in rats. *Toxicology and Industrial Health* 27, 279-288.

Heiman, M.L., Greenway, F.L., 2016. A healthy gastrointestinal microbiome is dependent on dietary diversity. *Molecular Metabolism* 5, 317-320.

Hirai, H., Takada, H., Ogata, Y., Yamashita, R., Mizukawa, K., Saha, M., Kwan, C., Moore, C., Gray, H., Laursen, D., Zettler, E.R., Farrington, J.W., Reddy, C.M., Peacock, E.E., Ward, M.W., 2011. Organic micropollutants in marine plastics debris from the open ocean and remote and urban beaches. *Marine Pollution Bulletin* 62, 1683-1692.

Horton, A.A., Vijver, M.G., Lahive, E., Spurgeon, D.J., Svendsen, C., Heutink, R., van Bodegom, P.M., Baas, J., 2018. Acute toxicity of organic pesticides to *Daphnia magna* is unchanged by co-exposure to polystyrene microplastics. *Ecotoxicology and Environmental Safety* 166, 26-34.

Hu, Z., Chen, X., Chang, J., Yu, J., Tong, Q., Li, S., Niu, H., 2018. Compositional and predicted functional analysis of the gut microbiota of *Radix auricularia* (Linnaeus) via high-throughput Illumina sequencing. *PeerJ* 6, e5537.

Imran, M., Saida, B., Ghadi, S.C., Verma, P., Shouche, Y.S., 2016. The gut-associated *Klebsiella* sp. of the apple snail produces multiple polysaccharide degrading enzymes.

Ji, K., Choi, K., Giesy, J.P., Musarrat, J., Takeda, S., 2011. Genotoxicity of several polybrominated diphenyl ethers (PBDEs) and hydroxylated PBDEs, and their mechanisms of toxicity. *Environmental Science & Technology* 45, 5003-5008.

Jin, Y., Xia, J., Pan, Z., Yang, J., Wang, W., Fu, Z., 2018. Polystyrene microplastics induce microbiota dysbiosis and inflammation in the gut of adult zebrafish. *Environmental Pollution* 235, 322-329.

Karapanagioti, H.K., Endo, S., Ogata, Y., Takada, H., 2011. Diffuse pollution by persistent organic pollutants as measured in plastic pellets sampled from various beaches in Greece. *Marine Pollution Bulletin* 62, 312-317.

Klein, S., Worch, E., Knepper, T.P., 2015. Occurrence and Spatial Distribution of Microplastics in River Shore Sediments of the Rhine-Main Area in Germany. *Environmental Science & Technology* 49, 6070-6076.

Koelmans, A.A., Bakir, A., Burton, G.A., Janssen, C.R., 2016. Microplastic as a Vector for Chemicals in the Aquatic Environment: Critical Review and Model-Supported Reinterpretation of Empirical Studies. *Environmental Science & Technology* 50, 3315-3326.

Kozich, J.J., Westcott, S.L., Baxter, N.T., Highlander, S.K., Schloss, P.D., 2013. Development of a dual-index sequencing strategy and curation pipeline for analyzing amplicon sequence data on the MiSeq Illumina sequencing platform. *Appl Environ Microbiol* 79, 5112-5120.

Li, C.Y., Dempsey, J.L., Wang, D., Lee, S., Weigel, K.M., Fei, Q., Bhatt, D.K., Prasad, B., Raftery, D., Gu, H., Cui, J.Y., 2018. PBDEs Altered Gut Microbiome and Bile Acid Homeostasis in Male C57BL/6 Mice. *Drug metabolism and disposition: the biological fate of chemicals* 46, 1226-1240.

Liang, X., Zhu, S., Chen, P., Zhu, L., 2010. Bioaccumulation and bioavailability of polybrominated diphenyl ethers (PBDEs) in soil. *Environmental Pollution* 158, 2387-2392.

Licht, T.R., Bahl, M.I., 2018. Impact of the gut microbiota on chemical risk assessment. *Current Opinion in Toxicology* In press.

Lozupone, C.A., Stombaugh, J.I., Gordon, J.I., Jansson, J.K., Knight, R., 2012. Diversity, stability and resilience of the human gut microbiota. *Nature* 489, 220-230.

Lu, L., Wan, Z., Luo, T., Fu, Z., Jin, Y., 2018. Polystyrene microplastics induce gut microbiota dysbiosis and hepatic lipid metabolism disorder in mice. *Science of the Total Environment* 631-632, 449-458.

Luo, Q., Cai, Z.W., Wong, M.H., 2007. Polybrominated diphenyl ethers in fish and sediment from river polluted by electronic waste. *Science of the Total Environment* 383, 115-127.

Magurran, A.E., 2004. *Measuring Biological Diversity*. Blackwell Publishing, Oxford, UK.

McMurdie, P.J., Holmes, S., 2013. phyloseq: an R package for reproducible interactive analysis and graphics of microbiome census data. *PLOS ONE* 8, e61217.

Mergaert, J., Swings, J., 1996. Biodiversity of microorganisms that degrade bacterial and synthetic polyesters. *Journal of industrial microbiology* 17, 463-469.

Mizukawa, K., Takada, H., Takeuchi, I., Ikemoto, T., Omori, K., Tsuchiya, K., 2009. Bioconcentration and biomagnification of polybrominated diphenyl ethers (PBDEs) through lower-trophic-level coastal marine food web. *Marine Pollution Bulletin* 58, 1217-1224.

Nogales, B., Lanfranconi, M.P., Piña-Villalonga, J.M., Bosch, R., 2011. Anthropogenic perturbations in marine microbial communities. *FEMS Microbiology Reviews* 35, 275-298.

Oberbeckmann, S., Kreikemeyer, B., Labrenz, M., 2018. Environmental Factors Support the Formation of Specific Bacterial Assemblages on Microplastics. *Frontiers in microbiology* 8, 2709-2709.

OECD, 2016. Test No. 243: *Lymnaea stagnalis* Reproduction Test. OECD Publishing.

Oksanen, J., Guillaume Blanchet, F., Kindt, R., Legendre, P., Minchin, P.R., O'Hara, R.B., Simpson, G.L., Solymos, P., Henry, M., Stevens, H., Wagner, H., 2013. Package vegan: community ecology package. R package version 2.0-7. R Foundation for Statistical Computing, Vienna, Austria.

Rahman, F., Langford, K.H., Scrimshaw, M.D., Lester, J.N., 2001. Polybrominated diphenyl ether (PBDE) flame retardants. *Science of the Total Environment* 275, 1-17.

Robrock, K.R., Coelhan, M., Sedlak, D.L., Alvarez-Cohen, L., 2009. Aerobic Biotransformation of Polybrominated Diphenyl Ethers (PBDEs) by Bacterial Isolates. *Environmental Science & Technology* 43, 5705-5711.

Rochman, C.M., Hoh, E., Kurobe, T., Teh, S.J., 2013a. Ingested plastic transfers hazardous chemicals to fish and induces hepatic stress. *Scientific Reports* 3, 3263.

Rochman, C.M., Manzano, C., Hentschel, B.T., Simonich, S.L., Hoh, E., 2013b. Polystyrene plastic: a source and sink for polycyclic aromatic hydrocarbons in the marine environment. *Environmental Science and Technology* 47, 13976-13984.

Rodrigues, M.O., Abrantes, N., Gonçalves, F.J.M., Nogueira, H., Marques, J.C., Gonçalves, A.M.M., 2018. Spatial and temporal distribution of microplastics in water and sediments of a freshwater system (Antuã River, Portugal). *Science of the Total Environment* 633, 1549-1559.

Rognes, T., Flouri, T., Nichols, B., Quince, C., Mahé, F., 2016. VSEARCH: a versatile open source tool for metagenomics. *PeerJ* 4, e2584.

SAPEA, 2019. A Scientific Perspective on Microplastics in Nature and Society. Science Advice for Policy by European Academies.

Sellström, U., Kierkegaard, A., de Wit, C., Jansson, B., 1998. Polybrominated diphenyl ethers and hexabromocyclododecane in sediment and fish from a Swedish River. *Environmental Toxicology and Chemistry* 17, 1065-1072.

Shade, A., Peter, H., Allison, S.D., Baho, D., Berga, M., Bürgmann, H., Huber, D.H., Langenheder, S., Lennon, J.T., Martiny, J.B., 2012. Fundamentals of microbial community resistance and resilience. *Frontiers in microbiology* 3, 417.

Shanmuganathan, D., Megharaj, M., Chen, Z., Naidu, R., 2011. Polybrominated diphenyl ethers (PBDEs) in marine foodstuffs in Australia: residue levels and contamination status of PBDEs. *Marine Pollution Bulletin* 63, 154-159.

Stecher, B., Denzler, R., Maier, L., Bernet, F., Sanders, M.J., Pickard, D.J., Barthel, M., Westendorf, A.M., Krogfelt, K.A., Walker, A.W., Ackermann, M., Dobrindt, U., Thomson, N.R., Hardt, W.-D., 2012. Gut inflammation can boost horizontal gene transfer between pathogenic and commensal *Enterobacteriaceae*. *Proceedings of the National Academy of Sciences* 109, 1269-1274.

Takahashi, S., Tomita, J., Nishioka, K., Hisada, T., Nishijima, M., 2014. Development of a prokaryotic universal primer for simultaneous analysis of bacteria and archaea using next-generation sequencing. *PLOS ONE* 9, e105592.

Teuten, E.L., Saquing, J.M., Knappe, D.R., Barlaz, M.A., Jonsson, S., Bjorn, A., Rowland, S.J., Thompson, R.C., Galloway, T.S., Yamashita, R., Ochi, D., Watanuki, Y., Moore, C., Viet, P.H., Tana, T.S., Prudente, M., Boonyatumanond, R., Zakaria, M.P., Akkavong, K., Ogata, Y., Hirai, H., Iwasa, S., Mizukawa, K., Hagino, Y., Imamura, A., Saha, M., Takada, H., 2009. Transport and release of chemicals from plastics to the environment and to wildlife. *Philosophical Transactions of the Royal Society B: Biological Sciences* 364, 2027-2045.

Tindall, B.J., Rosselló-Móra, R., Busse, H.-J., Ludwig, W., Kämpfer, P., 2010. Notes on the characterization of prokaryote strains for taxonomic purposes. *International Journal of Systematic and Evolutionary Microbiology* 60, 249-266.

Turnbaugh, P.J., Ridaura, V.K., Faith, J.J., Rey, F.E., Knight, R., Gordon, J.I., 2009. The effect of diet on the human gut microbiome: a metagenomic analysis in humanized gnotobiotic mice. *Science translational medicine* 1, 6ra14-16ra14.

Wang, D., Yan, J., Teng, M., Yan, S., Zhou, Z., Zhu, W., 2018. In utero and lactational exposure to BDE-47 promotes obesity development in mouse offspring fed a high-fat diet: impaired lipid metabolism and intestinal dysbiosis. *Archives of Toxicology* 92, 1847-1860.

Wang, Q., Garrity, G.M., Tiedje, J.M., Cole, J.R., 2007. Naive Bayesian classifier for rapid assignment of rRNA sequences into the new bacterial taxonomy. *Applied and Environmental Microbiology* 73, 5261-5267.

Wania, F., Dugani, C.B., 2003. Assessing the long-range transport potential of polybrominated diphenyl ethers: A comparison of four multimedia models. *Environmental Toxicology and Chemistry* 22, 1252-1261.

Watanabe, I., Sakai, S.-i., 2003. Environmental release and behavior of brominated flame retardants. *Environment International* 29, 665-682.

Wickham, H., 2016. *ggplot2: elegant graphics for data analysis*. Springer.

Yen, J.H., Liao, W.C., Chen, W.C., Wang, Y.S., 2009. Interaction of polybrominated diphenyl ethers (PBDEs) with anaerobic mixed bacterial cultures isolated from river sediment. *Journal of Hazardous Materials* 165, 518-524.

Yin, G., Zhou, Y., Strid, A., Zheng, Z., Bignert, A., Ma, T., Athanassiadis, I., Qiu, Y., 2017. Spatial distribution and bioaccumulation of polychlorinated biphenyls (PCBs) and polybrominated diphenyl ethers (PBDEs) in snails (*Bellamya aeruginosa*) and sediments from Taihu Lake area, China. *Environmental Science and Pollution Research* 24, 7740-7751.

Zhang, Y., Wang, W., Song, J., Ren, Z., Yuan, H., Yan, H., Zhang, J., Pei, Z., He, Z., 2016. Environmental Characteristics of Polybrominated Diphenyl Ethers in Marine System, with Emphasis on Marine Organisms and Sediments. *BioMed research international* 2016, 1317232-1317232.

Zhao, S., Rogers, M.J., Ding, C., He, J., 2018. Reductive Debromination of Polybrominated Diphenyl Ethers - Microbes, Processes and Dehalogenases. *Frontiers in microbiology* 9.

Zhu, B.-K., Fang, Y.-M., Zhu, D., Christie, P., Ke, X., Zhu, Y.-G., 2018a. Exposure to nanoplastics disturbs the gut microbiome in the soil oligochaete *Enchytraeus crypticus*. *Environmental Pollution* 239, 408-415.

Zhu, D., Chen, Q.-L., An, X.-L., Yang, X.-R., Christie, P., Ke, X., Wu, L.-H., Zhu, Y.-G., 2018b. Exposure of soil collembolans to microplastics perturbs their gut microbiota and alters their isotopic composition. *Soil Biology and Biochemistry* 116, 302-310.

Zonneveld, C., Kooijman, S., 1989. Application of a dynamic energy budget model to *Lymnaea stagnalis* (L.). *Functional Ecology*, 269-278.

Figure captions

Fig. 1. Measured PBDE concentrations in sediment, compared to the concentration within snails, for each BDE congener, with and without microplastics. 'No MP' = without microplastics, 'MP' = with microplastics. Fitted lines represent linear regression.

Fig. 2. Average snail wet weight change for all PBDE treatments (nominal concentration in ng g^{-1} within sediment) with and without microplastics, based on weight difference between 0 and 96 hours exposure. 'No MP' = without microplastics, 'MP' = with microplastics. Error bars represent standard deviation ($n = 6$).


Fig. 3: Comparison of changes in microbial community diversity of the snails (Fisher's Log alpha) in the blank and solvent controls, with and without microplastics. The boxes denote interquartile ranges (IQR) with the median as a black line. Whiskers represent lowest and highest values within 1.5 IQR ($n = 3$).

Fig. 4. Boxplots to show differences in microbial diversity in snails exposed to PBDEs in the absence and presence of microplastics, showing whole data for the whole microbial community, then subsequent separation into 'core' and 'non-core' community. The boxes denote interquartile ranges (IQR) with the median as a black line. Whiskers represent lowest and highest values within 1.5 IQR ($n = 3$).


Fig. 5. Order composition of bacterial communities at each nominal PBDE concentration (ng g^{-1}), with and without microplastics (present/absent). Relative abundance was calculated as rarefied number of sequences in OTU/total sequences in each sample (= 6359), relative abundances per treatment ($N=3$) are plotted on Y axis. For ease of representation taxa of an abundance of <0.02 (2%) from an individual

sample were excluded. Each column represents a PBDE concentration (0-3000 ng g⁻¹). *Note one sample was removed from this treatment due to inefficient sequencing, therefore $N=2$.


BDE 47



BDE 99



BDE 100



BDE 153

