

# Lichenometric dating: a commentary, in the light of some recent statistical studies

Tom Bradwell

*British Geological Survey, Murchison House, Edinburgh, UK*

## Abstract

This commentary article discusses the relative merits of new mathematical approaches to lichenometry. It highlights their strong reliance on complex statistics; their user unfriendliness; and their occasional mistreatment of existing lichenometric techniques. The article proposes that the success of lichenometric dating over the past 50 years has stemmed from its relative simplicity, transparency, and general field applicability. It concludes that any new techniques which ignore these principles are likely to be unjustified, unsuitable to the user community and inappropriate for the subject matter. Furthermore, the article raises a more general philosophical question: can statistical complexity and high precision in a 'geobotanical' dating technique, fraught with high degrees of environmental variability and in-built uncertainty, ever be scientifically valid?

## Introduction

Lichenometric dating has come a long way since its first use in the 1930s. Proposed as a relative dating technique by Knut Faegri (1934) and developed by Roland Beschel (1950, 1958, 1961, etc.), lichenometric dating has now been employed in over 600 studies worldwide and on all 7 continents. (See recent reviews by Noller and Locke 2001; Solomina and Calkin 2003; Muller 2006; Bradwell and Armstrong 2007; Benedict 2009). Various different methodologies and data collection techniques have been adopted – these range from measuring the single or several largest lichens on a surface to measuring whole populations of several thousand lichen thalli (Table 1). Measurement parameters also vary. The long axis, short axis, average diameter, the mean diameter of a number of lichens, the modal frequency of lichen sizes, and the percentage of lichen cover have all been used as metrics to estimate surface age. All of these sampling strategies have marked effects on the construction of lichenometric dating curves, the reported lichen 'growth' rate, and consequently the lichenometric age and precision of the surface being dated.

Lichenometry started out as a botanical science – field based in essence, primarily the domain of the ecologist or geographer. As its use as a dating technique became more established in the 1960s and 70s, lichens were measured more often by

42 geomorphologists and geologists eager to know the age of recent landforms,  
43 especially in high latitude and alpine settings. In the past decade, however, several  
44 papers have pushed lichenometry further towards the statistical sciences. Data  
45 collected in the field is now subjected to increasingly complex statistical procedures  
46 back in the office. In the past 3 years, 2 groups have presented lichen data using new  
47 and different statistical approaches: (1) The GEV (Generalized Extreme Value) group  
48 [Naveau *et al.* 2005, 2007; Cooley *et al.* 2006; Jomelli *et al.* 2007, 2008] and (2) The  
49  $U^2$  group [Orwin *et al.* 2008].

50

51 The GEV group aim to determine the age of a surface by modelling the lichen  
52 population distribution using a Bayesian treatment of Generalized Extreme Value  
53 (GEV) distribution theory. The authors go on to claim that each lichenometric surface  
54 is characterised in time by varying the GEV *location* and *scale* parameter functions,  
55 and is characterised in space by fixing the GEV *shape* parameter (Naveau *et al.* 2005).  
56 The whole process involves several complex steps, following collection of the field  
57 data, including: (1) generation of a statistical function considered to be a “growth  
58 curve”; (2) application of a Bayesian model; (3) many iterations using a Monte Carlo  
59 Markov Chain procedure to obtain parametric convergence; (4) computation of an  
60 expected ‘empirical’ distribution for each parameter; and finally (5) calculation of  
61 ‘surface-age’ and derivation of confidence intervals. In a recent assessment study of  
62 lichenometric dating techniques, Jomelli *et al.* (2007) find their GEV technique to be  
63 the best performing and most accurate method. The GEV group have repeated their  
64 statistical approach and their arguments several times in a number of recent similar  
65 publications (i.e. Naveau *et al.* 2007; Rabatel *et al.* 2007; Jomelli *et al.* 2007).

66

67 The second new approach is not a dating technique per se but a way to distinguish  
68 between lichen populations with different size-frequency distributions. The authors  
69 use the  $U^2$  statistic to group lichen populations and, after numerous statistical steps  
70 (e.g. observation ranking, cluster analysis and similarity matrices), to assign relative  
71 ages to recent glacial deposits and highlight complex depositional histories (Orwin *et*  
72 *al.* 2008).

73

74 Both new lichenometric approaches are novel and interesting but will probably be of  
75 limited use and applicability to the wider community. Essentially this is because they

76 are over-complicated and opaque to the non-statistician user. Unfortunately, both  
77 techniques also contain different flawed assumptions and inaccuracies. These are  
78 discussed within this article.

79

### 80 **Lichenometry as a dating technique**

81 Arguably, the beauty of lichenometry as a dating tool is its simplicity. It provides  
82 clear, powerful, quantitative results in a relatively quick, non-destructive and  
83 transparent way. It is particularly well suited to decoding Late Holocene glacial  
84 histories and has been used most often, and most successfully, in high latitude and  
85 alpine settings. In short, it has been demonstrated by many workers that a survey of  
86 largest-lichen diameters or lichen size-frequency distributions across recently  
87 deglaciated terrain will yield a good impression of the age of glacial landforms, whilst  
88 in the field. The size of the largest lichens acts as a good relative guide to the age of  
89 surfaces; which can be converted to absolute ages if a site-specific calibrated dating  
90 curve is available. It is this geobotanical phenomenon that was first noticed by Faegri  
91 and utilised by Beschel, and subsequently by many other workers in a wide range of  
92 settings. In its simplest form, lichenometry works well and can yield clear and  
93 meaningful results with very few intermediate steps or *a priori* assumptions. It is  
94 somewhat regrettable therefore that, in recent years, lichenometry has become  
95 removed from its humble origins and has started to lean too heavily on complex  
96 statistical approaches. It is particularly regrettable when these statistical approaches  
97 have not been shown to be appropriate to the lichenometric technique or to result in  
98 greater dating accuracy.

99

### 100 *Existing lichenometric techniques*

101 There are really only 4 different techniques in lichenometric dating:

- 102 1. The original approach of Beschel, often called the ‘traditional approach’ has  
103 been used to great effect many times since the 1950s. Beschel proposed that finding  
104 and measuring the largest lichen on a surface “growing under optimal environmental  
105 conditions” will result in the closest age-estimation (Beschel 1961: 1045).  
106 Consequently, this single largest lichen (LL) approach uses only the largest non-  
107 competing lichen of one species growing on an entire surface to derive a  
108 lichenometric age. The mean of the largest 5 lichens (5LL) on a surface was  
109 developed in the 1970s as a modification of the LL approach primarily to avoid

110 reliance on a single, potentially anomalous, lichen thallus. Others have chosen to use  
111 10 or more ‘largest lichens’, however several studies have shown that neither  
112 accuracy nor precision is improved by measuring more than the 5 largest lichens on a  
113 surface (e.g. Matthews 1975, 1994; Innes 1984). Some workers have chosen to use the  
114 LL or 5LL technique within a representative sample area (from 25-500 m<sup>2</sup>), when a  
115 whole-surface search is not practical. However, dating curves constructed using this  
116 fixed-area approach cannot be directly compared to those constructed using the LL on  
117 an entire surface, owing to the different sizes of the search areas (Innes 1983b, 1984).  
118 It is true that searching only part of a surface goes against the main assumption of the  
119 original LL technique, however as long as the same technique is used in the  
120 construction of the dating curve and for dating purposes the technique can be justified  
121 in most cases.

122 2. The fixed-area largest lichen (FALL) approach has been used, chiefly by Bull  
123 and co-workers, to ascertain the age and event history of diachronous surfaces.  
124 Essentially a development of the LL approach, this technique measures the single  
125 largest thallus of one species within a unit sample area. These sample areas, typically  
126 boulders, usually average ~1 m<sup>2</sup>. The measurements from one surface (c. 100-500) are  
127 pooled to allow statistical treatment and age projections. It is important to state that  
128 the FALL technique was specifically designed to study rockfall and talus  
129 accumulations where the age of the deposit may not be uniform (McCarroll 1993;  
130 Bull *et al.* 1994). Unlike the previous approaches, this technique is based on the  
131 assumption that lichen populations have a normal distribution of thallus sizes, and that  
132 the mean thallus size increases with surface age. Using the FALL technique, Bull and  
133 Brandon (1998) recorded an accuracy of +/-10 years on rockfall deposits up to 500  
134 years old in New Zealand.

135 3. The size-frequency approach (SF) was originally devised to identify multiple  
136 populations or anomalous, inherited or pre-existing, thalli growing on a single surface  
137 (Benedict 1967, 1985); but has since been used successfully as a relative and absolute  
138 dating technique (e.g. Caseldine 1991; Benedict 1999, 2009; Bradwell 2004; Bradwell  
139 *et al.* 2006). The SF approach has also been used to assess substrate stability, snow-  
140 kill frequency, lichen population structure and micro-environmental tolerance. It  
141 differs from the other techniques in that the operator records the long axis of all thalli  
142 of a single species growing within a representative sub-sample of the surface. Sample  
143 areas vary, but normally cover at least 25-50 m<sup>2</sup>, and may include between 200 to

144 5000 thalli. For best results, sample sizes of 1000 or more lichens are recommended  
145 (Benedict 2009). Whilst on smaller surfaces, every lichen should be measured.

146 4. The lichen cover approach (LC) is based on the assumption that the percentage  
147 of a rock surface covered by a single species of lichen will increase with time.  
148 Estimates of lichen cover are not common in lichenometric dating studies, although  
149 several authors have reported success in constructing relative chronologies using this  
150 technique (e.g. Birkeland 1973; Locke *et al.* 1979; Grab *et al.* 2005). The LC  
151 technique is the most subjective of the 4 lichenometric dating approaches (Innes  
152 1986a) and consequently is usually only used when the other 3 techniques are  
153 impractical. However, recent advances in digital image analysis may allow more  
154 quantitative lichen-cover studies to be performed (McCarthy and Zaniewski 2001).

155

156 All other lichenometric techniques are essentially modifications of one of these four  
157 methods. Most 'new' techniques merely use different statistical treatments of field  
158 data collected using one of the 4 techniques outlined above (i.e. LL/5LL, FALL, SF,  
159 LC). A powerful development of the LL technique was devised by Vanessa  
160 Winchester in the 1980s. She used multiple lichen species to derive several site-  
161 specific dating curves which, when used in combination, reduced uncertainty and  
162 improved accuracy (Winchester 1984). Using this multi-species approach, Winchester  
163 (1988) claimed precision of 1-2 years on stone monuments spanning the last 800 years  
164 in England. Surprisingly, few have adopted this technique to date recent glacial  
165 landforms – possibly owing to the lack of species diversity and the lack of control  
166 surfaces in many glacial environments.

167

168 Only the FALL approach makes assumptions about the size-frequency distribution of  
169 lichens on a surface. The SF approach measures, and therefore quantifies the precise  
170 size-frequency distribution of any given lichen population. The mathematical nature  
171 of the SF distribution on a specific surface, whether truncated log-normal, skewed,  
172 Poisson or otherwise, can only be determined from careful measurement of usually  
173 several hundred or more thalli. It is also worth stating that there is currently no  
174 consensus on the idealised nature of crustose lichen SF distributions (e.g. McCarthy  
175 1999). However, in young developmental populations, typical of those on Little Ice  
176 Age moraines, where space restriction is not a factor, statistical normality will

177 commonly apply (e.g. Innes 1983b, 1986b; Haines-Young 1988; McKinzey *et al.*  
178 2004).

179

180

### 181 **Recent statistical treatments of lichenometric data**

182 Processing lichenometric data and deriving absolute calendar ages for publication,  
183 with confidence intervals or error bars, is highly dependent on 2 things: the strength  
184 and validity of the dating-curve calibration; and the statistical treatment of the  
185 measurement data. Varying either of these 2 factors will produce widely differing  
186 results. The GEV group claim to build on a detailed statistical treatment published by  
187 McCarroll (1993, 1994). However, this lichenometric approach was principally  
188 devised to investigate geomorphic activity in multi-event deposits. Rather than using  
189 the size-frequency approach, which is best suited for dating single-event surfaces,  
190 McCarroll chose to modify the largest lichen approach to examine the age-frequency  
191 of avalanche boulders. As McCarroll (1993: 529) states in his study aims: “it is not  
192 the frequency distribution of lichens of different size that is of interest, but the  
193 frequency distribution of boulders of different age”. This study, and those of Bull and  
194 co-workers (1994, 1996, 1998) – who examined earthquake-generated rockfalls –  
195 have succeeded in using lichens to identify and date multi- and single-event deposits.  
196 But the GEV group go on to presume that all lichen-dating studies make the same  
197 assumptions made by McCarroll and Bull; whilst forgetting (or not recognising) that  
198 these authors were dealing with a specific modification of the lichenometric  
199 technique.

200

201 The GEV group criticise previous lichenometric techniques on the basis that “they  
202 assume that the largest lichens follow a Gaussian distribution” (Jomelli *et al.* 2007:  
203 137). However this is a misconception, and their statement may be based on a  
204 misunderstanding. The largest lichen in any population is by definition an extreme,  
205 hence why the largest lichens are far less numerous in any population, as found in  
206 many previous studies. But the “extreme” nature of the largest thalli does not require  
207 the statistical complexity of Generalized Extreme Value theory to calculate a  
208 lichenometric dating curve (or simply an age-size function) based on largest lichens.  
209 A calibrated age-size *dating curve* is simply an empirical relationship between the  
210 largest thallus (or mean of the 5 largest), assumed to be the oldest, and the surface age

211 of the feature, where the independent variable (x-axis) is time. There is no assumption  
212 of normal distribution in this procedure – Gaussian or otherwise. In its purest form,  
213 lichen dating curves can tell us, by interpolation, how old we should expect a certain-  
214 sized lichen to be. It is arguably this simplicity which has made the technique so  
215 useful to so many for so long.

216

217 The presentation of lichenometric dates has yet to be standardised, particularly  
218 regarding the calculation of confidence intervals. The GEV group claim (e.g. Cooley  
219 *et al.* 2006; Jomelli *et al.* 2007) that this as an inherent weakness in existing  
220 lichenometric approaches, and they attempt to devalue previous work which does not  
221 present the associated mathematical uncertainties. Jomelli *et al.* (2007: 140) criticize  
222 those studies which derive confidence intervals that “lack a mathematical foundation”.  
223 Instead they propose the use of their highly complex statistical approach (a Bayesian  
224 treatment of Generalized Extreme Value theory) in the perceived pursuit of greater  
225 precision and to calculate stronger mathematical confidence intervals (Cooley *et al.*  
226 2006; Naveau *et al.* 2006; Jomelli *et al.* 2007). They fail to recognise that  
227 uncertainties have been expressed quite succinctly and precisely in many ‘traditional’  
228 lichenometric studies (e.g. see Table 1). For dating curves constructed using the LL  
229 or 5LL, 2 standard deviations are preferred (95% confidence limits). The interpolated  
230 ages can be presented with the associated standard error, derived in the normal way,  
231 using (a) the lichen diameter, (b) the relevant calibration points, and (c) the value of  
232 the curve fitted through the calibration points at the relevant intersection. Any  
233 calibrated-age dating technique, such as lichenometry, will always be subject to the  
234 precision uncertainties of the field measurements combined with the construction of  
235 the calibration curve. These can be expressed and, in many cases, are incorporated  
236 into the derived lichenometric ages. If a new technique to derive mathematical  
237 uncertainty implies greater confidence than the original data warrants, regardless of its  
238 complexity, the technique risks serving no purpose. This is surely a major criticism of  
239 the new methodology proposed by the GEV group.

240

241 The SF technique makes use of a simple class-size statistical treatment in order to  
242 firstly determine the composition of the lichen population, whether it is unimodal or  
243 not, and secondly uses linear regression to determine the age of the population  
244 measured against a SF distribution ‘calibration curve’. This technique has had

245 considerable success both as a relative and an absolute dating technique, and is more  
246 statistically robust than the LL or 5LL techniques because of the large number of  
247 measurements which make up a single age-determination (Benedict 1985, 1999, 2009;  
248 Locke *et al.* 1979; Innes 1983b, 1986b; Caseldine 1991; Cook-Talbot 1991; Bradwell  
249 2004; Bradwell *et al.* 2006). It is not dependent on assumptions of statistical  
250 normality within lichen populations, although several studies have shown skewed  
251 normal distributions to be typical on young surfaces (e.g. Innes 1986b; Haines Young  
252 1988; Bradwell 2004; McKinzey *et al.* 2004). The SF approach is the least criticised  
253 by the GEV group in their assessment study of lichenometric dating techniques  
254 (Jomelli *et al.* 2007). However, they fail to see any advantages of the SF approach  
255 over their newly proposed GEV technique; and in conclusion Jomelli *et al.* (2007)  
256 omit the SF approach as a valid alternative to their own more statistically complex,  
257 and somewhat confusing, Bayesian GEV approach. The reason for this omission is  
258 not altogether clear, however it may be due to the construction of their experiment and  
259 a misunderstanding of the SF technique. Jomelli *et al.* (2007) could not perform the  
260 SF technique in one of their two test areas because they chose tombstones with small  
261 surface areas (typically  $<2 \text{ m}^2$ ). In the second test area, glacier forelands in the  
262 Bolivian Andes (Rabatel *et al.* 2006), the SF measurement data appear to have been  
263 collected unconventionally – possibly erroneously. Jomelli *et al.* (2007: 137) state that  
264 they measured at least 300 lichens “randomly selected” within a fixed area of  $50 \text{ m}^2$  –  
265 “1 lichen per block”. This is not the normal SF approach – which measures **all** thalli  
266 within a fixed area – and therefore their results cannot be compared with the  
267 conventional SF approach used by others (e.g. Benedict 1985, 1999, 2009; Innes  
268 1983b, 1986b; Bradwell 2004). This confused methodology, a mix of the SF and  
269 FALL techniques, may explain the apparent success of the GEV approach, as tested  
270 by Jomelli *et al.* (2007), over other more traditional lichenometric techniques such as  
271 the SF approach. Failure to recognise this flaw, along with the propagation of other  
272 false assumptions previously mentioned, seriously compromises the assessment study  
273 of Jomelli *et al.* (2007). Consequently, advocacy and adoption of the GEV method  
274 as the “most reliable” lichenometric dating technique (Jomelli *et al.* 2007: 131) is  
275 probably unjustified.

276

277 The complex statistical treatment proposed by Orwin *et al.* (2008) is not a dating  
278 method, but a technique which helps to identify lichen-colonized surfaces with similar

279 histories. Orwin *et al.* (2008) propose the use of the  $U^2$  statistic (Watson, 1961) to  
280 quantify the closeness of fit between any two lichen size-frequency distributions. The  
281  $U^2$  function has been used by statisticians for over 4 decades, but never before applied  
282 to lichenometry. Orwin *et al.*'s methodology is built around and based on the SF  
283 approach, and in fact uses the same dataset as the lichenometric study conducted by  
284 McKinzey *et al.* (2004).

285

286 The  $U^2$  technique may prove useful when examining lichen populations on multi-  
287 event surfaces. However, it is statistically cumbersome involving numerous steps,  
288 (observation ranking, cluster analysis and similarity matrices) whilst seeming to offer  
289 little in return. In many of the lichen populations from SE Iceland (Orwin *et al.* 2008;  
290 Fig. 3) visual inspection and simple statistics (i.e. mode, falling limb gradient or  
291 central tendency) easily describe their similarity or difference. Hence, the use of the  
292  $U^2$  statistic to distinguish between unimodal populations with markedly different SF  
293 gradients seems unnecessary and overcomplicated (e.g. HJ8708 & HJ8704 in Orwin  
294 *et al.* 2008; Fig. 3). The technique's ability to distinguish between complex or  
295 polymodal populations does represent a methodological advance. However, simple  
296 visual inspection can again prevent the inclusion of composite or polymodal  
297 populations in SF dating studies. This is important as older polymodal lichen  
298 populations cannot be dated with SF *age-gradient* curves (*sensu* Bradwell 2004) as  
299 they usually contain inherited thalli or multiple natality and mortality events (Innes  
300 1983b, 1986b; McCarthy 1999). Simply stated, the use of complex  $U^2$  statistics  
301 merely groups lichen populations with similar size-frequency distributions; it cannot  
302 decode moraine chronologies or the associated environmental conditions in any more  
303 detail than the lichen SF data itself. The use of this technique in "augmenting  
304 lichenometric surface dating" is suggested by Orwin *et al.* (2008: 151). However, it  
305 may offer little in uncomplicated, recessional moraine sequences; and it remains to be  
306 seen how the complex  $U^2$  statistics once generated can be applied to extract  
307 environmental information.

308

### 309 **Some ecological uncertainties**

310 Philosophically, it is hard to defend the use of high precision, highly complex  
311 statistics (such as those proposed by the GEV group) to solve what is essentially a  
312 simple problem: How can the size of lichens growing on a surface best inform us of

313 its age? Owing to the nature of the subject matter, uncertainty will always be high  
314 and hence dating precision will, in reality, always be low. Numerous ecological  
315 factors, central to the establishment and growth of the lichen thallus, determine this  
316 statement. A review of these factors, although probably timely, is far beyond the  
317 scope of this short article. However, it goes without saying that environmental  
318 conditions can vary greatly from site to site and even within sites. This can lead to  
319 problems when trying to calibrate or standardise field procedures, for instance when  
320 constructing a lichenometric dating curve. Uncertainties still surround the different  
321 growth rate of non-competing crustose lichens on surfaces with different aspect, slope  
322 angle, lithology, macro- and microclimate. Some of these topics remain largely  
323 unstudied, or are still being explored (e.g. Armstrong 1993, 2002, 2006, etc.). When  
324 combined with added uncertainties surrounding competition between thalli and  
325 between species (Armstrong and Welch 2007); differences in fungal (hypothallus)  
326 growth relative to algal (areolae) growth (Armstrong and Bradwell 2001); the impact  
327 and timing of mortality events (Loso and Doak 2006); and the importance of  
328 biological niches within certain environments (McCarthy 1999) – the range of factors  
329 likely to influence the growth rates of lichens becomes far greater. Even the exact  
330 nature of the growth curve in the most commonly used species in lichenometric dating  
331 (*R. geographicum*), although found to be non-linear over time, is still debated and in  
332 need of further study (cf. Proctor 1983; Matthews 1994; Bradwell and Armstrong  
333 2007). Careful research has shown that lichen growth is strongly controlled by  
334 moisture availability (Armstrong 1976, 2006; Benedict 1990). As a consequence,  
335 micro-environmental factors such as slope inclination (horizontal or vertical), surface  
336 orientation (to prevailing winds), surface texture and lithology may play an equally  
337 important role in determining growth rates alongside regional climatic conditions.  
338 Until the time when these key growth rate factors have been fully examined, and  
339 preferably quantified, in-built uncertainty will always surround the derivation of  
340 lichenometric dates even when local dating curves are used and field-measurement  
341 errors are minimised.

342

343

#### 344 **Summary**

345 This article attempts to dispel some of the current myths surrounding the statistical  
346 treatment of lichenometric data. Recent studies using complex statistics – most

347 notably by the GEV group – are attempting to overcomplicate what is a simple  
348 scientific technique. Many workers have successfully dated old surfaces using  
349 traditional lichenometric methods (see Table 1). In the North Atlantic region, the  
350 technique has enjoyed over 50 years of success and in this time has received several  
351 modifications and tweaks since the first studies by Roland Beschel. Importantly  
352 though, the long-established practice of comparing lichen sizes in the field with a  
353 carefully calibrated dating curve has proven effective for many workers in many  
354 countries over many years. So is it really time to adopt a new, considerably more  
355 complex, considerably less transparent technique, when the old one has not been  
356 found wanting? For this reason, advocacy of highly complex statistical techniques,  
357 such as the GEV approach, in pursuit of greater reliability or improved accuracy –  
358 over and above existing lichenometric techniques – seems premature and probably  
359 unjustified. These novel uses of statistics, whether to examine lichen populations  
360 growing on “similar historied surfaces” (Orwin *et al.* 2008), or to model uncertainties  
361 within idealised distributions (Naveau *et al.* 2007), leave the average potential user  
362 baffled by their complexity and inapplicability. Clearly, a good scientific technique is  
363 one which is only as complex as the subject matter warrants. In the case of  
364 lichenometry – a simple, user-friendly, field technique – the use of complex statistics  
365 is hard to support (e.g. Cooley *et al.* 2006; Naveau *et al.* 2007; Jomelli *et al.* 2007;  
366 Orwin *et al.* 2008) – particularly given the natural complexity and variability inherent  
367 within the lichen growing environment.

368

369 Whilst uncertainty still surrounds fundamental questions regarding lichen ecology,  
370 lichenometric dating will never be an exact science. In the meantime, any attempt to  
371 make it so should be viewed with caution and healthy scepticism. The lichen-dating  
372 community still awaits consensus on key questions relating to: the exact shape of the  
373 lichen growth curve; the typical size-frequency distribution for populations of  
374 different age; the effects of species competition; and the effects of temperature,  
375 precipitation and seasonality changes on lichen growth rates over many years. Lastly,  
376 on a more philosophical note (and maybe a suitable subtitle for this article), all this  
377 begs the question: can statistical complexity in pursuit of high precision ever be  
378 scientifically justified in a poorly understood ‘geobotanical’ dating technique?

379

380

381 **Acknowledgements**

382 I thank Richard Armstrong for discussions on lichen growth and lichen dating over  
383 the past 10 years. Communications with Richard have significantly shaped my  
384 thoughts on a wide range of lichen-related issues. Constructive and balanced reviews  
385 by Danny McCarroll, Olga Solomina and Wibjörn Karlén are gratefully  
386 acknowledged; although the views expressed in this paper are those of the author  
387 alone. Published with the permission of the Executive Director, BGS (NERC).

388

389 *Dr Tom Bradwell, British Geological Survey, Murchison House, West Mains Road,*  
390 *Edinburgh, EH9 3LA, UK*

391 *Email: tbrad@bgs.ac.uk*

392

393 **References**

394 *Anda, E. Orheim, O. and Mangerud, J., 1985: Late Holocene glacier variations and*  
395 *climate at Jan Mayen. Polar Research, 3: 129-145.*

396

397 *André, M.F., 1986: Dating slope deposits and estimating rates of rock wall retreat in*  
398 *Northwest Spitsbergen by lichenometry. Geografiska Annaler, 68(A): 65-75.*

399

400 *Armstrong, R.A., 1973: Seasonal growth and growth rate-colony size relationships in*  
401 *six species of saxicolous lichens. New Phytologist, 72: 1023-1030.*

402

403 *Armstrong, R.A., 1976: The influence of the frequency of wetting and drying on the*  
404 *radial growth of three saxicolous lichens in the field. New Phytologist, 77: 719-724.*

405

406 *Armstrong, R.A., 1983: Growth curve of the lichen Rhizocarpon geographicum.*  
407 *New Phytologist, 73: 913-918.*

408

409 *Armstrong, R.A., 1993: The growth of six saxicolous lichens transplanted to lime-rich*  
410 *and lime-poor substrates in south Gwynedd, Wales. Symbiosis, 15: 257-267.*

411

412 *Armstrong, R.A., 2002: The effect of rock surface aspect on growth, size structure and*  
413 *competition in the lichen Rhizocarpon geographicum. Environmental and*  
414 *Experimental Botany, 48: 187-194.*

415

416 *Armstrong, R.A., 2006: Seasonal growth of the crustose lichen Rhizocarpon*  
417 *geographicum (L.) DC. in South Gwynedd, Wales. Symbiosis, 41: 97-102.*

418

419 *Armstrong, R.A. and Bradwell, T., 2001: Variation in hypothallus width and the*  
420 *growth of the lichen Rhizocarpon geographicum (L.) DC. Symbiosis, 30: 317-328.*

421

422 *Armstrong, R.A. and Welch, AR., 2007: Competition in lichen communities.*  
423 *Symbiosis 43: 1-12.*

424

425 Bakke, J., Dahl, S.O., Paasche, O., Lovlie, R. and Nesje, A., 2005: Glacier  
426 fluctuations, equilibrium-line altitudes and palaeoclimate in Lyngen, northern  
427 Norway, during the Lateglacial and Holocene. *The Holocene*, 2005: 15: 518-540  
428

429 Ballantyne, C.K., 1990: The Holocene glacial history of Lyngshalvøya, northern  
430 Norway: chronology and climatic implications. *Boreas*, 19: 93-117.  
431

432 Benedict, J.B., 1967: Recent glacial history of an Alpine area in the Colorado Front  
433 Range, USA. 1: Establishing a lichen growth curve. *Journal of Glaciology*, 6: 817-  
434 832.  
435

436 Benedict, J.B., 1985: Arapaho Pass: Glacial geology and archaeology at the crest of  
437 the Colorado Front Range. *Center for Mountain Archaeology, Ward, Colorado,*  
438 *Research Reports*, 3: 1-197.  
439

440 Benedict, J.B., 1990: Experiments on lichen growth. I. Seasonal patterns and  
441 environmental controls. *Arctic and Alpine Research*, 22: 244-254.  
442

443 Benedict, J.B., 1999: Effects of changing climate on game-animal and human use of  
444 the Colorado High Country (U.S.A) since 1000BC. *Arctic, Antarctic and Alpine*  
445 *Research*, 31: 1-15.  
446

447 Benedict, J.B., 2009: Lichenometry for Archeologists: a Review. *American*  
448 *Antiquity*, 74: x-xx.  
449

450 Beschel, R.E., 1950: Flechten als Altersmassstab rezenter Moränen. *Zeitschrift für*  
451 *Gletscherkunde und Glazialgeologie*, 1: 152-161.  
452

453 Beschel, R.E., 1958: Lichenometrical studies in West Greenland. *Arctic*, 11: 254  
454

455 Beschel, R.E., 1961: Dating rock surfaces by lichen growth and its application to  
456 glaciology and physiography (lichenometry). In: G. O. Raasch (ed.): *Geology of the*  
457 *Arctic (Proceeding of the First International Symposium on Arctic Geology)*, Vol. 2.  
458 University of Toronto Press, Toronto: 1044-1062.  
459

460 Beschel, R.E., 1973: Lichens as a measure of the age of recent moraines. *Arctic and*  
461 *Alpine Research*, 5: 303-309. (Translation, by William Barr, of Beschel, R.E. 1950).  
462

463 Bickerton, R.W. and Matthews, J.A., 1992: On the accuracy of lichenometric dates:  
464 An assessment based on the 'Little Ice Age' moraine sequence at Nigardsbreen,  
465 southern Norway. *The Holocene*, 2: 227-237.  
466

467 Birkeland, P.W., 1973: The use of relative dating methods in a stratigraphic study of  
468 rock glacier deposits, Mt Sopris, Colorado. *Arctic and Alpine Research*, 5: 401-416.  
469

470 Bradwell, T., 2004: Lichenometric dating in southeast Iceland: the size-frequency  
471 approach. *Geografiska Annaler*, 86A: 31-41.  
472

473 *Bradwell, T., Dugmore, D.J. and Sugden, D.E., 2006: The Little Ice Age glacier*  
474 *maximum in Iceland and the North Atlantic Oscillation: evidence from*  
475 *Lambatungnajökull, southeast Iceland. *Boreas*, 35: 61-80.*  
476

477 *Bradwell, T. and Armstrong, R.A., 2007: Growth rates of *Rhizocarpon geographicum*:*  
478 *a review with new data from Iceland. *Journal of Quaternary Science*, 22: 311-320.*  
479

480 *Broadbent, N.D. and Berqvist, K.I., 1986: Lichenometric chronology and*  
481 *archaeological features on raised beaches: preliminary results from the Swedish north*  
482 *Bothnian coastal region. *Arctic and Alpine Research*, 18: 297-306.*  
483

484 *Bull, W.B., 1996: Dating San Andreas fault earthquakes with lichenometry. *Geology*,*  
485 *24: 111-114.*  
486

487 *Bull, W.B. and Brandon, M.T., 1998: Lichen dating of earthquake-generated regional*  
488 *rockfall events, Southern Alps, New Zealand. *Geological Society of America**  
489 *Bulletin*, 110: 60-84.  
490

491 *Bull, W.B., King, J., Kong, F., Moutoux, T. and Phillips, W.M., 1994: Lichen dating of*  
492 *coseismic landslide hazards in alpine mountains. *Geomorphology*, 10: 253-264.*  
493

494 *Caseldine, C.J., 1991: Lichenometric dating, lichen population studies and Holocene*  
495 *glacial history in Tröllaskagi, Northern Iceland. In: Maizels, J.K. and Caseldine, C.J.*  
496 *(Eds) *Environmental Change in Iceland: Past and Present*: 219-233. Kluwer,*  
497 *Dordrecht.*  
498

499 *Caseldine, C. and Baker, A., 1998: Frequency distribution of *Rhizocarpon**  
500 **geographicum* s.l., modeling, and climate variation in Tröllaskagi, northern Iceland.*  
501 **Arctic and Alpine Research*, 30: 175-183.*  
502

503 *Cook-Talbot, J.D., 1991: Sorted circles, relative-age dating and palaeoenvironmental*  
504 *reconstruction in an alpine periglacial environment, eastern Jotunheimen, Norway:*  
505 *lichenometric and weathering-based approaches. *The Holocene*, 1: 128-141.*  
506

507 *Cooley, D., Naveau, P., Jomelli, V., Rabatel, A. and Grancher, D., 2006: A Bayesian*  
508 *hierarchical extreme value model for lichenometry. *Environmetrics*, 17: 555-574.*  
509

510 *Evans, D.J.A., Butcher, C. and Kirthisingha, A.V., 1994: Neoglaciation and an early*  
511 *Little Ice Age in western Norway: lichenometric evidence from the Sandane area. *The**  
512 **Holocene*, 4: 278-289.*  
513

514 *Evans, D.J.A., Archer, S. and Wilson, D.J.H., 1999: A comparison of the*  
515 *lichenometric and Schmidt hammer dating techniques based on data from the*  
516 *proglacial areas of some Icelandic glaciers. *Quaternary Science Reviews*, 18: 13-41.*  
517

518 *Faegri, K., 1934: Über die Längenvariationen einiger Gletscher des Jostedalsbre und*  
519 *die dadurch bedingten Pflanzensukzessionen. *Bergens Museums Aarbog*, 1993: 137-*  
520 *142.*  
521

522 *Gordon, J.E. and Sharp, M., 1983: Lichenometry in dating recent glacial landforms*  
523 *and deposits, southeast Iceland. Boreas, 12: 191-200.*  
524

525 *Grab, S. van Zyl, C. and Mulder, N., 2005: Controls on basalt terrace formation in the*  
526 *eastern Lesotho Highlands. Geomorphology, 67: 473-485.*  
527

528 *Gudmundsson, H.J., 1998: Holocene glacier fluctuations of the Eiríksjökull ice cap,*  
529 *west central Iceland. Jökull, 46: 17-28.*  
530

531 *Haines-Young, R.H., 1983: Size variation of Rhizocarpon on moraine slopes in*  
532 *southern Norway. Arctic and Alpine Research, 15: 295-305.*  
533

534 *Haines-Young, R.H., 1988: Size-frequency and size-density relationships in*  
535 *populations from the Rhizocarpon sub-genus Cern. on moraine slopes in southern*  
536 *Norway. Journal of Biogeography, 15: 863-878.*  
537

538 *Hooker, T.N., 1980: Factors affecting the growth of Antarctic crustose lichens.*  
539 *British Antarctic Survey Bulletin, 50: 1-19.*  
540

541 *Innes, J.L., 1983a: The development of lichenometric dating curves for Highland*  
542 *Scotland. Transactions of the Royal Society of Edinburgh Earth Sciences, 74: 23-32.*  
543

544 *Innes, J.L., 1983b: Size-frequency distributions as a lichenometric technique: an*  
545 *assessment. Arctic and Alpine Research, 15: 285-294.*  
546

547 *Innes, J.L., 1984: The optimal sample size in lichenometric studies. Arctic and Alpine*  
548 *Research, 16: 233-244.*  
549

550 *Innes, J.L., 1985: Lichenometry. Progress in Physical Geography, 9: 187-254.*  
551

552 *Innes, J.L., 1986a: The use of percentage cover measurements in lichenometric*  
553 *dating. Arctic and Alpine Research, 18: 209-216.*  
554

555 *Innes, J.L., 1986b: The size-frequency distributions of the lichens Sporastatia*  
556 *testudinea and Rhizocarpon alpicola through time at Storbreen, south-west Norway.*  
557 *Journal of Biogeography, 13: 283-291.*  
558

559 *Innes, J.L., 1988: The use of lichens in dating. In: M. Galun (ed.) CRC Handbook of*  
560 *Lichenology. Volume III. CRC Press, Inc., Boca Raton: 75-91.*  
561

562 *Jomelli, V., Grancher, D., Naveau, P., Cooley, D. and Brunstein, D., 2007:*  
563 *Assessment study of lichenometric methods for dating surfaces. Geomorphology, 86:*  
564 *131-143.*  
565

566 *Jomelli, V., Grancher, D., Brunstein, D. and Solomina, O., 2008: Recalibration of the*  
567 *yellow Rhizocarpon growth curve in the Cordillera Blanca (Peru) and implications for*  
568 *LIA chronology. Geomorphology, 93: 201-212.*  
569

570 Jonasson, C., Kot, M. and Kotarba, A. 1991: Lichenometrical studies and dating of  
571 debris flow deposits in the High Tatra Mountains, Poland. *Geografiska Annaler*,  
572 73(A): 141-146.  
573

574 Kirkbride, M.P. and Dugmore, A.J., 2001: Can lichenometry be used to date the 'Little  
575 Ice Age' glacial maximum in Iceland? *Climatic Change*, 48: 151-167.  
576

577 Kugelmann, O., 1991: Dating recent glacier advances in the Svarfaðardalur–  
578 Skiðadalur area of northern Iceland by means of a new lichen curve. In: Maizels, J.K.  
579 and Caseldine, C. (Eds.) *Environmental Change in Iceland: Past and Present*: 203-  
580 217. Kluwer Academic Publishers.  
581

582 Locke, W.W., Andrews, J.T. and Webber, P.J., 1979. *A manual for lichenometry*.  
583 British Geomorphological Research Group Technical Bulletin 26.  
584

585 Loso, M.G. and Doak, D.F., 2006: The biology behind lichenometric dating curves.  
586 *Oecologia*, 147: 223–229.  
587

588 Macklin, M.G., Rumsby, B.T. and Heap, T., 1992: Flood alluviation and  
589 entrenchment: Holocene valley floor development and transformation in the British  
590 Uplands. *Geological Society of America Bulletin*, 104: 631-643.  
591

592 Matthews J.A., 1975: Experiments on the reproducibility and reliability of  
593 lichenometric dates, Storbreen gletschervorfeld, Jotunheimen, Norway. *Norsk*  
594 *Geografisk Tidsskrift*, 29: 97-109.  
595

596 Matthews, J.A., 1994: Lichenometric dating: A review with particular reference to  
597 'Little Ice Age' moraines in southern Norway. In: Beck, C. (ed.) *Dating in Surface*  
598 *Contexts*. New Mexico University Press: 185-212.  
599

600 Matthews J.A., 2005: 'Little Ice Age' glacier variations in Jotunheimen, southern  
601 Norway: a study in regionally controlled lichenometric dating of recessional moraines  
602 with implications for climate change and lichen growth rates. *The Holocene* 15: 1-19.  
603

604 McCarroll, D., 1993: Modelling late-Holocene snow-avalanche activity;  
605 incorporating a new approach to lichenometry. *Earth Surface Processes and*  
606 *Landforms*, 18: 527–539.  
607

608 McCarroll, D., 1994: A new approach to lichenometry: dating single-age and  
609 diachronous surfaces. *The Holocene*, 22: 383–396.  
610

611 McCarroll, D. Shakesby, R.A. and Matthews, J.A., 2001: Enhanced rockfall activity  
612 during the Little Ice Age: Further lichenometric evidence from a Norwegian talus.  
613 *Permafrost and Periglacial Processes*, 12: 157-164.  
614

615 McCarthy, D.P., 1999: A biological basis for lichenometry? *Journal of*  
616 *Biogeography*, 26: 379-386.  
617

- 618 McCarthy, D.P. and Zaniewski, K., 2001: Digital analysis of lichen cover: a technique  
619 for use in lichenometry and lichenology. *Arctic, Antarctic, and Alpine Research*, 33:  
620 107-113.  
621
- 622 McKinzey, K., Orwin, J. and Bradwell, T., 2004: Re-dating the moraines at  
623 Heinabergsjokull and Skalafellsjokull using different lichenometric methods:  
624 implications for the timing of the Icelandic Little Ice Age maximum. *Geografiska*  
625 *Annaler*, 86A: 319-336.  
626
- 627 Müller, G., 2006: Lichenometry and environmental history. *Environmental History*  
628 11: 604-609.
- 629 Naveau, P., Nogaja, M., Ammann, C., Yiou, P., Cooley, D. and Jomelli, V., 2005:  
630 Statistical methods for the analysis of Geophysical extreme events. *Comptes Rendus*  
631 *de l'Académie des Sciences*, 337: 1013–1022.
- 632 Naveau, P., Jomelli, V., Cooley, D. and Rabatel, A., 2007: Modeling uncertainties in  
633 lichenometry studies with an application: the Tropical Andes (Charquini Glacier in  
634 Bolivia). *Arctic, Antarctic, and Alpine Research*, 39: 277-285.
- 635 Noller J.S. and Locke W.W., 2001: Lichenometry. In *Quaternary Geochronology:*  
636 *Methods and Applications*, Noller, JS *et al.* (eds). American Geophysical Union:  
637 Washington DC; 261-272.  
638
- 639 Orwin, J.F., McKinzey, K.M., Stephens, M.A. and Dugmore, A.J., 2008: Identifying  
640 moraine surfaces with similar histories using lichen size distributions and the U<sup>2</sup>  
641 statistic, southeast Iceland. *Geografiska Annaler*, 90A: 151–164.  
642
- 643 Principato, S.M., 2008: Geomorphic evidence for Holocene glacial advances and sea  
644 level fluctuations on eastern Vestfirðir, northwest Iceland. *Boreas*, 37: 132–145.  
645
- 646 Proctor M.C.F., 1983: Sizes and growth-rates of thalli of the lichen *Rhizocarpon*  
647 *geographicum* on the moraines of the Glacier de Valsorey, Valais, Switzerland. *The*  
648 *Lichenologist* 15: 249-261.  
649
- 650 Rabatel, A., Jomelli, V., Francou, B., Naveau, P. and Grancher, D., 2005: Dating the  
651 Little Ice Age in the tropics from the moraines of Charquini Glaciers (Andes of  
652 Bolivia, 16°S). *Comptes Rendus de l'Académie des Sciences*, 337: 1311–1322.  
653
- 654 Rapp, A. and Nyberg, R., 1981: Alpine debris flows in Northern Scandinavia.  
655 *Geografiska Annaler*, 63: 183-196.  
656
- 657 Solomina, O. and Calkin, P.E., 2003: Lichenometry as applied to moraines in Alaska,  
658 USA and Kamchatka, Russia. *Arctic, Antarctic, and Alpine Research*, 35: 129-143.  
659
- 660 Thompson, A. and Jones, A., 1986: Rates and causes of proglacial river terrace  
661 formation in southwest Iceland: an application of lichenometric dating techniques.  
662 *Boreas*, 15: 231-246.  
663
- 664 Watson, G.S., 1961: Goodness-of-fit tests on a circle. *Biometrika*, 48: 109–114.

665

666 Winchester, V., 1984: A proposal for a new approach to lichenometry. *British*  
667 *Geomorphological Research Group, Technical Bulletin*, 33: 3-20.

668

669 Winchester, V., 1988: An assessment of lichenometry as a method for dating recent  
670 stone movements in two stone circles in Cumbria and Oxfordshire. *Botanical Journal*  
671 *of the Linnean Society*, 96: 57-68.

672

673 Winchester, V. and Chaujar R.K., 2002: Lichenometric dating of slope movements,  
674 Nant Ffrancon, North Wales. *Geomorphology* 47: 61-74.

675

676 Winkler S., Matthews J.A., Shakesby R.A. and Dresser P.Q., 2003: Glacier variations  
677 in Breheimen, southern Norway: dating Little Ice Age moraine sequences at seven  
678 low-altitude glaciers. *Journal of Quaternary Science*, 18: 359-413.

679

Table 1. A cross-section of lichenometric dating studies conducted in northern Europe since 1980.

author(s)	date <sup>1</sup> (AD)	location	lichen species <sup>2</sup>	technique <sup>3</sup>	lichen dimension	no. of lichens recorded <sup>4</sup>	survey area <sup>5</sup> (m <sup>2</sup> )	calibration surfaces <sup>6</sup>	uncertainty expressed
Rapp and Nyberg	1981	Abisko Mtns, Sweden	<i>R. geographicum</i> agg.	LL	long axis	1	variable	ex. curve	no
Innes	1983	Scottish Highlands	<i>R. section Rhizocarpon</i>	LL	long axis	1	entire	gravestones	no
Gordon and Sharp	1983	Breiðamerkurjökull and Skálafellsjökull, Iceland	<i>R. geographicum</i> agg. <i>R. geographicum</i> agg.	5LL 5LL	short axis long axis	5 5	1500 150	moraines moraines	yes yes
Anda <i>et al.</i>	1985	Jan Mayen	<i>R. geographicum</i>	LL	long axis	1	entire	moraines	no
Thompson and Jones	1986	Öræfi, SE Iceland	<i>R. geographicum</i> agg.	5LL	short axis	5	entire	moraines	yes
Broadbent and Bergqvist	1986	Bothnia coast, Sweden	<i>Rhizocarpon</i> subgenus	LL, SF	long axis	203	entire	raised beaches	yes
Andre	1986	NW Spitsbergen	<i>R. subgen. Rhizocarpon</i>	LL	long axis	1	variable	n/a	n/a <sup>7</sup>
Winchester	1988	Cumbria, England	<i>R. geographicum</i> subsp.	LL	long axis	1	entire	gravestones	no
Ballantyne	1990	Lyingshalvoya, Norway	<i>Rhizocarpon</i> subgenus	5LL, SF	long axis	100-400	variable	gravestones	no
Kugelmann	1991	Skiðadalur, Iceland	<i>R. geographicum</i> agg.	LL	long axis	1	entire	gravestones	yes
Cook-Talbot	1991	Jotunheimen, Norway	<i>R. geographicum</i> agg.	5LL, SF	long axis	300	variable	ex. curve	no
Jonasson <i>et al.</i>	1991	High Tatra Mtns, Poland	<i>R. geographicum</i>						
Caseldine	1991	Tröllaskagi, Iceland	<i>R. geographicum</i> s.l.	SF	long axis	1000	variable	debris flows	n/a <sup>7</sup>
Macklin <i>et al.</i>	1992	North Pennines, England	<i>R. geographicum</i> and <i>Huilia tuberculosa</i>	3LL	long axis	3	variable	gravestones, bridges	no
Bickerton and Matthews	1993	Jostedalsbreen, Norway	<i>Rhizocarpon</i> subgenus	LL, 5LL	long axis	5	c. 430	ex. curve	yes
McCarroll	1993	Jostedalen, W Norway	<i>R. geographicum</i> agg.	FALL	long axis	100	<2	ex. curve	yes
Evans <i>et al.</i>	1994	Sandane, W Norway	<i>R. section Rhizocarpon</i>	5LL	long axis	5	20	ex. curve	no
Gudmundsson	1998	Eiríksjökull, Iceland	<i>R. geographicum</i>	5LL	short axis	5	entire	ex. curve	no
Evans <i>et al.</i>	1999	Vatnajökull, Iceland	<i>R. geographicum</i> s.l.	5LL	long axis	5	entire	m, sh, br, g	no
McCarroll <i>et al.</i>	2001	Hurrungane, W Norway	genus <i>Rhizocarpon</i>	FALL	long axes	100	<2	ex. curve	yes
Kirkbride and Dugmore	2001	Eyjafjallajökull, Iceland	<i>R. geographicum</i>	LL, 5LL, SF	long axis	>250	50-100	m, fd	no
Winchester and Chaujar	2002	North Wales	<i>R. geographicum</i> subsp.	SF	long axis	100-500	variable	gravestones	no
Winkler <i>et al.</i>	2003	Breheimen, Norway	<i>Rhizocarpon</i> subgenus	LL, 5LL	long axis	5	variable	ex. curve	no
Bradwell	2004	SE Iceland	<i>R. section Rhizocarpon</i>	LL, SF	long axis	>250	30-100	m, rf, lf, fd	no
Matthews	2005	Jotunheimen, Norway	<i>Rhizocarpon</i> subgenus	LL, 5LL	long axis	5	200	moraines	no
Bakke <i>et al.</i>	2005	Lyngen, Norway	<i>R. geographicum</i>	5LL	long axis	5	30	ex. curve	no
Bradwell	2006	Lambatungnajökull, Iceland	<i>R. section Rhizocarpon</i>	LL, SF	long axis	>250	30-100	m, rf, lf, fd	yes
Principato	2008	Vestfirðir, Iceland	<i>R. geographicum</i>	5LL	mean diameter	5	entire	ex. curve	no

## Notes

1 – year of publication, not necessarily year of lichenometric survey.

2 – species, or taxonomic classification, as stated in publication.

3 – principal dating technique(s) used: LL (largest lichen); 3LL (3 largest lichens); 5LL (5 largest lichens); FALL (fixed-area largest lichen); SF (size-frequency distribution); see text for more details on different techniques.

4 – total number of lichens measured per surface in order to derive numerical age (1 = only largest-lichen used)

5 – average search area of lichenometric survey per surface, where stated. 'Entire' indicates the whole surface was searched. For FALL surveys, search areas are not recorded; a nominal value of  $<2 \text{ m}^2$  has been ascribed.

6 – surfaces used in calibration of dating curve, where applicable: moraines (m), gravestones (g), bridge (br), shoreline (sh), flood deposit (fd), rockfall (rf), lava flow (lf); ex.curve = existing (published) curve or modification of existing curve used to derive ages.

7 – relative ages only; uncertainty not applicable.