

# **Deep ocean exchange with west-European shelf seas**

**J. M. Huthnance, J. T. Holt and S. L. Wakelin**

Proudman Oceanographic Laboratory, 6 Brownlow Street, Liverpool L3 5DA, UK

Correspondence to: J. M. Huthnance ([jmh@pol.ac.uk](mailto:jmh@pol.ac.uk))

# Deep ocean exchange with west-European shelf seas

**J. M. Huthnance, J. T. Holt and S. L. Wakelin**

Proudman Oceanographic Laboratory, 6 Brownlow Street, Liverpool L3 5DA, UK

## **Abstract**

We review mechanisms and studies of exchange between the north-east Atlantic and the adjacent shelf sea. Mechanisms include: well-developed summer upwelling and associated filaments off Portugal and north-west Spain giving exchange  $O(3 \text{ m}^2/\text{s})$  per unit length of shelf; prevailing westerly winds further north driving exchange  $O(1 \text{ m}^2/\text{s})$ ; poleward flow along most of the upper slope with associated secondary circulation  $O(1 \text{ m}^2/\text{s})$ ; meanders and eddies in this poleward flow; eddies shed from slope waters into the Bay of Biscay; local exchanges at shelf spurs and depressions or canyons (e.g. dense-water cascading of order  $1 \text{ m}^2/\text{s}$ ). Tidal transports are larger; their reversal every six hours makes exchange largely ineffective except where internal tides are large and non-linear, as in the Celtic Sea where solitons carry water with exchange  $O(1 \text{ m}^2/\text{s})$ . These various physical exchanges amount to an estimated  $2\text{-}3 \text{ m}^2/\text{s}$  per unit length of shelf, between ocean and shelf; a numerical model estimate is comparable:  $2.5 \times 10^6 \text{ m}^3/\text{s}$  onto and off the shelf from Brittany to Norway. Mixing controls the seasonal thermocline, affecting primary production and hence fluxes and fate of organic matter. Specifically,  $\text{CO}_2$  take-up by primary production, settling below the thermocline before respiration, and then off-shelf transport, make an effective shelf-sea “pump” (for  $\text{CO}_2$  from the atmosphere to the deep ocean). However, knowledge of biogeochemical fluxes is generally sparse; there is scope for more measurements, model validation and estimates from models.

## **1 Introduction**

Ocean-shelf exchange has important consequences for shelf-sea currents, flushing and the supply of nutrients. Conversely, shelf processes impact on the open ocean: they appear to exert some control of circulation around major ocean basins (Hughes and Meredith, 2006); mixing over slopes is suggested to contribute to the main oceanic density structure (Munk

and Wunsch, 1998). Topography constrains large-scale (geostrophic) flow from crossing the slope, creating interest in special processes enabling exchange (Huthnance, 1995).

Transports of nutrients and carbon between shelf seas and the open ocean are critical parts of nutrient and carbon cycles, e.g. Wollast (1993), Liu et al. (2000), Muller-Karger et al., 2005). Although budgeting this part of their cycles is generally inadequate, there is evidence that the north-west European shelf can make a significant contribution to the oceanic up-take of carbon (Frankignoulle and Borges, 2001; Thomas et al., 2004). This “draw-down” is most efficient over an annual cycle if carbon taken up by plankton growth in spring and summer sinks below the seasonal thermocline before respiration and is exported to the deep ocean below the permanent thermocline.

Observational studies of ocean-shelf exchange have been carried out in several locations including the Middle Atlantic Bight (e.g. Biscaye et al., 1994), the US western shelf (e.g. Jahnke et al., 2008; Kudela et al., 2008; Hickey and Banas, 2008) and the Gulf of Lions (north-west Mediterranean; Monaco et al., 1990). Specific west-European margin studies (including multi-institutional experiments: MORENA, OMEX, SES) form the substance of review in Sect. 3.

Modelling of physical variables on kilometre scales is now possible in 3-D with fair realism for wind- and buoyancy-driven flow over realistic shelf and slope topography, albeit effects of smaller-scale internal waves and turbulence are parameterised (Samelson et al., 2008). Coupling with ecosystem models has progressed, especially for primary production; with higher trophic levels, there are increasing issues of parameter estimation and trade-offs (between number of variables, coverage, resolution and run duration). Modelling is also reviewed in Sect. 3.

These cited studies emphasise that net fluxes are hard to measure in unsteady conditions with along-shelf variations. An empirical approach to balancing a budget needs (in principle) sufficient spatial and temporal resolution covering all faces of a control volume. Typically this is not practical even for water; it is even harder to estimate net fluxes resulting from correlations of constituent concentrations with flow. Hence there is a need for theory and for models to give practical effect to theory. Models hold the prospect of 3-D representation of complex domains and processes, and inherent budgeting (if properly formulated). However, (expensive) measurements are necessary at least to test hypotheses and models. Such testing was the purpose of many of the studies cited. Then there may be more confidence in using

(tested) models to estimate exchanges where measurements are not dense enough. Confidence may be strengthened if models are shown to represent well the key processes contributing to ocean-shelf exchange in a region of interest.

The aim here, in the context of the “Deep Ocean Exchange with the Shelf” overview, is to review studies of exchange between the north-east Atlantic ocean and west-European shelf seas. In accord with the above discussion and issues, the emphasis is on a description of exchange processes occurring in this region, on their contributions to exchange in relation to overall estimates of exchange and on process representation in models which may give confidence to model estimates elsewhere.

We outline the west-European context (Sect. 2), review observational and modelling studies in this region (Sect. 3), summarise knowledge of exchange processes here (Sect. 4), give some overall estimates of exchange (Sect. 5), summarise by sector of shelf (Sect. 6) and discuss progress and needs for future work (Sect. 7).

## **2 Context; west-European margin**

In global terms, these shelf seas are quite broad and irregular (Fig. 1), especially compared with other eastern ocean margins. The shelf width varies; 10-60 km off Portugal and northern Spain, 50 km increasing to 100-150 km in eastern Biscay, approximately 500 km in the Celtic Sea, 50-150 km around Ireland and 100-200 km around Scotland with a very irregular coastal boundary and many islands. The Norwegian shelf broadens from about 20 km at 61°N to 200 km at 65°-68°N before narrowing again to about 50 km around 70°N.

Typical shelf depths are 100-150 m from Biscay to Scotland, increasing to more than 400 m in the Norwegian Trench. The North Sea shoals southwards, to 20-30 m depth between south-east England and the near continent. Depths on the Norwegian shelf are variable, extensively 200-300 m or more. The continental slope is steep from Portugal to north-west Scotland, and is indented with a few large canyons off Portugal and in southern Biscay; there are many smaller canyons north of Cape Ferret Canyon (44.7°N) as far as Goban Spur (49°N, 11°W). The Porcupine Sea Bight is a 2000 m deep intrusion; Porcupine Bank (depth shoaling to < 200 m) is a hydrographically semi-detached extension from the western Irish shelf. Around Scotland the upper slope is relatively smooth, and smaller (~ 0.02 or less) north-east of the Wyville-Thomson Ridge. Off Norway the shelf edge is less regular.

Rivers' inputs of freshwater, sediments and sediment-bound nutrients and carbon to west-European shelf seas are small in global terms; salinity in most areas is within 1 or 2 of open-ocean values (the Norwegian Trench is an important exception). Nevertheless, excepting west and north Scottish shelves with small-area river catchments, run-off adds more freshwater than precipitation-evaporation. In the adjacent upper ocean, warm North Atlantic water flows poleward past Ireland, Scotland and Norway; water formed by deep convection in northern Biscay flows slowly southwards to Iberia (van Aken, 2001); in 500-1500 m against the slope, northward flow (as far as the entrance to Rockall Trough) includes a Mediterranean Water fraction which increases the salinity.

Riverine input of phosphorus is much less than exchanges with the open ocean or transfers between shelf sectors. As losses are small, overall net export to the ocean is inferred. Nitrogen input from rivers and atmosphere is less than de-nitrification, and much less than exchanges with the open ocean or transfers between sectors.

Primary production in the north-east Atlantic is described by (e.g.) Pingree et al. (1976) and Longhurst (1998). A spring bloom begins when the light-determined critical depth for net algal growth descends to the mixed-layer depth, which shoals as heating increases and wind stress decreases. [The timing varies: inter-annually; also locally, for example if fresh-water inputs give a shallower surface layer, advancing the bloom, or sediment suspension limits light and delays growth]. The bloom becomes nutrient-limited when the initial near-surface charge is exhausted. Then summer growth depends on biologically regenerated nitrogen (as  $\text{NH}_3$ ),  $\text{NO}_3$  entrained through the thermocline by turbulence from winds, waves and internal waves, and Ekman suction from wind stress curl. Additionally on the shelf, tidal mixing can supply nutrients to the euphotic zone; fronts between summer-stratified and mixed waters particularly favour phytoplankton growth (Pingree et al., 1978). Mixed waters provide nutrients directly from benthic regeneration. An autumn bloom, usually weaker than the spring bloom, may be fuelled by nutrients entrained in the deepening mixed layer. Typical west-European shelf-sea production is  $O(100-250 \text{ g C m}^{-2} \text{ a}^{-1})$ , somewhat greater than in the adjacent north-east Atlantic. Ultimately, except in riverine-influenced coastal regions, production is fuelled mostly by nutrients from the open ocean.

### 3 Review of previous work

Off-shelf transport processes have yet to be explicitly investigated in the context of multi-annual and shelf wide-scales. Observational campaigns are necessarily limited to a few seasons and a particular area; in this region study areas have included the following: *Malin-Hebrides* shelf west of Scotland, regarding internal tides (Sherwin, 1988) and ocean-shelf interaction (LOIS/SES 1995-96, e.g. Souza et al., 2001); *Goban Spur* (OMEX I, 1993-1996; e.g. Wollast and Chou, 2001); *Celtic Sea* for internal tides (e.g. Pingree and New, 1989; New and Pingree, 1990), slope current (Pingree et al. 1999) and residual circulation (Pingree and Le Cann, 1989, 1990); the *Iberian Margin* where there have been several studies of upwelling as well as larger-scale studies of the current structure and biogeochemical exchanges: MORENA, e.g. Fiuza et al. (1998); OMEX II, e.g. Huthnance et al. (2002), Joint and Wassmann (2001), van Weering and McCave (2002). There have been *North Sea* studies of the seasonal cycle (Simpson, 1993) and subsequently more modelling (Lenhart et al., 1995; Radach and Lenhart, 1995; Iversen et al., 2002; Smith et al., 1996; Winther and Johannessen, 2006; Moll and Radach, 2003) and budgeting (Thomas et al., 2005) studies, albeit not directly addressing ocean-shelf exchange. Water, nutrient and carbon cycling along the sub-polar north-east Atlantic margin (Biscay to Norway) has been reviewed in the context of the JGOFS-LOICZ Continental Margin Task Team (Huthnance, 2009).

Holt et al. (2009) have used a 3-D physics-only model with ~ 12-km resolution to simulate 45 years exchange between the north-west European shelf and adjacent Atlantic (see Sect. 5). Proctor et al. (2003) modelled a year (1995) of nutrient and phytoplankton fluxes across the shelf edge west of Scotland in 2-D with resolution ~ 1.2 km in the horizontal and 60 layers in the vertical; the microbiological model represented nitrate, ammonium, oxygen, microplankton and detritus; there was fair agreement with 1995 measurements but indication of a need to improve bottom boundary-layer processes in the model. In general along-shelf variability influences ocean-shelf exchange (e.g. Kudela et al., 2008; Hickey and Banas, 2008) and calls for a 3-D model.

The 3-D POLCOMS Atlantic Margin Model as used in Holt et al. (2009) has been coupled with the European Regional Seas Ecosystem Model ERSEM (Allen et al., 2001). The 18 years 1988-2005 have been run for carbon cycling with ~12 km resolution, 42 s-levels, ERA40 and operational ECMWF surface forcing, about 300 river-flow inputs, time-varying (spatially-uniform) atmospheric pCO<sub>2</sub>; a mean annual cycle was used at ocean boundaries

and for optical attenuation and riverine concentrations of nutrient and dissolved organic carbon (DIC). Inputs to the model budget (largest first) are from rivers and the atmosphere (inorganic C) and advection of organic C. These simulations are presented in detail by Holt et al. (in preparation). A preliminary analysis shows that net burial  $0.02 \times 10^{12} \text{ mol C yr}^{-1}$  is the small difference between net settling of organic carbon and benthic respiration as inorganic carbon. Horizontal advection is the dominant loss; the net loss of (inorganic) carbon is  $0.9 \times 10^{12} \text{ mol C yr}^{-1}$ . The advective loss is an effective sink of carbon if it leaves the shelf to deep water below layers where it may re-equilibrate with the atmosphere.

#### 4 Exchange processes

Warm, saline North Atlantic Water (NAW) forms a *poleward current* along the continental slope. The current is approximately barotropic, centred at ~500m on the slope; it occurs off Portugal (e.g. Frouin et al., 1990) and again from Biscay past Britain to Norway (e.g. Pingree et al., 1999; Souza et al., 2001, documenting findings in the SES; McClimans, 1999; Skagseth et al., 2004). Drifter observations demonstrate its continuity (Pingree et al., 1999) and its dispersion (Burrows et al., 1999) onto the shelf. It is thought to be forced by the dynamic height of warmer sub-tropical waters (Huthnance, 1984; Hill et al., 1998). On the Celtic Sea shelf, mean flows are weak (Pingree and leCann, 1989) but supply ~0.1 Sv eastwards through the English Channel (Prandle et al., 1996;  $1 \text{ Sv} = 10^6 \text{ m}^3/\text{s}$ ) and ~0.1 Sv northwards through the Irish Sea (Knight and Howarth, 1999). Atlantic inflow to the northern North Sea, cyclonic North Sea circulation and fresher water from the Baltic all feed the northward-flowing Norwegian Coastal Current. Beneath the poleward slope current is a bottom *Ekman layer* (Fig. 2), modified by the slope and stratification (e.g. Trowbridge et al., 1998), where friction reduces the current to zero. This Ekman layer is expected (with some evidence from observations) to have off-shelf transport  $\tau/\rho f$  of order  $1 \text{ m}^2/\text{s}$  or  $1 \text{ Sv}/(1000 \text{ km})$  (Huthnance, 1995; Souza et al., 2001). Here  $\tau$  is the bottom stress arising from the current,  $\rho$  is the water density,  $f$  the Coriolis parameter. This Ekman transport is robust, being independent of the detailed form of stress, and forms the basis of an “Ekman drain” concept for off-shelf flux. Model results for the north-west European shelf (Holt et al., 2009) show a strong correlation ( $r \sim 0.7$ ) between the along-slope flux (across  $56.5^\circ\text{N}$ ) and the lower layer flux across the 200m contour in  $52.5^\circ\text{N}$  to  $57.5^\circ\text{N}$ , substantiating the “Ekman drain” concept.

The region is subject to strong *wind* forcing, at its most intense around 60°N. Here the winds are variable, usually associated with the passage of cyclones (depressions) driving storm surges (in the form of coastal Kelvin and continental shelf waves) rather than up- or downwelling. However, the relative orientation of the Irish-Norwegian shelves and prevailing westerlies tends to drive on-shelf surface flow and hence downwelling at the coast. Lack of an adjacent coast where the shelf is broad (North Sea, Celtic Sea, Irish Sea) allows more extensive net on-shelf flow, with scope for off-shelf return flow elsewhere (notably in the Norwegian Coastal Current). The surface flow can be estimated as an Ekman transport

$$\rho_a c_D (W^2 \cos\theta + w'^2) / (\rho_w f)$$

where  $\rho_a, \rho_w$  are the densities of air and water,  $c_D$  is a drag coefficient,  $W$  is the monthly mean wind speed at an angle  $\theta$  to the along-slope direction,  $w'$  is the wind's standard deviation (taken as isotropic) in the along-slope direction, and  $f$  is the Coriolis parameter.

Summer upwelling is forced by north-easterly trade winds over the north-south extent of the Iberian shelf and especially around Finisterre (e.g. Joint and Wassmann, 2001); less off the Algarve with different coast orientation. Well-developed upwelling can form filaments with associated off- and on-shelf fluxes, typically O(0.6 Sv) per filament (Barton et al., 2001) which exceeds the Ekman transport over the (semi-regular) spacing between filaments.

Table 1 gives annual averages of this cross-shelf edge Ekman flux, using wind speeds from Josey et al. (1998; 2002) and wind directions and standard deviations from Isemer and Hasse (1985). [In Table 1, W Iberia is given seasonally from Huthnance et al., (2002); the North Sea flux is not at the shelf edge but based on estimates of flow between Scotland and Norway (Huthnance, 1997; see also Sect. 5)].

Semi-diurnal *tides* are large in the north Atlantic generally. Tidal currents exceed 0.1 m/s across the wide shelves of most of the north-east Atlantic margin, creating significant turbulence and mixing. Locally in straits and around headlands, tidal currents often exceed 0.5 m/s. These currents are primarily barotropic. However, internal tides with comparable peak currents are generated over steep slopes in many locations: off Portugal (Jeans and Sherwin, 2001a,b); particularly strongly at the Celtic Sea shelf edge (e.g. Pingree and New, 1989); on the Malin Shelf (Sherwin, 1988) and over the Wyville-Thomson Ridge (Sherwin, 1991). Large-amplitude (non-linear) internal tides can transport water in their wave-forms on the summer thermocline, exceptionally as much as 1 m<sup>2</sup>/s at the Celtic Sea shelf edge. Locally, strong tidal currents may be rectified to along- and cross-slope flow, usually near

small-scale features (e.g. headlands) and the upper slope. Although generally small, such rectified flows may contribute significantly to long-term displacements (Holt and Proctor, 2008).

Shear dispersion,  $K$ , results from variations of tidal current with depth. Observed spreading of caesium-137 on the north-west European shelf can be modelled by a horizontal dispersion coefficient  $K = t_D U^2$ , where  $t_D \sim 10^3$  s and  $U$  (m/s) is the tidal current amplitude (Prandle, 1984). This approach suggests that shelf-edge exchange from tidal shear dispersion is relatively small unless  $U > 0.5$  m/s, as west of Brittany but rarely elsewhere.

*Buoyancy* inputs are the other main source of forcing. Although direct lateral inputs of freshwater from land and rivers are small in global terms, they drive anti-cyclonic flows around Ireland and Scotland. The largest flux of fresher water is the Norwegian Coastal Current (originally from rivers into the Baltic Sea). Surface fresh-water buoyancy input from precipitation minus evaporation is within  $\pm 200$  mm  $a^{-1}$  over much of the region. However, summer heating and winter cooling force a seasonal cycle of stratification in most areas.

*Stratification* is favoured by the buoyancy inputs but eroded by mixing due to tidal currents, winds and waves (surface and internal). Typically, in sequence from very shallow water out to the open shelf, there are: a nearshore region mixed by waves, a region of freshwater influence that may be stratified, shallow coastal water mixed by strong tidal currents and winds, and deeper shelf water which is thermally stratified in summer. The summer thermocline is at  $O(50$  m) or less. On the north-west European shelf, the “tidal mixing” front between the mixed and summer-stratified areas is inshore of the shelf edge (Simpson, 1998a,b); the outer shelf is stratified in summer.

*Inertial-internal* waves are particularly active at the shelf edge (a source of these waves). They are distinctive in causing mixing within the thermocline (via shear and large-wave breaking), and at the bottom for a favourable combination of slope, stratification and wave frequency. *Surface waves* are important for surface mixing and for air-sea exchange in this region of strong winds and rough seas.

The depth of *winter mixing* exceeds shelf depths in the whole region, being more than 500 m in Biscay and as deep as 750-900 m around  $50^\circ$ - $60^\circ$ N west of Ireland and Scotland. Water and its contents from below the seasonal thermocline are thus mixed to the surface during autumn and winter. Winter cooling in the Nordic seas occurs as deep convection, in small cells but probably extensively, forming a cold dense bottom layer, which spills southwards

over the sills between Greenland and Scotland. Dense water formed by winter cooling of shallow shelf seas may *cascade* down the slope under gravity, eventually leaving the sloping bottom at its density level. Typical values of such cascading fluxes are estimated in Shapiro et al. (2003) as  $0.5\text{-}1.6\text{ m}^2\text{s}^{-1}$ ; significant when and where they occur. There is evidence that cascading has occurred at the Celtic Sea shelf edge (Cooper and Vaux, 1949) and at the north end of Porcupine Bank (Hill et al., 1998).

The *irregular shelf*, with capes, canyons and varied shelf width, may cause locally-enhanced up-/down-welling and cross-slope flow (Trowbridge et al., 1998), e.g. at Goban Spur (Pingree et al., 1999). Large slope-current meanders and eddies are observed in the Faroe-Shetland Channel (Sherwin et al., 1999, 2006) and eddies are formed from the slope current off northern Spain (Pingree and LeCann, 1992)

## 5 Overall estimates of exchange

Empirical estimates of total exchange across some sectors of shelf edge have been made on the basis of (i) changing properties of water flowing along the slope, (ii) variance in cross-slope flow, (iii) cross-slope dispersion of drifters. Thus along the Iberian slope, northward-flowing Mediterranean water has lateral diffusivity  $\sim 500\text{ m}^2/\text{s}$  (Daniault et al., 1994; basis i). Off western Galicia, cross-slope current variance  $(19\text{mm/s})^2$  (basis ii) implies cross-slope exchange  $\sim 3.8\text{ m}^2/\text{s}$  in 200 m, or equivalent lateral diffusivity also  $\sim 500\text{ m}^2/\text{s}$  taking the offshore scale as the distance from the coast to the 200 m depth contour (26 km; Huthnance et al., 2002). Also off western Galicia, drifters deployed in August 1998 and winter 1998/99 dispersed with effective across-slope diffusivity  $870\text{ m}^2/\text{s}$  and  $190\text{ m}^2/\text{s}$  respectively, i.e. an average  $560\text{ m}^2/\text{s}$  (basis iii). West of Scotland, cross-slope current variance  $\sim (33\text{ mm/s})^2$  at the 200 m contour (Harikrishnan, 1998; basis ii) is equivalent to  $6.6\text{ m}^2/\text{s}$  exchange. North of Scotland and off Norway, cross-slope current variances are of order  $(0.1\text{ m/s})^2$  and there are estimates of lateral diffusivity from drogued-bouy dispersion (basis iii):  $\sim 700\text{ m}^2/\text{s}$  (Booth, 1988);  $\sim 360\text{ m}^2/\text{s}$  (Burrows and Thorpe, 1999). Analysis by process contributions is discussed in Sect. 6, initially for west Iberia, and presented in Table 1.

The period 1960-2004 was simulated for the west European shelf and nearby Atlantic by Holt et al. (2009) using surface forcing from ERA-40, COARE3 bulk formulae (Fairall et al., 2003), lateral boundary conditions from a  $\sim 1^\circ$  global model for 1958-2004 (NEMO; Smith and Haines, submitted) and river inputs (Young and Holt, 2007; [grdc.bafg.de/](http://grdc.bafg.de/)). The overall

pattern of across-shelf-edge transport is a net downwelling circulation of about 1.2 Sv. This is the net result of surface transport on to the shelf (variable, driven by prevailing winds) and an off-shelf transport in a thin near-bed Ekman layer under the slope current (little variability on monthly time scales). Total on-/off-shelf exchange (from Brittany to Norway) is larger, about 2.5 Sv, if all on-shelf flows are added and all off-shelf flows are added separately (Fig. 3). Across-shelf-edge transport in the American shelf and Celtic Sea is relatively weak and generally on-shelf, whereas there is strong off-shelf transport from the North Sea to the Norwegian Trench. These features are illustrated by tracer distributions in Fig. 4.

The fate of organic carbon settling to the bottom fraction of the water column was estimated by introducing tracer, each June, into the lowest model layer within the 200m depth contour, with uniform concentration / area. By the next March, typically 40% of tracer had left the shelf. Particularly for Irish and Scottish shelves, transport processes are very effective at removing material from the bottom layer on the shelf (Holt et al., 2009). This scenario forms the basis for extending the North Sea CO<sub>2</sub> “pump” concept (Sect. 6 and Thomas et al., 2004) to the wider west-European shelf.

## **6 Summary by sector**

This section summarises exchanges and some particular aspects of nutrients and carbon cycling for the west-European margin sectors in turn from south to north. Estimated process contribution to ocean-shelf exchange of water are summarised in Table 1.

### **6.1 West Iberia**

This sector has moderate shelf width, significant inputs of fresh water and very strong seasonality: in winter, (north-) westerly prevailing winds give mixing and downwelling overall; in spring, a shallow thermocline typically develops, overlying thicker remnants of the mixed layer from the previous winter (e.g. May 1993; Fiuza et al., 1998); in summer, long-periods of north-easterly winds give upwelling of cooler water rich in nutrients and filaments develop taking surface water off-shelf. Poleward along-slope flow comprises relatively warm and saline Eastern North Atlantic Central Water; it is manifested at the surface in winter, but is present as an undercurrent for much of the year, typically a few tens of kilometres offshore from the shelf break and reaching down to depths of some hundreds of metres. Deeper again, warm saline Mediterranean Water, often with two “cores” centred at levels above and below

1000m, flows generally northwards with progressive “dilution” by adjacent Atlantic Water (Fiuza et al., 1998).

Overall estimated exchange across the 200 m depth contour is  $3.1 \text{ m}^2/\text{s}$  on the basis of cross-slope current variance  $(15.5 \text{ mm/s})^2$  (Huthnance et al., 2002) but differs between summer and winter. Contributions (Table 1; Huthnance et al., 2002) come from up-/down-welling, secondary flow associated with the slope current ( $O(1 \text{ m}^2/\text{s})$  including the Ekman “drain”), non-linear internal tides on the summer thermocline  $O(1 \text{ m}^2/\text{s})$ , detaching eddies and cross-frontal flow  $O(0.6 \text{ m}^2/\text{s})$ . The overall estimate  $5.6 \text{ m}^2/\text{s}$  for summer exceeds that for winter ( $2.2 \text{ m}^2/\text{s}$ ), owing to non-linear internal tides and to filaments  $O(3 \text{ m}^2/\text{s})$  when upwelling is well-developed. The summer-and-winter average is in fair agreement with the overall  $3.1 \text{ m}^2/\text{s}$ , and the summer enhancement is in fair agreement with the summer-enhanced drifter dispersion ( $870 \text{ m}^2/\text{s}$  compared with  $190 \text{ m}^2/\text{s}$  in winter).

## 6.2 Southern Biscay

This north Spanish shelf is relatively narrow. Estimated ocean-shelf exchange  $\sim 1.6 \text{ m}^2/\text{s}$  is dominated by wind-forced flow. Winds drive summer upwelling around Cape Ortegal and enhance production on the Cantabrian shelf (OSPAR, 2000). Slope water eddies are shed into the ocean, especially around Cape Ortegal (Pingree and LeCann, 1992); Huthnance et al. (2002) estimated about five eddies per year in this sector, for a relatively small exchange  $0.16 \text{ m}^2 \text{ s}^{-1}$ .

Riverine freshwater input greatly exceeds net precipitation-evaporation over this small shelf area. Estimated ocean-shelf exchange is larger than in- and out-flows along the shelf. Production greatly exceeds what can be fuelled by riverine and atmospheric inputs; it depends on oceanic inputs.

## 6.3 Eastern Biscay

In this sector, an estimated excess of evaporation over precipitation only slightly offsets freshwater input from rivers, notably the Garonne, Dordogne and Loire. Salinity on the shelf is close to open-ocean values, indicating effective (mostly wind-forced) ocean-shelf exchange, larger than in- and out-flows along the shelf. Again, production greatly exceeds what can be supported by riverine and atmospheric inputs and is inferred to depend on oceanic inputs to the shelf.

## **6.4 Celtic Sea**

This is a very broad shelf sector between Brittany and south-west Ireland. The Ocean Margin Exchange (OMEX I) project studied biogeochemical fluxes in detail near Goban Spur during 1993-1997.

Low-frequency circulation here is generally weak (Pingree and le Cann, 1989) except along the upper slope and where channelled or accelerated around promontories. Tidal currents exceed 0.5 m/s at the shelf edge west of Brittany, where tidally-rectified flow is 0.1 m/s or more and large internal tides carry on-offshore exchange up to 1.3 m<sup>2</sup>/s (Huthnance et al., 2001). At the shelf edge, strong internal tides mix and diffuse the seasonal thermocline, and cooler water brought nearer to the surface is exposed by wind mixing as a cool surface band. There is evidence of dense winter-cooled water cascading. Pingree et al. (1999) show a satellite image suggesting slope-current "overshoot" into the ocean at Goban Spur. This localised exchange could be comparable with the slope current transport, O(1 Sv), when it occurs. These processes along with wind-forcing suggest a large ocean-shelf exchange O(3 m<sup>2</sup>/s), greatly exceeding in-/out-flows. Most nutrients come from the ocean but much is exported to the west of Ireland. Production at the shelf edge (Joint et al., 2001) is enhanced by mixing across the summer thermocline by internal waves. OMEX I estimated nutrient and organic carbon fluxes in a Goban Spur section (Wollast and Chou, 2001). Turbid down-slope Ekman flow on the slope (McCave et al., 2001) suggests a route for sediment transport off the shelf.

## **6.5 Western Ireland shelf**

Ocean-shelf exchange is predominantly wind-forced. Turbid down-slope Ekman flow (under the slope current) on the outer slope of Porcupine Bank (Dickson and McCave, 1986) again suggests a route for sediment transport off the shelf. River input of freshwater exceeds precipitation-evaporation, but large ocean-shelf exchange implies only a small salinity reduction, except in the coastal current. The nutrient budget is dominated by exchanges with the Celtic Sea and open ocean.

## **6.6 West Scotland (to Cape Wrath / Wyville-Thomson Ridge)**

Tidal currents include a notable internal component on the outer shelf. Irish Sea outflow forms the northward-flowing Scottish Coastal Current (SCC) with branches on both sides of the Outer Hebrides island chain. There is evidence of dense winter-cooled water cascading.

Harikrishnan (1998) estimated a downslope Ekman transport of  $0.46 \text{ m}^2/\text{s}$  below the slope current; again turbid down-slope flow near  $56.5^\circ\text{N}$  (McCandliss et al., 2000) suggests a route for sediment transport off the shelf. Wind forcing is the dominant cause of ocean-shelf exchange, which greatly exceeds in-/out-flows from/to adjacent shelf sectors.

As well as the Irish Sea outflow, river input and precipitation-evaporation are comparable in adding freshwater, but the effect on salinity is small except in the SCC. Riverine and atmospheric sources of nutrients are negligible compared with dominant oceanic supply. Nutrients from the Irish Sea also help to supply production and a large export to the north Scottish shelf. Balanced carbon fluxes have been estimated using a numerical ecological model of the cross-slope section (Proctor et al. 2003);  $\text{CO}_2$  uptake from the atmosphere is important therein. Results are validated by flux measurements in SES (Souza et al., 2001).

## **6.7 North Scotland**

This sector lies near the latitude of maximum wind forcing. The Wyville-Thomson Ridge is a source of internal tidal currents and waves. Prominent meso-scale activity in the Faroe-Shetland Channel is probably generated locally from slope-current instability or from the Iceland-Faroe front; sometimes eddies deflect much of the slope current North Atlantic Water into the central part of the Channel (2 Sv or more; Sherwin et al., 1999, 2006). Process-based estimates of ocean-shelf exchange accordingly show dominant wind-forced and eddy contributions to a large total  $3 \text{ m}^2/\text{s}$ . [Drifter-based estimates of dispersion as large as  $700 \text{ m}^2/\text{s}$  (Sect. 4) give estimated exchange  $7 \text{ m}^2/\text{s}$  across the 500 m contour (Huthnance, 1995)].

Dominant inflow from the ocean is required to supply the estimated flow to the North Sea. There is relatively little river inflow; precipitation-evaporation provides most of the freshwater input. The overall effect on salinity is small. Nutrient budgets are dominated by supply from the ocean (with some from the West Scottish shelf), inferred uptake by production and outflow to the North Sea.

## **6.8 North Sea**

Exchanges are dominated by “Atlantic Water” inflows to the northern North Sea and a similar flow along the continental slope into the Norwegian Trench (Huthnance, 1997). The northern inflow (by various routes around Scotland, notably the Fair Isle Current between Orkney and Shetland, and around Shetland) totals about 1.7 Sv; it greatly exceeds the Dover Strait inflow  $\sim 0.1 \text{ Sv}$  (Prandle et al., 1996) and combined river inputs (including those via

the Baltic). The “Atlantic Water” may have been significantly modified by its long transit across the shelf for the Dover Strait inflow; some of the inflow around Scotland may also have come from the Celtic Sea shelf edge (Pingree et al., 1999). The main transport out of the North Sea is northwards in the Norwegian Coastal current (NCC). River inputs in the southern and eastern North Sea add significantly to nutrient supply locally, although for the North Sea as a whole most nutrient input is via the “Atlantic Water”. In the southern North Sea, strong tidal currents in shallow water prevent summer stratification and induce turbidity sufficient to delay the spring bloom in places. However, the frontal regions and southern and eastern areas (off Belgium to Denmark) show large primary production (e.g. Howarth et al., 1993). The North Sea net carbon budget is dominated by carbon inputs from rivers, the Baltic Sea and the atmosphere; the North Sea acts as a sink for organic carbon with final export to the Atlantic Ocean accounting for more than 90% of the CO<sub>2</sub> taken up from the atmosphere (Thomas et al., 2005). The uptake and sink is aided by large pCO<sub>2</sub> draw-down; stratification separates production (upper layer) and respiration causing a carbon dioxide increase in the subsurface layer that is ultimately exported; this more than offsets emission of respired CO<sub>2</sub> in the vertically-mixed southern North Sea (Thomas et al., 2004).

## **6.9 Norway**

Tidal currents are strong with a significant diurnal component off northern Norway. The Norwegian Coastal Current (NCC) transports 1-2 Sv northwards from inflow across 61°N, with its variability also 1-2 Sv – current variance is 0.01-0.02 m<sup>2</sup> s<sup>-2</sup> (Poulain et al., 1996). NCC salinity (reduced by Baltic outflow) is less than 33 at 61°N. Despite much riverine freshwater (greatly exceeding precipitation-evaporation), the salinity increases to about 34 off northern Norway as Atlantic water is entrained. The NCC is unstable, forming large eddies, but there is no clear estimate of an eddy separation-rate to the ocean interior. The NCC also spreads under north/easterly winds, while south/westerlies confine it against the coast. Process-based estimates of ocean-shelf exchange totalling approximately 3 m<sup>2</sup>/s show dominant wind-forced and eddy contributions.

Nutrient input from the North Sea greatly exceeds riverine or atmospheric sources. By analogy with North Scotland, oceanic input is probably comparable but we lack an estimate. Estimated production would only use a small fraction of the nutrient supply, with the balance probably exported to the Arctic. Data and models suggest annual primary production of

about 120-160 g C m<sup>-2</sup> (Slagstad et al., 1999). Integrated along the shelf, average organic carbon production is small relative to import from the North Sea.

## **7 Discussion and future work**

The following characterise the west-European margin: poleward along-slope flow (not uniform, and may not be continuous; it may also lie under different surface flow, especially during upwelling); strong wind forcing with downwelling prevalent north of Biscay and summer upwelling off Iberia – when well developed, filaments increase ocean-shelf exchange; strong tidal currents and mixing on wide shelves; relatively small exchange in eddies; moderate freshwater and stratification except for the Norwegian Coastal Current; local rectified tides, solitons and cascading; overall exchange 2-3 m<sup>2</sup>/s.

The combination of estimated exchange-process contributions in Sect. 6 (for W Iberia; Table 1) is encouraging in relation to independent estimates of overall exchange and the summer enhancement. However, exchanges from different processes are not necessarily additive. For example, poleward flow along the slope may occur (as an undercurrent) with summer upwelling, but their effects on cross-slope flow will tend to cancel, or “at best” co-exist with upwelling above the undercurrent and hence correspondingly reduced vertical extent and transport. Or, poleward flow may be in part a response to wind forcing in a downwelling sense so that simple addition has an element of double-accounting.

The encouragement from combining exchange-process estimates nevertheless reinforces the use of models, when tested for their representation of processes. In most locations, measurements are insufficient to determine exchanges; some basis is needed to extrapolate from measured contexts. Numerical models are potentially a rational basis for extrapolation, giving practical effect to theory, especially for 3-D representation of complex domains and processes, with inherent budgeting (if properly formulated).

The model estimate 2.5 Sv (Sect. 5) for overall exchange from Brittany to Norway (about 2000 km of shelf edge) is about half of the amount suggested by the Table 1 values (1.7 to 4 m<sup>2</sup>/s per unit length of shelf) over this distance. This is consistent with averaging in the model estimate; entries in Table 1 but averaged out in the model estimate include temporal reversals in wind-driven transports, internal tide contributions on spatial scales unresolved in the model, and seasonal processes.

The overall circulation with downwelling  $O(1 \text{ Sv})$  north of Biscay has an important role in biogeochemical cycles and specifically the carbon pump. The along-slope current acts to replenish on-shelf nutrients aiding summer draw-down of  $\text{CO}_2$  in producing organic carbon. These tidally active shelf seas lack 'export production' or burial; hence (to conserve carbon) large air-sea  $\text{CO}_2$  flux implies divergent DIC transport; the overall downwelling allows transport of dissolved material from regions of high production to the deep ocean. In particular, near-bottom transport of DIC off the shelf, below the permanent thermocline on time scales less than a year, facilitates  $\text{CO}_2$  drawdown again in the next growing season. [The off-shelf flow of near-bottom water is equivalent to evacuating the bottom quarter of the whole shelf in less than one year; the whole shelf volume in  $\sim 3$  years]. This off-shelf flow is not uniform; the margin from Ireland to the Norwegian Trench and areas of the North Sea are most effective.

There is no simple relation between productivity and air-sea  $\text{CO}_2$  flux. Modelling this flux in shelf seas requires accurate representation of the circulation, mixing, chemistry and biology; these are all factors in the carbon "pump" concept. The near-coastal region is particularly important: it can act as either a  $\text{CO}_2$  sink or a source; however, it is also the most challenging in the need for model improvements to represent complex optics, fine horizontal resolution and uncertain land-sea fluxes.

The outcome is that the west-European shelf is a net sink of atmospheric  $\text{CO}_2$ . Within this, shelf edge regions tend to be strong sinks, open stratified regions are neutral or weaker sinks, coastal regions are either sources or sinks.

## **Acknowledgements**

This work is primarily funded under the UK Natural Environment Research Council Oceans 2025 programme. The JGOFS-LOICZ Continental Margin Task Team instigated much of this study. J.M. Huthnance thanks SCOR for funding his participation in the "Deep Ocean Exchange with the Shelf" workshop, Cape Town, October 2008, leading to this article.

## References

- Allen, J. I., Blackford, J., Holt, J. T., Proctor, R., Ashworth M., and Siddorn, J.: A highly spatially resolved ecosystem model for the North West European continental shelf. *Sarsia*, 86, 423-440, 2001.
- Barton, E. D., Inall, M. E., Sherwin, T. J., and Torres, R.: Vertical structure, turbulent mixing and fluxes during Lagrangian observations of an upwelling filament system off Northwest Iberia. *Prog. Oceanogr.*, 51, 249-267, 2001.
- Biscaye, P.E., Flagg, C. N., and Falkowski, P. G.: The Shelf Edge Exchange Processes experiment, SEEP-II: an introduction to hypotheses, results and conclusions. *Deep-Sea Res. II*, 41, 231-252, 1994.
- Booth, D.A.: Horizontal dispersion in the north east Atlantic. *Deep-Sea Res.*, 35, 1937-1951, 1988.
- Burrows, M., and Thorpe, S. A.: Drifter observations of the Hebrides slope current and nearby circulation patterns. *Ann. Geophys.*, 17, 280-302, 1999.
- Burrows, M., Thorpe, S. A., and Meldrum, D. T.: Dispersion over the Hebrides and Shetland shelves and slopes. *Continental Shelf Res.*, 19: 49-55, 1999.
- Cooper, L. N. H., and Vaux, D.: Cascading over the continental slope of water from the Celtic Sea. *J. Mar. Biol. Assoc. UK*, 28, 719-750, 1949.
- Daniault, N., Mazé J. P., and Arhan, M.: Circulation and mixing of Mediterranean water west of the Iberian peninsula. *Deep-Sea Res.*, 41, 1685-1714, 1994.
- Dickson, R. R., and McCave, I. N.: Nepheloid layers on the continental slope west of Porcupine Bank. *Deep-Sea Res.*, 33, 791-818, 1986.
- Fairall, C. W., Bradley, E. F., Hare, J. E., Grachev, A. A., and Edson, J. B.: Bulk parameterization of air-sea fluxes: updates and verification for the COARE algorithm. *J. Climate*, 16, 571-591, 2003.
- Fiuza, A. F. G., Hamann, M., Ambar, I., Del Rio, G. D., Gonzalez, N., Cabanas J. M., and Diaz Del Rio, G.: Water masses and their circulation off western Iberia during May 1993. *Deep-Sea Res.*, 1, 45, 1127-1160, 1998.
- Frankignoulle, M. and Borges, A. V.: European continental shelf as a significant sink for atmospheric carbon dioxide. *Global Biogeochem. Cycles*, 15, 569-576, 2001.

- Frouin, R., Fiuza, A. F. G., Ambar I., and Boyd, T. J.: Observations of a poleward surface current off the coasts of Portugal and Spain during winter. *J. Geophys. Res.*, 95, 679-691, 1990.
- Harikrishnan, M.: Flow and water column structure at the Hebridean shelf-edge. PhD thesis, University of Wales, Bangor, xi + 86pp + figures, 1998.
- Hickey, B. M. and Banas, N. S.: Why is the northern end of the California Current System so productive? *Oceanography*, 21(4), 90-107, 2008.
- Hill, A. E., Hickey, B. M., Shillington, F. A., Strub, P. T., Brink, K. H., Barton E. D., and Thomas, A. C.: Eastern Ocean Boundaries: Coastal Segment (E). *The Sea*, 11 (A. R. Robinson, K. H. Brink, eds.), 29-67, 1998.
- Hill, A. E., Souza, A. J., Jones, K., Simpson, J. H., Shapiro, G. I., McCandliss, R., Wilson, H., and Leftley, J.: The Malin cascade in winter 1996. *J. Mar. Res.*, 56, 87-106, 1998.
- Holt, J. T. and Proctor, R.: The seasonal circulation and volume transport on the northwest European continental shelf: a fine-resolution model study. *J. Geophys. Res.*, 113, C06021, doi:10.1029/2006JC004034, 2008.
- Holt, J. T., Wakelin, S. L., and Huthnance, J. M.: The downwelling circulation of the northwest European continental shelf: a driving mechanism for the continental shelf carbon pump. *Geophys. Res. Lett.*, accepted, 2009.
- Holt, J. T., Wakelin, S. L., Smyth, T., Blackford, J. C., Allen, J. I., Proctor, R., and Ashworth, M.: Modelling the carbon fluxes of the North West European continental shelf: validation and budgets, in preparation.
- Howarth, M. J., Dyer, K. R., Joint, I. R., Hydes, D. J., Purdie, D. A., Edmunds, H., Jones, J. E., Lowry, R. K., Moffatt, T. J., Pomroy A. J., and Proctor, R.: Seasonal cycles and their spatial variability. *Philos. Trans. Roy. Soc. London, A* 343, 383-403, 1993.
- Hughes, C. W., and Meredith, M.P.: Coherent sea-level fluctuations along the global continental slope. *Philos. Trans. Roy. Soc. London, A*364, 885-901, 2006.
- Huthnance, J. M.: Slope currents and JEBAR. *J. Phys. Oceanogr.*, 14, 795-801, 1984.
- Huthnance, J. M.: Circulation, exchange and water masses at the ocean margin: the role of physical processes at the shelf edge. *Prog. Oceanogr.*, 35, 353-431, 1995.

Huthnance, J. M.: North Sea interaction with the North Atlantic Ocean. *Deutsche Hydrogr. Zeit.*, 49, 153-162, 1997.

Huthnance, J. M.: The Northeast Atlantic Margins, Section 5.2 in: Carbon and nutrient fluxes in continental margins: a global synthesis, edited by: K. K. Liu and J. Sharples, 215-234, Springer, in press, 2009.

Huthnance, J. M., Coelho, H., Griffiths, C. R., Knight, P. J., Rees, A. P., Sinha, B., Vangriesheim, A., White, M., and Chatwin, P. G.: Physical structures, advection and mixing in the region of Goban Spur. *Deep-Sea Res. II*, 48, 2979-3021, 2001.

Huthnance, J. M., van Aken, H. M., White, M., Barton, E. D., Le Cann, B., Coelho, E. F., Fanjul, E. A., Miller, P., and Vitorino, J.: Ocean margin exchange – water flux estimates. *J. Mar. Systems*, 32, 107-137, 2002.

Isemer, H.-J., and Hasse, L.: The Bunker climate atlas of the North Atlantic Ocean. 1: Observations. Springer-Verlag, Berlin Heidelberg, vii + 218pp, 1985.

Iversen, S. A., Skogen, M. D., and Svendsen, E.: Availability of horse mackerel (*Trachurus trachurus*) in the north-eastern North Sea, predicted by the transport of Atlantic water. *Fisheries Oceanography*, 11, 245-250, 2002.

Jahnke, R. A., Roman, M. R., and Brink, K. H.: Coastal Ocean Processes program: advancing interdisciplinary research and technology development. *Oceanography*, 21, 18-21, 2008.

Jeans, D. R. G., and Sherwin, T. J.: The evolution and energetics of large amplitude nonlinear internal waves on the Portuguese shelf. *J. Mar. Res.*, 59, 327-353, 2001a.

Jeans, D. R. G., and Sherwin, T. J.: The variability of strongly non-linear solitary internal waves observed during an upwelling season on the Portuguese shelf. *Continental Shelf Res.*, 21, 1855-1878, 2001b.

Joint, I., and Wassmann, P.: Lagrangian studies of the Iberian upwelling system - an introduction. A study of the temporal evolution of surface production and fate of organic matter during upwelling on and off the NW Spanish continental margin. *Prog. Oceanogr.*, 51, 217-220, 2001.

Joint, I., Wollast, R., Chou, L., Batten, S., Elskens, M., Edwards, E., Hirst, A., Burkill, P., Groom, S., Gibb, S., Miller, A., Hydes, D., Dehairs, F., Antia, A., Barlow, R., Rees, A., Pomroy, A., Brockmann, U., Cummings, D., Lampitt, R., Loijens, M., Mantoura, F., Miller,

P., Raabe, T., Alvarez-Salgado, X., Stelfox, C., and Woolfenden, J.: Pelagic production at the Celtic Sea shelf break. *Deep-Sea Res. II*, 48, 3049-3081, 2001.

Josey, S. A., Kent, E. C., and Taylor, P. K.: The Southampton Oceanography Centre (SOC) Ocean-Atmosphere Heat, Momentum and Freshwater Flux Atlas. SOC Report 6, 30pp. and figures, 1998.

Josey, S. A., Kent, E. C., and Taylor, P. K.: Wind stress forcing of the ocean in the SOC Climatology: comparisons with the NCEP-NCAR, ECMWF, UWM/COADS and Hellerman and Rosenstein datasets. *J. Phys. Oceanogr.*, 32, 1993-2019, 2002.

Knight, P. J., and Howarth, M. J.: The flow through the north channel of the Irish Sea. *Continental Shelf Res.*, 19, 693-716, 1999.

Kudela, R. M., Banas, N. S., Barth, J. A., Frame, E. R., Jay, D. A., Largier, J. L., Lessard, E. J., Peterson, T. D., and Vander Woude, A. J.: New insights into the controls and mechanisms of plankton productivity in coastal upwelling waters of the northern California Current System. *Oceanography*, 21(4), 46-52, 2008.

Lenhart, H. J., Radach, G., Backhaus, J. O., and Pohlmann, T.: Simulations of the North Sea circulation, its variability, and its implementation as hydrodynamical forcing in ERSEM. *Netherlands J. Sea Res.*, 33, 271-299, 1995.

Liu, K.-K., Iseki, K., and Chao, S.-Y.: Continental margin carbon fluxes. Pp. 187-239 In: *The Changing Ocean Carbon Cycle* (Eds. R.B. Hanson, H.W. Ducklow and J.G. Field), Cambridge University Press, Cambridge, 2000.

Longhurst, A.: *The Ecological Geography of the Sea*, Academic Press, San Diego, xiii+398 pp, 1998.

McCandliss, R., Jones, S. E., and Jago, C. F.: Distribution and fluxes of suspended particulate matter at the Hebridean shelf edge. Abstract, Exchange Processes at the Ocean Margins. 32<sup>nd</sup> International Liège Colloquium on Ocean Hydrodynamics, Liège, 8-12 May, 2000.

McCave, I. N., Hall, I. R., Antia, A. N., Chou, L., Dehairs, F., Lampitt, R. S., Thomsen, L., van Weering, T. C. E., and Wollast, R.: Distribution, composition and flux of particulate material over the European margin at 47°-50°N: Results from OMEX I. *Deep-Sea Res. II*, 48, 3127-3139, 2001.

- McClimans, T. A., Johannessen, B. O., and Jenserud, T.: Monitoring a shelf edge current using bottom pressures or coastal sea-level data. *Continental Shelf Res.*, 19, 1265-1283, 1999.
- Moll, A., and Radach, G.: Review of three-dimensional ecological modelling related to the North Sea: Part 1 models and their results. *Prog. Oceanogr.*, 57, 175-217, 2003.
- Monaco, A., Biscaye, P. E., Soyer, J., Pocklington, R., and Heussner, S.: Particle fluxes and ecosystem response on a continental margin: the 1985-1988 Mediterranean ECOMARGE experiment. *Continental Shelf Res.*, 10, 809-839, 1990.
- Muller-Karger, F. E., Varela, R., Thunell, R., Luerssen, R., Hu, C., and Walsh, J. J.: The importance of continental margins in the global carbon cycle. *Geophys. Res. Lett.*, 32, L01602, doi:10.1029/2004GL021346, 4 pp, 2005.
- Munk, W., and Wunsch, C.: Abyssal recipes II: energetics of tidal and wind mixing. *Deep-Sea Res. I*, 45, 1977-2010, 1998.
- New, A. L., and Pingree, R. D.: Evidence for internal tidal mixing near the shelf break in the Bay of Biscay. *Deep-Sea Res.*, 37, 1783-1803, 1990.
- OSPAR: Quality Status Report 2000 for the North-East Atlantic. OSPAR Commission, <http://www.ospar.org/eng/html/qsr2000>, 2000.
- Pingree, R. D., and Le Cann, B.: Celtic and Armorican slope and shelf residual currents. *Prog. Oceanogr.*, 23, 303-338, 1989.
- Pingree, R. D., and Le Cann, B.: Structure, strength and seasonality of the slope currents in the Bay of Biscay region. *J. Mar. Biol. Assoc. UK*, 70, 857-885, 1990.
- Pingree, R. D., and Le Cann, B.: Three anticyclonic slope water ocean eddies (SWODDIES) in the southern Bay of Biscay in 1990. *Deep-Sea Res.*, 39, 1147-1175, 1992.
- Pingree, R. D., and New, A. L.: Downward propagation of internal tidal energy into the Bay of Biscay. *Deep-Sea Res.*, 36, 735-758, 1989.
- Pingree, R. D., Holligan, P. M., Mardell, G. T., and Head, R.N.: The influence of physical stability on spring, summer and autumn phytoplankton blooms in the Celtic Sea. *J. Mar. Biol. Assoc. UK*, 56, 845-873, 1976.

- Pingree, R. D., Holligan, P. M., and Mardell, G. T.: The effects of vertical stability on phytoplankton distributions in the summer on the northwest European shelf. *Deep-Sea Res.*, 25, 1011-1028, 1978.
- Pingree, R. D., Sinha, B., and Griffiths, C.R.: Seasonality of the European slope current (Goban Spur) and ocean margin exchange. *Continental Shelf Res.*, 19, 929-975, 1999.
- Poulain, P.-M., Warn-Varnas, A., and Niiler, P. P.: Near-surface circulation of the Nordic seas as measured by Lagrangian drifters. *J. Geophys. Res.*, 101, 18237-18258, 1996.
- Prandle, D.: A modelling study of the mixing of  $^{137}\text{Cs}$  in the seas of the European continental shelf. *Philos. Trans. Roy. Soc. London, A* 310, 407-436, 1984.
- Prandle, D., Ballard, G., Flatt, D., Harrison, A. J., Jones, S. E., Knight, P. J., Loch, S., McManus, J., Player, R., and Tappin, A.: Combining modelling and monitoring to determine fluxes of water, dissolved and particulate metals through the Dover Strait. *Continental Shelf Res.*, 16, 237-257, 1996.
- Proctor, R., Chen, F., and Tett, P. B.: Carbon and nitrogen fluxes across the Hebridean shelf break, estimated by a 2D coupled physical-microbiological model. *The Science of the Total Environment*, 314-316, 787-800, 2003.
- Radach, G., and Lenhart, H. J.: Nutrient dynamics in the North Sea: fluxes and budgets in the water column derived from ERSEM. *Netherlands J. Sea Res.*, 33, 301-335, 1995.
- Samelson, R. M., Allen, J. S., and MacCready, P.: Progress in coastal ocean modeling during CoOP. *Oceanography*, 21(4), 136-140, 2008.
- Shapiro, G. I., Huthnance, J. M., and Ivanov, V.V.: Dense water cascading off the continental shelf. *J. Geophys. Res.*, 108 (C12), 3390, doi:10.1029/2002JC001488, 2003.
- Sherwin, T. J.: Analysis of an internal tide observed on the Malin Shelf, north of Ireland. *J. Phys. Oceanogr.*, 18, 1035-1050, 1988.
- Sherwin, T. J.: Evidence of a deep internal tide in the Faeroe-Shetland Channel. Pp. 469-488 in *Tidal Hydrodynamics*, B.B. Parker, Ed., John Wiley & Sons, New York, 1991.
- Sherwin, T. J., Turrell, W. R., Jeans, D. R. G., and Dye, S.: Eddies and a mesoscale deflection of the slope current in the Faroe-Shetland Channel. *Deep-Sea Res. I*, 46, 415-438, 1999.

- Sherwin, T. J., Williams, M. O., Turrell, W. R., Hughes, S. L., and Miller, P. I.: A description and analysis of mesoscale variability in the Faroe-Shetland Channel. *J. Geophys. Res.*, 111, C03003, doi:10.1029/2005JC002867, 2006.
- Simpson, J. H.: The North Sea project: an overview and the way forward. *Philos. Trans. Roy. Soc. London*, A343, 585-596, 1993.
- Simpson, J. H.: Tidal processes in shelf seas. *The Sea*, 10 (K. H. Brink and A. R. Robinson, eds.), 113-150, 1998a.
- Simpson, J. H.: The Celtic Seas. *The Sea*, 11 (A. R. Robinson and K. H. Brink, eds.), 659-698, 1998b.
- Skagseth, Ø., Orvik, K. A., and Furevik, T.: Coherent variability of the Norwegian Atlantic Slope Current derived from TOPEX/ERS altimeter data. *Geophys. Res. Lett.*, 31, L14304, doi:10.1029/2004GL020057, 2004.
- Slagstad, D., Tande, K. S., and Wassmann, P.: Modelled carbon fluxes as validated by field data on the north Norwegian shelf during the productive period in 1994. *Sarsia*, 84, 303-317, 1999.
- Smith, J. A., Damm, P. E., Skogen, M. D., Flather, R. A., and Patsch, J.: An investigation into the variability of circulation and transport on the North-West European shelf using three hydrodynamic models. *Deutsche Hydrogr. Zeit.*, 48, 325-347, 1996.
- Souza, A. J., Simpson, J. H., Harikrishnan, M., and Malarkey, J.: Flow structure and seasonality in the Hebridean slope current. *Oceanol. Acta*, 24, S63-S76, 2001.
- Thomas, H., Bozec, Y., Elkalay, K., and de Baar, H. J. W.: Enhanced open ocean storage of CO<sub>2</sub> from shelf sea pumping. *Science*, 304, 1005-1008, 2004.
- Thomas, H., Bozec, Y., de Baar, H. J. W., Elkalay, K., Frankignoulle, M., Schiettecatte, L. S., Kattner, G., and Borges, A. V.: The carbon budget of the North Sea. *Biogeosciences*, 2, 87-96, 2005.
- Trowbridge, J. H., Chapman, D. C., and Candela, J.: Topographic effects, straits and the bottom boundary layer. *The Sea*, 10 (K. H. Brink and A. R. Robinson, eds.), 63-88, 1998.
- Van Aken, H. M.: The hydrography of the mid-latitude north-east Atlantic Ocean: III. The thermocline water mass. *Deep-Sea Res.*, I 48, 237-267, 2001.

Van Weering, T. C. E., and McCave, I. N.: Benthic processes and dynamics at the NW Iberian margin: an introduction. *Prog. Oceanogr.*, 52, 123-128, 2002.

Winther, N. G., and Johannessen, J. A.: North Sea circulation: Atlantic inflow and its destination. *J. Geophys. Res.-Oceans*, 111, C12018, doi:10.1029/2005JC003310, 2006.

Wollast, R.: Interactions of carbon and nitrogen cycles in the coastal zone. Pp. 195-201 in *Interactions of C, N, P and S biogeochemical cycles and global change* (ed. R. Wollast, F. T. Mackenzie and L. Chou), NATO ASI Series, I 4. Berlin: Springer-Verlag, 1993.

Wollast, R., and Chou, L.: The carbon cycle at the ocean margin in the northern Gulf of Biscay. *Deep-Sea Res. II*, 48, 3265-3292, 2001.

Young, E. F., and Holt, J. T.: Prediction and analysis of long-term variability of temperature and salinity in the Irish Sea. *J. Geophys. Res.*, 112, C01008, doi:10.1029/2005JC003386, 2007.

## Figure legends

Figure 1. Outline map of west-European shelf region, with 200m depth contour, 1000m contours thereafter and divisions (red) between shelf sectors (Sect. 6).

Figure 2. Schematic of cross-slope circulation associated with poleward along-slope flow.

Figure 3. Fluxes ( $S_v$ ) above 150m (blue) and below 150m depth (red). All fluxes are across the 200m contour shown; positive is onto the shelf except next to Norway (positive to north).

Figure 4. Evolution of modelled surface concentration of tracer during 2002-2003. Initial concentration (beginning of 2002) was  $100 \text{ m}^{-3}$  everywhere where total depth  $> 2000 \text{ m}$ , zero elsewhere. [Within and somewhat beyond the 150m isobath, concentrations homogenize through the water column each winter].

Table 1 Estimated process contributions to ocean-shelf water exchange ( $\text{m}^2/\text{s}$ ) by sector.  
Exchange time = shelf volume / total exchange rate.

Sector	Wind	Internal tide	Slope current	Eddies	Filament	~ Total	Exchange time (yr)
Norway	1.22		0.5	1		3	0.21
North Sea	2		2			4	1
N Scotland	1.45		0.5	1		3	0.08
W Scotland	1.17	0.25	0.5			2	0.20
W Ireland	1.19		0.5			1.7	0.23
Celtic Sea	0.91	1	0.5		1	3	0.28
E Biscay	0.97		0.2			1.2	0.34
S Biscay	0.99		0.5	0.16		1.6	0.05
W Iberia summer		1	1	0.6	3	5.6	0.02
W Iberia winter	0.6		1	0.6		2.2	0.05