

# Structural setting and U–Pb zircon geochronology of the Glen Scaddle Metagabbro: evidence for polyphase Scandian ductile deformation in the Caledonides of northern Scotland

R. A. STRACHAN\* & J. A. EVANS†

\*School of Earth & Environmental Sciences, University of Portsmouth, Burnaby Building,  
Burnaby Road, Portsmouth PO1 3QL, UK

†NERC Isotope Geosciences Laboratory, Kingsley Dunham Centre, Keyworth, Nottingham NG12 5GG, UK

(Received 18 January 2007; accepted 20 August 2007; First published online 6 March 2008)

**Abstract** – Within the Scottish Caledonides, the Glen Scaddle Metagabbro was intruded into the Moine Supergroup of the Northern Highland Terrane after Grampian  $D_2$  folding and prior to regional  $D_3$  and  $D_4$  upright folding and amphibolite-facies metamorphism. A U–Pb zircon age of  $426 \pm 3$  Ma obtained from the metagabbro is interpreted to date emplacement.  $D_3$ – $D_4$  folding is constrained to have occurred during the Scandian orogenic event. In contrast, polyphase folding and regional metamorphism of the Dalradian Supergroup southeast of the Great Glen Fault is entirely Grampian. These differences are consistent with published tectonic models that invoke a minimum of 700 km of post-Scandian sinistral displacements across the Great Glen Fault to juxtapose the Grampian and Northern Highland terranes.

Keywords: U–Pb zircon, Caledonides, Scandian, Scotland.

## 1. Introduction

The geological complexity of many Mesozoic–Cenozoic orogens often results from the progressive amalgamation of volcanic arcs and continental fragments during oceanic closure, as well as regionally significant strike-slip displacements that arise either from the oblique nature of a collision or lateral escape during orthogonal convergence (Dewey *et al.* 1986). Even in well-exposed recent orogens such as the Himalayas, the magnitude of such strike-slip displacements can be difficult to evaluate (e.g. Searle, Weinberg & Dunlap, 1998; Searle, 2006), and such difficulties are compounded in older, Palaeozoic and Precambrian orogens. The Lower Palaeozoic Caledonide orogen in Scotland and Ireland is an excellent ancient example of such complexity. It evolved as a result of the closure of the Iapetus Ocean and the convergence of three continental blocks, Laurentia, Baltica and Avalonia (Soper, 1988; Pickering, Bassett & Siveter, 1988; Soper *et al.* 1992; Dewey & Strachan, 2003). Early arc–continent collisions along the Laurentia and Baltica margins during the Ordovician were followed by final oblique continental collision during Silurian times. Orogen-parallel, sinistral displacements occurred during the final stages of collision along various structures, including the Great Glen Fault in Scotland (Fig. 1).

Estimates of displacements along the Great Glen Fault are controversial because of overprinting of

the palaeomagnetic record and the lack of any unambiguous correlation across the fault of pre-Devonian markers. An alternative way of evaluating likely displacements across such structures is by comparison of the varying intensity of individual Caledonian events across the orogen as a means of reconstructing original collisional templates. The results of isotopic dating have, for example, already provided indications of a significant contrast in the intensity of Scandian (Silurian) deformational and metamorphic events across the Great Glen Fault (Kinny *et al.* 2003). In this paper we investigate this further and provide additional evidence that the Great Glen Fault separates crustal blocks with contrasting Caledonian tectonothermal histories.

The Scottish Highland Caledonides comprise a series of fault-bounded terranes derived from the margin of eastern Laurentia. The Great Glen Fault separates the Northern Highland and Grampian terranes (Fig. 1). The Northern Highland terrane is mainly underlain by the early Neoproterozoic Moine Supergroup (Holdsworth, Strachan & Harris, 1994). In contrast, the Grampian terrane is dominated by the late Neoproterozoic to early Palaeozoic Dalradian Supergroup (Harris *et al.* 1994). A possible equivalent of the Moine Supergroup, the Dava succession, occurs in the Grampian Terrane (Highton, Hyslop & Noble, 1999; Smith, Robertson & Rollin, 1999), but otherwise there are no obvious lithostratigraphic correlations across the Great Glen Fault.

Both the Moine and Dalradian supergroups were deformed and metamorphosed during an Ordovician

\*Author for correspondence: rob.strachan@port.ac.uk




Figure 1. Sketch map of the southern outcrop of the Moine Supergroup (see inset for location) showing the main geological units, igneous bodies and structures mentioned in the text. Diagonal hatching in inset corresponds to the outcrop of the Moine Supergroup in northern Scotland. Numbers around edge of main map correspond to British National Grid. Abbreviations: A – Ardnish; Agg – Ardgour granite gneiss; CG – Carn Gorm; CS – Central Sutherland; FAgg – Fort Augustus granite gneiss; G – Glenfinnan; GD – Glen Doe; GDS – Glen Dessary syenite; GGF – Great Glen Fault; GM – Glen Moriston; GSM – Glen Scaddle metagabbro; GT – Grampian terrane; K – Knoydart; LE – Loch Eilt; LQ – Loch Quoich; LS – Loch Sunart; MTZ – Moine Thrust Zone; NHT – Northern Highland Terrane; NR – northern Ross-shire; S – Strontian; SB – Sgurr Breac; SBT – Sgurr Beag Thrust; SG – Strontian Granite.

(c. 480–465 Ma) Grampian orogenic event that resulted from the collision of the Laurentian margin with an intra-Iapetan volcanic arc (Dewey & Ryan, 1990; Dewey & Mange, 1999; Oliver *et al.* 2000; Strachan *et al.* 2002; Dewey, 2006). In addition, U–Pb zircon dating of syn-tectonic granites in east Sutherland (Fig. 1) has provided evidence for ductile reworking and associated amphibolite facies metamorphism of the Moine Supergroup during the Silurian (Kinny *et al.* 2003; Kocks, Strachan & Evans, 2006). This is attributed to the collision of the Northern Highland segment of Laurentia with Baltica and correlated with the Scandian event recognized in Norway (Dallmeyer *et al.* 2001). The lack of any evidence for a comparable Silurian tectonothermal event in the Grampian terrane has led to the suggestion that at that time it was situated further to the southwest along the Laurentian margin, remote from the Scandian collision (Dewey & Strachan, 2003; Kinny *et al.* 2003). It has been concluded that at least 700 km of subsequent sinistral displacement along the Great Glen Fault would have been necessary to bring the Grampian terrane to its

present location relative to the Northern Highland terrane (Dewey & Strachan, 2003).

In this paper we review briefly the Caledonian evolution of the Moine rocks of West Inverness-shire (Fig. 1) and present the results of a geochronological study of the Glen Scaddle Metagabbro (Bailey & Maufe, 1916; Drever, 1940). The structural setting of this intrusion was established by Stoker (1983; M. Stoker, unpub. Ph.D. thesis, Univ. Liverpool, 1980) and confirmed during recent fieldwork by one of us (RAS).

## 2. Geological setting

The Moine rocks of West Inverness-shire comprise the Morar, Glenfinnan and Loch Eil groups (Fig. 1; Holdsworth, Strachan & Harris, 1994). The Morar Group lies within the Moine Nappe, and the Glenfinnan and Loch Eil groups within the Sgurr Beag Nappe (Powell *et al.* 1981; Barr, Holdsworth & Roberts, 1986). The upper part of the Sgurr Beag Nappe is dominated by mainly unmigmatized psammites and quartzites of the Loch Eil Group (Stoker, 1983; Strachan, 1985). The base of the Loch Eil Group passes transitionally into the underlying migmatitic pelites and thinly interbanded psammites and semi-pelites of the Glenfinnan Group. A lower limit for deposition of the Glenfinnan and Loch Eil groups is provided by the c. 1000–950 Ma ages of the youngest detrital zircons incorporated within the metasediments (Friend *et al.* 2003; Cawood *et al.* 2004). The West Highland Granitic Gneiss suite, prominent members of which crop out in Ardgour, at Fort Augustus and at Glen Doe (Fig. 1), was emplaced at c. 870 Ma (Friend *et al.* 1997; Rogers *et al.* 2001). The current consensus is that this probably represents an early rift-related igneous suite, broadly coeval with intrusion of amphibolites and meta-gabbros that record all the tectonothermal events present in their Moine host rocks (Millar, 1999; Dalziel & Soper, 2001).

The Moine rocks of the Loch Eil Group in West Inverness-shire record four main phases of deformation (Strachan, 1985; see also Dalziel, 1966 and Stoker, 1983). Early minor isoclinal folding ( $D_1$ ) was associated with formation of an  $S_1$  schistosity that developed under amphibolite facies conditions. Elsewhere in West Inverness-shire (E. K. Hyslop, unpub. Ph.D. thesis, Univ. Hull, 1992; Rogers *et al.* 1998; Vance, Strachan & Jones, 1998; Tanner & Evans, 2003), a mid-Neoproterozoic (820–730 Ma) age has been demonstrated for early folding, amphibolite-facies metamorphism and segregation of dated syn-metamorphic pegmatites at Ardnish, Sgurr Breac, Knoydart, Loch Eilt and Carn Gorm (Fig. 1). A similar age is assumed for  $D_1$  in the Loch Eil Group. A subsequent phase of widespread minor tight to isoclinal folding ( $D_2$ ) resulted in formation of a composite  $S_0/S_1/S_2$  foliation (Strachan, 1985). The foliation carries a stretching and mineral lineation




Figure 2. Simplified map of the main D<sub>3</sub> and D<sub>4</sub> folds within the SW Moine (using data from Powell *et al.* 1981; Roberts, Smith & Harris, 1984; Stoker, 1983; Strachan, 1985). Abbreviations: SBT – Sgurr Beag Thrust; GS – Glenshian Synform; LEA – Loch Eilt Antiform; GDS – Glen Dessary Synform; SMA – Spidean Mialach Antiform; SGF – Sgurr Ghiubhsachain Fold; MCA – Meall na Cuartaige Antiform; BATS – Beinn an Tuim Synform; LES – Loch Eil Synform; SA – Stronchreggan Antiform; CA – Corrlarach Antiform; LCS – Lochan na Cruaich Synform; MFA – Meall an Feidh Antiform; GSS – Glen Scaddle Synform; GGA – Glen Gour Antiform; GGS – Glen Gour Synform.

(L<sub>2</sub>) that trends north–south where unaffected by later deformation. These structures are correlated with the D<sub>2</sub> folds and associated fabrics recognized in

the northern outcrop of the Loch Eil Group east of Loch Quoich (Fig. 1; Holdsworth & Roberts, 1984). Titanites aligned within the composite S<sub>1</sub>/S<sub>2</sub> foliation within the Fort Augustus Granite Gneiss (Fig. 1) have yielded a U–Pb age of 470 ± 1 Ma, suggesting that D<sub>2</sub> occurred during the Ordovician Grampian orogenic event (Rogers *et al.* 2001).

The large-scale structural geology of the southern Loch Eil Group is dominated by major upright D<sub>3</sub> and D<sub>4</sub> folds (Fig. 2; Strachan, 1985). D<sub>3</sub> folds are all upward-facing and plunge towards the Great Glen, with the result that younger rocks come on to the east. Folds are open to close in style, and an S<sub>3</sub> mica schistosity (locally with fibrolite) is locally developed axial-planar to D<sub>3</sub> folds. The refolding of D<sub>3</sub> folds during D<sub>4</sub> generated a regional-scale interference pattern (Fig. 2). D<sub>4</sub> folds are gentle to open, and vary in trend from north–south in the area north of Loch Eil to NNE–SSW in Ardgour; an axial-planar crenulation fabric (S<sub>4</sub>) is widespread. Fibrolite, hornblende and biotite all recrystallize around S<sub>4</sub> crenulations, and it is therefore concluded that D<sub>4</sub> was characterized by amphibolite-facies conditions. D<sub>4</sub> folding increases in intensity westwards to form the Northern Highland Steep Belt (Figs 2, 3; Roberts & Harris, 1983).

The prevailing view has been that the upright folds and associated metamorphic assemblages formed at a late stage of the Ordovician Grampian orogenic event (e.g. Kelley & Powell, 1985; Powell & Phillips, 1985; Harris, 1995). A lower limit for the formation of the steep belt is provided by the intrusion of the pre-D<sub>4</sub> Glen Dessary Syenite (Fig. 1) at 456 ± 5 Ma (U–Pb




Figure 3. Generalized geological cross-section across the SW Moine drawn approximately along line XY in Figure 1, showing the westerly increase in the intensity of D<sub>4</sub> upright folding to form the Northern Highland Steep Belt (modified from Powell & Glendinning, 1988). Abbreviations: MT – Moine Thrust; KNT – Knoydart Thrust; SBT – Sgurr Beag Thrust; GS – Glenshian Synform; LEA – Loch Eilt Antiform; GDS – Glen Dessary Synform; SMA – Spidean Mialach Antiform; LQL – Loch Quoich Line; GGF – Great Glen Fault. Note (a) the infold of Loch Eil Group in the core of the Glen Dessary Synform; (b) the local emergence of Glenfinnan Group rocks adjacent to the Great Glen Fault (see Fig. 1).




Figure 4. Composite cross-section across the Glen Scaddle Metagabbro and its Moine country rocks to illustrate the nature of the major  $D_3$  folds and the structural setting of the intrusion in the core of the Glen Scaddle Synform (modified from Stoker, 1983). The intrusion carries a moderately to steeply dipping foliation that is axial-planar to the Glen Scaddle Synform. Note that the Glen Gour Metagabbro to the southwest occupies a lower structural level, in the core of the  $D_3$  Glen Gour Antiform. The Moine succession is right-way-up. See Figure 2 for line of section; abbreviations for folds as in Figure 2.

zircon: van Breemen *et al.* 1979; Roberts, Smith & Harris, 1984). An upper limit is apparently provided by a U–Pb monazite age of  $450 \pm 10$  Ma obtained from a pegmatite sampled within the steep belt west of Glenfinnan and thought to pre-date upright folding (van Breemen, Pidgeon & Johnson, 1974). According to Powell & Phillips (1985), formation of the steep belt was followed by a prolonged period of post-tectonic cooling from *c.* 453 Ma to *c.* 405 Ma. The southern part of the steep belt is cut by the Strontian Granite that was intruded at  $425 \pm 3$  Ma (Fig. 1; Rogers & Dunning, 1991). Most of the intrusion is unmetamorphosed and dominated by pre-full-crystallization magmatic fabrics (Hutton, 1988).

Reappraisal of this interpretation is appropriate because the Grampian event is now thought to have been short-lived (*c.* 480–465 Ma) and followed by very rapid exhumation (Oliver *et al.* 2000). Furthermore, Kinny *et al.* (2003) have demonstrated widespread Silurian ductile deformation and amphibolite-facies metamorphism in the Moines of central and west Sutherland. Upright folds of the steep belt in northern Ross-shire (Fig. 1) deform lineations within the Morar Group that have been correlated with the Silurian fabrics identified in central Sutherland (Kinny *et al.* 2003). Either regional correlations of structures are in error or the published upper limit on the age of the steep belt (van Breemen, Pidgeon & Johnson, 1974) is less robust than thought previously. In an effort to place further constraints on the age of upright folding, we now focus on the geology of eastern Ardgour, with particular reference to the structural setting and U–Pb geochronology of the Glen Scaddle Metagabbro (Fig. 1).

### 3. Geology of the Glen Scaddle Metagabbro

#### 3.a. Structural setting and contact relationships

The Glen Scaddle Metagabbro intrudes the Moine rocks of the Loch Eil Group (Stoker, 1983; Strachan, 1985). In eastern Ardgour these comprise a varied succession of psammites and quartzites which are mostly right-way-up where sedimentary structures are present. The structure of eastern Ardgour is dominated by upright to moderately inclined, tight to close  $D_3$  folds (Figs 2, 4). The Glen Scaddle Metagabbro occupies the core of a major  $D_3$  fold, the Glen Scaddle Synform, flanked to the northeast by the  $D_3$  Meall an Feidh Antiform and to the southwest by the  $D_3$  Glen Gour Antiform (Figs 2, 4). A smaller intrusion, the Glen Gour Metagabbro, occupies the core of the Glen Gour Antiform (Fig. 4) and is apparently at a lower structural level than the Glen Scaddle Metagabbro.

The margins of the Glen Scaddle Metagabbro are broadly parallel to the regional composite  $S_0/S_1/S_2$  foliation within host Moine psammites. Contacts with Moine rocks are generally concordant and sharp [e.g. NN 0050 6920]. Small xenoliths (10–50 cm) of Moine lithologies are locally common within the marginal facies of the intrusion [e.g. NM 9662 7004]. Within a narrow (0.75–1 km) aureole developed within the Moine rocks, regional metamorphic assemblages are overprinted by contact metamorphic minerals (Drever, 1940; Ashworth & Chinner, 1978). Quartz–feldspar–biotite rocks that are interpreted as nebulitic migmatites formed from the partial melting and mobilization of Moine lithologies are locally developed on the decametre scale along the northeast contact of the Glen Scaddle body [e.g. NM 9656 7012]. These

are characterized by disoriented xenoliths of Moine psammities that apparently resisted melting, as well as zones of complex disharmonic folding.

### 3.b. Meta-igneous lithologies

The most common rock type is a medium- to coarse-grained, foliated metagabbro. Subordinate meta-igneous rock types range from ultrabasic to acidic in composition. Foliated hornblende peridotite and serpentinite were recorded by Drever (1940) from the margins of the Glen Gour and Glen Scaddle intrusions. Also present within the metagabbro are concordant metre-scale sheets of fine-grained hornblende schist and at least two phases of fine- to medium-grained granitic veins that are mainly developed on the centimetre–decimetre scale but can range up to 2–3 m in thickness. The earliest (meta)granitic veins are foliated and locally associated with hornblende schist; complex textures indicative of mingling between the two parent magmas are sometimes preserved [e.g. NM 9965 6856]. The later granitic veins are generally unfoliated and discordant, locally forming ramifying networks that cross-cut host meta-igneous lithologies.

### 3.c. Petrology and microfabrics

Relic igneous mineralogies and textures are best preserved in rare areas of low tectonic strain up to several hundred metres wide [e.g. River Scaddle, NM 9814 6818] and it is appropriate to describe these first before outlining the effects of superimposed deformation and metamorphism. In low strain zones, the metagabbro is essentially undeformed and typically consists of hypersthene, clinopyroxene, plagioclase ( $An_{35-40}$ ) and amphibole with minor quartz, biotite and accessory ilmenite, titanite and apatite. Feldspars are rectangular and randomly oriented. Pyroxene is mostly replaced partially to completely by uralitic amphibole, and this is interpreted to be the result of late stage deuteric alteration within the igneous protolith (Stoker, 1983). Feldspar is commonly highly altered and sericitized. Sub-ophitic textures are common, as are symplectic intergrowths between pyroxene and plagioclase. The effects of apparently static metamorphism are represented by the recrystallization of amphibole into granoblastic aggregates of hornblende, and the growth of radiating mats of biotite. M. Stoker (unpub. Ph.D. thesis, Univ. Liverpool, 1980) recorded a local crude rhythmic layering in metagabbro [e.g. NM 9945 6850] shown by gradation of melanocratic to leucocratic gabbroic types on scales of up to several metres. Low strain areas of metagabbro locally contain metre-scale mafic enclaves [e.g. NM 9852 6827] and are intruded by undeformed granitic sheets.

Areas of low tectonic strain pass transitionally into the foliated metagabbro that forms the bulk of the in-




Figure 5. Photomicrograph of typical foliated metagabbro (horizontal field of view, 4 mm). F – feldspar, H – hornblende; double-headed arrow indicates orientation of foliation.




Figure 6. Photomicrograph of mylonitic granite sheet within the Glen Scaddle Metagabbro (field of view, 4 mm). Q – aggregates of dynamically recrystallized quartz, F – feldspar porphyroclasts, M – myrmekite; double-headed arrow indicates orientation of foliation.

trusion. The relic igneous features described above are progressively replaced by a penetrative amphibolite-facies deformation fabric defined by aligned layers of intergrown hornblende and biotite that wrap augen of recrystallized plagioclase (Fig. 5). The mafic enclaves and the undeformed granites present in the low strain zones are interpreted as the protoliths of, respectively, the hornblende schists and the early foliated granitic veins within the foliated metagabbro. Within the hornblende schists, the foliation is defined by a penetrative alignment of fine-grained hornblende and recrystallized feldspar. The early granitic veins are characterized by an intense mylonitic fabric: classic mortar texture is displayed by ribbons of dynamically recrystallized quartz that wrap feldspar augen (Fig. 6). Feldspar porphyroclasts are sometimes fringed by myrmekite which is probably strain-induced (Fig. 6).

### 3.d. Structure and age of emplacement

Along the margins of the Glen Scaddle Metagabbro, the foliation described above is broadly parallel to the contact with host Moine rocks. However, within the intrusion the foliation is more uniform in orientation, trending approximately NW–SE and dipping moderately to steeply to the northeast. A moderately plunging mineral and extension lineation is defined locally by alignment of hornblende–biotite aggregates and plagioclase augen [e.g. NM 9965 6856].

The following two observations establish that intrusion post-dated the early fabric-forming events  $D_1$  and  $D_2$ . Firstly, within the aureole around the Glen Scaddle Metagabbro, syn- $D_1$  peak regional metamorphic assemblages are overprinted by contact metamorphic minerals, including sillimanite, biotite, plagioclase, quartz, K-feldspar, cordierite and andalusite which have been reported from various samples of pelite (Ashworth & Chinner, 1978; Stoker, 1983). Secondly, the intrusion locally incorporates xenoliths that contain isoclinal folds that are most probably  $D_2$  in age [e.g. NM 9680 6835].

A pre- $D_3$  age of emplacement is consistent with the broad parallelism of the metamorphic foliation in the intrusion to the axial plane of the  $D_3$  Glen Scaddle Synform, and the observation that contact metamorphic sillimanite is wrapped by the  $S_3$  fabric (Stoker, 1983). Furthermore, the Glen Scaddle Synform is defined within the intrusion by the orientation of primary layering which is folded by NW-trending upright folds that are equated with the  $D_3$  folds developed within the host Moine rocks (Stoker, 1983). Local structural complexity within the core of the synform is indicated by the observation that the foliation in the metagabbro and the mylonitic fabric within early granite veins are also locally deformed by tight, steeply plunging  $D_3$  folds [e.g. NM 9965 6856]. M. Stoker (unpub. Ph.D. thesis, Univ. Liverpool, 1980) interpreted this complexity as indicating that  $D_3$  was locally polyphase.

A pre- $D_4$  age of intrusion is certain because the foliation is deformed and crenulated by NE–SW-trending open folds that are correlated with regional folds of this age within the Moine country rocks. A mesoscopic  $D_4$  fold pair deforms the Glen Gour body and the southeasternmost part of the Glen Scaddle Metagabbro (Fig. 2; Stoker, 1983).

### 3.e. Summary

It is therefore concluded, following Stoker (1983), that the Glen Scaddle Metagabbro was emplaced after  $D_2$ , probably as a series of sills that were concordant with the regional  $S_0/S_1/S_2$  foliation within host Moine rocks. This foliation was probably flat-lying after  $D_2$ . The generalized gradation from ultrabasic rocks at the margins of the Glen Scaddle and Glen Gour intrusions to basic and acidic rocks centrally




Figure 7. Hand specimen of foliated metagabbro sampled for isotopic dating in this study.

suggests that they may represent differentiated layered sheet intrusions. The intrusion was then deformed and metamorphosed during  $D_3$  and  $D_4$ . The structural evidence therefore indicates that the Glen Scaddle Metagabbro is an important marker within the regional tectonometamorphic sequence; its age places a lower limit on the timing of regional upright folding and associated amphibolite-facies metamorphism.

## 4. U–Pb geochronology

### 4.a. Sample preparation procedure and analytical techniques

The sample analysed was a medium-grained, foliated metagabbro (Fig. 7) collected from the River Scaddle [NN 0004 6869]. A sample of 30 kg was crushed and milled, and the less than  $400 \mu$  fraction sieved out. Heavy mineral concentrations were obtained using a Gemini shaking table, followed by a superpanner. A separate, with specific gravity greater than  $3.3 \text{ gm ml}^{-1}$ , was recovered using Di-iodomethane. The minerals were then separated magnetically using a Frantz LB-1 magnetic separator. The recovered zircons from the non-magnetic  $\sim 1.8$  Amp fraction were hand-picked under alcohol and abraded. The majority of the zircons in the sample were fragments of quite large grains (about  $150 \mu$  length). The fragments were predominantly clear with sharp terminations and edges where visible. A few showed elongate melt inclusions and these were avoided during picking.

U and Pb separations followed the procedures of Krogh (1973) with minor modifications of Corfu & Ayres (1984). Zircon grains were analysed on a VG 354 mass spectrometer at the NERC Isotope Geosciences Laboratory following the procedures of Noble, Tucker & Pharaoh (1993). Chemistry blanks were  $\sim 5 \text{ pg}$ , and these were monitored in each batch of chemistry.




Figure 8. U–Pb concordant ages from the Glen Scaddle metagabbro. The MSWD is of concordance and equivalence at  $2\sigma$  and takes into account errors on the decay constant. The labelled ellipse is the weighted mean error ellipse of the three data points.

Uranium blanks contained  $< 0.1$  pg U. All results and errors were calculated following the methods of Ludwig (1993, 1994), and the Pb isotope ratios were corrected for initial common Pb in excess of laboratory blank using the model of Stacey & Kramers (1975). Ages were calculated using the decay constants of Jaffey *et al.* (1971).

**4.b. Results**

Three zircon fractions plot either on or close to concordia and give an age of  $426 \pm 3$  Ma (Fig. 8; Table 1). This age is interpreted to date closely igneous crystallization of the Glen Scaddle Metagabbro.

**5. Discussion**

**5.a. Age of polyphase upright folding in the SW Moine**

The U–Pb zircon age of  $426 \pm 3$  Ma reported here for the emplacement of the Glen Scaddle Metagabbro suggests that this intrusion is a basic member of the late Caledonian ‘Newer Granite’ suite (Read, 1961; Stephenson *et al.* 1999, and references therein). More importantly, in combination with the structural history outlined here, as well as the  $425 \pm 3$  Ma age for the Strontian Granite located only a few kilometres to the south, it implies a very different regional tectonometamorphic history for the Moine Supergroup in West Inverness-shire and Ross-shire to that currently published. Regional-scale  $D_3$ – $D_4$  upright folding, including the formation of the Northern Highland Steep Belt, must have occurred during a relatively short (Scandian) orogenic event. Western parts of the Strontian Granite are overprinted by locally developed steep, N–S-trending solid-state deformation fabrics (Hutton, 1988), and it is therefore possible that granite emplacement overlapped final stages of  $D_4$  folding.

Table 1. U–Pb data for zircon from the Glen Scaddle Metagabbro

Fraction code	$^1\text{Weight} (\mu\text{g})$	$^2\text{U} (\text{ppm})$	$^2\text{Pb} (\text{ppm})$	Total Pb (pg)	$^{206}\text{Pb}/^{204}\text{Pb}^3$	$^{208}\text{Pb}/^{206}\text{Pb}^3$	$^{206}\text{Pb}/^{238}\text{U}^3$	$\pm \%$	$^{207}\text{Pb}/^{235}\text{U}^3$	$\pm \%$	$^{207}\text{Pb}/^{206}\text{Pb}^4$	$\pm \text{Ma}$	Rho <sup>5</sup>
GSM-03-a	5.0	408	29.34	15	49225	0.1669	0.0683	0.69	0.52046	0.74	0.05528	6	0.94
GSM-03-b	8.0	307	24.02	17	5073	0.2543	0.0686	0.57	0.52189	0.64	0.05514	6	0.90
GSM-03-c	10.0	659	51.58	12	3800	0.2570	0.0683	0.59	0.52307	0.69	0.05557	8	0.86

(1) All analyses picked from non magnetic at 1.8 Amp fraction of zircons. (2) Sample weights and hence U and Pb concentrations are approximate. (3) Measured ratios are corrected for fractionation and common Pb spike. (4) Corrected for fractionation, spike, laboratory blank and initial common Pb calculated at 427 Ma (Stacey & Kramers, 1975). (5) Correlation coefficients of  $^{207}\text{Pb}/^{206}\text{Pb}$  and  $^{206}\text{Pb}/^{238}\text{U}$  are calculated using procedures and algorithms of (Ludwig, 1993). Errors for the measured ratios are propagated through data reduction and quoted at the 2 sigma level.

A younger age for regional upright folding than the Ordovician age proposed previously resolves the apparently contradictory structural relationships in Ross-shire, where steep belt folds deform structures assigned to the Scandian event (Kinny *et al.* 2003). However, a Silurian age for the formation of the Northern Highland Steep Belt is in contradiction to the U–Pb monazite age of  $450 \pm 10$  Ma obtained from a pegmatite west of Glenfinnan that was interpreted to post-date upright folding (van Breemen, Pidgeon & Johnson, 1974). The exact significance of this age is difficult to evaluate, partly due to the lack of a detailed field description of the dated pegmatite and its structural setting, but also because of the possibility that the dated monazites included an unrecognized inherited component. We consider the new age constraints reported here on the age of regional upright folding more reliable, although the structural setting and age of pegmatites within the steep belt clearly deserves reinvestigation.

#### 5.b. Implications for late-orogenic displacements along the Great Glen Fault

A Silurian age for regional, polyphase upright folding and associated amphibolite-facies metamorphism of the Moine Supergroup is consistent with the current model for Caledonian orogenic activity in the Northern Highland terrane, involving Grampian (470–460 Ma) and Scandian (435–425 Ma) events (Kinny *et al.* 1999, 2003; Friend, Jones & Burns, 2000; Dallmeyer *et al.* 2001). The new evidence reported here, in combination with the findings of Kinny *et al.* (2003), demonstrates widespread reworking of Grampian structures and metamorphic assemblages during the Scandian event. This culminated in development of the Moine Thrust Zone and emplacement of the Moine rocks onto the Laurentian foreland. The Northern Highland Steep Belt is thought to have formed at the same time and presumably detaches at depth on a ductile thrust zone, probably the Moine Thrust (Barr, Holdsworth & Roberts, 1986).

The tectonothermal history detailed above is in marked contrast to that recorded by the Dalradian Supergroup southeast of the Great Glen Fault, where the main phases of folding and associated regional metamorphism are assigned entirely to the Ordovician Grampian orogenic event. The present consensus is that the peak of Grampian metamorphism in the Dalradian rocks occurred at *c.* 470 Ma and was followed by cooling and uplift relatively soon thereafter at *c.* 460 Ma (Dempster, 1985; Dempster, Hudson & Rogers, 1995; Dempster *et al.* 2002; Soper, Ryan & Dewey, 1999; Oliver *et al.* 2000; Oliver 2001; Baxter, Ague & DePaulo, 2002). As yet, there is no evidence that the Dalradian Supergroup was affected by regionally significant Silurian deformation or metamorphism. The conclusions reported here therefore

reinforce tectonic models that invoke a minimum of 700 km of late Caledonian (425–390 Ma?) sinistral movement along the Great Glen Fault in order to juxtapose the Grampian and Northern Highland terranes following the Scandian collision between Baltica and the Northern Highland segment of eastern Laurentia (Dewey & Strachan, 2003; Kinny *et al.* 2003).

## 6. Conclusions

- (1) The Glen Scaddle Metagabbro forms a valuable time marker within the Northern Highland Terrane of the Scottish Caledonides because its age places a lower limit on the timing of regional upright folding and associated amphibolite-facies metamorphism.
- (2) The intrusion has yielded a U–Pb zircon age of  $426 \pm 3$  Ma that is interpreted to date its igneous crystallization.
- (3) Regional-scale D<sub>3</sub> and D<sub>4</sub> upright folding of the intrusion and its Moine host rocks, including formation of the Northern Highland Steep Belt, occurred during a relatively short Silurian (= Scandian) orogenic event. Final stages of D<sub>4</sub> deformation may have overlapped emplacement of the Strontian Granite, dated previously at  $425 \pm 3$  Ma.
- (4) The data reported here provide further evidence for widespread Silurian ductile deformation and amphibolite-facies metamorphism northwest of the Great Glen Fault; the apparent absence of this orogenic event within the adjacent Grampian Terrane reinforces arguments that this structure separates crustal blocks with contrasting Caledonian tectonothermal histories.
- (5) These differences support tectonic models that invoke a minimum of 700 km late Caledonian sinistral movement along the Great Glen Fault in order to juxtapose the Grampian and Northern Highland terranes following the Scandian collision.
- (6) This study therefore shows how comparison of the varying intensities of deformational events across an orogen can be used to reconstruct original collisional templates and constrain likely displacements along late-orogenic transcurrent faults.

**Acknowledgements.** Isotopic dating was funded by a grant from the Steering Committee of the NERC Isotope Geosciences Laboratory. Martyn Stoker is thanked for discussions about the geology of the Glen Scaddle area; Tony Harris and John Mendum contributed detailed reviews.



## References

- ASHWORTH, J. R. & CHINNER, G. A. 1978. Coexisting garnet and cordierite in migmatites from the Scottish Caledonides. *Contributions to Mineralogy and Petrology* **65**, 379–94.
- BAILEY, E. B. & MAUFE, H. B. 1916. *The geology of Ben Nevis, Glen Coe and the surrounding area*. Memoir of the Geological Survey of Great Britain.
- BARR, D., HOLDSWORTH, R. E. & ROBERTS, A. M. 1986. Caledonian ductile thrusting in a Precambrian metamorphic complex: the Moine of north-western Scotland. *Geological Society of America Bulletin* **97**, 754–64.
- BAXTER, E. F., AGUE, J. J. & DEPAULO, D. J. 2002. Prograde temperature–time evolution in the Barrovian type locality constrained by precise Sm/Nd garnet ages from Glen Clova, Scotland. *Journal of the Geological Society, London* **159**, 71–82.
- CAWOOD, P. A., NEMCHIN, A. A., STRACHAN, R. A., KINNY, P. D. & LOEWY, S. 2004. Laurentian provenance and tectonic setting for the upper Moine Supergroup, Scotland, constrained by detrital zircons from the Loch Eil and Glen Urquhart successions. *Journal of the Geological Society, London* **161**, 861–74.
- CORFU, F. & AYRES, L. D. 1984. U–Pb ages and genetic significance of heterogeneous zircon populations in rocks from the Favourable Lake area, north-western Ontario. *Contributions to Mineralogy and Petrology* **88**, 86–101.
- DALLMEYER, R. D., STRACHAN, R. A., ROGERS, G., WATT, G. R. & FRIEND, C. R. L. 2001. Dating deformation and cooling in the Caledonian thrust nappes of north Sutherland, Scotland: insights from  $^{40}\text{Ar}/^{39}\text{Ar}$  and Rb–Sr chronology. *Journal of the Geological Society, London* **158**, 501–12.
- DALZIEL, I. W. D. 1966. A structural study of the granitic gneiss of western Ardgour, Argyll and Inverness-shire. *Scottish Journal of Geology* **2**, 125–52.
- DALZIEL, I. W. D. & SOPER, N. J. 2001. Neoproterozoic extension on the Scottish promontory of Laurentia: paleogeographic and tectonic implications. *Journal of Geology* **109**, 299–317.
- DEMPSTER, T. J. 1985. Uplift patterns and orogenic evolution in the Scottish Dalradian. *Journal of the Geological Society, London* **142**, 111–28.
- DEMPSTER, T. J., HUDSON, N. F. & ROGERS, G. 1995. Metamorphism and cooling of the NE Dalradian. *Journal of the Geological Society, London* **152**, 383–90.
- DEMPSTER, T. J., ROGERS, G., TANNER, P. W. G., BLUCK, B. J., MUIR, R. J., REDWOOD, S. D., IRELAND, T. R. & PATERSON, B. A. 2002. Timing of deposition, orogenesis and glaciation within the Dalradian rocks of Scotland: constraints from U–Pb zircon ages. *Journal of the Geological Society, London* **159**, 83–94.
- DEWEY, J. F. 2006. Orogeny can be very short. *Proceedings of the National Academy of Sciences* **102**, 15286–93.
- DEWEY, J. F., HEMPTON, M. R., KIDD, W. S. F., SAROĞLU, F. & ŞENGÖR, A. M. C. 1986. Shortening of continental lithosphere: the neotectonics of Eastern Anatolia – a young collision zone. In *Collision Tectonics* (eds M. P. Coward & A. C. Ries), pp. 3–36. Geological Society of London, Special Publication no. 19.
- DEWEY, J. F. & MANGE, M. 1999. Petrography of Ordovician and Silurian sediments in the western Irish Caledonides: tracers of a short-lived Ordovician continent–arc collision orogeny and the evolution of the Laurentian–Appalachian–Caledonian collision. In *Continental Tectonics* (eds C. MacNiocall & P. D. Ryan), pp. 55–107. Geological Society of London, Special Publication no. 164.
- DEWEY, J. F. & RYAN, P. D. 1990. The Ordovician evolution of the South Mayo Trough, western Ireland. *Tectonics* **9**, 887–903.
- DEWEY, J. F. & STRACHAN, R. A. 2003. Changing Silurian–Devonian relative plate motion in the Caledonides: sinistral transpression to sinistral transtension. *Journal of the Geological Society, London* **160**, 219–29.
- DREVER, H. I. 1940. The geology of Ardgour, Argyllshire. *Transactions of the Royal Society of Edinburgh* **60**, 141–71.
- FRIEND, C. R. L., JONES, K. A. & BURNS, I. M. 2000. New high-pressure granulite facies event in the Moine Supergroup, northern Scotland: implications for Taconic (early Caledonian) crustal evolution. *Geology* **28**, 543–6.
- FRIEND, C. R. L., KINNY, P. D., ROGERS, G., STRACHAN, R. A. & PATERSON, B. A. 1997. U–Pb zircon geochronological evidence for Neoproterozoic events in the Glenfinnan Group (Moine Supergroup): the formation of the Ardgour granite gneiss, north-west Scotland. *Contributions to Mineralogy and Petrology* **128**, 101–13.
- FRIEND, C. R., STRACHAN, R. A., KINNY, P. D. & WATT, G. R. 2003. Provenance of the Moine Supergroup of NW Scotland: evidence from geochronology of detrital and inherited zircons from sediments, granites and migmatites. *Journal of the Geological Society, London* **160**, 247–57.
- HARRIS, A. L. 1995. The nature and timing of orogenesis in the Scottish Highlands and the role of the Great Glen Fault. In *Current Perspectives in the Appalachian–Caledonian Orogen* (eds J. Hibbard, C. R. van Staal & P. A. Cawood), pp. 65–79. Geological Association of Canada, Special Paper no. 41.
- HARRIS, A. L., HASELOCK, P. J., KENNEDY, M. J. & MENDUM, J. R. 1994. The Dalradian Supergroup in Scotland, Shetland and Ireland. In *A revised correlation of Precambrian rocks in the British Isles* (eds W. Gibbons & A. L. Harris), pp. 33–53. Geological Society of London, Special Report no. 22.
- HIGHTON, A. J., HYSLOP, E. K. & NOBLE, S. R. 1999. U–Pb zircon geochronology of migmatization in the northern Central Highlands: evidence for pre-Caledonian (Neoproterozoic) tectonometamorphism in the Grampian Block, Scotland. *Journal of the Geological Society, London* **156**, 1195–1204.
- HOLDSWORTH, R. E. & ROBERTS, A. M. 1984. A study of early curvilinear fold structures and strain in the Moine of the Glen Garry region, Inverness-shire. *Journal of the Geological Society, London* **141**, 327–38.
- HOLDSWORTH, R. E., STRACHAN, R. A. & HARRIS, A. L. 1994. Precambrian rocks in northern Scotland east of the Moine Thrust: the Moine Supergroup. In *A revised correlation of Precambrian rocks in the British Isles* (eds W. Gibbons & A. L. Harris), pp. 23–32. Geological Society of London, Special Report no. 22.
- HUTTON, D. H. W. 1988. Igneous emplacement in a shear zone termination: The biotite granite at Strontian, Scotland. *Geological Society of America Bulletin* **100**, 1392–9.

- JAFFEY, A. H., FLYNN, G. K. F., GLENDENIN, L. E., BENTLEY, W. C. & ESSLING, A. M. 1971. Precision measurements of half-lives and specific activities of  $^{235}\text{U}$  and  $^{238}\text{U}$ . *Physical Reviews C* **4**, 1889–1906.
- KELLEY, S. P. & POWELL, D. 1985. Relationships between marginal thrusting and movement on major, internal shear zones in the N. Highland Caledonides, Scotland. *Journal of Structural Geology* **7**, 43–56.
- KINNY, P. D., FRIEND, C. R. L., STRACHAN, R. A., WATT, G. R. & BURNS, I. M. 1999. U–Pb geochronology of regional migmatites, East Sutherland, Scotland: evidence for crustal melting during the Caledonian orogeny. *Journal of the Geological Society, London* **156**, 1143–52.
- KINNY, P. D., STRACHAN, R. A., FRIEND, C. R. L., KOCKS, H., ROGERS, G. & PATERSON, B. 2003. U–Pb geochronology of deformed meta-granites in central Sutherland, Scotland: evidence for widespread Silurian metamorphism and ductile deformation of the Moine Supergroup during the Caledonian orogeny. *Journal of the Geological Society, London* **160**, 259–69.
- KOCKS, H., STRACHAN, R. A. & EVANS, J. A. 2006. Heterogeneous reworking of Grampian metamorphic complexes during Scandian thrusting in the Scottish Caledonides: insights from the structural setting and U–Pb geochronology of the Strath Halladale Granite. *Journal of the Geological Society, London* **163**, 525–38.
- KROGH, T. E. 1973. A low contamination method for the hydrothermal decomposition of zircon and extraction of U and Pb for isotopic age determinations. *Geochimica et Cosmochimica Acta* **37**, 485–94.
- LUDWIG, K. R. 1993. *PBDAT: A computer program for processing Pb–U–Th isotope data, version 1.24*. U.S. Geological Survey Open-file Report 88-542.
- LUDWIG, K. R. 1994. *ISOPLOT: A plotting and regression program for radiogenic-isotope data, version 2.75*. U.S. Geological Survey Open-file Report 91-445.
- MILLAR, I. L. 1999. Neoproterozoic extensional basic magmatism associated with emplacement of the West Highland granite gneiss in the Moine Supergroup of NW Scotland. *Journal of the Geological Society, London* **156**, 1153–62.
- NOBLE, S. R., TUCKER, R. D. & PHARAOH, T. C. 1993. Lower Palaeozoic and Precambrian igneous rocks from eastern England and their bearing on Ordovician closure of the Tornquist Sea: constraints from U–Pb and Nd isotopes. *Geological Magazine* **130**, 835–46.
- OLIVER, G. J. H. 2001. Reconstruction of the Grampian episode in Scotland: its place in the Caledonian Orogeny. *Tectonophysics* **332**, 23–49.
- OLIVER, G. J. H., CHEN, F., BUCHWALDT, R. & HEGNER, E. 2000. Fast tectonometamorphism and exhumation in the type area of the Barrovian and Buchan zones. *Geology* **28**, 459–62.
- PICKERING, K. T., BASSETT, M. G. & SIVETER, D. J. 1988. Late Ordovician–Early Silurian destruction of the Iapetus Ocean: Newfoundland, British Isles and Scandinavia – a discussion. *Transactions of the Royal Society of Edinburgh: Earth Sciences* **79**, 361–82.
- POWELL, D., BAIRD, A. W., CHARNLEY, N. R. & JORDAN, P. J. 1981. The metamorphic environment of the Sgurr Beag Slide: a major crustal displacement zone in Proterozoic, Moine rocks of Scotland. *Journal of the Geological Society, London* **138**, 661–73.
- POWELL, D. & GLENDINNING, R. 1988. Excursion 4: Glenfinnan to Morar. In *An Excursion Guide to the Moine Geology of the Scottish Highlands* (eds I. Allison, F. May & R. A. Strachan), pp. 80–102. Edinburgh: Scottish Academic Press.
- POWELL, D. & PHILLIPS, W. E. A. 1985. Time of deformation in the Caledonide orogen of Britain and Ireland. In *The Nature and Timing of Orogenic Activity in the Caledonian Rocks of the British Isles* (ed. A. L. Harris), pp. 17–39. Geological Society of London, Memoir no. 9.
- READ, H. H. 1961. Aspects of Caledonian magmatism in Scotland. *Proceedings of the Liverpool and Manchester Geological Society* **2**, 653–83.
- ROBERTS, A. M. & HARRIS, A. L. 1983. The Loch Quoich Line – a limit of early Palaeozoic crustal reworking in the Moine of the northern Highlands of Scotland. *Journal of the Geological Society, London* **140**, 883–92.
- ROBERTS, A. M., SMITH, D. I. & HARRIS, A. L. 1984. The structural setting and tectonic significance of the Glen Dessary syenite, Inverness-shire. *Journal of the Geological Society, London* **141**, 1033–42.
- ROGERS, G. & DUNNING, G. R. 1991. Geochronology of appinitic and related granitic magmatism in the W Highlands of Scotland: constraints on the timing of transcurrent fault movement. *Journal of the Geological Society, London* **148**, 17–27.
- ROGERS, G., HYSLOP, E. K., STRACHAN, R. A., PATERSON, B. A. & HOLDSWORTH, R. E. 1998. The structural setting and U–Pb geochronology of Knoydartian pegmatites in W. Inverness-shire: evidence for Neoproterozoic tectonothermal events in the Moine of NW Scotland. *Journal of the Geological Society, London* **155**, 685–96.
- ROGERS, G., KINNY, P. D., STRACHAN, R. A., FRIEND, C. R. L. & PATERSON, B. A. 2001. U–Pb geochronology of the Fort Augustus granite gneiss: constraints on the timing of Neoproterozoic and Palaeozoic tectonothermal events in the NW Highlands of Scotland. *Journal of the Geological Society, London* **158**, 7–14.
- SEARLE, M. P. 2006. Role of the Red River Shear Zone, Yunnan and Vietnam, in the continental extrusion of SE Asia. *Journal of the Geological Society, London* **163**, 1025–36.
- SEARLE, M. P., WEINBERG, R. & DUNLAP, W. J. 1998. Transpressional tectonics along the Karakoram fault zone, northern Ladakh: constraints on Tibetan extrusion. In *Continental Transpressional and Transtensional Tectonics* (eds R. E. Holdsworth, R. A. Strachan & J. F. Dewey), pp. 345–66. Geological Society of London, Special Publication no. 160.
- SMITH, M., ROBERTSON, S. & ROLLIN, K. E. 1999. Rift basin architecture and stratigraphical implications for basement-cover relationships in the Neoproterozoic Grampian Group of the Scottish Caledonides. *Journal of the Geological Society, London* **156**, 1163–73.
- SOPER, N. J. 1988. Timing and geometry of collision, terrane accretion and sinistral strike-slip events in the British Caledonides. In *The Caledonian–Appalachian Orogen* (eds A. L. Harris & D. J. Fettes), pp. 481–92. Geological Society of London, Special Publication no. 38.
- SOPER, N. J., RYAN, P. D. & DEWEY, J. F. 1999. Age of the Grampian orogeny in Scotland and Ireland. *Journal of the Geological Society, London* **156**, 1231–6.
- SOPER, N. J., STRACHAN, R. A., HOLDSWORTH, R. E., GAYER, R. A. & GREILING, R. O. 1992. Sinistral transpression and the Silurian closure of Iapetus. *Journal of the Geological Society, London* **149**, 871–80.

- STACEY, J. S. & KRAMERS, J. D. 1975. Approximation of terrestrial lead isotope evolution by a two-stage model. *Earth and Planetary Science Letters*, **26**, 207–21.
- STEPHENSON, D., BEVINS, R. E., MILLWARD, D., HIGHTON, A. J., PARSONS, I., STONE, P. & WADSWORTH, W. J. 1999. *Caledonian Igneous Rocks of Great Britain*. Geological Conservation Review Series, **17**, Joint Nature Conservation Committee, 1–648.
- STOKER, M. S. 1983. The stratigraphy and structure of the Moine rocks of eastern Ardgour. *Scottish Journal of Geology* **19**, 369–85.
- STRACHAN, R. A. 1985. The stratigraphy and structure of the Moine rocks of the Loch Eil area, West Inverness-shire. *Scottish Journal of Geology* **21**, 9–22.
- STRACHAN, R. A., SMITH, M., HARRIS, A. L. & FETTES, D. J. 2002. The Northern Highland and Grampian terranes. In *Geology of Scotland* (4th edition) (ed. N. H. Trewin), pp. 81–147. Geological Society of London.
- TANNER, P. W. G. & EVANS, J. A. 2003. Late Precambrian U–Pb titanite age for peak regional metamorphism and deformation (Knoydartian orogeny) in the western Moine, Scotland. *Journal of the Geological Society, London* **160**, 555–64.
- VANCE, D., STRACHAN, R. A. & JONES, K. A. 1998. Extensional versus compressional settings for metamorphism: garnet chronometry and pressure–temperature–time histories in the Moine Supergroup, northwest Scotland. *Geology* **26**, 927–30.
- VAN BREEMEN, O., AFTALION, M., PANKHURST, R. J. & RICHARDSON, S. W. 1979. Age of the Glen Dessary syenite, Inverness-shire: diachronous Palaeozoic metamorphism across the Great Glen. *Scottish Journal of Geology* **15**, 49–62.
- VAN BREEMEN, O., PIDGEON, R. T. & JOHNSON, M. R. W. 1974. Precambrian and Palaeozoic pegmatites in the Moines of northern Scotland. *Journal of the Geological Society, London* **130**, 493–507.