




**British
Geological Survey**

NATURAL ENVIRONMENT RESEARCH COUNCIL

Model Metadata Report for the Glasgow Geothermal Energy Research Field Site Superficial Deposits Model

UK Geoenergy Observatories Programme

Open Report OR/18/064



BRITISH GEOLOGICAL SURVEY

UK GEOENERGY OBSERVATORIES PROGRAMME

OPEN REPORT OR/18/064

The National Grid and other
Ordnance Survey data © Crown
Copyright and database rights
2018. Ordnance Survey Licence
No. 100021290.

Keywords

UKGEOS, GGERFS, Glasgow,
Superficial, GSI3D.

National Grid Reference

SW corner 258000,660850
NE corner 265000,665000

Front cover

3D image of superficial deposits
beneath the UKGEOS Glasgow
area.

Bibliographical reference

S ARKLEY. 2018. Model
Metadata Report for the Glasgow
Geothermal Energy Research
Field Site Superficial Deposits
Model. *British Geological Survey
Open Report*, OR/18/064. 50pp.

Copyright in materials derived
from the British Geological
Survey's work is owned by UK
Research and Innovation (UKRI)
and/or the authority that
commissioned the work. You
may not copy or adapt this
publication without first
obtaining permission. Contact the
BGS Intellectual Property Rights
Section, British Geological
Survey, Keyworth,
e-mail ipr@bgs.ac.uk. You may
quote extracts of a reasonable
length without prior permission,
provided a full acknowledgement
is given of the source of the
extract.

3D images BGS©UKRI 2018
using GSI3D methodology and
software

Maps and diagrams in this book
use topography based on
Ordnance Survey mapping.

Model Metadata Report for the Glasgow Geothermal Energy Research Field Site Superficial Deposits Model

S Arkley

Contributor/editor

A A Monaghan

BRITISH GEOLOGICAL SURVEY

The full range of our publications is available from BGS shops at Nottingham, Edinburgh, London and Cardiff (Welsh publications only) see contact details below or shop online at www.geologyshop.com

The London Information Office also maintains a reference collection of BGS publications, including maps, for consultation.

We publish an annual catalogue of our maps and other publications; this catalogue is available online or from any of the BGS shops.

The British Geological Survey carries out the geological survey of Great Britain and Northern Ireland (the latter as an agency service for the government of Northern Ireland), and of the surrounding continental shelf, as well as basic research projects. It also undertakes programmes of technical aid in geology in developing countries.

The British Geological Survey is a component body of the Natural Environment Research Council.

British Geological Survey offices

**Environmental Science Centre, Keyworth, Nottingham
NG12 5GG**

Tel 0115 936 3100

BGS Central Enquiries Desk

Tel 0115 936 3143

email enquiries@bgs.ac.uk

BGS Sales

Tel 0115 936 3241

email sales@bgs.ac.uk

**The Lyell Centre, Research Avenue South, Edinburgh
EH14 4AP**

Tel 0131 667 1000

email scotsales@bgs.ac.uk

Natural History Museum, Cromwell Road, London SW7 5BD

Tel 020 7589 4090

Tel 020 7942 5344/45

email bgs-london@bgs.ac.uk

**Cardiff University, Main Building, Park Place, Cardiff
CF10 3AT**

Tel 029 2167 4280

**Macleon Building, Crowmarsh Gifford, Wallingford
OX10 8BB**

Tel 01491 838800

**Geological Survey of Northern Ireland, Department of
Enterprise, Trade & Investment, Dundonald House, Upper
Newtownards Road, Ballymiscaw, Belfast, BT4 3SB**

Tel 028 9038 8462

www.bgs.ac.uk/gsni/

Parent Body

**UK Research and Innovation, Polaris House, Swindon
SN2 1FL**

Tel 01793 444000

www.ukri.org

Website www.bgs.ac.uk

Shop online at www.geologyshop.com

Acknowledgements

This model builds on previous British Geological Survey (BGS) superficial models of the Glasgow area, developed as part of the Clyde Basin Urban Superproject (CUSP). A large number of individuals from that project contributed to creating those models and I would like to thank Diarmad Campbell, Andrew Finlayson, Tim Kearsey, Dave Lawrence, Sue Loughlin, Jo Merritt, Alison Monaghan, Ricky Terrington, Katie Whitbread and Charlotte Vye-Brown for their parts in creating previous versions the superficial geology models of Glasgow.

Glasgow City Council and Clyde Gateway Urban Regeneration Company have provided datasets and feedback on geological modelling over many years and are also thanked here.

In addition I would like to thank Tony Irving who has spent many years entering borehole data into the BGS corporate database, which underpins the geological model and interpretations presented here.

Contents

Acknowledgements.....	2
Contents.....	2
Summary	7
1 Modelled Volume, Purpose and Scale	7
2 Modelled Surfaces/Volumes	10
2.1 Sand and Gravel Unit (SUPD_XSV).....	13
2.2 Cadder Sand and Gravel Formation (CADR).....	13
2.3 Wilderness Till Formation (WITI)	13
2.4 Broomhouse Sand and Gravel Formation (BHSE).....	14
2.5 Clyde Clay Formation	16
2.6 Clydebank Clay Formation.....	18
2.7 Clyde Valley Formation	19
2.8 Peat	19
2.9 Made Ground (MGR)	19
2.10 Lenses.....	20
3 Model Workflow.....	21
4 Model Datasets	27
4.1 Map Data	27
4.2 GVS and GLEG Files	29
4.3 Digital Elevation Model	29
4.4 Borehole Data	30
5 Model Assumptions and Limitations.....	31
5.1 General Modelling Limitations	32

5.2	DEM	32
5.3	Borehole data	32
5.4	Map data	33
5.5	Modelled surfaces and volumes	33
6	Model Quality Assurance	33
7	Model Uncertainty	34
8	Model exports	35
9	Model Images.....	36
9.1	Cross-section images	36
9.2	Thickness maps.....	41
10	References	47

FIGURES

Figure 1 Location map of the Clyde Gateway regeneration area (blue outline) and GGERFS superficial model area (red outline) indicating the approximate location of the UKGEOS Glasgow boreholes. Includes mapping data licensed from Ordnance Survey. © Crown Copyright and/or database right 2019. Licence number 100021290 EUL..... 8

Figure 2 3D view of the Digital Terrain Model (5 m cell size) across the UKGEOS Glasgow area. The ground surface across the study area is between 2 m and 60 m above Ordnance Datum (OD). The course of the River Clyde can be clearly seen snaking its way north-westwards along the low ground of the Clyde valley (blue colours) with higher ground (orange colours) to the northeast and southwest. NEXTMap Britain elevation data from Intermap Technologies. 9

Figure 3 Location of the UKGEOS Glasgow study area (red rectangle) and the surrounding topography, depicted by the NEXTMap Surface model hillshade. Superficial deposits (DigMapGB-625k) are overlain as coloured tints: blue – glacial till, pink – glacial sand and gravel, orange – raised marine deposits, brown – peat, yellow – alluvium and green – blown sand. 10

Figure 4: Modelled distribution of Cadder Sand and Gravel Formation (orangey pink) and the Sand and Gravel unit (pink) across the UKGEOS Glasgow model area (red outline). 1 km grid squares shown. Includes mapping data licensed from Ordnance Survey. © Crown Copyright and/or database right 2019. Licence number 100021290 EUL. 13

Figure 5: Modelled distribution of Wilderness Till Formation (pale blue) across the UKGEOS Glasgow superficial geology model area (red outline). 1 km grid squares shown. Includes mapping data licensed from Ordnance Survey. © Crown Copyright and/or database right 2019. Licence number 100021290 EUL). 14

Figure 6: Modelled distribution of Broomhouse Sand and Gravel Formation (pink) across the UKGEOS Glasgow superficial geology model area (red outline). 1 km grid squares shown. Includes mapping data licensed from Ordnance Survey. © Crown Copyright and/or database right 2019. Licence number 100021290 EUL). 15

Figure 7: Modelled distribution of Broomhouse Sand and Gravel Formation (orange) and Ross Sand Member (pale pink), across the UKGEOS Glasgow superficial geology model area (red

outline). 1 km grid squares shown. Includes mapping data licensed from Ordnance Survey. © Crown Copyright and/or database right 2019. Licence number 100021290 EUL). 16

Figure 8: Modelled distribution of Paisley Clay Member (green) across the UKGEOS Glasgow superficial geology model area (red outline). 1 km grid squares shown. Includes mapping data licensed from Ordnance Survey. © Crown Copyright and/or database right 2019. Licence number 100021290 EUL). 17

Figure 9: Modelled distribution of Killearn Sand and Gravel Member (red) across the UKGEOS Glasgow superficial geology model area (red outline). 1 km grid squares shown. Includes mapping data licensed from Ordnance Survey. © Crown Copyright and/or database right 2019. Licence number 100021290 EUL). 18

Figure 10: Modelled distribution of Gourock Sand Member (pale yellow), Law Sand and Gravel Member (greeny yellow) and peat (brown), across the UKGEOS Glasgow superficial geology model area (red outline). 1 km grid squares shown. Includes mapping data licensed from Ordnance Survey. © Crown Copyright and/or database right 2019. Licence number 100021290 EUL). 19

Figure 11: Modelled distribution of made ground (MGR), across the UKGEOS Glasgow superficial geology model area (red outline). 1 km grid squares shown. Includes mapping data licensed from Ordnance Survey. © Crown Copyright and/or database right 2019. Licence number 100021290 EUL). 20

Figure 12 Screenshot of the GSI3D software showing the borehole, map and section window on the left-hand side and a view of the interlocking sections in the 3D window on the right-hand side. 21

Figure 13 Areas circled in red show where the UKGEOS model was updated/improved during phase 1, green lines indicate existing GSI3D section lines from the published Central Glasgow model, purple dots represent the location of boreholes entered into the BGS borehole database post-2005. 22

Figure 14 Example of a GSI3D section (Tester 1) which was significantly altered by including an additional 4 boreholes drilled since 2005. The upper image (original section) contained little borehole information to constrain the geology in this part of the section which has resulted in the construction of fairly simple geology. The lower image (updated section) indicates how additional borehole data suggests the presence of a further sand-body (pink), confirms the presence of till (blue) above rockhead across this part of the section, indicates that the Paisley Clay (green) has a patchy extent and provides improved constraint on the thickness and lateral extent of all the superficial deposits. 23

Figure 15 Example of a GSI3D section (NW_SE_5) which required little alteration when an additional 3 boreholes, drilled since 2005, were added: Upper image (original section) - little constraint in this part of the section – geology extrapolated from nearest borehole data and geological map at the surface. Lower image (updated section) – three additional boreholes confirm the superficial deposits present as predicted and provide improved constraint on the thickness of the deposits. 24

Figure 16 Summary map of proposed UKGEOS Glasgow borehole locations and sections constructed in GSI3D through the sites to indicate the nature and thickness of the superficial deposits beneath the surface. 25

Figure 17 Network of sections constructed in the vicinity of the UKGEOS Glasgow model area viewed in 3D. 25

Figure 18 Overview of the calculated superficial deposits model across the UKGEOS Glasgow model area, looking north-east, 3x vertical exaggeration. NEXTMap Britain elevation data from Intermap Technologies. 26

- Figure 19 Rockhead surface for the study area, generated in GSI3D by combining the bases of all the modelled superficial units. Low values are in yellow and brown and high values are in purple and white. Includes mapping data licensed from Ordnance Survey. © Crown Copyright and/or database right 2019. Licence number 100021290 EUL). 26
- Figure 20 1:50 000 scale (green) and 1:10 000 scale (blue) geological map sheets covering the UKGEOS Glasgow model area (pink). Includes mapping data licensed from Ordnance Survey. © Crown Copyright and/or database right 2019. Licence number 100021290 EUL).28
- Figure 21 UKGEOS Glasgow model area summary of superficial geology from BGS©UKRI 2016 DigMap 1:10 000 scale data, the Clyde Gateway urban regeneration area is shown in blue. 29
- Figure 22 Digital Terrain Model (50 m cell size) across the UKGEOS Glasgow superficial deposits model area (red outline), as used in the model. NEXTMap Britain elevation data from Intermap Technologies. Includes mapping data licensed from Ordnance Survey. © Crown Copyright and/or database right 2019. Licence number 100021290 EUL. 30
- Figure 23 Location of borehole records held in the BGS Single Onshore Borehole Index database. Red dots (more than 30 m deep), blue dots (between 10 m and 30 m), green dots (less than 10 m) and black dots (depth unknown). Includes mapping data licensed from Ordnance Survey. © Crown Copyright and/or database right 2019. Licence number 100021290 EUL. 31
- Figure 24 Distribution of boreholes used to construct the correlated cross-sections which constrain the superficial deposits model of the UKGEOS Glasgow area. 34
- Figure 25 Distribution of boreholes used to construct the correlated cross-sections which constrain the superficial deposits model around the proposed borehole sites for the UKGEOS Glasgow project. The River Clyde is shown in pale blue. 35
- Figure 26 Cross-section *GGERFS_slba_3*, orientated SSW-NNE through the superficial deposits of the UKGEOS Glasgow model area (see Figure 16 for location). Vertical exaggeration x 3. Borehole constraint points shown in red. Ground surface derived from NEXTMap Britain elevation data from Intermap Technologies. Height is in metres relative to Ordnance Datum.36
- Figure 27 Cross-section *GGERFS_slba_4*, orientated NW-SE through the superficial deposits of the UKGEOS Glasgow model area (see Figure 16 for location). Vertical exaggeration x 3. Borehole constraint points shown in red. Ground surface derived from NEXTMap Britain elevation data from Intermap Technologies. Height is in metres relative to Ordnance Datum.37
- Figure 28 Cross-section *GGERFS_slba_5*, orientated South-North through the superficial deposits of the UKGEOS Glasgow model area (see Figure 16 for location). Vertical exaggeration x 3. Borehole constraint points shown in red. Ground surface derived from NEXTMap Britain elevation data from Intermap Technologies. Height is in metres relative to Ordnance Datum. 37
- Figure 29 Cross-section *GGERFS_slba_7*, orientated NW-SE through the superficial deposits of the UKGEOS Glasgow model area (see Figure 16 for location). Vertical exaggeration x 3. Borehole constraint points shown in red. Ground surface derived from NEXTMap Britain elevation data from Intermap Technologies. Height is in metres relative to Ordnance Datum.38
- Figure 30 Cross-section *GGERFS_slba_8*, orientated North-South through the superficial deposits of the UKGEOS Glasgow model area (see Figure 16 for location). Vertical exaggeration x 3. Borehole constraint points shown in red. Ground surface derived from NEXTMap Britain elevation data from Intermap Technologies. Height is in metres relative to Ordnance Datum. 38
- Figure 31 Cross-section *GGERFS_slba_10*, orientated West-East through the superficial deposits of the UKGEOS Glasgow model area (see Figure 16 for location). Vertical exaggeration x 3.

Borehole constraint points shown in red. Ground surface derived from NEXTMap Britain elevation data from Intermap Technologies. Height is in metres relative to Ordnance Datum.39

- Figure 32 Cross-sections *GGERFS_slba_8* and *GGERFS_slba_10* as viewed in the 3D window of GSI3D. These two sections pass through the proposed site for borehole GGERFS10 at Dalmarnock (see Figure 16 for location and orientation of sections). The sections illustrate the calculated sequence and thickness of superficial deposits beneath this part of the UKGEOS Glasgow model area from available borehole and map data. Vertical exaggeration x 10. Ground surface derived from NEXTMap Britain elevation data from Intermap Technologies. Borehole constraint points along the sections and thickness of the geological units are indicated in Figures 24 and 25. 39
- Figure 33 Nine cross-sections as viewed in the 3D window of GSI3D. This network of sections pass through the proposed sites for the Cuningar Loop boreholes (see Figure 16 for location and orientation of sections). The sections illustrate the calculated sequence and thickness of superficial deposits beneath this part of the UKGEOS Glasgow model area from available borehole and map data. Vertical exaggeration x 10. Ground surface derived from NEXTMap Britain elevation data from Intermap Technologies. Borehole constraint points along some of the sections and thickness of the geological units are indicated in Figures 24 and 25..... 40
- Figure 34 Modelled distribution and thickness of Made Ground deposits (MGR) across the UKGEOS Glasgow superficial geology model area (red outline). 1 km grid squares shown. Includes mapping data licensed from Ordnance Survey. © Crown Copyright and/or database right 2019. Licence number 100021290 EUL. 41
- Figure 35 Modelled distribution and thickness of the Gourock Sand Member (GOSA) across the UKGEOS Glasgow superficial geology model area (red outline). These marine sands are extensive within the Clyde Valley and are over 15 m thick in places. 1 km grid squares shown. Includes mapping data licensed from Ordnance Survey. © Crown Copyright and/or database right 2019. Licence number 100021290 EUL. 42
- Figure 36 Modelled distribution and thickness of the Paisley Clay Member (PAIS) across the UKGEOS Glasgow superficial geology model area (red outline). These laminated marine clays and silts are extensive within the Clyde Valley and are over 30 m thick in places. 1 km grid squares shown. Includes mapping data licensed from Ordnance Survey. © Crown Copyright and/or database right 2019. Licence number 100021290 EUL. 43
- Figure 37 Modelled distribution and thickness of the Bridgeton Sand Member (BRON) across the UKGEOS Glasgow superficial geology model area (red outline). This deposit of marine sands is significant in the north-western corner of the modelled area where it is over 30 m thick in places. 1 km grid squares shown. Includes mapping data licensed from Ordnance Survey. © Crown Copyright and/or database right 2019. Licence number 100021290 EUL.44
- Figure 38 Modelled distribution and thickness of the Broomhouse Sand and Gravel Formation (BHSE) across the UKGEOS Glasgow superficial geology model area (red outline). This deposit of fluvioglacial sand and gravels generally occurs at a significant depth beneath the surface, as its known occurrence (shown above) is dependent on the presence of deep borehole records more extensive deposits may lie undetected in areas where there are a lack of data points. The unit is over 20 m thick in places. 1 km grid squares shown. Includes mapping data licensed from Ordnance Survey. © Crown Copyright and/or database right 2019. Licence number 100021290 EUL..... 45
- Figure 39 Modelled distribution and thickness of the Wilderness Till Formation (WITI) across the UKGEOS Glasgow superficial geology model area (red outline). This deposit of glacial till is widespread across the modelled area and beyond. It is thought to be almost 50 m thick in till-cored drumlins shaped by the over-riding Late Devensian glaciers. 1 km grid squares shown. Includes mapping data licensed from Ordnance Survey. © Crown Copyright and/or database right 2019. Licence number 100021290 EUL. 46

TABLES

Table 1 Summary of origin of superficial deposits stratigraphic units in the UKGEOS Glasgow model area: lithology, origin and age (Lac – lacustrine, Fluv – fluvial).	11
Table 2 The 15 lithostratigraphic units differentiated in the 3D superficial deposits model of the UKGEOS Glasgow area, in stratigraphic order.....	12
Table 3 Summary of the key datasets and associated files used in the construction of the UKGEOS Glasgow superficial deposits model.....	27
Table 4 Summary table of model exports	35

Summary

This report presents the metadata behind the superficial deposits model, developed by the British Geological Survey (BGS), for the Glasgow Geothermal Energy Research Field Site (GGERFS) of the UK Geoenergy Observatories Programme (UKGEOS). The site is located within the Clyde Gateway regeneration area of eastern Glasgow (Glasgow City Council) and Rutherglen (South Lanarkshire Council), central Scotland, UK.

The superficial deposits in the Clyde valley consist of a complex succession of glacial till, marine, lacustrine and fluvio-glacial deposits, overlain by fluvial deposits, recent alluvium and anthropogenic deposits, which locally reach around 50 m in thickness. There is widespread made, filled and landscaped ground mantling the natural superficial deposits.

The geological information in this report and accompanying 3D model characterise the various superficial deposits, and give an indication of their thickness and lateral extent. This has created a better understanding of the Quaternary (superficial) geology of the UKGEOS Glasgow area. The model allows users to visualise the likely subsurface sequence to be found beneath the UKGEOS Glasgow area and create synthetic borehole prognoses.

The model presented in this report builds on previous superficial deposit models of the Glasgow area, developed in BGS since 2005. A number of reports have been published describing previous models of the area, including Merritt et al. (2009), Monaghan et al. (2012, 2014) and Arkley et al. (2013). The model incorporates additional borehole data to increase the model resolution in the vicinity of the planned new borehole sites and incorporates any post-2005 borehole data more recently received by BGS than previous model versions. This version of the model represents a UKGEOS ‘pre-drill’ understanding of the superficial deposits.

The models are an interpretation of digital datasets held by the British Geological Survey. A summary of the construction method, limitations of the models and a brief description of the modelled units are given.

1 Modelled Volume, Purpose and Scale

This report provides an overview of the Glasgow Geothermal Energy Field Research Site superficial deposits model as part of the UK Geoenergy Observatories Programme (UKGEOS).

The model updates previous BGS superficial models in this part of the Clyde Gateway area.

The model was constructed to define the three dimensional spatial distribution of the superficial deposit succession from Devensian glacial till to recent made ground present in the UKGEOS Glasgow area, to help users with an understanding of the (Quaternary) superficial deposits of the area and as a basis for borehole prognoses, particularly those boreholes currently proposed as part of the UK Geoenergy Observatories Programme: the Dalmarnock borehole (GGERFS10) and the Cuningar Loop boreholes (GGERFS01, 02, 03 and 05).

The UKGEOS Glasgow area is located in the East End of Glasgow (Dalmarnock) and Rutherglen, straddling the boundary between Glasgow City and South Lanarkshire councils on the north and south sides of the River Clyde (Figure 1).




Figure 1 Location map of the Clyde Gateway regeneration area (blue outline) and GGERFS superficial model area (red outline) indicating the approximate location of the UKGEOS Glasgow boreholes. Includes mapping data licensed from Ordnance Survey. © Crown Copyright and/or database right 2019. Licence number 100021290 EUL.

The area is within the Clyde Gateway urban regeneration area, c.840 hectares, comprising sports facilities and housing for the 2014 Commonwealth Games, some pre-existing 19th and 20th century legacy housing, and vacant and derelict land with a history of multiple former industrial use and housing (Figure 1). Parts of the land are owned by [Clyde Gateway](#) URC (Urban Regeneration Company, a partnership of Glasgow City Council, South Lanarkshire Council and Scottish Enterprise backed by the Scottish Government). Some individual blocks of land have been acquired by private developers.

The aim of the study is to describe the distribution of the Quaternary-aged deposits based on a review and the interpretation of the available data held by BGS, largely comprising Site Investigation and exploration borehole data. This has been achieved by entering compositional and lithostratigraphic information into BGS databases and generating surfaces of geological units within the bespoke models using GSI3D software. This work allows the design and location of

UKGEOS boreholes to be decided with reference to the complex Quaternary geology in this area.

The ground surface across the study area is between 2 m and 60 m above Ordnance Datum (OD; Figure 2).




Figure 2 3D view of the Digital Terrain Model (5 m cell size) across the UKGEOS Glasgow area. The ground surface across the study area is between 2 m and 60 m above Ordnance Datum (OD). The course of the River Clyde can be clearly seen snaking its way north-westwards along the low ground of the Clyde valley (blue colours) with higher ground (orange colours) to the northeast and southwest. NEXTMap Britain elevation data from Intermap Technologies.

The UKGEOS Glasgow area falls within the previously published ‘Central Glasgow’ model (Merritt et al., 2009; Monaghan et al., 2014). This model has been updated between the coordinates SW corner 258000, 660850, NE corner 265000, 665000 to include boreholes drilled since 2005 and comprises an area of approximately 7 km x 4 km. The model extends to rockhead which lies between 36 m above OD and 40 m below OD. The UKGEOS Glasgow superficial deposits model is most suitable for use at scales between 1:10 000 to 1:50 000 scale, but is also useful for providing guidance at other scales. Users are advised against basing interpretations on extreme exaggeration of the vertical or horizontal scales of the model.

2 Modelled Surfaces/Volumes

Geological setting

The study area is located within the lower reaches of the Clyde valley which is thought to have been glaciated multiple times over the last 0.5 Ma, most recently during the Main Late Devensian glaciation when ice advanced into the Clyde basin. Glacial ice retreated from the Glasgow area by around 15,000 years BP, at a time when the relative sea level had risen locally to almost 40 m above current sea level and inundated low lying coastal areas. The relative sea level then fell during the Holocene and resulted in a series of raised estuarine flats around the Glasgow area at progressively lower elevations (approximately 12 m to 3 m above current sea level).

In the UKGEOS Glasgow area, the underlying Carboniferous rocks have been scoured by glacial, river and coastal erosion processes leaving an irregular undulating surface buried beneath the Quaternary sediments. Deposited on top of this is a complex succession of Quaternary superficial deposits, varying laterally and vertically across short distances, including widespread glacial till and marine, lacustrine and fluvio-glacial deposits, overlain by fluvial deposits, recent alluvium and anthropogenic (man-made) deposits (Figure 3).

The Quaternary deposits are of variable thicknesses, up to 50 m thick, with thicker accumulations of superficial deposits infilling a broadly NW–SE trending channel following the modern day River Clyde. There is also widespread made, filled and landscaped ground relating to a variety of prior industrial land use, which in some places reaches 10–15 m thick. Table 1 summarises the sequence. The Quaternary geology of the Clyde valley is described in detail in Browne and McMillan (1989), Forsyth et al. (1996), Finlayson et al. (2010) and Finlayson (2012).




Figure 3 Location of the UKGEOS Glasgow study area (red rectangle) and the surrounding topography, depicted by the NEXTMap Surface model hillshade. Superficial deposits (DigMapGB-625k) are overlain as coloured tints: blue – glacial till, pink – glacial sand and gravel, orange – raised marine deposits, brown – peat, yellow – alluvium and green – blown sand.

CLAY AND SILT		SAND AND GRAVEL		DIAMICTON	PEAT	LITHOLOGY
Marine	Lac/Fluv	Marine	Lac/Fluv	Glacial	Organic	ORIGIN
		Gourock	Law		Clippens	FLANDRIAN
		Killearn				
Paisley						DEVENSIAN
		Bridgeton				
	Bellshill					
			Ross			
			Broomhouse			
				Wilderness		
			Cadder			

Table 1 Summary of origin of superficial deposits stratigraphic units in the UKGEOS Glasgow model area: lithology, origin and age (Lac – lacustrine, Fluv – fluvial).

Code	Geological Unit	Equivalent description on 1: 10, 000 scale published map
Water	Water	Unattributed polygons or underlying sediments described
MGR-ARTDP	Made Ground (made and worked ground undifferentiated)	Made Ground (MGR), Made Ground and Worked Ground (WMGR), Infilled Ground (WMGR)
PEAT-P	Peat	Peat – blanket or basin peat, Flandrian (PEAT)
LAWSG-XCZSV	Law Sand and Gravel Member	Alluvium – modern river floodplains – located along the upper reaches and tributaries to the River Clyde, Flandrian (ALV). Also includes some Alluvial Fan Deposits, Flandrian (ALF) and some River Terrace Deposits, Flandrian (RTD1 and RTD2)
GOSA-XCZSV	Gourock Sand Member	Marine Deposits – located along the lower reaches of the River Clyde, Flandrian (MDU) and Alluvium – modern river floodplains – along the upper reaches of the River Clyde, Flandrian (ALV)
KARN-XSV	Killearn Sand and Gravel Member	Generally Raised Marine Deposits, Devensian (RMDV), Raised Marine Deltaic Deposits, Devensian (RMDDD) or Raised Marine Intertidal and

		Subtidal Deposits, Devensian (RMIS)
PAIS-XCZS	Paisley Clay Member	Generally Raised Marine Deposits, Devensian (RMDV) or Raised Marine Intertidal and Subtidal Deposits, Devensian (RMIS)
BRON-XSVZ	Bridgeton Sand Member	Largely concealed beneath younger deposits, where present, exposures usually represented as Raised Marine Deposits, Devensian (RMDV)
RSSA-XSV	Ross Sand Member	Glaciolacustrine Deposits, Devensian (GLLDD), Glaciolacustrine Deltaic Deposits, Devensian (GLDDD) or Glaciofluvial Deposits, Devensian (GFDUD)
RSSA-XSZ	Ross Sand Member (silt, sand)	Largely concealed beneath younger deposits, identified at depth from borehole data, rare exposures represented as Glaciolacustrine Deposits, Devensian (GLLDD) or Glaciolacustrine Deltaic Deposits, Devensian (GLDDD)
BHSE-XSV	Broomhouse Sand and Gravel Formation (sand and gravel)	Largely concealed beneath younger deposits, where present, exposures usually represented as Glaciofluvial Deposits, Devensian (GFDUD), but also as Glaciofluvial Ice-Contact Deposits, Devensian (GFICD)
BHSE-S	Broomhouse Sand and Gravel Formation (sand)	Not recorded on the maps in the UKGEOS Glasgow area (concealed beneath younger deposits), identified at depth from borehole data
WITI-DMTN	Wilderness Till Formation	Till - Devensian (TILLD)
CADR-XSV	Cadder Sand and Gravel Formation	Generally concealed beneath younger deposits, identified at depth from borehole data, rare exposures represented as Glaciofluvial Deposits, Devensian (GFDUD)
SUPD-XSV	Sand and gravel	Not recorded on the maps in the UKGEOS Glasgow area (concealed beneath younger deposits), identified at depth from borehole data

Table 2 The 15 lithostratigraphic units differentiated in the 3D superficial deposits model of the UKGEOS Glasgow area, in stratigraphic order.

Table 1 summarises the 15 lithostratigraphic units included in the superficial deposits model of the UKGEOS Glasgow area in stratigraphic order. Geological units are based on borehole and 1:10 000 scale geological map data., X indicates that each lithology is represented (e.g. XSV is a unit containing sand and gravel as opposed to SV which would be gravelly sand), where C=clay, Z=silt, S=sand, V=gravel, DMTN=diamicton, ARDP=artificial deposits, P=peat

A description of each of the modelled units is given below in approximately ascending (oldest/deepest first) order.

2.1 SAND AND GRAVEL UNIT (SUPD_XSV)

This small, un-named sand and gravel unit is thought to pre-date the Cadder Sand and Gravel Formation. The unit is generally thin (less than a few metres thick) and of limited lateral extent. It has been identified in a small number of boreholes in the northern part of the UKGEOS Glasgow area.

2.2 CADDER SAND AND GRAVEL FORMATION (CADR)

The Cadder Sand and Gravel Formation is found mainly to the north of the River Clyde (Figure 4), with deposits tens of metres thick associated with the deep bedrock depressions of the Kelvin buried valley-system. In the UKGEOS Glasgow area, the unit has largely been truncated by the emplacement of overlying tills and by modern river erosion, resulting in modification of the original depositional morphology of the unit. However a few boreholes in the north-eastern corner of the modelled area show sand and gravel deposits lying beneath glacial till up to 20 m thick.




Figure 4: Modelled distribution of Cadder Sand and Gravel Formation (orangey pink) and the Sand and Gravel unit (pink) across the UKGEOS Glasgow model area (red outline). 1 km grid squares shown. Includes mapping data licensed from Ordnance Survey. © Crown Copyright and/or database right 2019. Licence number 100021290 EUL.

2.3 WILDERNESS TILL FORMATION (WITI)

The Late Devensian Wilderness Till Formation is the most extensive unit in the Clyde area (Figure 5) and is named after temporary sections seen in the Wilderness Plantation area north of Bishopbriggs. It is characterised by a diamicton comprising isolated boulders, gravel and cobbles in a firm to stiff sandy, silty to clayey matrix (Browne and McMillan, 1989). Generally it rests directly on bedrock in the UKGEOS Glasgow area, but is underlain in places by older sand and gravel deposits (probably belonging to the Cadder Sand and Gravel Formation). Drumlins, which are large mounds formed during emplacement of the till below moving glacier ice, are a

characteristic landform associated with the Wilderness Till Formation. Drumlins constrain the hilly terrain of Central Glasgow and the surrounding areas where the Wilderness Till is exposed at the surface. Along the Clyde valley, buried drumlins cause considerable spatial variation in the thickness of overlying sediment deposits. The thickness of these glacial till deposits is highly variable across the UKGEOS Glasgow model area, reaching almost 50 m thick in some places whilst being absent in other areas.




Figure 5: Modelled distribution of Wilderness Till Formation (pale blue) across the UKGEOS Glasgow superficial geology model area (red outline). 1 km grid squares shown. Includes mapping data licensed from Ordnance Survey. © Crown Copyright and/or database right 2019. Licence number 100021290 EUL).

2.4 BROOMHOUSE SAND AND GRAVEL FORMATION (BHSE)

The Broomhouse Sand and Gravel is named after the Broomhouse area of eastern Glasgow where this unit generally overlies the Wilderness Till Formation. It comprises glaciofluvial ice-contact deposits, which produce features such as esker ridges, mounds, isolated flat-topped kames and kettleholes. Overall, the most abundant deposit is sand, except in esker ridges where gravel dominates. In some places this unit may also contain cobbles, clay and silt. The sands are planar and trough cross-bedded, ripple laminated and horizontally laminated; the gravels are typically massive or crudely bedded. Deposits are up to 25 m thick and flow directions were towards the east (Browne and MacMillan, 1989). The noted occurrences of Broomhouse Sand and Gravel Formation are coincident with areas of dense borehole data. This implies that there may be more extensive deposits in the area than have been modelled (Figure 6), but a lack of data means that further deposits remain undetected. Two components of the Broomhouse Sand and Gravel Formation have been modelled in this area: Broomhouse Sand and Gravel Member and the Ross Sand Member.




Figure 6: Modelled distribution of Broomhouse Sand and Gravel Formation (pink) across the UKGEOS Glasgow superficial geology model area (red outline). 1 km grid squares shown. Includes mapping data licensed from Ordnance Survey. © Crown Copyright and/or database right 2019. Licence number 100021290 EUL).

2.4.1 Ross Sand Member (RSSA)

The main lithologies of the Ross Sand Member, a member of the Broomhouse Sand and Gravel Formation, are sand or sand and silt, with clays at the base and thin local gravel layers. As noted above, the deposits are glacio-lacustrine in origin with deposits found in the south-east of Glasgow (Figure 7) interpreted to have formed in deltaic systems at the margins of glacial ‘Lake Clydesdale’ (Browne and MacMillan, 1989). For modelling purposes, the lithologies have been separated into two units, a main unit of sand with minor gravel (RSSA-XSV), and an underlying finer-grained sand with silt (RSSA-XSZ).

The Ross Sand Member is found draping the Wilderness Till Formation and is overlain by deposits of the Bridgeton Sand Member and Paisley Clay Member. Where it outcrops at the surface, the Ross Sand Member corresponds to glaciofluvial deltaic (and/or subaqueous fan) deposits (GFDD) on BGS DiGMapGB.

The Ross Sand Member has been mainly recognised at the eastern end of the UKGEOS Glasgow area, where it reaches up to 18 m thick in places.




Figure 7: Modelled distribution of Broomhouse Sand and Gravel Formation (orange) and Ross Sand Member (pale pink), across the UKGEOS Glasgow superficial geology model area (red outline). 1 km grid squares shown. Includes mapping data licensed from Ordnance Survey. © Crown Copyright and/or database right 2019. Licence number 100021290 EUL).

2.5 CLYDE CLAY FORMATION

The Clyde Clay Formation is part of the British Coastal Deposits Group and includes mainly Late Devensian deposits from marine isotope stages 2, 2a-b $\delta^{18}\text{O}$. Within the UKGEOS Glasgow area, the formation is modelled as three component members: the Bridgeton Sand Member, the Paisley Clay Member and the Killearn Sand and Gravel Member.

2.5.1 Bridgeton Sand Member (BRON)

The Bridgeton Sand Member is characterised by fine to medium, massive dense sand or silty sand. Locally, fine to coarse gravel and boulders occur in a sandy matrix. There is some flat bedding but generally the deposits are massive. The unit is largely confined to the Clyde valley, where it overlies the Wilderness Till and Broomhouse Sand and Gravel formations. The Bridgeton Sand Member is thought to be up to 32 m thick in some north-western parts of the UKGEOS Glasgow area.

Browne and MacMillan (1989) suggested that the sands were deposited as submarine outwash fans formed during catastrophic draining of pro-glacial Lake Clydesdale to the north-west, along the line of the Clyde valley, following breaching of the glacier dam.

2.5.2 Paisley Clay Member (PAIS)

The Paisley Clay Member typically comprises finely layered clay and silt-clay deposited in a glaciomarine setting. In borehole records it is often described as a grey and grey-brown, occasionally laminated, clayey silt and silty clay, often with a mottled appearance. The retreating glaciers are believed to have been to the north-west, in the sea lochs of the Southern Highlands. Relative sea level was high when deposition of the Paisley Clay Member commenced and some

clays were deposited at elevations up to 40 m above OD (Browne and McMillan, 1989). The Paisley Clay Member drapes the underlying topography, and appears to thin out over the tops of drumlins (Wilderness Till Formation) and bedrock ‘highs’. To the south-east of Glasgow, all clay units at the ground surface within lowland areas (max. 40 m to 45 m above OD) are assumed to be deposits of the Paisley Clay Member (Figure 8). All DiGMapGB polygons coded as raised marine deposits-Devensian (RMDV) have been interpreted as outcrops of the Paisley Clay Member. The Paisley Clay deposits within the UKGEOS Glasgow model area are thought to exceed 35 m thick in places.




Figure 8: Modelled distribution of Paisley Clay Member (green) across the UKGEOS Glasgow superficial geology model area (red outline). 1 km grid squares shown. Includes mapping data licensed from Ordnance Survey. © Crown Copyright and/or database right 2019. Licence number 100021290 EUL).

2.5.3 Killearn Sand and Gravel Member (KARN)

The Killearn Sand and Gravel Member is a patchy deposit comprising of varying proportions of sand and some clay layers. The deposits are commonly found in inter-drumlin areas along the Clyde Valley (Figure 9), and the distribution suggests that it formed as a result of marginal marine processes associated with the Late Devensian marine incursion responsible for the deposition of the Paisley and Linwood clays. The maximum level of the marine incursion thought to be responsible for the Killearn Member is approximately 34–36 m above OD (Rose, 1975; Browne and McMillan, 1989; Hall et al., 1998). Deposits are thought to reach almost 15 m thick in some eastern parts of the UKGEOS Glasgow model area.

The Killearn Sand and Gravel Member is thought to have been deposited in a range of environments, including beaches, and fluvial and deltaic systems. Rose (1975) describes temporary sections north of Erskine Bridge where Killearn Member sands and gravels overlie Broomhouse Sand and Gravel Formation. Here the Killearn Sand and Gravel Member is well bedded, with beds displaying shallow dip to the south-west. It is suggested that the sedimentary structures are consistent with shoreline processes, hence that the unit in this area is a beach

gravel. Rose (1975) also describes ice wedge casts suggesting that a periglacial environment persisted in the period following the deposition of the Killearn Sand and Gravel Member. It is suggested that these features may have formed during the cold period associated with the cooling event of the Younger Dryas. During this time, there is evidence glaciers readvanced in the Loch Lomond area, but the Clyde remained ice free.




Figure 9: Modelled distribution of Killearn Sand and Gravel Member (red) across the UKGEOS Glasgow superficial geology model area (red outline). 1 km grid squares shown. Includes mapping data licensed from Ordnance Survey. © Crown Copyright and/or database right 2019. Licence number 100021290 EUL).

2.6 CLYDEBANK CLAY FORMATION

2.6.1 Gourock Sand Member (GOSA)

The Gourock Sand Member forms extensive deposits in the Clyde valley, extending across the low ground either side of the River Clyde (Figure 10). The unit typically consists of fine to coarse sand with some gravel, silt and clay and organic detritus. The deposits are likely to have formed in estuarine environments, with a fluvial dominance in the east, becoming progressively more marine westwards with shallow channels linked by tidal flats (Browne and McMillan, 1989). The deposits have been modelled to be over 15 m thick in some central and north-western areas.




Figure 10: Modelled distribution of Gourrock Sand Member (pale yellow), Law Sand and Gravel Member (greenish yellow) and peat (brown), across the UKGEOS Glasgow superficial geology model area (red outline). 1 km grid squares shown. Includes mapping data licensed from Ordnance Survey. © Crown Copyright and/or database right 2019. Licence number 100021290 EUL).

2.7 CLYDE VALLEY FORMATION

2.7.1 Law Sand and Gravel Member (LAWSG)

The Law Sand and Gravel Member comprises fine to coarse sand with some silt, fine gravel and organic matter deposited in river channels and associated floodplains. The Law Sand and Gravel Member includes recent (currently accumulating) river deposits; in the UKGEOS Glasgow model area these deposits are mainly found associated with the White Cart Water in the south-western corner of the modelled area. Small alluvial fans and deposits associated with minor streams are also included in this unit.

2.8 PEAT

Small discrete deposits of peat, consisting of dark, organic rich, humic material formed by the accumulation of partially decomposed vegetation, are found in places in the UKGEOS Glasgow model area. Peat accumulations are generally associated with lacustrine and alluvial sediments, where they may form surface deposits, or form bands or lenses within the strata (discussed below).

2.9 MADE GROUND (MGR)

Made ground in the 3D model represents a combination of made and worked ground including filled and partially back-filled pits and quarries - hence it comprises all anthropogenic deposits.

Glasgow has a long and varied industrial past and these deposits are what remains of the former land uses and developments in the area (Figure 11).

Areas of made and worked ground were primarily identified using DiGMapGB 1:10 000 polygons. These were subsequently altered to encompass areas where boreholes reported additional areas of artificial ground (Figure 11). Alterations were made using the Ordnance Survey maps to identify the extent of industrial areas, housing developments and other information. It is likely that more extensive but thinner deposits of made ground occur within the model that are not currently represented. Artificial (made) ground has not been subdivided. Made ground is thought to be up to 27 m in places in the UKGEOS Glasgow model area and have a variable composition. Its distribution in the model is patchy (Figure 11) and is likely to be much more extensive than modelled.




Figure 11: Modelled distribution of made ground (MGR), across the UKGEOS Glasgow superficial geology model area (red outline). 1 km grid squares shown. Includes mapping data licensed from Ordnance Survey. © Crown Copyright and/or database right 2019. Licence number 100021290 EUL).

2.10 LENSES

Many of the units described above contain considerable variation in lithologies within their strata. Where thin deposits of a contrasting lithology to the main unit are recorded within borehole descriptions, these have been depicted by the modelling of a lens contained within the 'parent' unit. In general, only lithological sub-units with thicknesses greater than approximately 2 m have been identified as separate lenses.

The only lens modelled in the UKGEOS Glasgow superficial model area is a small, thin peat_lense; located in the south-western corner of the model.

3 Model Workflow

The superficial deposits model was constructed using GSI3D[®] software and the standard modelling workflow was followed (Kessler et al. 2009). The software utilises a range of data such as boreholes, digital terrain models (DTM) and geological linework to enable the geologist to construct a 3D model.

The workflow initially involves correlating the various geological units through strings of borehole logs in 2D cross-sections, the cross-sections intersect to form an interlocking network across the model area (Figures 12-17). Construction of the cross-sections is intuitive and flexible; combining borehole and outcrop data with the geologist's experience to refine the interpretation. The correlated sections and geological map data provide information on the lateral extent of each geological unit which is saved as an 'envelope' for the geological units.

Using both the information from the cross-sections and the geographical extent of each unit (envelope) a calculation algorithm creates the triangulated surfaces for the top and base of each unit. In order to control the relative vertical ordering of the calculation, a generalised vertical section file (.gvs) is established. A proprietary legend file (.gleg) is created to control symbolisation of the cross-section and model. The modeller can view all the units in 3D and iteratively return to the cross-section to make amendments or add further cross-sections to refine the model. This process is a standard methodology for modelling Quaternary and simple bedrock horizons and is fully documented in Kessler et al. (2009).




Figure 12 Screenshot of the GSI3D software showing the borehole, map and section window on the left-hand side and a view of the interlocking sections in the 3D window on the right-hand side.

The UKGEOS Glasgow superficial deposits model was updated in two phases:

Phase 1: Incorporated boreholes drilled since 2005 into the superficial model, in areas where it was felt the new data would significantly improve the resolution and accuracy to the model to better fit with available geological data (Figure 13).




Figure 13 Areas circled in red show where the UKGEOS model was updated/improved during phase 1, green lines indicate existing GSI3D section lines from the published Central Glasgow model, purple dots represent the location of boreholes entered into the BGS borehole database post-2005.

In detail, in Phase 1 model updating:

The approach taken was to focus on updating existing section line interpretations and adding new section lines in the areas with new borehole data (Figure 13). This improved the detail and certainty in parts of the model.

The post-2005 boreholes were used to update the model in the following ways:

- **Added into existing sections** – existing sections were re-routed to include additional boreholes where possible, this was the neatest approach if a section was already present nearby (Figures 14, 15).
- **Used to create a new section** – new sections were created or existing sections extended if there wasn't an existing section close to the additional boreholes. New sections were always extended to cross existing sections so the geological units could be tied-in with the existing network of sections.
- **Projected into existing sections** – where an existing section passed through a large number of additional boreholes spread along its length, the most appropriate approach was to project close boreholes into the existing section and adjust the geological linework as necessary. This approach quickly improved existing sections without the need to re-draw the section.
- **Informed the presence or absence of a unit across the model** – away from the lines of section the additional boreholes provide a constraint when drawing the geological unit 'envelopes' which show the lateral extent of a unit

Changes were also made to any crossing sections where necessary and edits were made to the associated geological envelopes, where required.




Figure 14 Example of a GSI3D section (Tester 1) which was significantly altered by including an additional 4 boreholes drilled since 2005. The upper image (original section) contained little borehole information to constrain the geology in this part of the section which has resulted in the construction of fairly simple geology. The lower image (updated section) indicates how additional borehole data suggests the presence of a further sand-body (pink), confirms the presence of till (blue) above rockhead across this part of the section, indicates that the Paisley Clay (green) has a patchy extent and provides improved constraint on the thickness and lateral extent of all the superficial deposits.




Figure 15 Example of a GSI3D section (NW_SE_5) which required little alteration when an additional 3 boreholes, drilled since 2005, were added: Upper image (original section) - little constraint in this part of the section – geology extrapolated from nearest borehole data and geological map at the surface. Lower image (updated section) – three additional boreholes confirm the superficial deposits present as predicted and provide improved constraint on the thickness of the deposits.

Phase 2 model updating:

Once provisional borehole sites for the UKGEOS Glasgow boreholes had been chosen (GGERFS01, 02, 03, 05 and 10), an additional 16 sections were completed passing through these points (Fig. 16) to illustrate what sequence of superficial deposits were expected at the sites.

Section lines were chosen to include as many additional boreholes in the vicinity of the UKGEOS proposed borehole sites to provide the highest resolution as possible in the site area. Sections were also placed to cross existing lines of section so the geology could be tied in with the existing network of sections.

The sections were created and geological linework drawn to increase the resolution of the model in the vicinity of the proposed sites to give our best indication of what lies beneath the site for boreholes prognoses.

Following the editing of sections and envelopes across the UKGEOS Glasgow area, the model was calculated, grids for each geological unit were exported (25m cell size) into an ArcGIS project along with information such as boreholes used, and lines of section. A rockhead grid was also calculated and exported for use in the bedrock model.




Figure 16 Summary map of proposed UKGEOS Glasgow borehole locations and sections constructed in GSI3D through the sites to indicate the nature and thickness of the superficial deposits beneath the surface.




Figure 17 Network of sections constructed in the vicinity of the UKGEOS Glasgow model area viewed in 3D




Figure 18 Overview of the calculated superficial deposits model across the UKGEOS Glasgow model area, looking north-east, 3x vertical exaggeration. NEXTMap Britain elevation data from Intermap Technologies.

The 3D superficial deposits model contains 15 superficial units (Figure 18) and is of use in borehole prognosis, site planning and can be used in hydrogeological conceptual models. For example, the calculated rockhead surface shows significant variability across the area from -40 m to +36 m relative to OD (Figure 19).




Figure 19 Rockhead surface for the study area, generated in GSI3D by combining the bases of all the modelled superficial units. Low values are in yellow and brown and high values are in purple and white. Includes mapping data licensed from Ordnance Survey. © Crown Copyright and/or database right 2019. Licence number 100021290 EUL).

4 Model Datasets

The model was constructed using a variety of datasets including NEXTMap® Digital Elevation Model ©Intermap Technologies, the BGS digital borehole database, BGS 1:10 000 scale digital maps (DiGMapGB, 2009), BGS 1:50 000 maps (DiGMapGB, 2008), field slip scans, historic maps and scanned geological cross-sections. Other literature such as BGS regional geological guides and scientific papers influenced the correlation of geological units.

Published BGS superficial model of Glasgow (2013) and basis for UKGEOS Glasgow model	CentralGlasgow_v3_7.gsipr
GVS	Glasgow_master_LEX_RCS_GVS_Jan2013v3.gvs
Legend	Glasgow_master_LEX_RCS_GLEG_Jan2013v3.gleg
BOREHOLE_INDEX	Porcupine_ESIOS_all.bid
BOREHOLE_LOGS	Porcupine_ESIOS_all.blg
DEM, NextMap	CentralGlasgow_dtm_50m
1:10 000 scale map sheets	NS56SE and NS66SW

Table 3 Summary of the key datasets and associated files used in the construction of the UKGEOS Glasgow superficial deposits model.

4.1 MAP DATA

BGS published maps of the superficial deposits over the UKGEOS Glasgow model area includes map sheets NS66SW (2007) and NS56SE (2007) at 1:10 000 scale, and map sheets 30E Glasgow (1994) and 31W Airdrie (1992) at 1:50 000 scale (Figure 20). The associated BGS Memoirs form a definitive reference source for the geology of this area (Hall et al., 1998, Forsyth et al., 1996). Map data is available as GIS shapefiles (DigMapGB) or as scans ([Mapviewer](#)).



Figure 20 1:50 000 scale (green) and 1:10 000 scale (blue) geological map sheets covering the UKGEOS Glasgow model area (pink). Includes mapping data licensed from Ordnance Survey. © Crown Copyright and/or database right 2019. Licence number 100021290 EUL).

The UKGEOS Glasgow model broadly matches the corresponding 1:10 000 scale digital geological map data of the area (Figure 21), some differences are present as a result of incorporating new (post-2005) borehole data which post-date the map construction. The map shows an extensive area of alluvium (river deposits) through the centre, representing the floodplain of the River Clyde. A variety of raised marine, lacustrine and glaciofluvial deposits have been mapped between areas of alluvium and the glacial till that is only seen at the surface over higher ground. Bedrock is mapped 'at or close' to the surface in a few areas where superficial deposits are absent. Bedrock units and faults are represented in separate maps and models (see Burkin and Kearsey, 2018).




Figure 21 UKGEOS Glasgow model area summary of superficial geology from BGS©UKRI 2016 DigMap 1:10 000 scale data, the Clyde Gateway urban regeneration area is shown in blue.

4.2 GVS AND GLEG FILES

The GSI3D geological vertical sequence (.gvs) file contains all the geological units in their correct and unique super-positional (stratigraphical) order. The geological legend (.gleg) file assigns colours and textures to the different geological units. The files were assembled in a combination of Notepad, Wordpad and Excel and iterated as the model expanded and new units were encountered. The GVS was created using DiGMapGB-50 data. The .gleg file was created by using data from the BGS 1:10 000 and 1:50 000 DiGMapGB datasets.

4.3 DIGITAL ELEVATION MODEL

The UKGEOS Glasgow superficial deposits model is capped by an extract of the NEXTMap® Digital Elevation Model from ©Intermap Technologies. The maximum horizontal NEXTMap® resolution of 5 m was subsampled to 50 m for the previous ‘Central Glasgow’ model to reduce the file size and allow model calculation. The same extract was used for the UKGEOS Glasgow model to avoid any ‘re-snapping’ of the existing geological linework to a new surface in the cross-sections. The elevation range of the ground surface in the UKGEOS Glasgow area is +2 m to +59 m above Ordnance Datum (OD). Figure 22 clearly shows the sinuous course of the River Clyde along the low ground of the Clyde valley (blue colours) with higher ground (yellow-red colours) to the northeast and southwest.




Figure 22 Digital Terrain Model (50 m cell size) across the UKGEOS Glasgow superficial deposits model area (red outline), as used in the model. NEXTMap Britain elevation data from Intermap Technologies. Includes mapping data licensed from Ordnance Survey. © Crown Copyright and/or database right 2019. Licence number 100021290 EUL.

4.4 BOREHOLE DATA

Boreholes, along with geological map data were the primary data sources for the construction of cross-sections across the UKGEOS Glasgow superficial deposits model area. BGS holds approximately 10 000 borehole records (including site investigations and trial pits) across the area (Figure 23). Although the majority of these (approximately three-quarters) penetrate to depths of less than 30 metres, they define the complex artificial and superficial deposit succession.

Borehole records are held in the BGS *Single Onshore Borehole Index* (SOBI) database and downhole geological information is held in the BGS *Borehole Geology* (BoGe) database. All borehole records are entered into the SOBI database at acquisition, however geological information is entered into BoGe as required. The bulk of geological information was entered for the construction of the Central Glasgow and earlier models, however additional geological information for new borehole data received post-2005 was entered for the UKGEOS Glasgow work. Boreholes were coded with bedrock units from the BGS Lexicon <http://www.bgs.ac.uk/lexicon/> and the Rock Classification Scheme (RCS) <http://www.bgs.ac.uk/bgsrscs/> as appropriate to log lithology

For the model updates described as ‘phase 1’ in section 3 (above), all borehole data was recalled from the database for the UKGEOS Glasgow superficial deposits model area, this was then filtered to only include boreholes entered/drilled in 2005 or later. Where multiple interpretations existed for a single borehole, priority was given to AAMI, then ECAL, then SJSE.

Generally, the boreholes selected for use in the GSI3D cross-sections were those which intersected geological rockhead and therefore recorded the full sequence of superficial deposits and/or those which gave detailed descriptions of the geological units. The spread of borehole data was variable (Figure 23) from extremely closely-spaced at site investigation locations to

more widely spaced and isolated boreholes. Therefore the borehole location was also of consideration during selection for inclusion in cross-sections, to ensure an even spatial distribution of borehole data as far as possible. As the UKGEOS Glasgow superficial deposits model is based on the earlier Central Glasgow model which covers a much larger area, cross-sections extend beyond the boundaries of the model area. This reduces any edge effects.

Recorded borehole start heights are often different to the ground level shown by the digital elevation model (DEM). As the boreholes range in age to over 100 years old, it was generally assumed that the differences were either due to ground level changes as a result of landscaping and development, or due to the relatively coarse 50 m resolution of the DEM. Recorded borehole start heights were used in preference, as opposed to ‘hanging’ the boreholes on the DEM.




Figure 23 Location of borehole records held in the BGS Single Onshore Borehole Index database. Red dots (more than 30 m deep), blue dots (between 10 m and 30 m), green dots (less than 10 m) and black dots (depth unknown). Includes mapping data licensed from Ordnance Survey. © Crown Copyright and/or database right 2019. Licence number 100021290 EUL.

5 Model Assumptions and Limitations

Some key assumptions and rules used in the modelling process include:

- Where no boreholes were present, basal units were modelled to the BGS rockhead model (see <http://www.bgs.ac.uk/products/onshore/superficialThickness.html>). Where the depth of rockhead in boreholes differed from the rockhead model, the borehole depth was taken in preference.

- The depth of water within the tidal limit of the River Clyde was based approximately on a UK Hydrographic Office Admiralty Chart (International Chart Series c.2000). Where no information is available a sub-rectangular river channel cross-section has been assumed.
- Water body extents were extracted from the OS 1:10 000 scale Open data digital map. The River Clyde and lakes/reservoirs exceeding approximately 150 m in width/length are included in the 3D model. During modelling, the water bodies were generally fitted to the mapped water extent and not to the simplified DEM. In places this leads to artefacts where the water surface does not appear to be horizontal.

5.1 GENERAL MODELLING LIMITATIONS

- Geological interpretations are made according to the prevailing understanding of the geology at the time. The quality of such interpretations may be affected by the availability of new data, by subsequent advances in geological knowledge, improved methods of interpretation, improved databases and modelling software, and better access to sampling locations. The 3D geological model represents an individual interpretation of the data available; other interpretations may be valid.
- This model builds upon the earlier models described in Merritt et al. (2009), Monaghan et al. (2012, 2014) and Arkley et al. (2013). Over this period of time, a number of different geologists have constructed and updated models. Although efforts have been made to retain consistency of methodologies, some geologists may have interpreted the borehole data in different ways.
- Best endeavours (detailed quality checking procedures) are employed to minimise data entry errors but given the diversity and volume of data used, it is anticipated that occasional erroneous entries will still be present (e.g. boreholes locations, elevations etc.) Any raw data considered when building geological models may have been transcribed from analogue to digital format. Such processes are subjected to quality control to ensure reliability; however undetected errors may exist.
- The model does not reflect the full complexity of the superficial deposits geology. In reality, surfaces have been subjected to more glaciectonic deformation than is represented in the model. It is also known that made, worked and artificial ground is more widespread than is shown by the model, and could be subdivided into more detail than the ‘made ground’ currently used.
- Smaller rivers, streams and water bodies have not been included in the model.
- The cross-section density and therefore model certainty is variable across the model, and is based on complexity and type of geology, borehole density etc.

5.2 DEM

- The Digital Elevation Model (DEM) was subsampled from a 5 m resolution to 50 m resolution which means the surface distribution and geometry of a geological unit does not reflect the highest resolution possible. This resolution was chosen based on the size of the area, the resolution of the original modelling and the software capability. Some minor mismatches between geomorphological features and modelled units (including water) and the DEM occur due to the coarse resolution of the DEM.
- The DEM may contain artefacts such as trees or artificial structures such as pylons. The majority of these have been stripped out before modelling. If any of these artefacts were found during the modelling then the effects of these were minimised in the model as much as possible.

5.3 BOREHOLE DATA

- The spatial distribution of available borehole data is highly variable with concentrations and gaps. For example there are concentrations along infrastructure routes such as the M74 and

the site of the Commonwealth Games, and scattered areas of very sparse data, particularly in the southern part of the model around Rutherglen.

- The precise relationships between the Quaternary deposits are complex, varying laterally and vertically across short distances, and in areas where borehole data are sparse or absent extrapolation is difficult.
- A subset of the most reliable borehole data has been included to constrain the cross-sections within the model. However there is also a large subset of borehole data that has not been included within the model. Some of the boreholes are more than 100 years old and although often deep, rockhead and location details are often vague, start heights are not recorded and descriptions of the superficial deposits are brief.

Factors affecting the usefulness of the borehole include drill depth, the amount of detail described in the log and whether they have been logged to industry standards. The drilled depth is a limitation because not all of the boreholes used in the model reach rockhead. Where clusters of boreholes occur only the deepest boreholes with the most geological information were selected for use in the model. Out of the 10,000 boreholes in the study area, approximately 4500 boreholes have been included in the network of sections with additional boreholes being projected into the sections to inform the geological linework.

- Borehole start heights are obtained from the original records, Ordnance Survey mapping or a digital terrain model. Where borehole start heights look unreasonable, they are checked and amended if necessary in the index file. In some cases, the borehole start height may be different from the ground surface, if for example, the ground surface has been raised or lowered since the borehole was drilled, or if the borehole was not originally drilled at the ground surface.

5.4 MAP DATA

- The geological map linework in the model has been modified during the modelling process to modify the interpretation where new data is available. The most common mapped areas requiring revision relate to worked/made ground activity post-mapping. This model supersedes the published BGS 1: 10 000 scale maps of the Glasgow area. Updates in line with advances made in geological understanding of the area during 3D modelling are yet to be made to the maps

5.5 MODELLED SURFACES AND VOLUMES

- The thin nature of made ground, and the thin draped form of some areas e.g. deposits of the Killearn Sand and Gravel Member, means that these units are poorly shown in visualisations of the 3D model.
- A known limitation is that for some thin units close to DEM surface and over topographically variable ground the superficial deposits output grids contain small patches of no modelled surface within the unit extent.
- The modelled volumes representing some elongate units such as water, made ground along road or rail embankments and alluvium are in places spiky/angular due to a combination of steep edges, DEM resolution and limited constraining cross-sections. The size of the angularity is in proportion to the unit and is accepted as a known limitation.

6 Model Quality Assurance

In order for a geological model to be approved for publication or delivery to a client a series of quality assurance checks is carried out. This includes visual examination of the modelled cross-sections to ensure that they match each other at cross-section intersections and fit the borehole and geological map data used. The model calculation is checked to ensure that all units calculate to their full extent within the area of interest and the modelled geological surfaces are checked for artefacts such as spikes and thickness anomalies. The naming convention of the modelled

geological units is checked to ensure that recognised entries in the BGS Lexicon of Named Rock Units (<http://www.bgs.ac.uk/lexicon/home.html>) and the BGS Rock Classification Scheme (<http://www.bgs.ac.uk/bgsrscs/>) are used as far as possible.

Any issues found in the QA checking process are recorded and addressed before delivery/publication of the model.

7 Model Uncertainty

Quantitative model uncertainty studies have not been undertaken. However the cross-sections and borehole data points constraining the UKGEOS Glasgow superficial deposits model shown in Figures 24 and 25 give an indication of the most certain areas of the model, those with the most cross-sections and data points (e.g. around Cuningar Loop), and the least certain areas (e.g. around Rutherglen). These maps do not qualify increasing uncertainty with depth below the surface, noting that more than half of the boreholes in the model area reached less than 10 m below the surface.




Figure 24 Distribution of boreholes used to construct the correlated cross-sections which constrain the superficial deposits model of the UKGEOS Glasgow area.




Figure 25 Distribution of boreholes used to construct the correlated cross-sections which constrain the superficial deposits model around the proposed borehole sites for the UKGEOS Glasgow project. The River Clyde is shown in pale blue.

8 Model exports

Modelled tops, bases, thicknesses and extents have been exported as (Table 4). Files are named with the Lexicon and lithological description from the GVS file (e.g. WITI_DMTN_BASE). Z values are in metres relative to Ordnance Datum (positive upwards).

File format	Description
ASCII grids .asc	Top, base and thickness (shell) of each modelled unit
ESRI grids	Top, base and thickness (shell) of each modelled unit
Gocad .ts surface	Top, base and thickness (shell) of each modelled unit
ASCII.asc and ERSI grid format	Rockhead grid calculated in GSI3D
ESRI .shp	Extent of geological units and cross-section locations

Table 4 Summary table of model exports

9 Model Images

9.1 CROSS-SECTION IMAGES




Figure 26 Cross-section *GGERFS_slba_3*, orientated SSW-NNE through the superficial deposits of the UKGEOS Glasgow model area (see Figure 16 for location). Vertical exaggeration x 3. Borehole constraint points shown in red. Ground surface derived from NEXTMap Britain elevation data from Intermap Technologies. Height is in metres relative to Ordnance Datum.




Figure 27 Cross-section *GGERFS_slba_4*, orientated NW-SE through the superficial deposits of the UKGEOS Glasgow model area (see Figure 16 for location). Vertical exaggeration x 3. Borehole constraint points shown in red. Ground surface derived from NEXTMap Britain elevation data from Intermap Technologies. Height is in metres relative to Ordnance Datum.




Figure 28 Cross-section *GGERFS_slba_5*, orientated South-North through the superficial deposits of the UKGEOS Glasgow model area (see Figure 16 for location). Vertical exaggeration x 3. Borehole constraint points shown in red. Ground surface derived from NEXTMap Britain elevation data from Intermap Technologies. Height is in metres relative to Ordnance Datum.




Figure 29 Cross-section *GGERFS_slba_7*, orientated NW-SE through the superficial deposits of the UKGEOS Glasgow model area (see Figure 16 for location). Vertical exaggeration x 3. Borehole constraint points shown in red. Ground surface derived from NEXTMap Britain elevation data from Intermap Technologies. Height is in metres relative to Ordnance Datum.




Figure 30 Cross-section *GGERFS_slba_8*, orientated North-South through the superficial deposits of the UKGEOS Glasgow model area (see Figure 16 for location). Vertical exaggeration x 3. Borehole constraint points shown in red. Ground surface derived from NEXTMap Britain elevation data from Intermap Technologies. Height is in metres relative to Ordnance Datum.




Figure 31 Cross-section *GGERFS_slba_10*, orientated West-East through the superficial deposits of the UKGEOS Glasgow model area (see Figure 16 for location). Vertical exaggeration x 3. Borehole constraint points shown in red. Ground surface derived from NEXTMap Britain elevation data from Intermap Technologies. Height is in metres relative to Ordnance Datum.




Figure 32 Cross-sections *GGERFS_slba_8* and *GGERFS_slba_10* as viewed in the 3D window of GSI3D. These two sections pass through the proposed site for borehole GGERFS10 at Dalmarnock (see Figure 16 for location and orientation of sections). The sections illustrate the calculated sequence and thickness of superficial deposits beneath this part of the UKGEOS Glasgow model area from available borehole and map data. Vertical exaggeration x 10. Ground surface derived from NEXTMap Britain elevation data from Intermap Technologies. Borehole constraint points along the sections and thickness of the geological units are indicated in Figures 24 and 25.




Figure 33 Nine cross-sections as viewed in the 3D window of GSI3D. This network of sections pass through the proposed sites for the Cuningar Loop boreholes (see Figure 16 for location and orientation of sections). The sections illustrate the calculated sequence and thickness of superficial deposits beneath this part of the UKGEOS Glasgow model area from available borehole and map data. Vertical exaggeration x 10. Ground surface derived from NEXTMap Britain elevation data from Intermap Technologies. Borehole constraint points along some of the sections and thickness of the geological units are indicated in Figures 24 and 25.

9.2 THICKNESS MAPS




Figure 34 Modelled distribution and thickness of Made Ground deposits (MGR) across the UKGEOS Glasgow superficial geology model area (red outline). 1 km grid squares shown. Includes mapping data licensed from Ordnance Survey. © Crown Copyright and/or database right 2019. Licence number 100021290 EUL.

Figure 34 highlights the variable made ground thickness. The thickest deposits are shown in red and reach over 20 m thick in places. These are a relic of the area's industrial past and they often represent formerly worked areas such as the infilled sand and gravel pits at Cuningar Loop, areas of former heavy industry such as the Clydebridge Steel Works at the east side of the study area or sites of significant infrastructure such as that at the former Polmadie Mineral Depot at the west end of the study area. Additional made ground, associated with former land use and development is likely to be present across most of the modelled area, mantling the natural superficial deposits.




Figure 35 Modelled distribution and thickness of the Gourrock Sand Member (GOSA) across the UKGEOS Glasgow superficial geology model area (red outline). These marine sands are extensive within the Clyde Valley and are over 15 m thick in places. 1 km grid squares shown. Includes mapping data licensed from Ordnance Survey. © Crown Copyright and/or database right 2019. Licence number 100021290 EUL.




Figure 36 Modelled distribution and thickness of the Paisley Clay Member (PAIS) across the UKGEOS Glasgow superficial geology model area (red outline). These laminated marine clays and silts are extensive within the Clyde Valley and are over 30 m thick in places. 1 km grid squares shown. Includes mapping data licensed from Ordnance Survey. © Crown Copyright and/or database right 2019. Licence number 100021290 EUL.




Figure 37 Modelled distribution and thickness of the Bridgeton Sand Member (BRON) across the UKGEOS Glasgow superficial geology model area (red outline). This deposit of marine sands is significant in the north-western corner of the modelled area where it is over 30 m thick in places. 1 km grid squares shown. Includes mapping data licensed from Ordnance Survey. © Crown Copyright and/or database right 2019. Licence number 100021290 EUL.




Figure 38 Modelled distribution and thickness of the Broomhouse Sand and Gravel Formation (BHSE) across the UKGEOS Glasgow superficial geology model area (red outline). This deposit of fluvioglacial sand and gravels generally occurs at a significant depth beneath the surface, as its known occurrence (shown above) is dependent on the presence of deep borehole records more extensive deposits may lie undetected in areas where there are a lack of data points. The unit is over 20 m thick in places. 1 km grid squares shown. Includes mapping data licensed from Ordnance Survey. © Crown Copyright and/or database right 2019. Licence number 100021290 EUL.




Figure 39 Modelled distribution and thickness of the Wilderness Till Formation (WITI) across the UKGEOS Glasgow superficial geology model area (red outline). This deposit of glacial till is widespread across the modelled area and beyond. It is thought to be almost 50 m thick in till-cored drumlins shaped by the over-riding Late Devensian glaciers. 1 km grid squares shown. Includes mapping data licensed from Ordnance Survey. © Crown Copyright and/or database right 2019. Licence number 100021290 EUL.

10 References

British Geological Survey holds most of the references listed below, and copies may be obtained via the library service subject to copyright legislation (contact libuser@bgs.ac.uk for details). The library catalogue is available at: <https://envirolib.apps.nerc.ac.uk/olibcgi>

- ARKLEY, S, WHITBREAD, K, MONAGHAN, A. 2013. Clyde superficial deposits and bedrock models released to the ASK Network 2013: a guide for users. *British Geological Survey Open Report*, OR/13/002. 29pp.
- BRITISH GEOLOGICAL SURVEY. 2007. Glasgow, SE. Sheet NS56SE. Superficial Deposits. 1:10 000 scale. [Keyworth, Nottingham: British Geological Survey].
- BRITISH GEOLOGICAL SURVEY. 2007. Rutherglen. Sheet NS66SW. Superficial Deposits. 1:10 000 scale. [Keyworth, Nottingham: British Geological Survey].
- BRITISH GEOLOGICAL SURVEY. 1994. Glasgow, Scotland Sheet 30E. Drift Geology. 1:50 000 scale. [Keyworth, Nottingham: British Geological Survey].
- BRITISH GEOLOGICAL SURVEY. 1992. Airdrie. Scotland Sheet 31W. Drift Geology. 1:50 000 scale. [Keyworth, Nottingham: British Geological Survey].
- BRITISH GEOLOGICAL SURVEY. 2008. *Digital Geological Map of Great Britain 1:50 000 scale (DiGMapGB-50) data. Version 5.18*. [Keyworth, Nottingham: British Geological Survey]. Release date 20-05-2008.
- BRITISH GEOLOGICAL SURVEY [DIGMAPGB]. 2009. Digital Geological Map of Great Britain 1:10 000 scale (DiGMapGB-10) data. Version 2.18. [Keyworth, Nottingham: British Geological Survey]. Release date 15-01-2009.
- BROWNE, M A E AND McMILLAN, A A. 1989. Quaternary geology of the Clyde valley. *Research Report of the British Geological Survey* SA/89/1.
- BURKIN, J N AND KEARSEY, T. 2018. Model Metadata report for Glasgow Geothermal Energy Research Field Site Bedrock Model. *British Geological Survey Open Report*, OR/10/053. 17pp
- FINLAYSON, A, MERRITT, J, BROWNE, M, MERRITT, J, McMILLAN, A, WHITBREAD, K. 2010. Ice sheet advance, dynamics, and decay configurations: evidence from west central Scotland. *Quaternary Science Reviews*, 29 (7–8), 969–988.
- FINLAYSON, A. 2012. Ice dynamics and sediment movement: last glacial cycle, Clyde basin, Scotland. *Journal of Glaciology*, 58, 487–500.
- FORSYTH, I H, HALL, I H S, McMILLAN, A A, ARTHUR, M J, BRAND, P J, GRAHAM, D K, and ROBINS, N S. 1996. Geology of the Airdrie district: *Memoir of the British Geological Survey*, Sheet 31W (Scotland). ISBN 011884508X
- HALL, I H S, BROWNE, M A E AND FORSYTH, I H. 1998. Geology of the Glasgow district. *Memoir of the British Geological Survey*, Sheet 30E (Scotland). ISBN 0118845349.
- KESSLER, H, MATHERS, S J, AND SOBISCH, H-G. 2009. GSI3D – The capture and dissemination of integrated 3D geospatial knowledge at the British Geological Survey using GSI3D software and methodology. *Computers and Geosciences*, Vol 35, Issue 6, pp 1311–1321.
- MERRITT, J E, MONAGHAN, A A, LOUGHLIN, S C, MANSOUR, M, Ó DOCHARTAIGH, B É, AND HUGHES, A G. 2009. Clyde Gateway Pilot 3D Geological and Groundwater Model. *British Geological Survey, Commissioned Report*, CR/09/005N. <http://nora.nerc.ac.uk/id/eprint/14773/>
- MONAGHAN, A A, TERRINGTON, R L, AND MERRITT, J E. 2012. Clyde Gateway Pilot 3D Geological Model, Version 2. *British Geological Survey Commissioned Report*, CR/12/010N. 10pp <http://nora.nerc.ac.uk/id/eprint/21118/>
- MONAGHAN, A A, ARKLEY, S L B, WHITBREAD, K AND McCORMAC, M. 2014. Clyde superficial deposits and bedrock models released to the ASK Network 2014: a guide for users Version 3. *British Geological Survey Open Report*, OR/14/013. 31pp. <http://nora.nerc.ac.uk/id/eprint/505554/>
- MONAGHAN, A A, STARCHER, V, O DOCHARTAIGH, B, SHORTER, K AND BURKIN, J N. 2018. UK Geoenergy Observatories: Glasgow Geothermal Energy Research Field Site: science infrastructure. Nottingham, UK, British Geological Survey, 46pp. (OR/18/037) (Unpublished) <http://nora.nerc.ac.uk/id/eprint/521444/>
- ROSE, J. 1975. Raised beach gravels and ice wedge casts at Old Kilpatrick, near Glasgow. *Scottish Journal of Geology*. Vol 11. p 15–21.