



**Effect of heavy metals and other xenobiotics on biodegradation of waste canola oil by cold-adapted *Rhodococcus* sp. strain AQ5-07**

**Efecto de metales pesados y otros xenobioticos en la biodegradación de residuos de aceite de canola mediante la cepa adaptada al frío *Rhodococcus* sp. AQ5-07**

S. Ibrahim<sup>1,2</sup>, A. Zulkharnain<sup>3</sup>, K.N.M. Zahri<sup>1</sup>, G.L.Y. Lee<sup>1</sup>, P. Convey<sup>4</sup>, C. Gomez-Fuentes<sup>5</sup>, S. Sabri<sup>6</sup>, K.A. Khalil<sup>7</sup>, S.A. Alias<sup>8</sup>, G. Gonzalez-Rocha<sup>9</sup> and S.A. Ahmad<sup>1,8\*</sup>

<sup>1</sup>Department of Biochemistry, Faculty of Biotechnology and Biomolecular Sciences, Universiti Putra Malaysia, 43400 UPM Serdang, Selangor, Malaysia.

<sup>2</sup>Centre for Biotechnology Research, Bayero University PMB 3011 Kano, Nigeria.

<sup>3</sup>Department of Bioscience and Engineering, College of Systems Engineering and Science, Shibaura Institute Technology, 307 Fukasaku, Minuma-ku, Saitama, 337-8570, Japan.

<sup>4</sup>British Antarctic Survey, NERC, High Cross, Madingley Road, Cambridge CB3 0ET, United Kingdom.

<sup>5</sup>Department of Chemical Engineering, Universidad de Magallanes, Avda. Bulnes 01855, Punta Arenas, Región de Magallanes y Antártica Chilena, Chile.

<sup>6</sup>Department of Microbiology, Faculty of Biotechnology and Biomolecular Sciences, Universiti Putra Malaysia, 43400 UPM Serdang, Selangor, Malaysia.

<sup>7</sup>Department of Biomolecular Sciences, Faculty of Applied Sciences, Universiti Teknologi MARA, 40450, Shah Alam, Selangor, Malaysia.

<sup>8</sup>National Antarctic Research Centre, B303 Level 3, Block B, IPS Building, Universiti Malaya, 50603 Kuala Lumpur, Malaysia.

<sup>9</sup>Laboratorio de Investigacion en Agentes Antibacterianos, Facultad de Ciencias Biologicas, Universidad de Concepcion, Concepcion, Chile.

Received: October 9, 2019; Accepted: November 19, 2019

**Abstract**

The Antarctic is generally considered to be one of the most pristine areas in the world. However, both long and short-range pollutants are now known to be present in the Antarctic environment. Canola oil is an example of a polluting hydrocarbon that can be accidentally released into the Antarctic environment in oil wastewater treatment plants. The Antarctic soil bacterial strain *Rhodococcus* sp. AQ5-07, known to be capable of using waste canola oil (WCO) as its sole source of carbon, was tested for its ability to degrade canola oil in the presence of different heavy metals and xenobiotics. *Rhodococcus* sp. AQ5-07 was grown on minimum salt media containing different heavy metals (Zn, Co, Ni, Ag, Pb, Cu, Cr, Hg, Cd and As), xenobiotics (acrylamide and phenol) supplemented with 3% WCO. Three out of the 10 heavy metals tested (Hg, Cd and Ag) led a significant reduction in canola oil degradation at a concentration of 1 ppm. The IC<sub>50</sub> values of Hg, Cd and Ag were 0.38, 0.45 and 0.32 ppm, respectively. The strain could also withstand 10 mg/L acrylamide, 50 mg/L phenol and 0.5% (v/v) diesel. This study confirmed the ability of *Rhodococcus* sp. AQ5-07 to degrade canola oil in the presence of various heavy metals and other xenobiotics, supporting its potential use in bioremediation of vegetable oil and wastewater treatments in low temperature environments.

**Keywords:** Arctica, biodegradation, canola oil, heavy metals, *Rhodococcus*, xenobiotics.

**Resumen**

La Antártida se considera generalmente una de las áreas más prístinas del planeta. Sin embargo, es sabido que contaminantes de largo y corto alcance están presentes en el medio ambiente antártico. El aceite de canola es un ejemplo de un hidrocarburo contaminante que puede ser liberado accidentalmente al medio ambiente antártico por las plantas de tratamiento de aguas residuales. La cepa bacteriana de suelo antártico *Rhodococcus* sp. AQ5-07, de la cual se sabe que es capaz de usar aceite de canola de desecho (ACD) como única fuente de carbono, fue testeada en su capacidad para degradar el aceite de canola en presencia de diferentes metales pesados y xenobióticos. *Rhodococcus* sp. AQ5-07 se cultivó en medio de cultivo mínimo de sales conteniendo diferentes metales pesados (Zn, Co, Ni, Ag, Pb, Cu, Cr, Hg, Cd y As), xenobióticos (acrilamida y fenol) suplementado con 3% de ACD. Tres de los 10 metales pesados probados (Hg, Cd y Ag) produjeron una reducción significativa en la degradación del aceite de canola a una concentración de 1 ppm. Los valores de CI<sub>50</sub> de Hg, Cd y Ag fueron 0,38, 0,45 y 0,32 ppm, respectivamente. La cepa también pudo soportar 10 mg/L de acrilamida, 50 mg/L de fenol y 0.5% (v/v) de diesel. Este estudio confirmó la capacidad de *Rhodococcus* sp. AQ5-07 para degradar el aceite de canola en presencia de varios metales pesados y otros xenobióticos, apoyando su potencial uso en la biorremediación de aceites vegetales y tratamientos de aguas residuales en ambientes de baja temperatura.

**Palabras clave:** Antártica, biodegradación, aceite de canola, metales pesados, *Rhodococcus*, xenobióticos.

\* Corresponding author. E-mail: aqlima@upm.edu.my

<https://doi.org/10.24275/rmiq/Bio917>

issn-e: 2395-8472

## 1 Introduction

---

Antarctica is a unique region globally, and a key natural laboratory devoted to scientific research. It is often assumed to be one of the most pristine regions in the world (Xie and Sun, 2008), but it is not sealed from the rest of the world and is reached by global contaminants (Bargagli, 2005). Since the early 20th century, coastal regions have suffered the direct impacts of human presence (Vodopivec *et al.*, 2015). Anthropogenic pollution can reach Antarctica areas through atmospheric circulation and deposition. Over the last 200 years, human activities in Antarctica have changed focus from initial exploration towards scientific research and, more recently, tourism. Since the International Geophysical Year of 1957/1958, scientific research programmes have developed to a level where now approximately 5000 researchers and support staff work in Antarctica annually (Tiwari, 2017; COMNAP, 2012). A further surge in activity took place with the International Polar Year of 2007/2008 (Convey *et al.*, 2012). At the same time, by the 2015/16 austral summer, tourist numbers increased to almost 40,000 (IAATO, 2017), a trend that is expected to continue. Today, nationals of more than 30 countries are actively involved in scientific research on the continent (Williams *et al.*, 2017; Rack, 2015). The increasing levels of human activity in the region inevitably are associated with risks of anthropogenic pollution events that have become a significant threat to the natural Antarctic environment.

Heavy metals occur naturally in the Earth's crust. They can represent a serious environmental threat due to accumulation, as they cannot be broken down like other contaminant types (Buendía-González *et al.*, 2019; Corral-Escárceg *et al.*, 2017; Chu *et al.*, 2019). Elements such as copper, lead and arsenic are often present in limited quantities but the threat also extends to other elements, metalloids and metallic elements such as Sb, Tl, Mo, Rh, Ir, Pt, U, Bi, Ba, Cd, Ag, Co, Zn, Mn, Cr and V (Hong *et al.*, 2012; Soyol-Erdene *et al.*, 2011). The presence of heavy metals in contaminated areas can therefore be a major challenge for bioremediation of other pollutants such as oils and other hydrocarbons (Alcázar-Medina *et al.*, 2020; Hernández-Martínez *et al.*, 2019; Interiano-López *et al.*, 2019; Suárez-García *et al.*, 2019; Subramaniam *et al.*, 2019; Karamba *et al.*, 2016; Ruiz-Marín *et al.*, 2013).

Remediation of lipid-contaminated areas through biodegradation provides an important example of recovery from pollution. Canola oil, a vegetable oil, is a major cooking oil widely used in food preparation both industrially and at domestic scale. Fats, fatty acids (Fas), oil, and grease are released into the natural environment together with industrial and municipal wastewater (Čipinyte *et al.*, 2009; Ibrahim *et al.*, 2018). These polluting compounds can lead to serious anthropogenic problems such as odour, pipe and filter blockage, oil film formation and oxygen depletion, with deleterious effects on aquatic organisms and ecosystems (Fadile *et al.*, 2011). Metabolism of lipids involves emulsification and degradation. Free fatty acid(s) and glycerol are generated during the degradation process. The free fatty acid(s) are converted to acetyl-CoA via the beta-oxidation pathway and finally citric acid cycle (Phong *et al.*, 2014), while the glycerol generated is converted to glucose via a series of chemical reactions.

Many aerobic and anaerobic microorganisms capable of degrading lipids have been isolated from water and soil samples (Ibrahim *et al.*, 2018). These are commonly bacterial and fungal strains from the genera *Rhodococcus*, *Staphylococcus*, *Arthrobacter*, *Pseudomonas*, *Enterobacter*, *Bacillus*, *Penicillium*, *Serratia*, *Aspergillus*, *Mucor*, *Burkholderia*, *Acinetobacteria* and *Lactobacillus* (Danikuu and Sowley, 2014; Nagarajan *et al.*, 2014; Phong *et al.*, 2014; Bharathi *et al.*, 2012; Kumar *et al.*, 2012; Čipinyte *et al.*, 2009; Matsumiya *et al.*, 2007). These organisms have the capacity to release extracellular lipases that are responsible for vegetable oil degradation (Serikovna *et al.*, 2013; Pereira *et al.*, 2003).

Considerable attention has been given in microbial biotechnology to the development of enzymatic lipases (Dors *et al.*, 2013). Several studies have shown that lipolytic aerobic treatment techniques do not develop odour problems and lead to better effluent quality (Chan *et al.*, 2009). The use of microbial enzymes in lipid degradation methodologies is considered to improve efficiency, as they are eco-friendly, specific, reduce time investment, are easy to use, disease-free and cost-effective, as well as reducing chemical oxygen demand as well as the quantity of suspended lipid solids and colour (Dors *et al.*, 2013).

The previously isolated strain *Rhodococcus* sp. AQ5-07, a known phenol-degrading and cold-adapted Antarctic soil bacterium (Lee *et al.*, 2018), was tested here for its ability to degrade waste canola oil (WCO) as a sole source of carbon in the presence of different

types of heavy metal and other xenobiotic compounds. To our knowledge, no studies to date have examined the effects of heavy metals and other xenobiotics on the biodegradation activity of WCO using cold-adapted bacteria.

## 2 Materials and methods

### 2.1 Chemicals

Waste canola oil was collected from the Chilean Bernardo O'Higgins Riquelme Station, northern Antarctic Peninsula, in February 2018. Analytical grade chemicals were obtained from Fisher Scientific (Malaysia), Sigma (USA) and Merck (Darmstadt, Germany).

### 2.2 Microorganism culture and media preparation

Previously isolated *Rhodococcus* sp. strain AQ5-07 originally obtained from King George Island (South Shetland Islands, Antarctica) was cultured at 10°C in sterilised canola oil liquid medium. The liquid medium is used to enrich the lipid-degrading organism contained 3% canola oil and the following compositions (mg/L): 100 (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub>, 30 yeast extract, 20 MgSO<sub>4</sub>, 60 KH<sub>2</sub>PO<sub>4</sub>, 90 K<sub>2</sub>HPO<sub>4</sub>. For solid medium, Tween-peptone agar was prepared containing the following (mg/L): 10 CaCl<sub>2</sub>, 50 NaCl, 100 peptone, 180 agar and 5 mL/L Tween 80 (Aktas *et al.*, 2002). The media were autoclaved for 20 min at 121°C. The isolate was maintained and sub-cultured every two weeks in Tween-peptone agar medium.

### 2.3 Flask culture experiments

A single loop of a freshly grown culture from a Tween-peptone agar plate was transferred to 10 mL nutrient broth medium and incubated on a rotary shaker at 150 rpm and 10°C for 24 hr. Ten percent (v/v) of the culture was transferred to 100 mL of the canola oil medium in 250 mL Erlenmeyer flasks in triplicates and incubated for 72 hr at 150 rpm and 10°C. After 72 hr, the bacterial growth was measured according to colony forming units (CFU/mL) procedure of Hazan *et al.* (2012). Dilutions of 10<sup>-1</sup>, 10<sup>-2</sup>, 10<sup>-3</sup>, 10<sup>-4</sup>, 10<sup>-5</sup>, 10<sup>-6</sup> and 10<sup>-7</sup> made with 900 µL normal saline were carried out. Then, 100 µL of the suspension from 10<sup>-2</sup> to 10<sup>-7</sup> dilutions were aseptically transferred to petri

dishes followed by spreading the bacterial suspension to allow proper mix between the agar and bacterial suspension. The plates were then incubated at 15°C for 48 hr.

After the incubation, petri dishes containing between 30 and 300 colonies were collected and counted. Colony that forms the units per millilitre were calculated by multiplying the number of colonies by the dilution factor and divide by the volume of bacterial suspension plated on the agar (Equation 1) (Sieuwerts *et al.*, 2008).

$$CFU/mL = \frac{\text{No of colonies counted}}{\text{Volume of sample plated}} \times \text{dilution factor} \quad (1)$$

Control tubes containing the media and oil without the bacterial inoculation were used to ensure the dependency of observed growth on the inoculated bacteria.

### 2.4 Determination of canola oil degradation by gravimetric method

The amount of residual oil was determined using a gravimetric method following Abubakar *et al.* (2019) and Sihag and Pathak (2016). Fresh bacterial culture (OD<sub>600</sub> 1.3 - 1.4) was inoculated into 100 mL MSM media supplemented with 3% WCO. After 72 hr incubation, 1 mL was removed and bacterial growth was measured by assessment of colony forming units (CFU/mL).

For extracting the residual oil, 10 mL of n-hexane was added to the medium and the mixture was separated using a separating funnel. Two layers were formed; the lower layer that included the medium and the upper layer including the oil. The oil from the upper layer was collected in a pre-weighed Petri dish. One millilitre of filtered residual oil was used for GC-MS analysis. The percentage of canola oil degradation was calculated using Equation 2.

$$\text{Canola oil}(\%) = \frac{(a - b)}{a} \times 100\% \quad (2)$$

where a is the mass of canola oil added to the medium and b is the mass of residual canola oil.

### 2.5 Screening for heavy metal inhibition

A batch of experiment was conducted in a conical flask with initial WCO concentration of 3% (v/v) and the strain AQ5-07 was cultivated in nutrient broth (NB). About 10% (v/v) of the seed cultures were transferred

to 100 mL of WCO liquid media. Assessment of bacterial growth and oil degradation was carried out at every 12 hr intervals. The effect of different heavy metals on WCO degradation was studied. Heavy metals ions, namely zinc (Zn), cobalt (Co), nickel (Ni), silver (Ag), lead (Pb), copper (Cu), chromium (Cr), mercury (Hg), cadmium (Cd) or arsenic (As) at a concentration of 1 ppm, were added into the canola oil media. The media was incubated in an optimised condition for of  $(\text{NH}_4)_2\text{SO}_4$  1.0 g/L, yeast extract 0.3 g/L, temperature of 10°C, pH 7.5 and substrate concentration of 3% v/v. The amount of bacterial growth and degradation were then measured. Heavy metal ions that inhibited bacterial growth and oil-degrading ability were selected for further studies. All the experiments were conducted in triplicate and the results obtained for WCO were compared to confirm whether the metals caused a reduction in oil degradation. Media without the addition of heavy metals served as controls.

## 2.6 Effects of xenobiotics

The effect of xenobiotics (acrylamide, phenol and diesel) on WCO degradation was studied. Different concentrations were selected based on the literature (Lee et al., 2018; Rajasekar et al., 2007), and the media were incubated under optimised conditions for 72 hr at 10°C on a rotary shaker at 150 rpm.

## 2.7 Standardisation of parameters

The concentration of the selected heavy metals was manipulated, ranging between 0.0, 0.2, 0.4, 0.6, 0.8 and 1.0 ppm, while keeping all other parameters constant. Aliquots of *Rhodococcus* sp. AQ5-07 (10%) were transferred to 100 mL of each canola oil medium flask containing different concentrations of the heavy metals.

## 2.8 Inhibitory concentration 50 ( $IC_{50}$ )

The inhibitory concentration ( $IC_{50}$ ) was defined as the concentration that inhibited oil degradation by 50%. This test is used to determine best fit for a particular application from a list of various heavy metals concentrations. Tubes were set sequentially in decreasing order of concentration and incubated for 72 hr and the  $IC_{50}$  was calculated. To determine the  $IC_{50}$  values, the concentration range used of each heavy metals were from 0 - 1.0 ppm, and the data were analysed using Graphpad Prism v 8

software, by plotting % degradation against heavy metals concentration in ppm. All the analyses were conducted in triplicate and the mean  $\pm$  standard deviation values were taken.

## 2.9 Statistical analyses

Analysis of variance, using SPSS statistics V. 24 software package (SPSS Inc., Chicago, Illinois, USA), was used to compare results within treatment groups, and Tukey's test was used to conduct *post hoc* pairwise tests if significant differences were obtained. All experiments were carried out in triplicate and data are presented as a mean  $\pm$  standard deviation.

# 3 Results and discussion

## 3.1 Canola oil degradation

Cold-adapted *Rhodococcus* sp. AQ5-07 was previously tested in canola oil media for its ability to degrade WCO in the absence of heavy metals (Ibrahim et al. in review). In this study, 3% WCO was used as it is the optimum initial oil concentration exhibited by the strain (Fig. 1). Canola oil, as with vegetable oil generally at high concentrations (Shon et al., 2002), is known to be toxic to microorganisms, and efficient degradation demands the selection of a threshold level of canola oil that can be tolerated and degraded efficiently.




Fig. 1. Degradation and bacterial growth with a 3% initial canola oil concentration by *Rhodococcus* sp. AQ5-07 over 72 hr incubation. Error bars represent mean  $\pm$  standard deviation,  $n = 3$ .




Fig. 2. Effect of 1 ppm heavy metal ion concentrations on canola oil degradation and bacterial growth of *Rhodococcus* sp. AQ5-07 after 72 hr incubation. Error bars represent mean  $\pm$  standard deviation,  $n = 3$ .

### 3.2 Effects of heavy metal on canola oil degradation

The effects of different heavy metals on WCO degradation by *Rhodococcus* sp. AQ5-07 were compared after 72 h incubation. Three of the 10 heavy metals tested had an inhibitory effect on WCO degradation (Fig. 2), Ag, Hg, and Cd. Hg had the strongest inhibition effect with only 3.88% of WCO being degraded, followed by Ag (6.15%) and Cd (10.90%). One-way ANOVA confirmed significant differences between the different heavy metals tested in terms of bacterial growth and oil degradation ( $F(10, 22) = 176.51$ ,  $p < 0.001$  and  $F(10, 22) = 48.61$ ,  $p < 0.001$ , respectively). *Post hoc* tests identified no significant differences in degradation between control and Zn, Pb, Cu, Cr and As (all  $p > 0.05$ ), while the remaining heavy metals showed significant differences with the control (all  $p < 0.01$ ). None of the metals tested enhanced the degradation rate, and control medium without the addition of any heavy metals showed the highest rate.

Ahmad *et al.* (2018) and Zakaria *et al.* (2018) reported similar findings in studies of Antarctic *Arthrobacter bambusae* AQ5-003 and *Rhodococcus baikonurensis* AQ5-001, with Hg, Ag and Cd again inhibiting bacterial growth and phenol degradation. Abou-Shanab *et al.* (2007) reported that Zn, Ni, Cr and Hg inhibited *Arthrobacter* sp. AY509239. More widely, the detrimental effect of heavy metals on aquatic vertebrates include increased respiration, impaired fertilisation and reduced survival (Stebbing, 2002).




Fig. 3. Effect of a) Hg b) Cd and c) Ag concentration on canola oil degradation and growth of *Rhodococcus* sp. AQ5-07 after 72 hr incubation. Error bars represent mean  $\pm$  standard deviation,  $n = 3$ .

The pattern of microbial growth inhibition by heavy metals is often specific for each organism and each metal and, for any given metal, the toxic concentration may vary between different organisms (Sadler and Trudinger, 1967). Rajendran *et al.* (2003) reported that many heavy metals could affect microorganisms even at low concentration. As with most antimicrobial substances, the effects of heavy metals depend on their concentration which may be either bacteriostatic, where growth is inhibited but the microorganisms are not destroyed, or bactericidal, where the microorganisms are destroyed and will not recover even if the inhibitors are removed. It has been reported that both biochemical and



morphological changes in microorganisms can be observed at sub-lethal heavy metal concentrations (Sadler and Trudinger, 1967). Biochemical changes include changes in the proportion of some molecular components, particularly nucleic acids, which are stimulated in some organisms by cobalt, while the activities of certain oxidative enzymes are decreased (Nies, 1999). Morphological changes include the abnormal production of long filamentous forms when exposed to platinum and the conversion of bacterial rods to spherical forms. These may result in the inhibition of normal cell division and cell wall synthesis processes (Sharma et al., 2014).

The effect of Hg on WCO degradation was studied in more detail at concentrations between 0 and 1 ppm. Hg concentrations from 0.2 to 1 ppm reduced canola oil-degrading ability (Fig. 3a). Increasing Hg concentration above 0.4 ppm led to a significant decrease in canola oil degradation of 30.1%. One-way ANOVA revealed an overall significant difference between different Hg concentrations in both bacterial growth and oil degradation ( $F(5, 12) = 79.95, p < 0.001$  and  $F(5, 12) = 64.57, p < 0.001$ , respectively). *Post hoc* comparisons indicated that there were no significant differences in terms of bacterial growth between the control ( $M = 7.59, SD = 0.04$ ), and 0.2 ppm ( $M = 7.42, SD = 0.05$ ) and 0.4 ppm ( $M = 7.27, SD = 0.04$ ). In terms of oil degradation, there were significant differences between control and all Hg concentrations (all  $p < 0.001$ ). Hg is the most toxic non-radioactive metal in the natural environment and is toxic in any form (Abdel-Salam et al., 2010; Ross, 2004). Hg exerts its inhibitory effects by binding to sulfhydryl, phosphoryl, amide, carboxylic and amine groups of proteins. Robinson and Tuovinen (1984) reported that Hg is soluble in lipids and can easily bind to proteins containing sulfhydryl groups in the cell membrane, as well as to enzymes, causing toxicity. Mercury toxicity to bioremediation activity can be reduced by the addition of metal immobilising agents (Yusuf et al., 2019). Ibrahim et al. (2016) reported that immobilised *Leifsonia* sp. strain SIU can withstand up to 0.4 ppm Hg in caffeine degradation, helping to reduce Hg toxicity and maintaining the function of the bioremediation agent even in a polluted environment.

The effects of cadmium concentration on WCO degradation were studied at concentrations between 0 and 1 ppm. Cd concentrations from 0.4 to 1 ppm reduced canola oil-degrading ability (Fig. 3b). One-way ANOVA again identified overall significant differences in bacterial growth and oil degradation between the different concentrations ( $F(5, 12) =$

$62.84, p < 0.001$  and  $F(5, 12) = 62.00, p < 0.001$ , respectively). *Post hoc* comparisons indicated that there were no significant differences ( $p = 0.595$ ) between the degradation achieved between the control ( $M = 74.34, SD = 9.42$ ) and 0.2 ppm concentration ( $M = 66.60, SD = 5.97$ ), but there were significant differences between the control and all other Cd concentrations (all  $p < 0.001$ ). Cadmium occurs naturally in the soil environment but, due to human activities, it is more widely present. Ahmad et al. (2018) reported that cadmium inhibited phenol degradation by Antarctic *Arthrobacter bambusae* strain AQ5-003, being toxic even at 0.1 ppm. In contrast, Cd can also enhance bacterial growth and degradation in another Antarctic bacterium, *Rhodococcus baikonurensis* strain AQ5-001 (Zakaria et al., 2018). Cd also inhibits the biodegradation of other xenobiotics (Hoffman et al., 2005; Sandrin and Maier, 2003). The inhibitory effect of Cd can be lessened by immobilising the bacteria (Ahmad et al., 2017; Ibrahim et al., 2016). Furthermore, the addition of chemical additives can reduce the toxicity of cadmium ions (El-Deeb and Altalhi, 2009). Finally, the effects of various Ag concentrations on WCO degradation were studied from 0 to 1 ppm. The results obtained demonstrated that concentrations from 0.6 to 1 ppm reduced canola oil-degrading ability by 31.49, 19.23 and 13.14%, respectively (Fig. 3c). A significance difference was observed between different Ag concentrations in both analyses as shown by analysis of variance ( $F(5, 12) = 51.14, p < 0.001$  and  $F(5, 12) = 75.18, p < 0.001$ , respectively). In *post hoc* tests the degradation achieved under control conditions ( $M = 74.34, SD = 9.42$ ) and 0.2 ppm concentration ( $M = 70.77, SD = 3.09$ ) did not differ significantly, but there were significant differences between the control and all other Ag concentrations (all  $p < 0.001$ ). The strain AQ5-07 degraded 13.14% WCO in the presence of 1.0 ppm Ag, thus confirming silver as a strong inhibitor of canola oil degradation.

Silver can be activated in the presence of cytochrome b and exerts its toxicity to bacteria by penetration of the silver ions through the plasma membrane and the bacterial cell wall, thus inactivating membrane-bound proteins. The ion also binds to bacterial DNA, interrupting replication. Silver ions can also damage the ability of ribosomes to translate mRNA into proteins (Chudobova et al., 2013). Nies (1999) reported that Ag in coins and medicine can act as an anti-microbial agent.

### 3.3 Inhibitory concentration 50 (IC<sub>50</sub>)

The half maximal inhibitory concentration (IC<sub>50</sub>) is an effective measurement of the activity of a substance in preventing a particular biochemical or biological function. In the current study, the IC<sub>50</sub> is the concentration that reduces canola oil degradation by 50%. The IC<sub>50</sub> values of Hg, Cd and Ag for canola oil degradation by *Rhodococcus* sp. AQ5-07 are shown in Fig. 4a, b and c. The IC<sub>50</sub> values were 0.3866 ppm (Hg), 0.4539 ppm (Cd) and 0.3217 ppm (Ag). The IC<sub>50</sub> value for Hg inhibition by Antarctic *Rhodococcus baikonurensis* strain AQ5-001 was 0.5 ppm (Zakaria et al., 2018). Ahmad et al. (2018) reported IC<sub>50</sub> values for *Arthrobacter bambusae* strain AQ5-003 of 0.04 ppm for Cd and 0.53 ppm for Ag and concluded that Cd was the most toxic metal of those tested.




Fig. 4. The IC<sub>50</sub> of a) Hg b) Cd and c) Ag on canola oil degradation by *Rhodococcus* sp. AQ5-07. Error bars represent mean  $\pm$  standard deviation,  $n = 3$ .

### 3.4 Effects of xenobiotics (acrylamide, phenol and diesel) on canola oil degradation

The effects of different acrylamide concentrations on WCO degradation by *Rhodococcus* sp. AQ5-07 were studied at concentrations from 0 - 200 mg/L over 72 hr incubation. Acrylamide concentrations from 10 - 200 mg/L reduced canola oil degradation (Fig. 5a). One-way ANOVA showed overall significant differences in oil degradation and bacterial growth between different acrylamide concentrations ( $F(5, 12) = 28.63, p < 0.001$  and  $F(5, 12) = 69.09, p < 0.001$ , respectively).

There were no significant differences between the degradation achieved by the control ( $M = 68.50, SD = 7.90$ ) and 10.00 mg/L ( $M = 60.30, SD = 6.81$ ), but there were significant differences between the control and all other acrylamide concentrations (all  $p < 0.001$ ). At 200 mg/L, there was a significant decrease in oil degradation (19.20%), which may be due to the inhibitory effect of acrylamide on thiol groups of proteins (Buranasilp and Charoenpanich, 2011). These data suggest that, at high concentrations, acrylamide is toxic to this bacterial strain. Shukor et al. (2009a) similarly reported a decrease in bacterial growth at high acrylamide concentration above 500 mg/L in Antarctic *Pseudomonas* sp. strain DRYJ7. Nawaz et al. (1998) and Buranasilp and Charoenpanich (2011) compared acrylamide degradation by free and immobilised cells of *Rhodococcus* sp. and *Enterobacter aerogenes*, respectively, and found that immobilised cells degraded acrylamide to acrylic acid and ammonia faster than free cells.

The effects of different phenol concentrations on WCO degradation by *Rhodococcus* sp. AQ5-07 were studied from 0 - 500 mg/L over a 72 hr incubation. Phenol concentrations from 50 - 500 mg/L affected canola oil degradation (Fig. 5b). One-way ANOVA confirmed that the effects of different phenol concentrations on bacterial growth and oil degradation were significant ( $F(5, 12) = 25.77, p < 0.001$  and  $F(5, 12) = 20.77, p < 0.001$ , respectively). *Post hoc* tests indicated that there were no significant differences in the growth and degradation achieved in the control ( $M = 68.50, SD = 7.90$ ), 10 mg/L ( $M = 70.44, SD = 3.90$ ) and 50 mg/L ( $M = 51.02, SD = 8.30$ ) phenol concentrations, while the remaining pairwise comparisons with the control were significantly different. At 10 mg/L, there was a slight increase in WCO degradation, which is due

to the effect of the phenol. At 500 mg/L, there was a significant decrease in oil degradation (27.40%), which may be due to the toxic effect of phenol on the bacterium. According to Wang *et al.* (2007) maximum phenol degradation by *Acinetobacter* sp. strain PD 12 was achieved at 500 mg/L. It has also been reported that, at high phenol concentrations, the compound inhibits bacterial growth. Lee *et al.* (2018) recorded maximum phenol degradation at 500 mg/L by the Antarctic strains *Arthrobacter* sp. AQ5-05 and AQ5-06 and *Rhodococcus* sp. AQ5-07.

The effects of different initial concentrations of diesel on WCO degradation by *Rhodococcus* sp. AQ5-07 were studied from 0 - 2.5% after 72 hr incubation. Diesel concentrations from 1 - 2.5% reduced canola oil degradation (Fig. 5c). This suggests that the optimum carbon source (WCO and diesel) initial concentrations for the growth of strain AQ5-07 were 3% (v/v) and 0.5% (v/v) diesel, respectively. One-way ANOVA identified overall significant differences in bacterial growth and oil degradation between different diesel concentrations ( $F(5, 12) = 30.04$ ,  $p < 0.001$  and  $F(5, 12) = 31.84$ ,  $p < 0.001$ , respectively). *Post hoc* comparisons showed that there were significant differences between degradation achieved between the control and all diesel concentrations (all  $p < 0.001$ ).

Canola oil in the presence of diesel is required as a carbon source at certain concentrations. According to Shukor *et al.* (2009b), diesel can be toxic to microorganisms due to the solvent effect of diesel, which destroys the bacterial cell membrane. Therefore, many studies on microbial degradation of diesel have been carried out using lower diesel concentrations ranging from 0.5 - 1.5% (v/v) (Rajasekar *et al.*, 2007). This result is consistent with that reported by Shukor *et al.* (2009b) which found that, at high diesel concentrations of 1 or 1.5% (v/v), degradation is generally slow and toxic to bacteria. Diesel degradation at higher concentrations greater than 5.5% (v/v) has been reported, but requires additional yeast extract (1 g/L) and glucose (2 g/L) (Kwapisz *et al.*, 2008). Since *Rhodococcus* sp. AQ5-07 was able to withstand a high concentration of diesel, this implies that the strain is a good candidate for canola oil bioremediation when diesel is present as a co-contaminant.

Due to increased in human activities like tourism and scientific research have led to the risk of diesel pollution. The spilled diesel may serve as a substrate for microbial growth. Oil and other xenobiotics pollutants in Antarctica tends to be persistent in the region due to extremely low temperatures and often

dryness limit the rates of bioprocesses in addition to abiotic degradation (Lee *et al.*, 2018).

## Conclusions

The present study revealed the effect of heavy metals and other xenobiotics on canola oil biodegradation using a recently isolated cold-adapted Antarctic bacterium, *Rhodococcus* sp. AQ5-07. Three of the ten heavy metals tested, namely Ag, Cd and Hg, had a strongly inhibitory effect on bacterial growth and oil degradation at concentrations of 1 ppm. Hg and Cd showed significant inhibition at 0.2 and 0.4 ppm with reduction in canola oil degradation of 43.63% and 50.1%, respectively, while Ag showed an effect at 0.6 ppm with reduction in degradation of 31.49%. The  $IC_{50}$  values of Hg, Cd and Ag were 0.39, 0.45 and 0.32 ppm, respectively. The strain can also withstand 10 mg/L acrylamide, 50 mg/L phenol and 0.5 % (v/v) diesel respectively. The present information regarding the effects of heavy metals and xenobiotics on canola oil degradation by this cold-tolerant bacterium can be useful to the real state for oil degradation in Antarctic environments exposed to multiple pollutants under low temperature.

## Acknowledgements

This work was supported by Matching Grant PUTRA (UPM 9300436), PUTRA Berimpak (9660000 and 9678900). We acknowledge Tertiary Education Trust Fund (Tetfund) and Bayero University, Kano for sponsorship and Universiti Putra Malaysia for providing an enabling environment to conduct the research. P. Convey is supported by NERC core funding to the British Antarctic Survey 'Biodiversity. Evolution and Adaptation' Team. The authors also would like to thank Centro de Investigacion y Monitoreo Ambiental Antártico (CIMAA), Chilean Army and the Antarctic General Bernardo O'Higgins Station staff especially the Comandante de la Base O'Higgins; Teniente Coronel Jose Ignacio Alvarado Camps, Chef; Suboficial Juan David Sandoval Navarrete and Sargento Juan Eduardo Cortínez Padovani, Nancy Calisto-Ulloa, Yayasan Penyelidikan Antartika Sultan Mizan (YPASM), National Antarctic Research Centre (NARC) and Instituto Antártico Chileno (INACH).



## References

---

- Abdel-Salam, A.M., Al-Dekheil, A., Babkr, A., Farahna, M. and Mousa, H.M. (2010). High fiber probiotic fermented mares milk reduces the toxic effects of mercury in rats. *North American Journal of Medical Sciences* 2, 569-575.
- Abou-Shanab, R.A.I., Van Berkum, P. and Angle, J.S. (2007). Heavy metal resistance and genotypic analysis of metal resistance genes in gram-positive and gram-negative bacteria present in Ni-rich serpentine soil and in the rhizosphere of *Alyssum murale*. *Chemosphere* 68, 360-367.
- Abubakar, M., Abdul Habib, N.M.S., Manogaran, M., Yasid, N.A., Alias, S.A., Ahmad, S.A., Smykla J., Hassan M.A., and Abd Shukor M.Y. (2019). Response surface-based optimization of the biodegradation of a simulated vegetable oily ballast wastewater under temperate conditions using the Antarctic bacterium *Rhodococcus erythropolis* ADL36. *Desalination and Water Treatment* 144, 129-137.
- Ahmad, S.A., Shamaan, N.A., Syed, M.A., Dahalan, F.A., Abdul Khalil, K., Ab Rahman, N.A. and Shukor, M.Y. (2017). Phenol degradation by *Acinetobacter* sp. in the presence of heavy metals. *Journal of the National Science Foundation of Sri Lanka* 45, 247-253.
- Ahmad, S.A., Asokan, G., Yasid, N.A., Nawawi, N.M., Subramaniam, K., Zakaria, N.N. and Shukor, M.Y. (2018). Effect of heavy metals on biodegradation of phenol by Antarctic: *Arthrobacter bambusae* strain AQ5-003. *Malaysian Journal of Biochemistry and Molecular Biology* 21, 47-51.
- Aktas, E., Yigit, N. and Ayyildiz, A. (2002). Esterase activity in various *Candida* species. *Journal of International Medical Research* 30, 322-324.
- Alcázar-Medina, F., Núñez-Núñez, C., Rodríguez-Rosales, M., Valle-Cervantes, S., Alarcón-Herrera, M. and Proal-Nájera, J. (2019). Lead removal from aqueous solution by spherical agglomeration using an extract of *Agave lechuguilla* Torr. as biosurfactant. *Revista Mexicana De Ingeniería Química* 19, 71-84.
- Bargagli, R. (2005). *Antarctic Ecosystems: Environmental Contamination, Climate Change, and Human Impact*. Berlin: Springer.
- Bharathi, P., Elavarasi, N. and Mohanasundaram, S. (2012) Studies on rate of biodegradation of vegetable (coconut) oil by using *Pseudomonas aureginosa*. *International Journal of Environmental Biology* 2, 12-19.
- Buendía-González, L., Cruz-Sosa, F., Rodríguez-Huezo, M., Barrera-Díaz, C., Hernández-Jaimes, C. and Orozco-Villafuerte, J. (2019). *In vitro* simultaneous accumulation of multiple heavy metals by *Prosopis laevigata* seedlings cultures. *Revista Mexicana de Ingeniería Química* 18, 1167-1177.
- Buranasilp, K. and Charoenpanich. J. (2011) Biodegradation of acrylamide by *Enterobacter aerogenes* isolated from wastewater in Thailand. *Journal of Environmental Sciences* 23, 396-403.
- Chan, Y.J., Chong, M.F., Law, C.L. and Hassell, D.G. (2009) A review on anaerobic-aerobic treatment of industrial and municipal wastewater. *Chemical Engineering Journal* 155, 1-18.
- Chu, W-L., Dang, N-L., Kok, Y-Y., Yap K-S. I., Phang, S-W. and Convey, P. (2019) Heavy metal pollution in Antarctica and its potential impacts on algae. *Polar Science* 20, 75-83.
- Chudobova, D., Maskova, D., Nejdil, L., Kopel, P., Merlos-Rodrigo, M.A., Adam, V. and Kizek, R. (2013). The effect of silver ions and silver nanoparticles on *Staphylococcus aureus*. In: *Microbial Pathogens and Strategies for Combating them: Science, Technology and Education* (A. Méndez-Vilas, ed.), 728-735.
- Čipinyte, V., Grigiškis, S. and Baškys, E. (2009) Selection of fat-degrading microorganisms for the treatment of lipid-contaminated environment. *Biologija* 55, 84-92.
- Council of Managers of National Antarctic Programs (COMNAP) (2012). Main Antarctic facilities operated by national Antarctic programs in the Antarctic Treaty Area (south of 60' latitude south). <https://www.comnap.aq/Information/SitePages/Home.aspx>
- Convey, P., Hughes K.A. and Tin, T. (2012). Continental governance and environmental

- management mechanisms under the Antarctic Treaty System: Sufficient for the biodiversity challenges of this century?. *Biodiversity* 13, 234-248.
- Corral-Escárcega, M.C., Ruíz-Gutiérrez, M.G., Quintero-Ramos, A., Meléndez-Pizarro, C.O., Landizabal-Gutiérrez, D. and Campos-Venegas, K. (2017). Use of biomass-derived from pecan nut husks (*Carya illinoensis*) for chromium removal from aqueous solutions. Column modeling and adsorption kinetic studies. *Revista Mexicana de Ingeniería Química* 13, 939-953.
- Danikuu F.M. and Sowley E.N.K. (2014) Biodegradation of shea nut cake by indigenous soil bacteria. *Journal of Medical and Biomedical Sciences* 3, 9-15.
- Dors, G., Mendes, A.A., Pereira, E.B., de Castro, H.F. and Furigo, A. (2013) Simultaneous enzymatic hydrolysis and anaerobic biodegradation of lipid-rich wastewater from poultry industry. *Applied Water Science* 3, 343-349.
- El-Deeb, B. and Altalhi, A.D. (2009) Degradative plasmid and heavy metal resistance plasmid naturally coexist in phenol and cyanide assimilating bacteria. *American Journal of Biochemistry and Biotechnology* 5, 84-93.
- Fadile A., El-Hassani, F.Z., Aissam, H., Merzouki, M. and Benlemlih, M. (2011). Aerobic treatment of lipid-rich wastewater by a bacterial consortium. *African Journal of Microbiology Research* 5, 5333-5342.
- Hazan, R., Que, Y.A., Maura, D. and Rahme, L.G. (2012). A method for high throughput determination of viable bacteria cell counts in 96-well plates. *BMC Microbiology* 12, 1-7.
- Hernández-Martínez, R., Valdivia-Rivera, S., Betto-Sagahon, J., Coreño-Alonso, A., Tzintzun-Camacho, O. and Lizardi-Jiménez, M. (2018). Solubilization and removal of petroleum hydrocarbons by a native microbial biomass in a bubble column reactor. *Revista Mexicana de Ingeniería Química* 18, 181-189.
- Hoffman, D.R., Okon J.L. and Sandrin T.R. (2005) Medium composition affects the degree and pattern of cadmium inhibition of naphthalene biodegradation. *Chemosphere* 59, 919-927.
- Hong, S., Soyol-Erdene, T.O., Hwang, H.J., Hong, S.B., Hur, S.D. and Motoyama, H. (2012). Evidence of global-scale As, Mo, Sb, and Tl atmospheric pollution in the Antarctic Snow. *Environmental Science and Technology* 46, 11550-11557.
- IAATO (2017) Tourism Statistics. Available at: [https://iaato.org/en\\_GB/tourism-statistics](https://iaato.org/en_GB/tourism-statistics)
- Ibrahim, S., Muhammad, A., Tanko, A.S., Abubakar, A., Ibrahim, H., Shukor, M.Y. and Ahmad, S.A. (2016). Studies of action of heavy metals on caffeine degradation by immobilised *Leifsonia* sp. strain SIU. *Bayero Journal of Pure and Applied Sciences* 8, 138-144.
- Ibrahim, S., Shukor, M.Y.A., Yazid N.A. and Ahmad S.A. (2018) Microbial degradation of vegetable oils: A review. *Malaysian Journal of Biochemistry and Molecular Biology* 28, 45-55.
- Ibrahim, S., Zahri, K.N.M., Convey, P., Khalil, K.A., Gomez-Fuentez, C., Zulkarnain, A., Alias, S.A., Gonzalez-Rocha, G. and Ahmad, S.A. (2019). Optimisation of biodegradation conditions for waste canola oil by cold-adapted *Rhodococcus* sp. AQ5-07 from Antarctica. *Electronic Journal of Biotechnology* (Under Review).
- Interiano-López, M., Ramírez-Coutiño, V., Godínez-Tovar, L., Zamudio-Pérez, E. and Rodríguez-Valadez, F. (2019). Bioremediation methods assisted with humic acid for the treatment of oil-contaminated drill cuttings. *Revista Mexicana de Ingeniería Química* 18, 929-937.
- Karamba, K.I., Ahmad, S.A., Zulkarnain, A., Syed, M.A., Khalil, K.A., Shamaan, N.A., Dahalan, F.A. and Shukor, M.Y. (2016). Optimisation of biodegradation conditions for cyanide removal by *Serratia marcescens* strain AQ07 using one-factor-at-a-time technique and response surface methodology. *Rendiconti Lincei* 27, 533-545.
- Kumar, S., Mathur, A., Singh, V., Nandy, S., Khare, S.K. and Negi, S. (2012). Bioremediation of waste cooking oil using a novel lipase produced by *Penicillium chrysogenum* SNP5 grown in solid medium containing waste grease. *Bioresource Technology* 120, 300-304.

- Kwapisz, E., Wszelaka, J., Marchut, O. and Bielecki, S. (2008). The effect of nitrate and ammonium ions on kinetics of diesel oil degradation by *Gordonia alkanivorans* S7. *International Biodeterioration and Biodegradation* 61, 214-222.
- Lee, G.L.Y., Ahmad, S.A., Yasid, N.A., Zulkharnain, A., Convey, P., Johari, W.L.W., Alias, S.A., Gonzalez-Rocha, G. and Shukor, M.Y. (2018). Biodegradation of phenol by cold-adapted bacteria from Antarctic soils. *Polar Biology* 3, 553-562.
- Matsumiya, Y., Wakita, D., Kimura, A., Sanpa, S. and Kubo, M. (2007). Isolation and characterization of a lipid-degrading bacterium and its application to lipid-containing wastewater treatment. *Journal of Bioscience and Bioengineering* 103, 325-330.
- Nagarajan, J., Nawawi, N.M. and Ibrahim, A.L. (2014). Rhodococcus UKMP-5M, an endogenous lipase producing actinomycete from Peninsular Malaysia. *Biologia* 69, 123-132.
- Nawaz, M.S., Billedeau, S.M. and Cerniglia, C.E. (1998) Influence of selected physical parameters on the biodegradation of acrylamide by immobilized cells of Rhodococcus sp. *Biodegradation* 9, 381-387.
- Nies, D.H. (1999) Microbial heavy-metal resistance. *Applied Microbiology and Biotechnology* 51, 730-750.
- Pereira, M.G., Mudge, S.M. and Latchford, J. (2003) Vegetable oil spills on salt marsh sediments; comparison between sun- flower and linseed oils. *Marine Environmental Research* 56, 367-385.
- Phong, N.T., Duyan, N.T. and Diep, C.N. (2014) Isolation and characterization of lipid-degrading bacteria in wastewater of food processing plants and restaurants in Can Tho City, Vietnam. *American Journal of Life Sciences* 2, 382-388.
- Rack, U. (2015). Polar expeditions. In: *Exploring the Last Continent*, (Liggett, D., Storey, B., Cook, Y. and Meduna, V. eds), Cham: Springer.
- Rajasekar, A., Babu, T.G., Pandian, S.T., Maruthamuthu, S., Palaniswamy, N. and Rajendran, A. (2007). Role of *Serratia marcescens* ACE2 on diesel degradation and its influence on corrosion. *Indian Journal of Microbiology and Biotechnology* 34, 589-598.
- Rajendran, P., Muthukrishnan, J. and Gunasekaran, P. (2003). Microbes in heavy metal remediation. *Indian Journal Experimental Biology* 41, 935-944.
- Robinson, J.B. and Tuovinen, O.H. (1984). Mechanisms of microbial resistance and detoxification of mercury and organomercury compounds: physiological, biochemical, and genetic analyses. *Microbiological Reviews* 48, 95-124.
- Ross, D.A. (2004). The toxicology of mercury. *New England Journal of Medicine* 350, 945-947.
- Ruiz-Marín, A., Zavala-Loria, J.C., Canedo-López, Y. and Cordova-Quiroz, A.V (2013). Tropical bacteria isolated from oil-contaminated mangrove soil: bioremediation by natural attenuation and bioaugmentation. *Revista Mexicana de Ingeniería Química* 12, 553-560.
- Sadler, W.R. and Trudinger, P.A. (1967). The inhibition of microorganisms by heavy metals. *Mineralium Deposita* 2, 158-168.
- Sandrin, T.R. and Maier, R.M. (2003) Impact of metals on the biodegradation of organic pollutants. *Environmental Health Perspectives* 111, 1093-1101.
- Serikovna, S.Z., Serikovich, K.S., Sakenovna, A.S., Murzakhmetovich, S.S. and Khamitovich, A.K. (2013). Screening of lipid degrading microorganisms for wastewater treatment. *Malaysian Journal of Microbiology* 9, 219-226.
- Sharma, B., Singh S. and Siddiqi N.J. (2014). Biomedical implications of heavy metals induced imbalances in redox systems. *Biomed Research International* 2014, doi:10.1155/2014/640754.
- Shon, H.K., Tian, D., Kwon, D.Y., Jin, C.S., Lee, T.J. and Chung, W.J. (2002). Degradation of fat, oil, and grease (FOGs) by lipase-producing bacterium *Pseudomonas* sp. strain D2D3. *Journal of Microbiology and Biotechnology* 12, 583-591.

- Shukor, M.Y., Gusmanizar, N., Ramli, J., Shamaan, N.A., MacCormack, W.P. and Syed, M.A. (2009a). Isolation and characterization of an acrylamide-degrading Antarctic bacterium. *Journal of Environmental Biology* 30, 107-112.
- Shukor, M.Y., Dahalan, F.A., Jusoh, A.Z., Muse, R., Shamaan, N.A. and Syed, M.A. (2009b). Characterization of diesel-degrading strain isolated from hydrocarbon-contaminated site. *Journal of Environmental Biology* 30, 145-150.
- Sihag, S. and Pathak, H. (2016). Biodegradation of 2T engine oil using soil microbe and gravimetric analysis. *International Journal of Scientific Engineering Research* 7, 1286-1295.
- Soyol-Erdene, T.O., Huh, Y., Hong, S. and Hur, S.D. (2011). A 50-year record of platinum, iridium, and rhodium in Antarctic Snow: volcanic and anthropogenic sources. *Environmental Science and Technology* 45, 5929-5935.
- Stebbing, A.R.D (2002). Tolerance and hormesis - increased resistance to copper in hydroids linked to hormesis. *Marine Environmental Research* 54, 805-809.
- Suárez-García, L., Cuervo-López, F., & Texier, A.-C. (2019). Biological removal of mixtures of ammonium, phenol, cresol isomers, and sulfide in a sequencing batch reactor. *Revista Mexicana de Ingeniería Química* 18, 1189-1202.
- Subramaniam, K., Mazuki, T.A.T., Shukor, M.Y., and Ahmad, S.A. (2019). Isolation and optimisation of phenol degradation by Antarctic isolate using one factor at time. *Malaysian Journal of Biochemistry and Molecular Biology* 22, 79-86.
- Tiwari, A.K. (2017). Environmental monitoring around Indian Antarctic Stations. *Proceedings of Indian National Science Academy* 83, 399-413.
- Vodopivec, C., Curtosi, A., Villaamil, E., Smichowski, P., Pelletier, E. and Mac Cormack, W.P. (2015) Heavy metals in sediments and soft tissues of the Antarctic clam *Laternula elliptica*: More evidence as a? Possible biomonitor of coastal marine pollution at high latitudes?. *Science of Total Environment* 502, 375-384.
- Williams, L., Borchhardt, N., Colesie, C., Baum, C., Komsic-Buchmann, K., Rippin, M., Becker, B., Karsten, U. and Büdel, B. (2017) Biological soil crusts of Arctic Svalbard and of Livingston Island, Antarctica. *Polar Biology* 40, 399-411.
- Wang, Y., Tian, Y., Han, B., Zhao, H., Bi, J. and Cai, B. (2007). Biodegradation of phenol by free and immobilized *Acinetobacter* sp. strain PD12. *Journal of Environmental Sciences* 19, 222-225.
- Xie, Z. and Sun, L. (2008). A 1,800-year record of arsenic concentration in the penguin dropping sediment, Antarctic. *Environmental Geology* 55, 1055-1059.
- Yusuf, I., Ahmad, S.A., Phang, L.Y., Yasid, N.A. and Shukor, M.Y. (2019). Effective production of keratinase by gellan gum-immobilised *Alcaligenes* sp. AQ05-001 using heavy metal-free and polluted feather wastes as substrates. *3 Biotech* 9, 32. Doi:[org/10.1007/s13205-018-1555-x](https://doi.org/10.1007/s13205-018-1555-x)
- Zakaria, N.N., Ahmad, S.A., Yasid, N.A., Yin, G.L.L., Manogaran, M., Subramaniam, K., Mazuki, T.T.A., Nawawi, N.M. and Shukor, M.Y. (2018). Biodegradation of phenol by Antarctic bacterium *Rhodococcus baikonurensis* strain AQ5-001 in the presence of heavy metals. *Malaysian Journal of Biochemistry and Molecular Biology* 21, 29-36.