

Geochemistry, Geophysics, Geosystems®



TECHNICAL REPORTS: DATA

10.1029/2021GC010154

Key Points:

- PetroChron Antarctica is a new relational database containing petrological, geochemical, and geochronological data sets from sampled rocks across Antarctica
- Lithology and age of geolocated samples, along with computed chemical and physical rock properties, facilitate quantitative analysis and data integration for interdisciplinary use (e.g., geodynamics, oceanography, ice sheet dynamics, biodiversity, and soil studies)
- The PetroChron Antarctica database is accessible online via a web portal, where data can be freely downloaded as comma-separated text flat files or individual tables to be used in a relational database system

Supporting Information:

Supporting Information may be found in the online version of this article.

Correspondence to:

J. A. Halpin,
jacqueline.halpin@utas.edu.au

Citation:

Sanchez, G., Halpin, J. A., Gard, M., Hasterok, D., Stål, T., Raimondo, T., et al. (2021). PetroChron Antarctica: A geological database for interdisciplinary use. *Geochemistry, Geophysics, Geosystems*, 22, e2021GC010154. <https://doi.org/10.1029/2021GC010154>

Received 8 SEP 2021
Accepted 21 NOV 2021

Author Contributions:

Conceptualization: G. Sanchez, J. A. Halpin, M. Gard, D. Hasterok, T. Stål, T. Raimondo, S. Peters, A. Burton-Johnson
Data curation: G. Sanchez, J. A. Halpin, M. Gard, D. Hasterok, T. Stål, T. Raimondo, S. Peters

© 2021 The Authors.

This is an open access article under the terms of the [Creative Commons Attribution-NonCommercial License](https://creativecommons.org/licenses/by-nc/4.0/), which permits use, distribution and reproduction in any medium, provided the original work is properly cited and is not used for commercial purposes.

PetroChron Antarctica: A Geological Database for Interdisciplinary Use

G. Sanchez^{1,2} , J. A. Halpin¹ , M. Gard^{2,3} , D. Hasterok^{3,4} , T. Stål^{1,5} , T. Raimondo^{6,7} , S. Peters⁶ , and A. Burton-Johnson⁸ 

¹Institute for Marine and Antarctic Studies, University of Tasmania, Hobart, TAS, Australia, ²Geoscience Australia, Canberra, ACT, Australia, ³Department of Earth Sciences, University of Adelaide, Adelaide, SA, Australia, ⁴Mawson Centre for Geoscience, University of Adelaide, Adelaide, SA, Australia, ⁵School of Natural Sciences (Physics), University of Tasmania, Hobart, TAS, Australia, ⁶UniSA STEM, University of South Australia, Adelaide, SA, Australia, ⁷Future Industries Institute, University of South Australia, Adelaide, SA, Australia, ⁸British Antarctic Survey, Cambridge, UK

Abstract We present PetroChron Antarctica, a new relational database including petrological, geochemical and geochronological data sets along with computed rock properties from geological samples across Antarctica. The database contains whole-rock geochemistry with major/trace element and isotope analyses, geochronology from multiple isotopic systems and minerals for given samples, as well as an internally consistent rock classification based on chemical analysis and derived rock properties (i.e., chemical indices, density, p -velocity, and heat production). A broad range of meta-information such as geographic location, petrology, mineralogy, age statistics and significance are also included and can be used to filter and assess the quality of the data. Currently, the database contains 11,559 entries representing 10,056 unique samples with varying amounts of geochemical and geochronological data. The distribution of rock types is dominated by mafic (36%) and felsic (33%) compositions, followed by intermediate (22%) and ultramafic (9%) compositions. Maps of age distribution and isotopic composition highlight major episodes of tectonic and thermal activity that define well known crustal heterogeneities across the continent, with the oldest rocks preserved in East Antarctica and more juvenile lithosphere characterizing West Antarctica. PetroChron Antarctica allows spatial and temporal variations in geology to be explored at the continental scale and integrated with other Earth-cryosphere-biosphere-ocean data sets. As such, it provides a powerful resource ready for diverse applications including plate tectonic reconstructions, geological/geophysical maps, geothermal heat flow models, lithospheric and glacial isostasy, geomorphology, ice sheet reconstructions, biodiversity evolution, and oceanography.

Plain Language Summary On a continent with less than 0.18% of outcrop, information such as the rock type, chemistry and age of Antarctic rock samples are critical inputs for understanding complex interactions between the lithosphere, cryosphere, biosphere, and ocean. We have created PetroChron Antarctica, a relational database containing a compilation of petrological, geochemical and geochronological data from geological samples across Antarctica. The database contains more than 10,000 samples, along with chemical indices and rock properties calculated from chemical analyses. PetroChron Antarctica contains spatial meta-information to enable visualization and analysis of the database using an online interactive map, which highlights the variability in crustal geology at the continental scale and can be used for interdisciplinary scientific studies. PetroChron Antarctica is freely available through Zenodo and an ESRI Web Feature Service (<http://bit.ly/petrochron>).

1. Introduction

The Antarctic lithosphere was built over billions of years (e.g., Boger, 2011; Harley et al., 2013), and it is increasingly clear that this long and complex lithospheric evolution both records and influences interactions with the oceans and cryosphere (e.g., Burton-Johnson et al., 2020; Hochmuth et al., 2020; Paxman et al., 2020; Whitehouse et al., 2019). Understanding these interrelated processes critically depends on the ability to integrate large heterogeneous data sets from regional to continental scale (Stål et al., 2020). Antarctic data sets are typically poorly represented in global databases. In the Antarctic geosciences, data set hosting and dissemination are mainly supported through the Scientific Committee on Antarctic Research (SCAR; <https://www.scar.org/resources/data/>) and NASA's Earth Science Data Systems Program (<https://search.earthdata.nasa.gov/search>). However, geological data sets are poorly resolved compared with the burgeoning geophysical data streams. Where available,

Funding acquisition: J. A. Halpin
Investigation: G. Sanchez, J. A. Halpin, A. Burton-Johnson
Methodology: G. Sanchez, J. A. Halpin, M. Gard, D. Hasterok, T. Stål, T. Raimondo, S. Peters
Project Administration: J. A. Halpin, T. Raimondo
Resources: G. Sanchez, J. A. Halpin, M. Gard, D. Hasterok, A. Burton-Johnson
Software: G. Sanchez, M. Gard, D. Hasterok, T. Stål, T. Raimondo, S. Peters
Supervision: J. A. Halpin
Validation: G. Sanchez, J. A. Halpin, T. Stål
Visualization: G. Sanchez, J. A. Halpin, T. Stål, T. Raimondo, S. Peters
Writing – original draft: G. Sanchez, J. A. Halpin
Writing – review & editing: G. Sanchez, J. A. Halpin, M. Gard, D. Hasterok, T. Stål, T. Raimondo, S. Peters, A. Burton-Johnson

geological data are typically hosted within national databases (e.g., OZCHEM; Champion et al., 2007; Petlab; Strong et al., 2016) or individual publications and are therefore difficult to utilize.

Here, we present PetroChron Antarctica, a new geological database that includes geochemical, geochronological and petrological data sets from Antarctic rock samples, compiled from existing databases and individual publications. We also generate compositionally based classifications, geochemical indices and physical properties derived from the geochemical data where possible. This database builds upon the global whole-rock geochemistry compilation developed by Gard et al. (2019). A newly generated schema implemented to account for the newly incorporated data types and associated meta-information is described, including the data integration procedure. Finally, we relate some applications to highlight potential future uses of the database.

2. Existing Initiatives and Motivation for Data Augmentation and Integration

The PetroChron Antarctica database incorporates various geochemical and geochronological data sets, together with related petrological information, from both global and national initiatives (Table 1). Whereas these collections are a valuable asset for the geoscience community and are incorporated in numerous regional and global studies, they are mostly organized around data types of interest (Figure 1a) or localized in specific geographic areas where national campaigns have focused mapping and sampling efforts on accessible outcrop (Figures 1b and 1c). This lack of integration between geochemical and geochronological data (and other rock-based data), along with a strong asymmetry in data density from these existing databases, demonstrates the need to augment and integrate additional Antarctic geological data streams. PetroChron Antarctica, therefore, incorporates standardized peer-reviewed academic publications and some unpublished data (Figure 1d). Currently, the PetroChron Antarctica database contains 10,056 rock samples representing 11,559 data entries, of which around 40% are compiled from existing data repositories spanning over 80 years of research (Table 1). Whereas the existing databases are mostly located in West Antarctica, the distribution of geological data incorporated from individual publications is more widespread, and mostly located in East Antarctica (Figure 1d; 72%). These data can be integrated with geological map information (e.g., GeoMAP <https://www.scar.org/science/geomap/geomap/> and OneGeology <http://www.onegeology.org/>) to extract further geological information from the Antarctic lithosphere. Ideally, it could be linked to Antarctic sample collection information in the spirit of the Polar Rock Repository (<https://prr.osu.edu/>), enabling further data discovery and sample sharing for future research.

3. Database Foundational Framework

3.1. Data Model

The database architecture follows the key concept described in Figure 2. We decided to use a simplified relational database structure including only five sub-tables (metadata, petrology, geochemistry, geochronology, and rock properties) representing the core elements of sample-related information (Table 2). In an effort to meet the FAIR (findable, accessible, interoperable, and reusable) data standard for inter- and intra-disciplinary studies, we organize the different sub-tables around subdomains of knowledge used across the research community.

The minimalist relational model simplifies maintenance and minimizes file size. Indeed, complex relational models are usually not sustainable in the long term to support the expansion of data sets or fields to track provenance and modification. Our approach facilitates the extraction of the data from the database, the incorporation of the data into other databases with different schemas, and enables its use in various scientific workflows.

3.2. Data Compilation Workflow

To ensure data consistency and enhance database reliability over PetroChron Antarctica's lifetime, we implemented several procedures written in a com-

Table 1
Number of Sample Entries Per Data Source

Data source	No. entries
Others (publications, unpublished ...)	5,266
OZCHEM (Champion et al., 2007)	1,792
Petlab (Strong et al., 2016)	1,819
GEOROC (http://georoc.mpch-mainz.gwdg.de)	1,464
Burton-Johnson BAS compilation (Gard et al., 2019)	1,074
DateView (Eglington, 2004)	144
Total	11,559

Note. The Geochemistry of Rocks of the Oceans and Continents (GEOROC) data compilation contains chemical, isotope and limited age data for igneous rocks. National government collections include the Australian national whole-rock geochemical database (OZCHEM; Champion et al., 2007), the New Zealand national rock, mineral and geoanalytical database (Petlab; Strong et al., 2016) and the whole-rock geochemical data compilation from Burton-Johnson, British Antarctic Survey (BAS; included in Gard et al., 2019). Part of the geochronological database DateView (Eglington, 2004) is also included, but are not cited as such when the data have been modified or independently entered from individual publications.




Figure 1. Spatial distribution of PetroChron Antarctica samples categorized by (a) data type, and data source including (b) national databases, (c) international compilations, and (d) international peer-reviewed publications.

bination of programming languages (i.e., Python and PostgreSQL) for data standardization to create a common data schema (Figure 2, Table 2).

Collecting a useful Antarctic geological data set starts with accurate sample location information. Historically (i.e., prior to GPS), this information was not readily recorded in a useful format, or it may have been lost in the process of transcribing notes or maps. In the case, where accurate absolute spatial information is not provided in the original paper or data set, geographic locations along with latitude and longitude from the SCAR Gazetteer is used (Secretariat SCAR, 1992 updated 2014). For each entry, an attribute identifies the source of the geographic coordinate (i.e., Geographic Coordinate Information). This approach allows us to retain 45% of the ge-




Figure 2. The PetroChron Antarctica data model using a simple five-table structure representing metadata information and sub-domains of knowledge (petrology, geochemistry, geochronology, and rock properties). Text in blue represents computed data based on chemical analyses. For readability purposes, chemical and isotopic elements are grouped by element types (i.e., major elements, trace elements, and isotopes) as shown by the asterisks.

ological samples in PetroChron Antarctica that would previously have been excluded due to the lack of location information.

Lithology has a dominant control over the physical and chemical properties of rocks. We therefore categorize the database according to rock group (i.e., igneous, sedimentary, metamorphic) and rock type (e.g., plutonic, clastic, and metavolcanic) where known or inferred. However, there are a variety of lithology names based on different criteria (mineralogical, textural, chemical). Thus, to achieve consistency and reproducibility and avoid any subjectivity in assigning rock names to samples, we include a computed lithology based on whole-rock geochemical data as described by Hasterok et al. (2018) and Gard et al. (2019). Note that these classifications are purely chemical, and do not reflect the mineralogy, grain size, texture, and/or metamorphic grade of the sample.

The database structure is then focused on the integration of geochronological data sets with geochemical data. In the global whole-rock geochemical database of Gard et al. (2019), petrological information and geochemical analyses were linked to “estimated” crystallization ages of the sample as presented in the original paper. A key difference in PetroChron Antarctica is that geochronological information is stored as a set of parameters including the age type (i.e., isotopic system), the mineral isotope (i.e., analyzed mineral and/or whole rock), the age significance (e.g., crystallization age), the age approach (e.g., concordia age), and the analytical technique (e.g., SHRIMP; Figure 2). This configuration significantly increases the flexibility to support geochronological data from multiple isotopic systems and minerals for a given sample, which potentially have different geological significances. The inclusion of age-related statistical information if applicable (e.g., mean squared weighted deviation—MSWD, and probability of fit) enables data to be manipulated through more complex statistical analyses and could also be useful for data quality assessment. Geochronological parameters generally follow the schema of the established geochronological database DateView (Eglington, 2004) for consistency and easy transfer between databases.

Table 2
Description of Table Contents and Detailed Information of Key Field Attributes

Table name	Table content description	Field attribute	Field description
Metadata	Contains metadata information related to the recorded data including the approximative location and spatial reference (name, geographic coordinates, datum ...) of the sample, the source of the data (existing database, original paper reference), the type of sample and the technique used to collect it, the sample name and other geological information related to the terrane and/or stratigraphic unit the sample may belong to	geolocation	Information on the sample location (geographic area, place name). Additional information may be included, such as sites number, distance. Note that SCAR Gazetteer place names were used in most cases to consistently populate location names
		coordinfo	Indicates the technique used to flag how geographic coordinates were recorded in the database
		data_source	Source of the data if the record was extracted from an existing database or data compilation
		sample_type	Type of the sample collected—for example, veins, dyke, xenolith ...
		sampling	Sampling technique used to collect sample—for example, outcrop, dredge, core ...
		sample_name	Sample name as recorded by the author in the publication or existing database. Duplicate number may occur
Petrology	Comprises rock group, type, name, description, facies, mineralogy of the sample. Additional information are in chemical based classification (TAS, SIA granite type, frost classification). For further explanation, the reader is referred to Hasterok et al. (2018)	rock_group	High-level rock group of the sample (igneous, metamorphic and sedimentary rocks) assigned by original author/database
		rock_type	Standardized rock type—for example, plutonic, volcanic, metavolcanic, metaplutonic, metasedimentary, clastic, assigned, or inferred by the original author/database
		rock_name	Non-standardized rock name designated by the original author/database
		rock_description	Non-standardized detailed description of the rock sample from the original author/database
		rock_facies	Metamorphic facies information
		mineralogy_major	List of major minerals present in the rock sample
		mineralogy_minor	List of minor minerals present in the rock sample
		lithology	Chemical based rock type following methods described in Hasterok et al. (2018)
		qap_name	Computed rock names based on the TAS igneous classification (Middlemost, 1994), including high-Mg volcanics (Le Bas & Streckeisen, 1991)
		sia_scheme	S-, I-, and A-type granite classification
		frost_class1	Magnesian or Ferroan (Frost et al., 2001)
		frost_class2	Calcic, calc-alkalic, alkali-calcic, and alkalic (Frost et al., 2001)
		frost_class3	Metaluminous, peraluminous, and peralkaline (Frost et al., 2001)
Geochemistry	Sets of major, trace and isotope analyses. It also includes a set of chemical based indices computed from major element normalised (LOI-free) geochemical composition	geochem_mineral	Mineral/fraction analyzed—for example, whole rock, zircon ...
		geochem_tech_analysis	Analytical technique used for geochemical measurements
		geochem_major	Major element analyses—includes major element oxides as well as volatile, carbonate and LOI content where available
		geochem_trace	Trace element analyses
		geochem_isotopes	Isotopic ratio analyses, including initial ratio

Table 2
Continued

Table name	Table content description	Field attribute	Field description
		mg_number	Magnesium number. Fe ²⁺ estimated using $0.85 \times \text{FeOT}$
		fe_number	Iron number (Frost et al., 2001)
		mali	Modified alkali–lime index (Frost et al., 2001)
		asi	Alumina Saturation Index (ASI; Frost et al., 2001)
		maficity	$n\text{Fe} + n\text{Mg} + n\text{Ti}$
		cia	Chemical index of alteration (Nesbitt & Young, 1989)
		wip	Weathering index of Parker (1970)
		spar	Modified from Debon and Le Fort (1983) to remove apatite
		qtzindex	Quartz Index (Debon & Le Fort, 1983)
		r1	R1R2 chemical variation diagram (De la Roche et al., 1980)
		r2	R1R2 chemical variation diagram (De la Roche et al., 1980)
Geochronology	Includes age, age uncertainty and associated statistics of the age calculation (if provided in original reference/database). A set of metadata information related to the type of radiochronometer, the mineral dated, the approach and analytical technique used and the significance of the age are populated	age_type	Radiochronometer used to estimate the rock sample age—Ar–Ar, U–Pb ...
		age_mineral	Mineral used for dating—for example, mica, zircon ...
		age_significance	Significance of the calculated age—for example, Crystallization, Cooling ...
		age_approach	The approach used to calculate an age—for example, Regression, Concordia, Discordia, Ar Plateau
		age_techgeochem	The technique used to measure isotopic ratio used for dating—for example, TIMS (single grain and multigrain), SHRIMP, Laser ...
		age_ma	Radiometric age in Ma
		age_2SD_ma	Standard deviation—95% or 2 sigma—in Ma
		age_mswd	The calculated MSWD
		age_probffit	The calculated probability of fit
		age_probchi2	The calculated probability of Chi ² test
		age_analyse_n	Total number of analyses used to calculate an age
Rock properties	List of physical rock properties including heat production, seismic velocity and density estimation computed from geochemical analysis. For further information on the computation, see Hasterok et al. (2018)	p_velocity	Empirically calculated seismic velocity based on chemical composition. The compositional empirical model used was $V_p \text{ (km s}^{-1}\text{)} = 6.9 - 0.011\text{CSiO}_2 + 0.037\text{CMgO} + 0.045\text{CCaO}$. For further discussion on the computation, the reader can refer to Hasterok and Webb (2017)
		s_velocity	Empirically calculated seismic velocity based on chemical composition. For further discussion on the computation, the reader can refer to Jennings et al. (2019)
		density_model	Rock density computed from chemical analyses using linear regression as described in Hasterok et al. (2018)
		thermal_conductivity	Empirically calculated thermal conductivity based on chemical composition. For further discussion on the computation, the reader can refer to Jennings et al. (2019)
		heat_production	Heat production mass multiplied by the density estimate (in kg m^{-3}) (Rybach, 1988)

Table 2
Continued

Table name	Table content description	Field attribute	Field description
		heat_production_mass	Estimated from the chemical rock composition using the empirical formula $HP_{mass} = 10^{-5} \times (9.67CU + 2.56CTh + 2.89CK2O)$ where C are the concentrations of the heat producing elements in ppm except K2O in wt.% (Rybach, 1988)

Note. MSWD, mean squared weighted deviation; SCAR, Scientific Committee on Antarctic Research.

4. PetroChron Antarctica Data and Applications

4.1. Data Statistics

Igneous rocks included in PetroChron Antarctica correspond to 60% of the total entries, followed by 39% for metamorphic rocks (Figures 3a and 3b). Sedimentary rocks are poorly represented at only 1%. Igneous rocks are mainly represented by plutonic rocks (42%), whereas metamorphic rocks are dominated by metaplutonic varieties (20%). A large proportion (38%) of igneous rocks are mafic in composition, followed by those of felsic (29%) and intermediate (24%) compositions (Figure 3d). Metamorphic and sedimentary rocks are dominated by felsic compositions (42% and 39%, respectively). Overall, the compositional range across all sampled rocks recorded in PetroChron Antarctica compared with the global whole-rock geochemical database (Gard et al., 2019) is similar (Figures 3c and 3d), when excluding samples marked as oceanic from the global data set.

Computed properties in PetroChron Antarctica include lithology based on chemical classification (Figure S1 in Supporting Information S1). There is a clear dominance of granitoid (32%) and gabbroic rocks (22%). Dioritic and syenitoid compositions (including geochemically equivalent volcanic rocks) are also a significant proportion of the igneous rocks (19% and 16%, respectively). Other computed geochemical indices include ASI, WIP, CIA, or CPA that are often used in soil science as a proxy for alteration/weathering conditions of sampled rocks (see the full list of computed indices in Table 2). Petrophysical properties (density, p- and s-wave velocity, thermal conductivity, and heat production) were computed from geochemical data, following the method described in Hasterok et al. (2018) and Jennings et al. (2019).

4.2. Visualizations and Applications

To illustrate the versatility and the utility of PetroChron Antarctica, we describe below some applications that could use interrelated data sets (i.e., geological, geochemical, and geochronological data associated with rock properties) to gain insights through map visualization.

Figure 4 shows a set of maps illustrating some of the geochronological components of PetroChron Antarctica. For example, the “crystallization age” map (Figure 4a), based on zircon U-Pb isotopic data and typically interpreted to date high-temperature magmatic processes, highlights the dominance of Phanerozoic crust-forming events in the Antarctic Peninsula and Transantarctic Mountains (e.g., Allibone & Wysoczanski, 2002; Burgess et al., 2015; Goodge et al., 2012; Hagen-Peter & Cottle, 2016; Pankhurst et al., 1998; Riley et al., 2017; Zheng et al., 2018). In contrast, the majority of East Antarctic crust formed during the Proterozoic and Archean (Figure 4a; e.g., Adachi et al., 2013; Boger et al., 2006; Corvino et al., 2008; Elburg et al., 2015, 2016; Goodge & Fanning, 2010; Grew et al., 2012; Hokada et al., 2019; Liu et al., 2016; Maritati et al., 2019; Mikhalsky et al., 2017; Morrissey et al., 2017; Tsunogae et al., 2016; Tucker et al., 2017; Zhang et al., 2012), including some of the oldest rocks on Earth (c. 3.9 Ga in Enderby Land; e.g., Black et al., 1986).

A “metamorphic age” map (Figure 4b) based on U-Pb and Sm-Nd isotopic data from zircon, monazite, garnet and whole-rock samples, show the predominance of late Neoproterozoic–Cambrian (~630–500 Ma) ages in the Transantarctic Mountains, Dronning Maud Land, MacRobertson Land, and Princess Elizabeth Land (e.g., Baba et al., 2015; Bisnath et al., 2006; Board et al., 2005; De Vries Van Leeuwen et al., 2019; Goodge & Fanning, 2016; Halpin et al., 2007; Jacobs et al., 2003; Kawakami et al., 2017; Liu et al., 2018; Mikhalsky et al., 2013; Morrissey et al., 2016; Wang et al., 2016). These tectonothermal events record prolonged ocean




Figure 3. Sample rock type and composition. (a) Sample distribution colored by rock type. (b) Bar chart representing rock type. (c) Compositional distribution colored by SiO₂ wt.% content. (d) Comparison of SiO₂ wt.% content between the global whole-rock geochemical database (Gard et al., 2019) and PetroChron Antarctica.

closure, terrane accretion and collision-related processes related to Gondwana amalgamation and active margin tectonics (e.g., Boger, 2011; Fitzsimons, 2003; Goodge, 2020; Harley et al., 2013; Jacobs et al., 2015; Jordan et al., 2020; Mulder et al., 2019).

A map of “cooling ages” (Figure 4c), recorded by low-temperature thermochronology across numerous minerals and whole-rock samples, is dominated by ages <600 Ma (84% of ages recorded by fission track, Ar-Ar, He). The youngest cooling ages (~140–30 Ma with a larger proportion of Paleogene ages) are located along the elevated Transantarctic Mountains (e.g., Fitzgerald & Stump, 1997; Foland et al., 1993; Gleadow & Fitzgerald, 1987; Prenzel et al., 2018; Zattin et al., 2014), whereas East Antarctica records a predominance of late Carboniferous–Triassic (~340–200 Ma) ages and to a lesser extent Cretaceous ages (e.g., Rolland et al., 2019; Sirevaag et al., 2018). The variability in spatial and temporal cooling patterns across Antarctica, although poorly documented, has fueled debate about whether topographic relief evolved




Figure 4. PetroChron Antarctica isotopic age/composition distributions. Maps of (a) zircon crystallization ages; (b) metamorphic ages for different minerals/whole rock; and (c) cooling ages for different minerals/whole rock. The color scale follows the GeoMAP (Cox et al., 2019) chronostratigraphic chart and highlights the variability in isotopic age within the mapped geological units. Dashed rectangles show the location of inset maps.




Figure 5. Computed physical property estimate distributions including (a) density; (b) P-wave velocity; and (c) heat production. Histograms compare distributions for the global whole-rock geochemical database (Gard et al., 2019) and PetroChron Antarctica.

via continental-scale tectonic and/or climatic processes during the Phanerozoic (e.g., Maritati et al., 2020; Rolland et al., 2019).

Collectively, geochronological and isotopic data from across Antarctica reveal major episodes of tectonic and thermal activity, as well as denudation and deposition associated with complex crustal forming processes operating during at least three supercontinent cycles (i.e., Nuna, Rodinia, and Gondwana/Pangea). As such, this database provides a valuable resource for testing possible links between plate tectonic configurations, major climatic and paleoenvironmental change and Antarctic landscape evolution.

Figure 5 shows a map of rock properties computed from geochemical data across Antarctica. Density estimates peak at $\sim 2,630$ and $\sim 2,930$ kg m^{-3} , and P-wave seismic velocity estimates peak at ~ 6.2 and ~ 7.0 km s^{-1} , corresponding to felsic and mafic rock compositions, respectively. These values agree with the densities (2,690 and 2,950 kg m^{-3}) and velocities (~ 6.1 and 7.1 km s^{-1}) recorded in the global whole-rock geochemical database (Gard et al., 2019), when calculated from the same bin size. Antarctic heat production has a median value of ~ 1.3 $\mu\text{W m}^{-3}$, with first and third quartiles at 0.6 and 2.4 $\mu\text{W m}^{-3}$ (Figure 5c), which is higher than the value of 1.0 $\mu\text{W m}^{-3}$ estimated by Gard et al. (2019), who included oceanic samples. At a regional and local scale, crustal heat production shows a high degree of heterogeneity (Figure 5c) due to the high variability of Antarctic local geology (Carson et al., 2014; Goode, 2018) that can be integrated into geothermal heat flow models (Stål et al., 2021). This compositional variability clearly highlights the need to include robust and petrologically valid constraints from direct measurements in geophysical interpretations and numerical computations (Stål et al., 2020).

4.3. Accuracy and Ownership

Although we have made every effort to ensure accuracy when collating information from databases and individual publications, we have undoubtedly inherited or introduced some errors. There are certainly omissions. For example, for any reference or sample, whereas geochemical information may be included in PetroChron Antarctica, accompanying geochronological/isotopic data may not (and vice versa). We strongly advise researchers to revisit the original publications to validate the data for their own use. We encourage users to contact us when they find errors or omissions. Ownership of these data remains with the original authors, and users must cite the relevant original reference(s) and/or data sources as appropriate. In addition to the summary information in the “reference” and “data_source” fields, we provide a list of references in Supporting Information S1.

5. Future Work

We hope the PetroChron Antarctica database can be applied and integrated across Antarctic Earth-cryosphere-biosphere-ocean research. Future work will aim at expanding the database by incorporating not yet considered and newly published data, as well as correcting any errors and adding new data types including metamorphism, protolith, and data-quality parameters. We also invite researchers to collaborate on our data compilation using the user input XLSX template (Table S1 in Supporting Information S1), or by contacting the corresponding author directly.

Data Availability Statement

The PetroChron Antarctica database is available on Zenodo (<https://doi.org/10.5281/zenodo.5032026>) and through the PetroChron Antarctica web portal (an ESRI Web Feature Service; <http://bit.ly/petrochron>). Future versions of the database will be updated at both these locations. The service copies the current data model and helps visualize the distribution of the data. The complete database file in a CSV format can be directly downloaded from the PetroChron Antarctica web portal and Zenodo, or as subset data tables that can be used in any Relational Database Management System (RDBMS) through Zenodo. Code to reproduce figures in this paper is available here: <https://github.com/TobbeTripitaka/PetroChron>.

Acknowledgments

This research was supported by the Australian Research Council Discovery Project DP180104074 and the Australian Research Council Special Research Initiative for Antarctic Gateway Partnership (SR140300001). The authors thank Chris Carson (GA), Marlina Elburg (UJ), Simon Cox (GNS), and Laura Morrissey (UniSA) for providing data sets and/or personal compilations, Sally Watson (NIWA) for early contributions to the database, and Josh Gore (UniSA) for early contributions to the web portal. The authors thank Adam Martin and an anonymous reviewer for their encouraging and constructive feedback. The authors also acknowledge the support and motivation of the SCAR Geosciences Group, including the INSTANT, SERCE, and PAIS Scientific Research Programmes and the GHF subgroup.

References

- Adachi, T., Osanai, Y., Hokada, T., Nakano, N., Baba, S., & Toyoshima, T. (2013). Timing of metamorphism in the central Sør Rondane Mountains, eastern Dronning Maud Land, East Antarctica: Constraints from SHRIMP zircon and EPMA monazite dating. *Precambrian Research*, 234, 136–160. <https://doi.org/10.1016/j.precamres.2012.11.011>
- Allibone, A., & Wysoczanski, R. (2002). Initiation of magmatism during the Cambrian–Ordovician Ross orogeny in southern Victoria Land, Antarctica. *GSA Bulletin*, 114(8), 1007–1018. [https://doi.org/10.1130/0016-7606\(2002\)114<1007:iomdtc>2.0.co;2](https://doi.org/10.1130/0016-7606(2002)114<1007:iomdtc>2.0.co;2)
- Baba, S., Horie, K., Hokada, T., Owada, M., Adachi, T., & Shiraishi, K. (2015). Multiple Collisions in the East African–Antarctica Orogen: Constraints from Timing of Metamorphism in the Filchnerfjella and Hochlinfjellet Terranes in Central Dronning Maud Land. *The Journal of Geology*, 123(1), 55–77. <https://doi.org/10.1086/679468>
- Bisnath, A., Frimmel, H. E., Armstrong, R. A., & Board, W. S. (2006). Tectono-thermal evolution of the Maud Belt: New SHRIMP U-Pb zircon data from Gjelsvikfjella, Dronning Maud Land, East Antarctica. *Precambrian Research*, 150(1–2), 95–121. <https://doi.org/10.1016/j.precamres.2006.06.009>
- Black, L. P., Williams, I. S., & Compston, W. (1986). Four zircon ages from one rock: The history of a 3930 Ma-old granulite from Mount Sones, Enderby Land, Antarctica. *Contributions to Mineralogy and Petrology*, 94, 427–437. <https://doi.org/10.1007/BF00376336>
- Board, W. S., Frimmel, H. E., & Armstrong, R. A. (2005). Pan-African tectonism in the Western Maud Belt: P-T-t path for high-grade gneisses in the H.U. Sverdrupfjella, East Antarctica. *Journal of Petrology*, 46(4), 671–699. <https://doi.org/10.1093/ptrology/egh093>
- Boger, S. D. (2011). Antarctica – Before and after Gondwana. *Gondwana Research*, 19(2), 335–371. <https://doi.org/10.1016/j.gr.2010.09.003>
- Boger, S. D., Wilson, C. J. L., & Mark Fanning, C. (2006). An Archaean province in the southern Prince Charles Mountains, East Antarctica: U-Pb zircon evidence for c. 3170 Ma granite plutonism and c. 2780 Ma partial melting and orogenesis. *Precambrian Research*, 145(3–4), 207–228. <https://doi.org/10.1016/j.precamres.2005.12.003>
- Burgess, S. D., Bowring, S. A., Fleming, T. H., & Elliot, D. H. (2015). High-precision geochronology links the Ferrar large igneous province with early-Jurassic ocean anoxia and biotic crisis. *Earth and Planetary Science Letters*, 415, 90–99. <https://doi.org/10.1016/j.epsl.2015.01.037>
- Burton-Johnson, A., Dziadek, R., & Martin, C. (2020). Review article: Geothermal heat flow in Antarctica: Current and future directions. *The Cryosphere*, 14(11), 3843–3873. <https://doi.org/10.5194/tc-14-3843-2020>
- Carson, C. J., McLaren, S., Roberts, J. L., Boger, S. D., & Blankenship, D. D. (2014). Hot rocks in a cold place: High sub-glacial heat flow in East Antarctica. *Journal of the Geological Society*, 171(1), 9–12. <https://doi.org/10.1144/jgs2013-030>
- Champion, D. C., Budd, A. R., Hazell, M. S., & Sedgmen, A. (2007). *OZCHEM National Whole Rock Geochemistry Dataset*. Retrieved from <http://pid.geoscience.gov.au/dataset/ga/65464>; <https://researchdata.andcs.org.au/ozchem-national-rock-geochemistry-dataset>
- Corvino, A. F., Boger, S. D., Henjes-Kunst, F., Wilson, C. J. L., & Fitzsimons, I. C. W. (2008). Superimposed tectonic events at 2450 Ma, 2100 Ma, 900 Ma and 500 Ma in the North Mawson Escarpment, Antarctic Prince Charles Mountains. *Precambrian Research*, 167(3–4), 281–302. <https://doi.org/10.1016/j.precamres.2008.09.001>
- Cox, S. C., Smith Lyttle, B., & the GeoMAP team. (2019). *SCAR GeoMAP dataset*. GNS Science. Release v.201907. <https://doi.org/10.21420/7SH7-6K05>
- Debon, F., & Le Fort, P. (1983). A chemical–mineralogical classification of common plutonic rocks and associations. *Transactions of the Royal Society of Edinburgh Earth Sciences*, 73(3), 135–149. <https://doi.org/10.1017/S0263593300010117>
- De la Roche, H., Leterrier, J., Grandclaude, P., & Marchal, M. (1980). A classification of volcanic and plutonic rocks using R1R2-diagram and major-element analyses — Its relationships with current nomenclature. *Chemical Geology*, 29(1), 183–210. [https://doi.org/10.1016/0009-2541\(80\)90020-0](https://doi.org/10.1016/0009-2541(80)90020-0)
- De Vries Van Leeuwen, A. T., Morrissey, L. J., Kelsey, D. E., & Raimondo, T. (2019). Recognition of Pan-African-aged metamorphism in the Fisher Terrane, central Prince Charles Mountains, East Antarctica. *Journal of the Geological Society*, 176(4), 785–798. <https://doi.org/10.1144/jgs2018-146>
- Eglinton, B. M. (2004). DateView: A windows geochronology database. *Computers & Geosciences*, 30(8), 847–858. <https://doi.org/10.1016/j.cageo.2004.06.002>
- Elburg, M. A., Andersen, T., Jacobs, J., Läufer, A., Ruppel, A., Krohne, N., & Damaske, D. (2016). One hundred fifty million years of intrusive activity in the Sør Rondane Mountains (East Antarctica): Implications for Gondwana Assembly. *The Journal of Geology*, 124(1), 1–26. <https://doi.org/10.1086/684052>
- Elburg, M. A., Jacobs, J., Andersen, T., Clark, C., Läufer, A., Ruppel, A., et al. (2015). Early Neoproterozoic metagabbro-tonalite-trondhjemite of Sør Rondane (East Antarctica): Implications for supercontinent assembly. *Precambrian Research*, 259, 189–206. <https://doi.org/10.1016/j.precamres.2014.10.014>
- Fitzgerald, P. G., & Stump, E. (1997). Cretaceous and Cenozoic episodic denudation of the Transantarctic Mountains, Antarctica: New constraints from apatite fission track thermochronology in the Scott Glacier region. *Journal of Geophysical Research*, 102(B4), 7747–7765. <https://doi.org/10.1029/96jb03898>
- Fitzsimons, I. C. W. (2003). Proterozoic basement provinces of southern and southwestern Australia, and their correlation with Antarctica. *Geological Society, London, Special Publications*, 206(1), 93–130. <https://doi.org/10.1144/GSL.SP.2003.206.01.07>
- Foland, K. A., Fleming, T. H., Heimann, A., & Elliot, D. H. (1993). Potassium-argon dating of fine-grained basalts with massive Ar loss: Application of the ⁴⁰Ar/³⁹Ar technique to plagioclase and glass from the Kirkpatrick Basalt, Antarctica. *Chemical Geology*, 107(1), 173–190. [https://doi.org/10.1016/0009-2541\(93\)90109-V](https://doi.org/10.1016/0009-2541(93)90109-V)
- Frost, B. R., Barnes, C. G., Collins, W. J., Arculus, R. J., Ellis, D. J., & Frost, C. D. (2001). A geochemical classification for granitic rocks. *Journal of Petrology*, 42(11), 2033–2048. <https://doi.org/10.1093/ptrology/42.11.2033>
- Gard, M., Hasterok, D., & Halpin, J. A. (2019). Global whole-rock geochemical database compilation. *Earth System Science Data*, 11(4), 1553–1566. <https://doi.org/10.5194/essd-11-1553-2019>
- Gleadow, A. J. W., & Fitzgerald, P. G. (1987). Uplift history and structure of the Transantarctic Mountains: New evidence from fission track dating of basement apatites in the Dry Valleys area, southern Victoria Land. *Earth and Planetary Science Letters*, 82(1), 1–14. [https://doi.org/10.1016/0012-821X\(87\)90102-6](https://doi.org/10.1016/0012-821X(87)90102-6)
- Goode, J. W. (2018). Crustal heat production and estimate of terrestrial heat flow in central East Antarctica, with implications for thermal input to the East Antarctic ice sheet. *The Cryosphere*, 12(2), 491–504. <https://doi.org/10.5194/tc-12-491-2018>
- Goode, J. W. (2020). Geological and tectonic evolution of the Transantarctic Mountains, from ancient craton to recent enigma. *Gondwana Research*, 80, 50–122. <https://doi.org/10.1016/j.gr.2019.11.001>
- Goode, J. W., & Fanning, C. M. (2010). Composition and age of the East Antarctic Shield in eastern Wilkes Land determined by proxy from Oligocene–Pleistocene glaciomarine sediment and Beacon Supergroup sandstones, Antarctica. *The Geological Society of America Bulletin*, 122(7–8), 1135–1159. <https://doi.org/10.1130/B30079.1>

- Goodge, J. W., & Fanning, C. M. (2016). Mesoarchean and paleoproterozoic history of the Nimrod Complex, central Transantarctic Mountains, Antarctica: Stratigraphic revisions and relation to the Mawson Continent in East Gondwana. *Precambrian Research*, 285, 242–271. <https://doi.org/10.1016/j.precamres.2016.09.001>
- Goodge, J. W., Fanning, C. M., Norman, M. D., & Bennett, V. C. (2012). Temporal, isotopic and spatial relations of early Paleozoic Gondwana-margin arc magmatism, central Transantarctic Mountains, Antarctica. *Journal of Petrology*, 53(10), 2027–2065. <https://doi.org/10.1093/ptrology/egs043>
- Grew, E. S., Carson, C. J., Christy, A. G., Maas, R., Yaxley, G. M., Boger, S. D., & Fanning, C. M. (2012). New constraints from U–Pb, Lu–Hf and Sm–Nd isotopic data on the timing of sedimentation and felsic magmatism in the Larsemann Hills, Prydz Bay, East Antarctica. *Precambrian Research*, 206–207, 87–108. <https://doi.org/10.1016/j.precamres.2012.02.016>
- Hagen-Peter, G., & Cottle, J. M. (2016). Synchronous alkaline and subalkaline magmatism during the late Neoproterozoic–early Paleozoic Ross orogeny, Antarctica: Insights into magmatic sources and processes within a continental arc. *Lithos*, 262, 677–698. <https://doi.org/10.1016/j.lithos.2016.07.032>
- Halpin, J. A., Clarke, G. L., White, R. W., & Kelsey, D. E. (2007). Contrasting P–T–t paths for Neoproterozoic metamorphism in MacRobertson and Kemp Lands, east Antarctica. *Journal of Metamorphic Geology*, 25, 683–701. <https://doi.org/10.1111/j.1525-1314.2007.00723.x>
- Harley, S. L., Fitzsimons, I. C. W., & Zhao, Y. (2013). Antarctica and supercontinent evolution: Historical perspectives, recent advances and unresolved issues. *Geological Society, London, Special Publications*, 383, 1–34. <https://doi.org/10.1144/SP383.9>
- Hasterok, D., Gard, M., & Webb, J. (2018). On the radiogenic heat production of metamorphic, igneous, and sedimentary rocks. *Geoscience Frontiers*, 9(6), 1777–1794. <https://doi.org/10.1016/j.gsf.2017.10.012>
- Hasterok, D., & Webb, J. (2017). On the radiogenic heat production of igneous rocks. *Geoscience Frontiers*, 8(5), 919–940. <https://doi.org/10.1016/j.gsf.2017.03.006>
- Hochmuth, K., Gohl, K., Leitchenkov, G., Sauermilch, I., Whittaker, J. M., Uenzelmann-Neben, G., et al. (2020). The evolving paleobathymetry of the circum-Antarctic Southern Ocean Since 34 Ma: A key to understanding past cryosphere-ocean developments. *Geochemistry, Geophysics, Geosystems*, 21(8). <https://doi.org/10.1029/2020GC009122>
- Hokada, T., Grantham, G. H., Arima, M., Saito, S., Shiraishi, K., Armstrong, R. A., et al. (2019). Stenian A-type granitoids in the Namaqua-Natal Belt, southern Africa, Maud Belt, Antarctica and Nampula Terrane, Mozambique: Rodinia and Gondwana amalgamation implications. *Geoscience Frontiers*. <https://doi.org/10.1016/j.gsf.2019.04.003>
- Jacobs, J., Bauer, W., & Fanning, C. M. (2003). Late Neoproterozoic/Early Palaeozoic events in central Dronning Maud Land and significance for the southern extension of the East African Orogen into East Antarctica. *Precambrian Research*, 126(1–2), 27–53. [https://doi.org/10.1016/S0301-9268\(03\)00125-6](https://doi.org/10.1016/S0301-9268(03)00125-6)
- Jacobs, J., Elburg, M., Läufer, A., Kleinhanns, I. C., Henjes-Kunst, F., Estrada, S., et al. (2015). Two distinct Late Mesoproterozoic/Early Neoproterozoic basement provinces in central/eastern Dronning Maud Land, East Antarctica: The missing link, 15–21°E. *Precambrian Research*, 265, 249–272. <https://doi.org/10.1016/j.precamres.2015.05.003>
- Jennings, S., Hasterok, D., & Payne, J. (2019). A new compositionally based thermal conductivity model for plutonic rocks. *Geophysical Journal International*, 219(2), 1377–1394. <https://doi.org/10.1093/gji/ggz376>
- Jordan, T. A., Riley, T. R., & Siddoway, C. S. (2020). The geological history and evolution of West Antarctica. *Nature Reviews Earth & Environment*, 1(2), 117–133. <https://doi.org/10.1038/s43017-019-0013-6>
- Kawakami, T., Higashino, F., Skrzypek, E., Satish-Kumar, M., Grantham, G., Tsuchiya, N., et al. (2017). Prograde infiltration of Cl-rich fluid into the granulitic continental crust from a collision zone in East Antarctica (Perlebandet, Sør Rondane Mountains). *Lithos*, 274–275, 73–92. <https://doi.org/10.1016/j.lithos.2016.12.028>
- Le Bas, M. J., & Streckeisen, A. L. (1991). The IUGS systematics of igneous rocks. *Journal of the Geological Society*, 148(5), 825–833. <https://doi.org/10.1144/gsjgs.148.5.0825>
- Liu, X., Ling, X., & Jahn, B. M. (2018). U–Th–Pb monazite and Sm–Nd dating of high-grade rocks from the Grove Mountains, East Antarctica: Further evidence for a Pan-African-aged monometamorphic terrane. *Advances in Polar Science*, 29(2), 108–117. <https://doi.org/10.13679/j.advps.2018.2.00108>
- Liu, X., Wang, W., Zhao, Y., Liu, J., Chen, H., Cui, Y., & Song, B. (2016). Early Mesoproterozoic arc magmatism followed by early Neoproterozoic granulite facies metamorphism with a near-isobaric cooling path at Mount Brown, Princess Elizabeth Land, East Antarctica. *Precambrian Research*, 284, 30–48. <https://doi.org/10.1016/j.precamres.2016.08.003>
- Maritati, A., Danišik, M., Halpin, J. A., Whittaker, J. M., & Aitken, A. R. A. (2020). Pangea Rifting Shaped the East Antarctic Landscape. *Tectonics*, 39(8). <https://doi.org/10.1029/2020TC006180>
- Maritati, A., Halpin, J. A., Whittaker, J. M., & Daczko, N. R. (2019). Fingerprinting Proterozoic Bedrock in Interior Wilkes Land, East Antarctica. *Scientific Reports*, 9(1), 10192. <https://doi.org/10.1038/s41598-019-46612-y>
- Middlemost, E. A. K. (1994). Naming materials in the magma/igneous rock system. *Earth-Science Reviews*, 37(3), 215–224. [https://doi.org/10.1016/0012-8252\(94\)90029-9](https://doi.org/10.1016/0012-8252(94)90029-9)
- Mikhalsky, E. V., Krylov, D., Rodionov, N., Presnyakov, S., Skublov, S., & Myasnikov, O. (2017). Refined geological history of the polyphase plutonometamorphic complex in the Thala Hills area (Enderby Land, East Antarctica) from zircon SHRIMP dating and implications for Neoproterozoic amalgamation of Gondwanaland. *Geological Society, London, Special Publications*, 457(1), 7–36. <https://doi.org/10.1144/SP457.2>
- Mikhalsky, E. V., Sheraton, J. W., Kudriavtsev, I. V., Sergeev, S. A., Kovach, V. P., Kamenev, I. A., & Laiba, A. A. (2013). The Mesoproterozoic Rayner Province in the Lambert Glacier area: Its age, origin, isotopic structure and implications for Australia–Antarctica correlations. *Geological Society, London, Special Publications*, 383, 35–57. <https://doi.org/10.1144/SP383.1>
- Morrissey, L. J., Hand, M., Kelsey, D. E., & Wade, B. P. (2016). Cambrian high-temperature reworking of the Rayner-Eastern Ghats terrane: Constraints from the Northern Prince Charles Mountains region, East Antarctica. *Journal of Petrology*, 57(1), 53–91. <https://doi.org/10.1093/ptrology/egv082>
- Morrissey, L. J., Payne, J. L., Hand, M., Clark, C., Taylor, R., Kirkland, C. L., & Kylander-Clark, A. (2017). Linking the Windmill Islands, east Antarctica and the Albany–Fraser Orogen: Insights from U–Pb zircon geochronology and Hf isotopes. *Precambrian Research*, 293, 131–149. <https://doi.org/10.1016/j.precamres.2017.03.005>
- Mulder, J. A., Halpin, J. A., Daczko, N. R., Orth, K., Meffre, S., Thompson, J. M., & Morrissey, L. J. (2019). A Multiproxy provenance approach to uncovering the assembly of East Gondwana in Antarctica. *Geology*, 47(7), 645–649. <https://doi.org/10.1130/G45952.1>
- Nesbitt, H. W., & Young, G. M. (1989). Formation and diagenesis of weathering profiles. *The Journal of Geology*, 97(2), 129–147. <https://doi.org/10.1086/629290>
- Pankhurst, R. J., Weaver, S. D., Bradshaw, J. D., Storey, B. C., & Ireland, T. R. (1998). Geochronology and geochemistry of pre-Jurassic superterranes in Marie Byrd Land, Antarctica. *Journal of Geophysical Research*, 103(B2), 2529–2547. <https://doi.org/10.1029/97jb02605>
- Parker, A. (1970). An index of weathering for silicate rocks. *Geological Magazine*, 107(6), 501–504. <https://doi.org/10.1017/S0016756800058581>

- Paxman, G. J. G., Gasson, E. G. W., Jamieson, S. S. R., Bentley, M. J., & Ferraccioli, F. (2020). Long-term increase in Antarctic ice sheet vulnerability driven by bed topography evolution. *Geophysical Research Letters*, *47*(20). <https://doi.org/10.1029/2020GL090003>
- Prenzel, J., Lisker, F., Monsees, N., Balestrieri, M. L., Läufer, A., & Spiegel, C. (2018). Development and inversion of the Mesozoic Victoria Basin in the Terra Nova Bay (Transantarctic Mountains) derived from thermochronological data. *Gondwana Research*, *53*, 110–128. <https://doi.org/10.1016/j.gr.2017.04.025>
- Riley, T. R., Flowerdew, M. J., Pankhurst, R. J., Curtis, M. L., Millar, I. L., Fanning, C. M., & Whitehouse, M. J. (2017). Early Jurassic magmatism on the Antarctic Peninsula and potential correlation with the Subcordilleran plutonic belt of Patagonia. *Journal of the Geological Society*, *174*(2), 365–376. <https://doi.org/10.1144/jgs2016-053>
- Rolland, Y., Bernet, M., van der Beek, P., Gautheron, C., Duclaux, G., Bascou, J., et al. (2019). Late Paleozoic Ice Age glaciers shaped East Antarctica landscape. *Earth and Planetary Science Letters*, *506*, 123–133. <https://doi.org/10.1016/j.epsl.2018.10.044>
- Rybach, L. (1988). Determination of heat production rate. In R. Hänel, L. Rybach, & I. Stegena (Eds.), *Handbook of terrestrial heat flow density determination* (Vol. 486, pp. 125–142). Kluwer Academic Publishers.
- Secretariat SCAR. (1992, updated 2014). Composite Gazetteer of Antarctica, Scientific Committee on Antarctic Research. GCMD Metadata. Retrieved from http://gcmd.nasa.gov/records/SCAR_Gazetteer.html
- Sirevaag, H., Jacobs, J., Ksienzyk, A. K., Dunkl, I., & Marshall, H. R. (2018). Extent, thickness and erosion of the Jurassic continental flood basalts of Dronning Maud Land, East Antarctica: A low-T thermochronological approach. *Gondwana Research*, *61*, 222–243. <https://doi.org/10.1016/j.gr.2018.04.017>
- Stål, T., Reading, A. M., Halpin, J. A., Phipps, S. J., & Whittaker, J. M. (2020). The Antarctic crust and upper mantle: A flexible 3D model and software framework for interdisciplinary research. *Frontiers of Earth Science*, *8*(447). <https://doi.org/10.3389/feart.2020.577502>
- Stål, T., Reading, A. M., Halpin, J. A., & Whittaker, J. M. (2021). Antarctic geothermal heat flow model: Aq1. *Geochemistry, Geophysics, Geosystems*, *22*(2), e2020GC009428. <https://doi.org/10.1029/2020GC009428>
- Strong, D. T., Turnbull, R. E., Haubrock, S., & Mortimer, N. (2016). Petlab: New Zealand's national rock catalogue and geoanalytical database. *New Zealand Journal of Geology and Geophysics*, *59*(3), 475–481. <https://doi.org/10.1080/00288306.2016.1157086>
- Tsunogae, T., Yang, Q.-Y., & Santosh, M. (2016). Neoproterozoic–Early Paleoproterozoic and Early Neoproterozoic arc magmatism in the Lützow–Holm Complex, East Antarctica: Insights from petrology, geochemistry, zircon U–Pb geochronology and Lu–Hf isotopes. *Lithos*, *263*, 239–256. <https://doi.org/10.1016/j.lithos.2016.02.010>
- Tucker, N. M., Payne, J. L., Clark, C., Hand, M., Taylor, R. J., et al. (2017). Proterozoic reworking of Archean (Yilgarn) basement in the Bunge Hills, East Antarctica. *Precambrian Research*, *298*, 16–38. <https://doi.org/10.1016/j.precamres.2017.05.013>
- Wang, W., Liu, X., Zhao, Y., Zheng, G., & Chen, L. (2016). U–Pb zircon ages and Hf isotopic compositions of metasedimentary rocks from the Grove Subglacial Highlands, East Antarctica: Constraints on the provenance of protoliths and timing of sedimentation and metamorphism. *Precambrian Research*, *275*, 135–150. <https://doi.org/10.1016/j.precamres.2015.12.018>
- Whitehouse, P. L., Gomez, N., King, M. A., & Wiens, D. A. (2019). Solid Earth change and the evolution of the Antarctic Ice Sheet. *Nature Communications*, *10*(1), 503. <https://doi.org/10.1038/s41467-018-08068-y>
- Zattin, M., Pace, D., Andreucci, B., Rossetti, F., & Talarico, F. M. (2014). Cenozoic erosion of the Transantarctic Mountains: A source-to-sink thermochronological study. *Tectonophysics*, *630*, 158–165. <https://doi.org/10.1016/j.tecto.2014.05.022>
- Zhang, S. H., Zhao, Y., Liu, X. C., Liu, Y. S., Hou, K. J., Li, C. F., & Ye, H. (2012). U–Pb geochronology and geochemistry of the bedrocks and moraine sediments from the Windmill Islands: Implications for Proterozoic evolution of East Antarctica. *Precambrian Research*, *206*–207, 52–71. <https://doi.org/10.1016/j.precamres.2012.02.019>
- Zheng, G.-G., Liu, X., Liu, S., Zhang, S.-H., & Zhao, Y. (2018). Late Mesozoic–early Cenozoic intermediate–acid intrusive rocks from the Gerlache Strait area, Antarctic Peninsula: Zircon U–Pb geochronology, petrogenesis and tectonic implications. *Lithos*, *312*–313, 204–222. <https://doi.org/10.1016/j.lithos.2018.05.008>