

Article (refereed)

Thomas, Philippe J.; Mineau, Pierre; **Shore, Richard F.**; Champoux, Louise; Martin, Pamela A.; Wilson, Laurie K.; Fitzgerald, Guy; Elliott, John E.. 2011 Second generation anticoagulant rodenticides in predatory birds: Probabilistic characterisation of toxic liver concentrations and implications for predatory bird populations in Canada. *Environment International*, 37 (5). 914-920. [10.1016/j.envint.2011.03.010](https://doi.org/10.1016/j.envint.2011.03.010)

Copyright © 2011 Elsevier Ltd.

This version available <http://nora.nerc.ac.uk/14168/>

NERC has developed NORA to enable users to access research outputs wholly or partially funded by NERC. Copyright and other rights for material on this site are retained by the authors and/or other rights owners. Users should read the terms and conditions of use of this material at <http://nora.nerc.ac.uk/policies.html#access>

This document is the author's final manuscript version of the journal article prior to the peer review process. Some differences between this and the publisher's version may remain. You are advised to consult the publisher's version if you wish to cite from this article.

www.elsevier.com/

Contact CEH NORA team at
noraceh@ceh.ac.uk

Title: Second generation anticoagulant rodenticides in predatory birds: probabilistic characterisation of toxic liver concentrations and implications for predatory bird populations in Canada.

Authors: Philippe J. Thomas¹, Pierre Mineau¹, Richard F. Shore², Louise Champoux³, Pamela A. Martin⁴, Laurie K. Wilson⁵, Guy Fitzgerald⁶, John E. Elliott⁷

¹ National Wildlife Research Centre, Science & Technology Branch, Environment Canada, Ottawa, Ontario, K1A 0H3, Canada, philippe.thomas@ec.gc.ca (corresponding author), pierre.mineau@ec.gc.ca,

Tel: (613) 998-0511 Fax: (613) 998-0458

² Centre for Ecology & Hydrology, Lancaster Environment Centre, Library Avenue, Bailrigg, Lancaster LA1 4AP, UK, rfs@ceh.ac.uk

³ Science & Technology Branch, Environment Canada, Ste-Foy, QC, G1V 4H5, Canada, louise.champoux@ec.gc.ca

⁴ Science & Technology Branch, Environment Canada, Burlington, ON, L7R 4A6, Canada, pamela.martin@ec.gc.ca

⁵ Canadian Wildlife Service, Environment Canada, Delta, BC, V4K 3N2, Canada, laurie.wilson@ec.gc.ca

⁶ Université de Montréal, Saint-Hyacinthe, QC, J2S 7C6, Canada, guy.fitzgerald@umontreal.ca

⁷ Science & Technology Branch, Environment Canada, Delta, BC, V4K 3N2, Canada, john.elliott@ec.gc.ca

Abstract:

Second-generation anticoagulant rodenticides (SGARs) are widely used to control rodent pests but exposure and poisonings occur in non-target species, such as birds of prey. Liver residues are often analyzed to detect exposure in birds found dead but their use to assess toxicity of SGARs is problematic. We analyzed published data on hepatic rodenticide residues and associated symptoms of anticoagulant poisoning from 270 birds of prey using logistic regression to estimate the probability of toxicosis associated with different liver SGAR residues. We also evaluated exposure to SGARs on a national level in Canada by analyzing 196 livers from great horned owls (*Bubo virginianus*) and red-tailed hawks (*Buteo jamaicensis*) found dead at locations across the country. Analysis of a broader sample of raptor species from Quebec also helped define the taxonomic breadth of contamination. Calculated probability curves suggest significant species differences in sensitivity to SGARs and significant likelihood of toxicosis below previously suggested concentrations of concern (<0.1 mg/kg). Analysis of birds from Quebec showed that a broad range of raptor species are exposed to SGARs, indicating that generalized terrestrial food chains could be contaminated in the vicinity of the sampled areas. Of the two species for which we had samples from across Canada, great horned owls are exposed to SGARs to a greater extent than red-tailed hawks and liver residue levels were also higher. Using our probability estimates of effect, we estimate that a minimum of 11% of the sampled great horned owl population is at risk of being directly killed by SGARs. This is the first time the potential mortality impact of SGARs on a raptor population has been estimated.

Keywords: rodent, exposure, liver residues, toxicity threshold, anticoagulant rodenticide

SGAR – second generation anticoagulant rodenticide

GHOW – great horned owl

RTHA – red-tailed hawk

BAOW – barred owl

BNOW – barn owl

ALL – all bird species pooled

EIIS – ecological incident information system

EPA – environmental protection agency

1 **1. Introduction**

2 Introduced in the 1970s, second-generation anticoagulant rodenticides (SGARs) were
3 developed to combat the reported development of rodent resistance to first-generation
4 compounds (Buckle et al. 1994). These newer anticoagulant poisons differ from their
5 first-generation counterparts in that they are more acutely toxic at lower doses (often
6 allowing a lethal dose to be obtained in a single feeding), and are more persistent in
7 vertebrate livers (Parmar et al. 1987, Stone et al. 1999, Newton et al. 1999, Erickson and
8 Urban 2004). Greater acute toxicity increases the potential for primary poisoning
9 amongst non-target species while the longer tissue half-lives of SGARs enhance the
10 potential for bioaccumulation in non-target predators in particular, and so may increase
11 the risk of secondary poisoning. Furthermore, rodents survive for several days after
12 consuming a lethal dose of SGARs and often will continue feeding on the bait (Cox and
13 Smith 1992). That increases the likelihood that the body burden in poisoned rodents may
14 significantly exceed the LD50 or even LD100 dose, and poisoned animals may remain
15 active and available for capture by predators for some period after ingestion of the
16 rodenticide. Additionally, poisoned rodents exhibit an altered state of behaviour, such as
17 spending more time in open areas in a lethargic state, and this may further predispose
18 them to predation (Cox and Smith 1992).

19 SGARs bind and inhibit vitamin K epoxide reductase and persist for at least six
20 months in organs and tissues containing this enzyme such as the liver (Stone et al. 1999,
21 Eason et al. 2002). In an attempt to monitor exposure in non-target wildlife, the presence
22 of detectable SGAR residues as well as the magnitude of concentrations has been
23 measured in the livers of some Canadian, American and European predatory birds and

24 scavengers (Albert et al. 2010, Newton et al. 1990, Shore et al. 1999, Shore et al. 2006).
25 There was a common trend among those studies for most SGARs, namely brodifacoum,
26 bromadiolone, difenacoum and difethialone being detected at an increasing frequency in
27 numerous predators and scavengers. Species most commonly monitored in North
28 America are great horned owls (*Bubo virginianus*) and red-tailed hawks (*Buteo*
29 *jamaicensis*) (Albert et al. 2010, Erickson and Urban 2004).

30 It is still uncertain what SGAR liver concentration is diagnostic of a potentially lethal
31 dose and, indeed Erickson and Urban (2004) have questioned whether such a cause-effect
32 relationship is appropriate. A sometimes cited –toxicity threshold is given as –greater
33 than 0.1 – 0.2 mg/kg wet weight (Newton et al. 1998, Newton et al. 1999). This was, in
34 fact, described as a –potentially lethal range and was derived for a single species, the
35 barn owl (*Tyto alba*); it stems from two sets of observations (Shore et al. 2001). Firstly,
36 barn owls diagnosed post-mortem as having died from rodenticides had liver
37 concentrations > 0.1 mg/kg. Those owls exhibited classical toxicosis signs such as
38 haemorrhaging from organs such as the heart, lungs, liver, brain and/or subcutaneous
39 areas (Newton et al. 1998). Secondly, owls that were experimentally poisoned had liver
40 residues in the range of 0.2 – 1.72 mg/kg (Newton et al. 1999). However, it is uncertain
41 whether these barn owl criteria would apply to other species. Liver residues associated
42 with SGAR poisonings in various species typically range over two orders of magnitude
43 and were reported to be as low as 0.01 mg/kg wet wt in one great horned owl that was
44 examined (Stone et al. 1999). Thus, liver SGAR concentrations associated with toxicity
45 vary markedly among both individuals and species. This suggests a probabilistic
46 approach; which we adopt to review the evidence pertaining to how liver residues are

47 related to toxicity. Our principal objectives are: i) to determine SGAR liver
48 concentrations that may be associated with mortality in birds (ie- to quantify the -toxicity
49 threshold) and ii) using the threshold values, assess the extent and severity of exposure
50 in Canadian birds of prey.

51

52 **2. Methods**

53 **2.1. Toxicity Threshold**

54 2.1.1. Literature Search

55 Recently published (~ last 10 years) peer-reviewed publications as well as the United
56 States Environmental Protection Agency (EPA)'s Ecological Incident Information
57 System (EIS) were surveyed in order to locate liver residue data sets for birds of prey.
58 The EIS is the EPA's database managing information on incidents linked to the exposure
59 of non target plants and animals to pesticides. It is currently managed by the Office of
60 Pesticide Programs (Mastrota 2007). Data were retained for our assessment if they met a
61 set of pre-determined conditions. These conditions included:

- 62 i) SGAR detection limits in liver were under 0.02 mg/kg wet wt;
- 63 ii) post-mortem evaluations were conducted prior to liver extraction and analysis;
64 pathophysiological signs of rodenticide poisoning were included.
- 65 iii) post-mortem evaluations were conducted by a reputable professional such as a
66 doctor of veterinary medicine (DVM); and
- 67 iv) adequate sample sizes were available ($n > 15$) for any given species (in order to
68 have greater statistical power).

69 2.1.2. Data Analysis

70 Raptor necropsies with attending SGAR liver analyses were collected and compiled
71 in database software, and each case was given a binary code as positive (1) or negative
72 (0) for pathophysiological signs of poisoning. A positive coding meant that, after a
73 detailed post-mortem evaluation, an anticoagulant was diagnosed as being the cause of
74 death or a significant contributory factor (ie- when necropsies showed hemorrhage or
75 anemia in the absence of traumatic injury or infectious or parasitic diseases and an
76 anticoagulant residue was detected in the liver). A negative coding represented cases
77 where the cause of death was deemed to be natural or accidental (for example incidental
78 take, hunting, motor vehicle collisions, starvation).

79 The binary dataset was imported into SAS/STAT (version 9.2 TS2M0). Residue
80 concentrations of all SGAR compounds were summed for the logistic regression.
81 Concentrations were log transformed to meet the assumption of normality and re-tested.
82 The PROC LOGISTIC macro was invoked to determine how liver residues affected
83 presence or absence of poisoning symptoms. An effects plot was generated to illustrate
84 the relationship and equations were built for every species with sufficient data ($n \geq 15$).
85 Using these equations, liver residue levels (in mg/kg wet weight (ww)) were determined
86 for probabilities of 5%, 10%, 15% and 20% of exhibiting pathologies consistent with
87 rodenticides exposure. Species comparisons were completed using analysis of variance
88 (ANOVA) in conjunction with Tukey's Studentized Range test. Because all birds were
89 found dead or moribund, there was a logical inference that those pathologies
90 (haemorrhaging of the heart, lungs, liver, brain and/or subcutaneous areas) were
91 responsible for, or strongly contributed to, the mortality of the individual.

92 **2.2. Exposure extent in Canada**

93 2.2.1. Sample Collection

94 To obtain a cross-Canada survey of residue levels, liver samples of birds were
95 selected, irrespective of the cause of death, from British Columbia, the prairie provinces,
96 Ontario and Quebec. The birds were collected near agricultural and urban areas of the
97 country where SGAR use was thought to be common. They were typically submitted to
98 rehabilitation or veterinary centres either dead or in a moribund state. Initial diagnosis
99 frequently involved car strike or other obvious ‘mishap’. They were not chosen because
100 they showed signs of anticoagulant poisoning, but rather reflect the population of
101 reported birds of prey dying from a multitude of causes. The subsequent liver samples
102 were harvested initially as part of previous investigations of exposure to heavy metals or
103 other toxicants, and then rodenticides residues were determined in later years. Three main
104 collections were sampled. These included an Ontario/prairie sample of red-tailed hawks
105 and great-horned owls, two common species known to scavenge; a broader phylogenetic
106 collection from Quebec and a collection of three owl species from British Columbia (barn
107 owl, barred owl [*Strix varia*] and great-horned owl). Those owl species are less mobile
108 than most of the hawk species and were chosen to help identify geographical patterns of
109 contamination and hence, potential sources of rodenticides residues. Results from the
110 latter have already been reported (Albert et al. 2010).

111 2.2.2. Chemical Analysis

112 Chemical analysis was conducted at the National Wildlife Research Center in Ottawa,
113 Ontario, Canada. Methods were similar to those reported by Albert et al. (2010). 50 mg of
114 liver was ground in a mortar with about 5 g anhydrous sodium sulphate (Fisher no. S420-
115 3). The resulting mixture was transferred to an amber glass septum bottle and acetonitrile

116 (EMD Omnisolv, AX0142-1, HPLC grade; 1 x 7 mL and 2 x 5 mL) was used for
117 extraction. The extract was shaken for 2 minutes by hand and 15 minutes mechanically.
118 After centrifuging for 15 minutes at 1000 rpm, the supernatant was removed and
119 transferred into a 40 mL conical tube. The supernatant of the two subsequent extractions
120 were combined with the first supernatant. The total product was evaporated to 10 mL
121 under a stream of nitrogen in a water bath kept at 40°C.

122 In order to clean up liver extract, a 2 mL portion was transferred into a test tube and
123 heated to dryness. The sample was reconstituted in acetonitrile and cleaned by solid-
124 phase extraction. After the introduction of the sample into the SPE cartridge, the tube
125 containing the sample was rinsed with acetonitrile and added to the SPE cartridge
126 solution. The eluate was then evaporated to dryness and reconstituted in MeOH and
127 filtered through an Acrodisc® syringe filter with a polyvinylidene fluoride (PVDF)
128 membrane. A volume of 10 µL of the diluted filtered extract was analyzed by liquid
129 chromatography-mass spectrometry (LC-MSMS). Some of the owl samples analysed
130 (mainly from British Columbia) were not cleaned using an SPE cartridge. However,
131 limits of detection were calculated for the procedure with and without an SPE sample
132 cleaning phase and were found to be identical. For this reason, both SPE-cleaned data and
133 non-SPE data were pooled for our analysis.

134 Brodifacoum, bromadiolone and difethialone were detected with a triple quadrupole
135 mass Quatro-Ultima (Waters) with negative electrospray ionization (ESI) in multiple
136 reaction monitoring scanning mode (MRM). LC-MSMS, MRM parameters and triple
137 quadrupole settings were identical as the ones reported in Albert et al. 2010.

138 The method's detection limit was 0.005 mg/kg for difethialone and 0.002 mg/kg for
139 brodifacoum and bromadiolone. The standards were all analytical grade (>98% purity). A
140 calibration curve was built with five levels of concentrations ranging from 2.5 to 80 pg
141 with an $r^2 > 0.99$. Samples were diluted in order to fit within the limits of the calibration
142 curve. Recoveries at low and high level were >70% for all compounds. Known amounts
143 of coumatetralyl (5 pg/L; transition 291.00>140.90) and flocoumafen (1 pg/L; transition
144 541.40>382.00) were added to each sample prior to the injection allowing ion
145 suppression monitoring. Methanol was injected between each sample to monitor any
146 possible contamination.

147 2.2.3. Statistical Analysis

148 Since great horned owls and red-tailed hawks represented the two species consistently
149 found across Canada (no red-tailed hawk samples were submitted from British Columbia,
150 however) and for which we had a large enough sample size to warrant a meaningful
151 analysis, cumulative frequency distribution graphs were constructed for these species.
152 The graphs were generated through a bootstrapping procedure (501 samples) using
153 BurrliOZ (version 1.0.14, © Commonwealth Scientific and Industrial Research
154 Organisation, Australia 2000). Using the values identified in our toxicity threshold
155 analysis, it was possible to identify the percentage of the sampled population exposed to
156 SGARs belonging to a certain risk category (5%, 10%, 15% and 20% risk of becoming
157 symptomatic).

158

159 **3. Results**

160 **3.1. Toxicity Threshold**

161 Five sources of data matched our criteria and were used in the analysis. Data
162 published by Newton et al. (1990, 1998, 2000; n=45), Albert et al. (2010; n=164) as well
163 as data from the Ecological Incident Information System (EIIS; n=61). All but four of the
164 EIIS cases were submitted by the State of New York and several of the values were
165 published in Stone et al. (1999, 2003). Barn owl samples were collected from localized
166 areas across Canada and the United Kingdom (UK) with a few individuals from the
167 United States (USA). Barred owl samples were mostly collected in Canada with only one
168 from the USA while red-tailed hawk samples were obtained from the USA only. Great
169 horned owl samples were collected from across both Canada and the USA. Samples were
170 often collected from relatively developed areas or areas where the public was likely to
171 report and submit carcasses.

172 There were significant differences between species in liver SGAR concentrations
173 ($F_{(4,535)}=12.68$, $p<0.0001$). Post hoc-tests (Tukey's Studentized Range test, $\alpha = 0.05$)
174 revealed that, on average, red-tailed hawks (n=32) were the species with the highest liver
175 concentrations of SGARs (Figure 1). All three owl species (great horned owl [n=86],
176 barred owl [n=26] and barn owl [n=126]) had SGAR liver residues that were comparable.

177 Logistic regression plots were calculated to predict the probability of a bird being
178 symptomatic as a function of SGAR liver residues (Figure 2). This was done for each
179 species separately and for all species combined (total of 270 individuals). Only the
180 predicted probability curve for the great horned owl (GHOW) was located inside the 95%
181 confidence limits for the pooled data and the estimated probability of becoming
182 symptomatic differed significantly between species ($F_{(1,4)} = 82.9$, $p<0.0001$). The curve
183 for the red-tailed hawk curve differed from those of the three owl species and the curves

184 for the great horned owl and the barn owl also differed from each other (Tukey's
185 Studentized Range post-hoc test, $P < 0.05$).

186 Using the probability curves, we calculated the predicted SGAR liver residue levels
187 for different probability risk thresholds for different species (Table 1), although this was
188 not possible for red-tailed hawks, as the data for this species could not be significantly
189 modeled by a logistic regression. The majority of the calculated values are under the
190 >0.1 - 0.2 mg/kg threshold suggested by Newton et al. (1999) and all are below 0.2 mg/kg.
191 If the lower range of 0.1 mg/kg and 0.2 mg/kg from the potentially lethal range suggested
192 for barn owls is applied to the barn owl probability curve, they correspond to toxicity
193 probabilities of 11% and 22% , respectively. The higher 0.7 mg/kg level proposed by the
194 Rodenticide Registrants Task Force (Erickson and Urban 2004) corresponds to a 54%
195 probability of effect in barn owls.

196 Although the differences among the species curves indicate that probabilities of
197 toxicity should be considered on a species-by-species basis, that is not possible where
198 data for species are lacking. In such cases, it may be necessary to estimate toxicity
199 probabilities on the basis of pooled data for other species. The probability curve for the
200 pooled data in our study predicts that one in 20 birds with detectable residues would
201 become symptomatic with SGAR liver residues of 0.02 mg/kg and one in five when
202 residue levels reach 0.08 mg/kg.

203 **3.2. The extent of SGAR exposure in Canada**

204 Of the two species sampled over a relatively broad area of Canada (great horned owl,
205 red-tailed hawk), great horned owls were most consistently exposed to SGARs (Figure
206 3). Roughly 65% of great horned owls across Canada had detectable levels of SGARs in

207 their liver (detection limit of 0.005 mg/kg ww). Frequency of exposure in red-tailed
208 hawks seemed to increase eastward from the Prairie Provinces to Ontario and Quebec.
209 The frequency of exposed birds was the lowest (~20%) in the Prairie and Northern
210 provinces (and territories), increased to ~70% in Ontario and reached the highest in
211 Quebec (~90% of red-tailed hawks found with detectable SGAR liver residues), although
212 the sample size in Quebec was smaller than in the other regions. However, as sampling
213 was fortuitous and sampling effort was not uniform, these spatial comparisons must be
214 considered preliminary.

215 Great horned owls and red-tailed hawks were exposed to a number of SGARs (Figure
216 4). The majority of great-horned owls had multiple compounds in the liver; it was the
217 only species with detectable levels of all three registered compounds. Sixty percent of
218 red-tailed hawks had detectable liver residues of one or two compounds (Figure 4).

219 Although the proportion of great horned owls with detectable residues was greater than
220 for red-tailed hawks, this difference was not significant when data were compared for
221 those provinces from which carcasses of both species were collected (Prairie Provinces,
222 Ontario and Quebec; (paired t-test, $t_{(2)} = -0.78$, $p = 0.26$; Figure 4). Brodifacoum and
223 bromadiolone were both detected in great horned owls and red-tailed hawks.

224 Difethialone was only ever detected in great horned owls (Table 2) but has only been
225 registered in Canada relatively recently.

226 When the liver SGAR concentrations in great horned owls measured in the present
227 study were plotted as a cumulative frequency graph (Figure 5; birds with detectable
228 residues only), it was apparent that approximately 25% had liver SGARs that exceeded
229 the 20% probability level for effect (0.07mg/kg; Table 1). The lack of a probability curve

230 for red-tailed hawks precludes making a similar calculation for that species, but it is
231 evident that liver residue levels were much lower than for great-horned owls (Figure 5
232 and 6). For-example, 50% of great horned owls with detectable residues had liver
233 concentrations greater than 0.05 mg/kg ww compared with only 10% of red-tailed hawks.
234 Comparison of liver concentrations in the two species in which birds were matched by
235 province confirmed that liver residues were significantly higher in the owls than in the
236 hawks (paired t-test; $t_{(2)} = -4.0$, $p=0.03$). This finding is in contrast to the previously
237 published literature (Figure 1) where liver residues were higher in red-tailed hawks than
238 in great-horned owls.

239 Of the small number of individuals from 13 other species analyzed from Quebec,
240 eight of those had at least one individual with detectable liver SGAR residues (Figure 7).
241 That indicates that a wide breadth of species is probably also exposed to these
242 compounds elsewhere in Canada.

243

244 **4. Discussion**

245 **4.1. Toxicity Threshold**

246 Critical SGAR liver concentrations associated with adverse effects and/or mortality
247 have not been defined for most raptor species (Walker et al. 2008a), and establishing liver
248 -toxicity thresholds for SGARs is problematic (Stone et al. 2003). This is partly
249 because there are a number of factors that contribute uncertainty. For instance, the limit
250 of quantification used to measure the liver SGAR residues can vary widely with the
251 analytical method. That can lead to underestimates of the extent of contamination but,
252 conversely, inflation of residue magnitude if residues which were detected but were

253 below the level of quantification using older analytical methodology were assigned an
254 inflated limit value (Taylor et al. 2009). Species also vary markedly in their sensitivity to
255 SGARs. This is known for laboratory mammals (World Health Organisation 1995) but
256 almost nothing is known about the relative sensitivity of different avian species (Walker
257 et al. 2008a). Our risk probability curves strongly suggest significant differences exist
258 among raptor species.

259 To date, the only residue toxicity threshold for SGARs in raptors that has been
260 suggested is the >0.1-0.2 mg/kg –potentially lethal range for barn owls (Newton et al.
261 1998, 1999). At best, that provides a range of concern for potential toxicity, and gives no
262 indication of likelihood of effects. The approach described in the current study offers a
263 major advance in our ability to assess risk from SGAR residues in that it proposes
264 quantitative toxicity thresholds for different probability levels of dying from SGAR
265 intoxication for three species, including the barn owl. If sufficient data were available, it
266 should be possible to extend this approach to other species. That, in turn, would help to
267 identify raptor species that may be more sensitive to SGAR toxicity. Overall, on the basis
268 of the probability curves defined so far, it would seem that the >0.1-0.2 mg/kg level for
269 barn owls already carries considerable risk of acute intoxication (> 10-20% of barn owls
270 with this residue being likely to suffer mortality). Clearly, the probability of acute
271 poisoning associated with the 0.7 mg/kg residue level proposed by the Rodenticide
272 Registrants Task Force (Erickson and Urban 2004) is worse still.

273 The probabilistic methods described here are, as with all predictive methods, subject
274 to biases and uncertainties. Of these, perhaps two of the most important are likely to be
275 underestimation of non-lethal residues, because birds characterised as –zeros in the

276 probabilistic plot may have metabolised some of their non-lethal SGAR residues before
277 dying [from non-SGAR related causes], and over-estimation of residues associated with
278 mortality because birds ingest more than a lethal dose before they die; animals typically
279 die some 5-7 days after ingestion of a lethal dose (Meehan 1984). Both biases would
280 have the effect of flattening the probability curve.

281 **4.2. Exposure extent in Canada**

282 4.2.1. Spatial extent

283 Stone et al. (2003) stated that, at the time, SGARs appeared to be present in the
284 majority of great horned owls and in roughly half of the red-tailed hawks from the
285 sampled areas of the State of New York. That conclusion can be directly applied to our
286 situation in Canada. Furthermore, a substantial fraction of a number of other raptors in
287 Quebec (from the western half of the province including areas surrounding Gatineau,
288 Montreal, Sherbrooke, Quebec and as far north as Obedjiwan) were also exposed to
289 SGARs (43% – or 13 of 30 birds tested), supporting the notion that other avian species
290 are also being impacted by SGAR use. This wider exposure in Quebec suggests a broad
291 contamination of terrestrial food chains as Accipiters, such as the Cooper’s hawk, as well
292 as other species such as the merlin and the American kestrel, feed predominantly on small
293 birds and occasionally on insects (Ehrlich et al. 1988). Small birds, if the source of
294 rodenticides, are most likely being exposed to SGARs from insects or other invertebrates,
295 and possibly through direct uptake of grain-based baits.

296 In our study, great horned owls were consistently exposed to SGARs across the
297 country. In apparent contrast, their daytime ecological counterpart, the red-tailed hawk,
298 showed an increasing frequency of exposure eastward from the Prairie Provinces. This

299 difference could be explained by the lower dietary diversity of owls than hawks. Marti
300 and Kochert (1995) showed that, on a finer scale, food-niche breadth became narrower
301 along an eastward transect from the west coast of North America. This may reflect
302 greater diversity of available prey in the west that could permit local populations of those
303 two raptors to increase their diet segregation in western regions (Marti and Kochert
304 1995). Houston et al. (1998) lists the main prey of great horned owls as including rabbits
305 and hares, coots and other waterfowl and mice. While snowshoe hares (*Lepus*
306 *americanus*), black-tailed jackrabbits (*Lepus californicus*), and ground squirrels
307 (*Spermophilus spp.*) dominate the hawk's diet in western and northern parts of North
308 America (Preston and Beane 2009). The bulk of their diet in eastern and midwestern
309 North America includes voles (*Microtus*), mice (*Peromyscus spp.*, *Reithrodontomys spp.*,
310 *Mus musculus*), rats (*Sigmodon hispidus*, *Oryzomys palustris*), and cottontails (*Sylvilagus*
311 *spp.*) (Preston and Beane, 2009). Thus, it may be that in eastern areas that are more
312 agricultural and urban (and subject to a higher degree of SGAR use), red-tailed hawks are
313 exposed more frequently to SGARs through their increased feeding on rodents and
314 reduced predation on other prey.

315 To obtain a more reliable estimate on actual exposure in Canada, we examined the
316 livers of birds found dead from all causes. Our data indicate that, despite a smaller human
317 population and the harsher climate in Canada (albeit some south-western regions of the
318 country are characterised by milder weather), both of which should limit the need for
319 rodenticides, the scale of exposure reported in our study are comparable to those in
320 Europe. In the French Department of Loire Atlantique, 73% of a sample consisting of
321 common kestrels (*Falco tinnunculus*), common buzzards (*Buteo buteo*), barn owls and

322 tawny owls (*Strix aluco*) had detectable SGAR liver residues (Lambert et al. 2007). In the
323 UK, between 40% and 74% of barn owls, kestrels, and avian scavengers such as buzzards
324 and red kites (*Milvus milvus*) found dead from various causes had detectable liver SGAR
325 residues (Newton et al. 1999; Shore et al., 1999, 2006; Walker 2008b). However, it
326 should be noted that the sampled areas of Canada were those with higher population
327 densities and where landscape features are not greatly dissimilar from Europe. That may
328 at least in part account for the apparent similarity in the frequency of contamination.

329 The widespread exposure in Canada in part most likely reflects the increase in sales
330 and use of SGARs in the last few decades (Albert et al. 2010), and the use of persistent
331 compounds that remain detectable in the liver long after the exposure event (Laas et al.
332 1985). However, it is also clear from our data that multiple exposures, as detected by the
333 presence of multiple compounds in the liver, are common. Although SGARs cannot be
334 used legally on crops or orchards in Canada and are labelled for ‘_indoor uses’ only,
335 ‘_indoor’ is defined to include use of baits outside farms and food establishments. This is
336 likely to increase the exposure of non-target organisms. SGARs in Canada are currently
337 labelled for domestic use although this is likely to change soon. Proposed regulatory
338 actions relating to exposure risks for wildlife includes (amongst others), prohibiting use
339 of SGAR compounds in residential settings or outdoor areas where wildlife may be
340 exposed. In the case of commercial applications, bait stations would be required where
341 wildlife could be exposed. Furthermore, labels of commercial class products would be
342 amended to state that those products could be used only by certified operators, farmers
343 and persons authorized in government-approved pest control programs (Pest Management

344 Regulatory Agency 2009). Those risk mitigation measures should have an overall
345 positive impact on reducing unnecessary exposure risks to wildlife.

346 Regarding the impact of SGARs, we must be cautious in extrapolating from our data
347 to predict likely mortality. However, if the probability of mortality is applied to each
348 residue value in our dataset for great horned owls, this equates to an estimated predicted
349 mortality of 11% (calculated by multiplying the probability of being exposed to SGARs
350 [65% in GHOW] by the mean probability of exhibiting signs of intoxication [17% in
351 GHOW]). This is the first time that the scale of potential mortality from SGARs has been
352 estimated for any wild raptor population. That estimate may well be too low, as some
353 proportion of poisoned birds likely die out of sight (Shore et al. 2005) and so be under-
354 represented in our sample. Furthermore, our estimates of the scale of mortality do not
355 account for any indirect effects that SGARs may have. Sub-lethal exposures may
356 indirectly increase mortality associated with natural or accidental events. For instance,
357 SGARs may hinder the recovery of birds from non-fatal collisions or accidents. They
358 may also impair hunting ability through behavioural changes such as lethargy, thus
359 increasing the probability of starvation. Intoxication with rodenticides has been shown to
360 alter behaviour in rodents (Cox and Smith 1992) but there is no evidence to date of
361 indirect effects in free-ranging raptors (Shore et al. 2005).

362 The lack of a probability plot for red-tailed hawks means that a comparable estimate
363 for SGAR-induced mortality in Canada cannot be made for this species. The available
364 data suggest that red-tailed hawks may be more sensitive to SGARs than great horned
365 owls (Figure 2) but red-tailed hawks generally had lower liver SGAR concentrations in
366 Canada, and, it is notable that in New York, great horned owls are poisoned more

367 frequently than red-tailed hawks (Stone et al. 1999, 2003). Additional studies and
368 monitoring of red-tailed hawk SGAR residues would strengthen our ability to estimate
369 the risk of toxicosis following exposure to SGARs.

370 4.2.2. Future directions

371 Most studies that investigate exposure of non-target species to SGARs have focused
372 on the uptake of poisoned rodents by various predators (Newton et al.1990, 1999; Berny
373 et al.1997; McDonald et al.1998; Howald et al. 1999; Shore et al. 1999, 2003). The
374 finding that falcons and accipiters were also exposed in Quebec suggests that terrestrial
375 food chains are broadly contaminated by SGARs despite their very restricted use.
376 Invertebrates represent another route of exposure, especially in insectivorous avian
377 species (Dowding et al. 2006). Some potential routes of exposure to aerial insectivores
378 include the consumption of invertebrates that previously fed on rodent faeces or carcasses
379 and even the consumption of ground-dwelling earthworms and beetles that ingested
380 residues or actual rodent bait (Spurr and Drew 1999; Dunlevy et al. 2000). Clearly, given
381 the fact that many ecosystems contain a larger proportion of insectivorous vertebrates
382 relative to higher trophic predators, exposure could even be greater in those taxa
383 (Dowding et al. 2010). Developing probability curves or even metabolism studies for a
384 wider range of species would provide us with insight into the relative sensitivities and
385 risks to other species (Watanabe et al 2010). Finally, researching further indirect effects
386 of SGARs on survival would refine current risk assessments of direct and indirect
387 mortalities in wildlife.

388 **4.3. Conclusion**

389 Our results continue to support recommendations that persistent SGARs such as
390 brodifacoum, bromadiolone and difethialone should be used with caution (or not at all in
391 some circumstances) given that it appears difficult to eliminate the risk of exposure to
392 non-target wildlife. The results presented will hopefully aid policy-makers in refining
393 risk-assessments of SGARs on non-target wildlife.

394 Our results can also help regulatory agencies worldwide provide guidance on both
395 commercial and residential use of SGARs and enforce appropriate risk mitigation as
396 needed. In this context, the extent of non-target exposure to SGARs may not always
397 depend on the amount of bait used, but also on the way it is used (Shore et al. 2006).
398 Focusing on improving application methods, such as baiting in areas of high rat activity
399 only, conducting periodic and frequent searches for dead or dying rodents, enclosing the
400 bait in a fashion that reduces invertebrate uptake may help reduce exposure of SGARs to
401 predatory birds and other non-target species. Whether or not rodenticide resistance is
402 common, an Integrated Pest Management (IPM) approach, that seeks to combine
403 mechanical, biological and chemical controls, should be favoured as opposed to relying
404 on a purely chemical mode of control.

405
406
407
408
409
410
411
412
413
414
415
416

417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460

Acknowledgements

We wish to thank all the wildlife agencies, provincial ministries, Conservation Officers, wildlife rehabilitation centers and shelters and the various wildlife rescue associations who submitted raptor carcasses for this study. National Wildlife Research Centre staff are thanked for specimen bank archiving and rodenticide residue analysis. Thanks are extended to Nick Mastrota for access to the US EPA EIIS, Dr. Malcolm McAdie and Dr. Craig Stephens and staff at the Centre for Coastal Health for conducting post mortem examinations and Dr. Ward Stone for his unflagging and inspirational work on contaminant and pesticide issues in New York State. Funding was from the Canadian Wildlife Service and the Pesticide Science Fund of Environment Canada.

References:

Albert, C.A., Wilson, L.K., Mineau, P., Trudeau, S. and Elliott, J.E. 2010. Anticoagulant rodenticides in three owl species from western Canada, 1998-2003. Arch. Environ. Contam. Toxicol. 163,451-459.

Berny, P.J., Buronfosse, T., Buronfosse, F., Lamarque, F. and Lorgue, G. 1997. Field evidence of secondary poisoning of foxes (*Vulpes vulpes*) and buzzards (*Buteo buteo*) by bromadiolone, a 4-year survey. Chemosphere. 35,1817–1829.

Buckle, A.P., Prescott, C.V. and Ward, K.J. 1994. Resistance to the first and second generation anticoagulant rodenticides – a new perspective. In: Halverson, W.S., Marsh, R.E. (eds) Proc. 16th Vert. Pest Conf. pp 138-144.

Cox, P. and Smith, R.H. 1992. Rodenticide ecotoxicology: pre-lethal effects of anticoagulants on rat behaviour. In: Borreco JE, Marsh RE (eds) Proc. 15th Vert. Pest Conf. University of California, Davis, California, USA. pp 165-170.

Dowding, J.E., Lovegrove, T.G., Ritchie, J., Kast, S.N. and Puckett, M. 2006. Mortality of northern New Zealand dotterels (*Charadrius obscurus aquilonius*) following an aerial poisoning operation. Notornis. 53,235–239.

Dowding, C.V., Shore, R.F., Worgan, A., Baker, P.J. and Harris, S. 2010. Accumulation of anticoagulant rodenticides in a non-target insectivore, the European hedgehog (*Erinaceus europaeus*). Environ. Pollut. 158,161-66.

Dunlevy, P.A., Campbell, E.W. and Lindsey, G.D. 2000. Broadcast application of a placebo rodenticide bait in a native Hawaiian forest. Int. Biodeterior. Biodegradation. 45,199–208.

Eason, C.T., Murphy, E.C., Wright, G.R.G., Spurr, E.B. 2002. Assessment of risks of brodifacoum to non-target birds and mammals in New Zealand. Ecotoxicology. 11,35-48.

461 Ehrlich, P.R., Dobkin, D.S. and Wheye, D. 1988. The birder's handbook – a field guide
462 to the natural history of North American birds. Simon & Schuster Inc. New York, New
463 York.

464
465 Erickson, W. and Urban, D. 2004. Potential risks of nine rodenticides to birds and
466 nontarget mammals: a comparative approach. United States Environmental Protection
467 Agency, Washington, DC, USA.

468
469 Hedgal, P.L. and Colvin, B.A. 1988. Potential hazard to eastern screech-owls and other
470 raptors of brodifacoum bait used for vole control in orchards. Environ. Toxicol. Chem.
471 7,245-260.

472
473 Houston, C.S., Dwight, G.S. and Christoph, R. 1998. Great Horned Owl (*Bubo*
474 *virginianus*), The Birds of North America Online (A. Poole, Ed.). Ithaca: Cornell Lab of
475 Ornithology; Retrieved from the Birds of North America Online:
476 <http://bna.birds.cornell.edu/bna/species/372>

477
478 Howald, G.R., Mineau, P., Elliott, J.E., and Cheng, K.M. 1999. Brodifacoum poisoning
479 of avian scavengers during rat control on a seabird colony. Ecotoxicology 8,429-446.

480
481 Laas, F.Y., Forss, D.A., Godfrey, M.E.R. 1985. Retention of brodifacoum in sheep
482 tissues and excretion in faeces. N.Z. J. Agric. Res. 28,357–359.

483
484 Lambert, O., Pouliquen, H., Larhantec, M., Thorin, C. and L'Hostis, M. 2007. Exposure
485 of raptors and waterbirds to anticoagulant rodenticides (difenacoum, bromadiolone,
486 coumatetralyl, coumafen, brodifacoum): epidemiological survey in Loire Atlantique
487 (France). Bull. Environ. Contam. Toxicol. 79,91-94.

488
489 McDonald, R.A., Harris, S., Turnbull, G., Brown, P. and Fletcher, M. 1998.
490 Anticoagulant rodenticides in stoats (*Mustela erminea*) and weasels (*Mustela nivalis*) in
491 England. Environ. Pollut. 103,17–23.

492
493 Marti, C.D. and Kochert, M.N. 1995. Are Red-tailed Hawks and Great Horned Owls
494 diurnal-nocturnal dietary counterparts? Wilson Bull. 107,615-628.

495
496 Mastrota, N.F. 2007. EIIS Version 2.0 – User Manual. EFED Information and Support
497 Branch, Environmental Fate and Effects Division, Office of Pesticide Programs, USA.

498
499 Meehan, A.P. 1984. Rats and Mice: their Biology and Control. Rentokil Limited, East
500 Grinstead, Sussex, UK.

501
502 Newton, I., Wyllie, I. and Freestone, P. 1990. Rodenticides in British barn owls. Environ.
503 Pollut. 68,101-117.

504
505 Newton, I., Dale, L., Finnie, J.K., Freestone, P., Wright, J., Wyatt, C. and Wyllie, I. 1998.
506 Wildlife and Pollution: 1997/98 Annual Report. JNCC Report No. 285

507
508 Newton, I., Shore, R.F., Wyllie, I., Birks, J.D.S., Dale, L. 1999. Empirical evidence of
509 side-effects of rodenticides on some predatory birds and mammals. In: Cowan DP, Feare
510 CJ (eds) Advances in vertebrate pest management. Filander Verlag, Furth, pp 347–367.
511
512 Newton, I., Afsar, A., Dale, L., Finnie, J., Shore, R.F., Wright, J., Wyatt, C. and Wyllie, I.
513 2000. Wildlife and Pollution: 1998/99 Annual Report. JNCC Report No.305
514
515 Parmar, G., Bratt, H., Moore, R. and Batten, P.L. 1987. Evidence for a common binding-
516 site in vivo for the retention of anticoagulants in rat liver. Hum. Toxicol. 6,431–432.
517
518 Pest Management Regulatory Agency. 2009. Proposed Risk Mitigation Measures for
519 Eight Rodenticides. Re-evaluation Note REV2009-05. Health Canada. Ottawa, Ontario.
520 22 June 2009. 23 pages.
521
522 Preston, C.R. and Beane, R.D. 2009. Red-tailed Hawk (*Buteo jamaicensis*), The Birds of
523 North America Online (A. Poole, Ed.). Ithaca: Cornell Lab of Ornithology; Retrieved
524 from the Birds of North America Online: <http://bna.birds.cornell.edu/bna/species/052>
525
526 Shore, R.F., Birks, J.D.S. and Freestone, P. 1999. Exposure of non-target vertebrates to
527 second-generation rodenticides in Britain, with particular reference to the polecat *Mustela*
528 *putorius*. N.Z. J. Ecol. 23,199-206.
529
530 Shore, R.F., Malcolm, H.M., Wienburg, C.L., Turk, A., Horne, J.A., Dale, L., Wyllie, I.
531 and Newton, I. 2001. Wildlife and pollution: 1999/2000 Annual report. JNCC Report No.
532 321
533
534 Shore, R.F., Birks, J.D.S., Afsar, A., Wienburg, C.L. and Kitchener, A.C. 2003. Spatial
535 and temporal analysis of second-generation anticoagulant rodenticide residues in polecats
536 (*Mustela putorius*) from throughout their range in Britain, 1992–1999. Environ. Pollut.
537 122,183–193.
538
539 Shore, R.F., Malcolm, H.M., Wienburg, C.L., Walker, L.A., Turk, A. and Horne, J.A.
540 2005. Wildlife and Pollution: 2001/02 Annual Report. JNCC Report No. 352
541
542 Shore, R.F., Malcolm, H.M., McLennan, D., Turk, A., Walker, L.A., Wienburg, C.L. and
543 Burn, A.J. 2006. Did foot-and-mouth disease-control operations affect rodenticide
544 exposure in raptors? J. Wildl. Manage. 70,588-593.
545
546 Spurr, E.B. and Drew, K.W. 1999. Invertebrates feeding on baits used for vertebrate pest
547 control in New Zealand. N.Z. J. Ecol. 23,167–173.
548
549 Stone, W.B., Okoniewski, J.C. and Stedelin, J.R. 1999. Poisoning of wildlife with
550 anticoagulant rodenticides in New York. J. Wildl. Dis. 35,187-193.
551

552 Stone, W.B., Okoniewski, J.C. and Stedelin, J.R. 2003. Anticoagulant rodenticides and
553 raptors: recent findings from New York, 1998-2001. Bull. Environ. Contam. Toxicol.
554 70,34-40.
555
556 Taylor, M.J., Sharp, E.A., Melton, L.M., Giela, A. and Watson, J.E. 2009. Walk on the
557 wild side: an investigation into secondary poisoning of wild birds using LC-MS. British
558 Mass Spectrometry Society. 60,12-14.
559
560 Walker, L.A., Turk, A., Long, S.M., Wienburg, C.L., Best, J. and Shore, R.F. 2008a.
561 Second generation anticoagulant rodenticides in tawny owls (*Strix aluco*) from Great
562 Britain. Sci. Total Environ. 392,93-98.
563
564 Walker, L.A., Shore, R.F., Turk, A., Pereira, M.G., and Best, J. 2008b. The Predatory
565 Bird Monitoring Scheme (PBMS) — identifying chemical risks to top predators in
566 Britain. Ambio. 37,466-471.
567
568 Walker, L.A., Lawlor, A.J., Llewellyn, N., Peréira, M.G., Potter, E., Townsend, J., Turk,
569 A., and Shore, R.F. 2010. The Predatory Bird Monitoring Scheme (PBMS) Report 2006-
570 7. A contract report from the Centre for Ecology & Hydrology to Natural England, 65
571 pages. Retrieved from the website at
572 http://pbms.ceh.ac.uk/docs/AnnualReports/PBMSreport2006_2007_FINAL.pdf
573
574 Watanabe, K.P., Saengtienchai, A., Tanaka, K.D., Ikenaka, Y. and Ishizuka, M. 2010.
575 Comparison of warfarin sensitivity between rat and bird species. Comp. Biochem. Phys.
576 C. 152,114-119.
577
578 World Health Organisation. 1995. Environmental Health Criteria 175: Anticoagulant
579 Rodenticides. World Health Organisation (WHO), Geneva.
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597

598 **Figure captions:**

599

600 **Figure 1:** Published liver SGAR residues (combined concentrations of bromadiolone,
601 brodifacoum and difethialone) in barred owl (BAOW), barn owl (BNOW), great horned
602 owl (GHOW) and red-tailed hawk (RTHA). Total number of birds = 270 and do not
603 include birds with non-detected residues. Diamond in the center of the box represents
604 average, line is the median, box is the upper and lower quartiles and the whiskers are the
605 standard deviation. Sources of the data are: Newton et al. 1990, 1998, 2000; Stone et al.
606 1999, 2003; Albert et al. 2010; EIIS 2010 download.

607

608 **Figure 2:** Effect plot of the probability of becoming symptomatic (0,1) as a function of
609 \log_{10} [mg/kg]. ALL represents pooled data (n=270), BAOW represents barred owls
610 (n=26), BNOW represents barn owls (n=126), GHOW represents great horned owls
611 (n=86) and RTHA represents red-tailed hawks (n=32). Shading represents 95%
612 confidence limits for ALL birds. Curves were drawn using the formula $y(\text{probability}) =$
613 $1/(1+\exp(-(\text{int} + b*x)))$ where int is the intercept and b is the parameter estimate for X
614 (concentration).

615

616 **Figure 3:** Percentage of great horned owls (GHOW) and red-tailed hawks (RTHA)
617 across Canada sampled in our study that had detectable (≥ 0.005 mg/kg ww) liver SGAR
618 residues. No RTHA samples were collected from PYR. PYR stands for the Pacific and
619 Yukon region of Canada and PNR is the Prairie and Northern Region.

620

621 **Figure 4:** Percentage of great horned owls (GHOW) and red-tailed hawks (RTHA) with
622 0, 1, 2 and 3 different SGARs detected in the liver. Tested compounds were brodifacoum,
623 bromadiolone and difethialone.

624

625 **Figure 5:** Cumulative frequency graph for liver SGAR residues in 79 great horned owls.
626 Red line represents the 20% probability level for effect (0.07 mg/kg; Table 1).

627

628 **Figure 6:** Cumulative frequency graph for liver SGAR residues in 42 red-tailed hawks.

629

630 **Figure 7:** Numbers of birds of prey from Québec that contained detectable and non-
631 detectable liver SGAR residues (13/30 samples tested positive or 43%).

632

633 **Table 1:** Toxicity threshold values (mg/kg ww) for 5%, 10%, 15% and 20% probability
 634 risk levels. For-example, in barred owls (BAOW), an owl with 0.06mg/kg SGAR
 635 residues in the liver would have a 5% chance of showing signs of toxicosis. Sample sizes
 636 (n) as well as the number of positive (1) and negative (0) cases are presented. P value
 637 representing binary logit model fit is also showed. BNOW stands for barn owl, GHOW is
 638 the great horned owl, RTHA the red-tailed hawk and ALL represents the pooled data for
 639 all birds.
 640

Probability	BAOW n=26 0=22 1=4 p=0.008	BNOW n=126 0=114 1=12 p=<0.0001	GHOW n=86 0=62 1=24 p=<0.0001	RTHA n=32 0=3 1=29 p=0.37	ALL n=270 0=201 1=69 p=<0.0001
0.05	0.06	0.05	0.02	---	0.02
0.10	0.09	0.09	0.03	---	0.04
0.15	0.13	0.13	0.05	---	0.06
0.20	0.16	0.18	0.07	---	0.08

641 --- - values not presented if binary logit model fit was not statistically significant
 642
 643
 644
 645
 646
 647
 648
 649
 650
 651
 652
 653
 654
 655
 656
 657
 658
 659
 660
 661
 662
 663
 664
 665
 666
 667
 668
 669


670 **Table 2:** Geometric mean (range) liver SGAR concentrations [mg/kg ww] for great
 671 horned owls (GHOW) and red-tailed hawks (RTHA) from the Pacific and Yukon region
 672 of Canada (PYR), the prairie and northern region (PNR), Ontario and Quebec.
 673

	PYR		PNR		Ontario		Quebec		Pooled – all provinces	
	GHOW	RTHA	GHOW	RTHA	GHOW	RTHA	GHOW	RTHA	GHOW	RTHA
Brodifacoum	0.04 (0.003- 0.61) n=28	N/A	0.008 (0.001- 0.016) n=6	0.004 (0.001- 0.02) n=3	0.007 (0.001- 0.05) n=17	0.006 (0.001- 0.17) n=18	0.013 (0.003- 0.08) n=7	0.01 (0.008- 0.04) n=5	0.017 (0.001- 0.61) n=58	0.006 (0.001- 0.17) n=26
Bromadiolone	0.03 (0.005- 0.57) n=33	N/A	0.007 (0.001- 0.07) n=7	0.004 (0.001- 0.008) n=3	0.01 (0.001- 0.07) n=15	0.004 (0.001- 0.06) n=25	0.01 (0.003- 0.14) n=6	0.003 (0.002- 0.006) n=4	0.018 (0.001- 0.57) n=61	0.004 (0.001- 0.064) n=32
Difethialone	0.02 (0.013- 0.03) n=3	N/A	ND	ND	0.003 (0.003- 0.003) n=1	ND	ND	ND	0.013 (0.003- 0.03) n=4	0
Pooled - all compounds	0.03 (0.003- 0.609) n=64	N/A	0.007 (0.001- 0.07) n=13	0.004 (0.001- 0.017) n=6	0.008 (0.001- 0.07) n=33	0.005 (0.001- 0.17) n=43	0.012 (0.003- 0.14) n=13	0.006 (0.002- 0.04) n=9	0.016 (0.001 - 0.61) n=123	0.005 (0.001 - 0.064) n=58


674 N/A = no samples obtained; ND = no detectable residue in any livers; n= number of birds with detectable
 675 residues.

676
 677
 678
 679

680
681
682



683
684 **Figure 1**
685
686
687
688
689
690
691
692
693
694
695
696



697
 698
 699
 700
 701
 702
 703
 704
 705
 706
 707
 708
 709
 710
 711

Figure 2

712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757




Figure 3

758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803




Figure 4

804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849




Figure 5





Figure 6

850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890



891
 892
 893
 894
 895
 896
 897
 898
 899
 900
 901
 902
 903
 904
 905
 906
 907
 908
 909
 910
 911
 912
 913
 914
 915

Figure 7

916
917
918
919
920