

## **LOCH DUNTELCHAIG (NH 600 285 – 600 291)**

*D. Stephenson*

### **Introduction**

At the SW limit of the Orcadian Basin, the outcrop of Middle Old Red Sandstone rocks extends down the SE side of the Great Glen as far as Foyers (Stephenson, 1972, 1977; Mykura, 1982). Within this outcrop 'extension', the basal beds are characterized by coarse-grained breccias, breccio-conglomerates and conglomerates, resting on a highly irregular paleo-land surface of Grampian Group (Dalradian) metasedimentary rocks and late-Caledonian granitic rocks. The unconformity is well exposed in several places, the most dramatic and instructive being on the SE side of Dun Chia hill, between Loch Ruthven and Loch Duntelchaig. There, the unconformable junction is exposed continuously for a distance of some 1300 m and the overlying breccio-conglomerates form cliffs from Tom Mor (591 280) in the SW, through Craig na h-Iolaire (597 282), to the spectacular, 50 m-high Creag nan Clag (Crag of the Bells; 601 289) in the NE (Figure 2.xx). The effects of pre-MORS weathering and erosion on the underlying Grampian Group rocks are well seen and the breccio-conglomerates yield much information about sedimentary processes and palaeogeography of the region in Mid Devonian time. Similar basal beds are also well exposed in cliffs between Creag Dhearg (618 294) and Carn Mor (627 303), 2 to 3 km to the ENE of Craig nan Clag, but there the unconformity is less well exposed.

The Old Red Sandstone rocks of the area were first described by Wallace (1880), who described the unconformity and the overlying beds at Dun Chia and Creag Dhearg in some detail. Further descriptions and interpretations of these localities were given by Mould (1946) in her account of the nearby Foyers granitic pluton, but the most extensive investigation was that of Mykura (1982), which was used in the compilation of the BGS 1:50 000 geological map, Sheet 73E (Foyers, 1996) and is the basis for much of the following account.

### **Description**

Between Loch Duntelchaig and Loch Ness in the Great Glen, there is a cross-strike outcrop width of about 5 km of Middle Old Red Sandstone strata. This represents a succession over 2 km thick that includes coarse- and fine-breccias, conglomerates, gritty and pebbly feldspathic sandstones and fine sandstones, with subordinate mudstones, siltstones and calcareous beds. In general, the lower parts of the sequence (up to 900 m) are of breccio-conglomerate and coarse bimodal conglomerate overlain by planar-bedded medium- to fine-grained sandstones. The sequence is interdigitated along strike to the NE with that of the Inverness and Nairnside districts, where fragmentary fish and plant spores of Mid Devonian age have been found (Horne and Hinxman, 1914; Horne, 1923; Fletcher *et al.*, 1996). There is a regional dip of between 10° and 50° to the NW.

Throughout the area in general the pre-MORS topography was undulating with some steep slopes locally that probably reflected active fault scarps (Stephenson, 1972; Mykura, 1982). Within the GCR site, between Tom Mor (591279) and Creag

nan Clag (600 287), the basal unconformity is exposed almost continuously for 1300 m at the foot of steep cliffs (Figure 2.xx). It is undulating, but is generally inclined in a NNW to westerly direction at angles ranging from 20° to 65°, with an average of 45°. Locally it is almost vertical on Creag na h-Iolaire (at 5985 2835).

The underlying rocks are platy psammites (metasandstones), gneissose in parts, of the Dalradian, Grampian Group, which are generally vertical or steeply inclined, but exhibit some quite large scale folding. Directly beneath the unconformity at the southern end of Creag nan Clag (5988 2857 – 6000 2870), a monocline, overturned towards the west, has a near-horizontal upper limb. Locally this limb is broken into small blocks, some rotated, but others still in their original position and enclosed by sandstone. Sandstone veins also penetrate downwards through the horizontal limb and axial plane of the monocline into the vertical limb beneath. A small inlier of psammite on the NW slope of Dun Chia, some 40 m below the summit (Figure 2.xx), probably represents the peak of a buried hillock or ridge crest. Here, just below the unconformity, is another monocline overturned to the SW.

The psammites are intruded by dykes, up to 10 m thick, of microgranitic rock ('felsite') and microdiorite. Many of the felsites are intensely shattered at the unconformity and are overlain by a deposit composed almost entirely of felsite clasts.

The MORS rocks of the Moray Firth area are formally assigned to the Inverness Sandstone Group, and the basal beds at the GCR site are all part of the Bochruben Formation (Mykura, 1982). These are mainly breccio-conglomerates and conglomerates in which almost all of the clasts are locally derived from the Dalradian, Grampian Group or from the nearby Foyers Pluton. Horizontal and vertical variations in the basal beds of the formation are well displayed in the cliffs on the east side of Dun Chia.

At the south end of Craig nan Clag, the lowest 5 – 10 m are composed entirely of angular, locally derived clasts, with plates of psammite and felsite in roughly equal proportions; in the vicinity of felsite dykes, felsite clasts predominate. Elsewhere, for example at the eastern end of the inlier on the NW flank of Dun Chia, felsite clasts commonly constitute 40% of the basal deposits and in some restricted areas they constitute up to 90%.

The basal breccias of almost in-situ angular material are overlain by massive, very crudely bedded breccio-conglomerates, composed of ungraded subangular clasts up to 60 cm in diameter. The larger clasts are not in contact and are supported in a sparse red gritty matrix. Their long axes are randomly orientated, some being almost perpendicular to the bedding. Most are psammite and quartzite of local origin, with subordinate granite-gneiss and gneissose semipelite, but angular felsite clasts constitute up to 30% in places. Rounded boulders of biotite granite also occur sporadically. The breccio-conglomerates vary in thickness, being thickest in the cliffs of Creag nan Clag and thinning to the NE and SW, though they still form the greater part of all the cliff sections.

The NW flanks of Dun Chia and parts of the summit ridge are composed of bimodal conglomerates with a high proportion of leucogranite clasts. On Creag na h-Iolaire, coarse conglomerates with rounded granite boulders up to 60 cm in diameter

rest in places almost directly on the Dalradian 'basement', though residual patches of local breccia only a few metres thick, and some breccio conglomerates are also present. Sandstone lenses are intercalated locally with the conglomerates, such as those dipping at 15° to the WNW, between 6 and 10 m above the unconformity at the SW end of Creag nan Clag.

## **Interpretation**

The prominent monocline in the Dalradian 'basement' at Creag nan Clag has an axial plane almost parallel to the unconformity and was interpreted by Mykura (1982) as flexuring due to hill creep prior to deposition of the MORS sediments. However, this coherent fold is on a remarkably large scale to have been formed in such a manner and subsequent visitors to the site have rejected the interpretation, observing that the geometry of the fold is of a tectonic style and conforms with others in the surrounding psammites. At one locality a dyke, truncated by the unconformity, is intruded along the vertical limb of the monocline but is not bent over as it enters the horizontal limb. Mykura explained this by suggesting that the dyke was intruded after the hill creep but before the sedimentation. This somewhat implausible scenario does fit a further suggestion by Mykura (1982), concerning the origin of local concentrations of felsite in the basal MORS. Observing that, in several places on Dun Chia and Creag Dearg, the percentage of felsite clasts (40 – 90%) is much higher than the percentage of felsite dykes seen in the Dalradian beneath, Mykura suggested that the dykes may have been feeders to small extrusive domes on the pre-MORS land surface. Such domes are usually unstable and would have crumbled to breccia shortly after extrusion; hence no definite domes have been identified. However, in the absence of any other evidence for volcanism in the Grampian Highlands in Mid Devonian time, this scenario is also regarded as highly tenuous.

The local nature of most of the clasts in the basal MORS and the low degree of sorting indicate that many of the deposits were formed quite near to their source of material. Much of the angular material in the basal breccias is virtually *in situ* and is in effect a regolith or, if on a steep original slope, a 'fossil scree'. The lack of movement is particularly noticeable above some felsite dykes where it is difficult to separate the in-situ felsite from 'felsite scree'. This effect is even more characteristic of basal MORS deposits to the SW of the GCR site, where 'granitic scree' are indistinguishable in places from underlying in-situ granitic rocks (Stephenson, 1972; Mykura, 1982). The freshness of many of the clasts indicates rapid accumulation in a climate in which chemical weathering was minimal. However, oxidation is commonly intense, imparting a red or purple hue to the matrix of the rocks, so climatic conditions during deposition were probably arid or semiarid.

The breccio-conglomerates that dominate the GCR site merge with the basal breccias and have no marked basal erosion surfaces. They are unstratified and poorly sorted, with mainly subangular clasts of local material 'floating' in a matrix, usually of mud or silt grade. These are the characteristics of debris flows that moved as dense, viscous masses in which waterlogged matrix was able to support and transport clasts up to boulder size. Such deposits form typically in areas of high relief and rapid erosion, in which deposition on steeply sloping alluvial fans results from occasional sheet floods during sudden violent rainstorms in an otherwise dry area. The thickest development of breccio-conglomerate, in the 50 m-high cliffs of Creag nan Clag, has

no horizontal breaks and it is possible that the entire section represents a single debris flow, possibly filling a palaeovalley. Lenses of conglomerate and sandstone within the sequence probably originated as channel-fill deposits within the alluvial fans and some of these have cut down to rest directly on basement in places.

Higher in the sequence, reworking by water becomes apparent as the breccias become slightly better sorted with the development of some bedding and clasts become more rounded. Debris of a less local nature appears, including leucogranites that must have been transported for some distance. Bimodal clast-supported conglomerates, which occur locally as channel-fills within the breccio-conglomerates, become dominant higher in the succession, from the summit of Dunchia north-westwards. These higher conglomerates are of fluvial origin and probably originated as braided river deposits on the distal parts of fans and on the piedmont plain. Beyond the GCR site, still higher parts of the succession record the eventual burial of the fans by finer-grained clastic deposits accumulating on a broad flood plain (Mykura, 1982).

By the time of deposition of the MORS sediments, the Grampian Highlands had already experienced considerable rapid uplift at the end of the Caledonian Orogeny. An estimate of the scale of uplift in the northern Grampians can be obtained from petrological evidence in the aureole of the Foyers Pluton, which Marston (1971) suggested had a possible cover of 7 km at its time of emplacement in the Early Devonian. This cover was removed in a maximum of c. 20 Ma to expose the pluton in the Mid Devonian, representing an erosion rate of at least 3.5 cm per 100 years, comparable with rates in present day newly formed mountain areas.

The Great Glen Fault was already in existence in the Mid Devonian and could well have been the locus of a major valley that drained NE and was bounded to the SE by high mountains (Stephenson, 1972, 1977). The MORS basal sediments were deposited predominantly in coalescing alluvial fans along the foot of these mountains and in the valley bottom, which widened out north-eastwards into a piedmont plain on the edge of the main Orcadian Basin and probably extended across the Great Glen Fault into the area now occupied by the Black Isle and Tarbat Ness (Mykura, 1982, figure 11). Close to the Great Glen Fault, in the Foyers – Inverfarigaig area, the highlands to the SE were probably bounded by major active NE – SW fault scarps, related to the Great Glen Fault and with a component of downthrow to the NW. However, around Loch Duntelchaig, the edge of the hills probably swung to the ENE; here major fault scarps were probably absent and the boundary zone between plain and mountain may have been a belt of foothills with several small active local fault scarps.

## **Conclusions**

The spectacular cliffs on the SE flank of Dun Chia hill, at the SW end of Loch Duntelchaig, provide a representative section of the basal beds of the Middle Old Red Sandstone at the southern limit of the Orcadian Basin. The unconformable junction beneath these beds is very well exposed almost continuously over a distance of 1300 m and is probably one of the best preserved ‘buried landscape’ unconformities in Great Britain. It is certainly of national importance and visitors cannot fail to be impressed by the scale of the exposure and by the many exceptional features that are exhibited. Beneath the unconformity, folded Neoproterozoic flaggy metasandstones,

cut by felsitic dykes, are disaggregated and shattered close to an undulating former land surface. The overlying angular debris can be matched precisely to the immediately underlying rock in many places and probably accumulated as scree. Poorly sorted mixtures of angular fragments and rounded boulders (breccio-conglomerates) comprise the majority of the section and probably originated as debris flows or sheet floods associated with alluvial fans that debouched from mountainous terrain to the SE and spread out over a broad piedmont plain to the north on the edge of the alluvial plain of the main Orcadian Basin.

## References

Fletcher, T.P., Auton, C.A., Highton, A.J., Merritt, J.W., Robertson, S. and Rollin, K.E. 1996. Geology of the Fortrose and eastern Inverness district. *Memoir of the British Geological Survey*, Sheet 84W (Scotland).

Horne, J. 1923 The geology of the Lower Findhorn and Lower Strath Nairn. *Memoir of the Geological Survey of the U.K.*, Sheets 84 and part of 94 (Scotland).

Horne, J. and Hinxman, L.W. 1914. The geology of the country around Beaully and Inverness: including part of the Black Isle. *Memoir of the Geological Survey of the U.K.*, Sheet 83 (Scotland).

Marston, R.J. 1971. The Foyers granitic complex, Inverness-shire, Scotland. *Quarterly Journal of the Geological Society of London*, Vol. 126, 331-368.

Mould, D.D.C.P. 1946. The geology of the Foyers 'granite' and the surrounding country. *Geological Magazine*, Vol. 83, 249-265.

Mykura, W. 1982. The Old Red Sandstone east of Loch Ness, Inverness-shire. *Report of the Institute of Geological Sciences*, No. 82/13.

Stephenson, D. 1972. Middle Old Red Sandstone alluvial fan and talus deposits at Foyers, Inverness-shire. *Scottish Journal of Geology*, Vol. 8, 121-127.

Stephenson, D. 1977. Intermontane basin deposits associated with an early Great Glen feature in the Old Red Sandstone of Inverness-shire. 35-45 In Gill, G. (editor) *The Moray Firth Area Geological Studies*. (Inverness: Inverness Field Club.)

Wallace, T. 1880. The structural geology of Strathnairn. *Transactions of the Edinburgh Geological Society*, Vol. 3, 204-219.

## Figure Captions

Figure 2.xx The cliffs of Creag nan Clag, on the eastern flank of Dun Chia hill, Loch Duntelchaig GCR site. The undulating unconformity between Middle Old Red Sandstone massive breccio-conglomerates and underlying Grampian Group (Dalradian) flaggy psammities can be traced for some 1300 m along the base of the steep upper cliffs. (Photo: BGS No. D1813.)

Figure 2.xx Map of the area around Dun Chia hill, at the SW end of Loch Duntelchaig, based upon BGS 1:50 000 Sheet 73W, Scotland (Foyers, 1996).

**Site Selection Criteria** (for table)

Representative of the basal MORS at the southern edge of the Orcadian Basin; exceptional example of a 'buried landscape' type of unconformity.