

Geology as a control on landslides on the Isle of Wight: an overview

G. O. Jenkins^{a*}, C. Foster^a & P. M. Hopson^a

^aBritish Geological Survey, Keyworth, Nottingham, NG12 5GG

* Corresponding author, Email: gjenkins@bgs.ac.uk, Tel: +44(0)115 9363128, Fax: +44(0)115 9363200

Abstract

The Cretaceous and Palaeogene sedimentary rocks that crop out on the Isle of Wight are highly prone to landsliding and the island offers an important field laboratory wherein to investigate a number of the different types of failure. Many of these landslides represent a significant engineering hazard, with several urban areas requiring remedial work and planning constraints (e.g. The Undercliff and Seagrove Bay) to aid development. Previous studies have thoroughly investigated the major landslides in the Undercliff area around Ventnor and presented a mechanism for that massive failure. This overview of the landslides throughout the Isle of Wight by the British Geological Survey was completed as part of the multidisciplinary survey of the surface geology, structure, geophysical response and offshore interpretations of the island between 2007 and 2010. The survey has collected new observational data on the extensive coastal landslides, as well as the distribution, nature and mechanism of failure of the lesser-studied inland examples.

Keywords: Landslides, Isle of Wight, Cretaceous, Palaeogene, coastal erosion.

1 Introduction

The Cretaceous and Palaeogene strata of the Isle of Wight are mainly composed of relatively soft, often poorly lithified sedimentary rocks, the nature of which makes them highly susceptible to landsliding. These susceptible units are interbedded with more competent layers that affect the style of the failure by providing a natural 'reinforcement' of the stratal units during failure. Slope instability is exacerbated at the coast where marine erosion unloads the toes of cliffs leading to cliff failure and sequential failure within existing landslides. Extensive landslides occurring inland, for example on the slopes of the southern chalk downs, are the result of a complex inter-relationship between the strata, geomorphological development, palaeoclimate and climate, and groundwater. Here the removal of toe debris is slow and sporadic and the slides have a natural angle of stability. Together, these areas present a significant hazard to infrastructure on the island.

The studies of landslides affecting the Undercliff in Ventnor, Luccombe, and Seagrove Bay (e.g. Hutchinson, 1991; Hutchinson et al., 1991a; Hutchinson et al., 1991b; Moore et al., 2007; Winfield et al., 2007; Moore et al., 2010) have significantly improved the understanding of the nature and mechanisms of these landslides. However, there is much less discussion of significant failures elsewhere. Thus the purpose of the interdisciplinary survey of the island described in this paper has been to provide an overview of all other major failures.

1 The existing British Geological Survey (BGS) 1:50 000 geological map of the Isle of
2 Wight (British Geological Survey, 1976) shows extensive landslides around the coast;
3 one continuous landslide is shown extending approximately 25 km from Brook Bay
4 on the south-west coast to Luccombe on the SE coast. These published landslide
5 polygons were principally derived from inspection of aerial photographs and limited
6 field inspections in the mid 1970s. The landslide survey by the BGS in 2007 and 2008
7 re-examined these large landslide areas and mapped them with greater precision, more
8 accurately reflecting the type of instability that is present as well as identifying
9 additional areas affected by landslides. The survey has accounted for over 90
10 landslide affected areas. This paper highlights the widespread distribution of
11 landslides on the Isle of Wight, documenting the extensive coastal landslides other
12 than those around Ventnor, including the Palaeogene landslides along the northwest
13 coast, and the lesser-studied inland landslides.
14
15

16
17 The exposed geology of the Isle of Wight spans a total of about 135 million years,
18 including sediments deposited during the Cretaceous, Palaeogene and the Quaternary.
19 No sediments are thought to have been deposited during the Neogene (Insole et al.,
20 1998). A detailed account of the lithology, stratigraphy and depositional environment
21 is beyond the scope of this paper, and the reader is referred to Insole et al., (1998) and
22 the overview provided in Hopson (this issue). A simplified geological map, based on
23 the current published map of the Isle of Wight is presented in Fig. 1 incorporating the
24 landslides identified during the most recent survey.
25
26

27
28 In general flat-lying Cretaceous and Palaeogene strata are found to the south and
29 north, respectively, of the central spine of the island within which both are
30 incorporated into the complex steeply-dipping Isle of Wight structure. The internal
31 architecture and lithology of these bedrock units underpin their response to erosion
32 and slope stability and this will be discussed below. Thin, patchy and widely
33 dispersed outcrops of Quaternary deposits overlie the bedrock, providing evidence for
34 the complex climatic history that has affected the island. However, these limited
35 outcrops belie the widespread affect of periglaciation on the both bedrock and
36 superficial deposits alike. In some cases, the effects of periglaciation may also be
37 significant in the development of slope failures, but this aspect of landslide
38 development has not been well documented on the island.
39
40
41

42
43 The Isle of Wight has a varied geomorphology, including chalk downlands, coastal
44 plains and large drowned estuaries (rias) (Booth & Brayson, this issue) founded
45 principally on the Palaeogene. As noted above the island is divided from east to west
46 by a prominent chalk ridge that is the expression of the monocline, formed along the
47 northern limbs of the asymmetric Brighstone and Sandown anticlines (which Hopson
48 describes in detail in this issue). The structure forms the foundation of most of the
49 inland steep slopes and also controls the formation of steep chalk cliffs at both ends of
50 the island. With the exception of the low-lying estuaries the majority of the coastal
51 sections around the island present cliffs of variable height reflecting the variably
52 resistant strata and their response to coastal erosion. The majority of these cliffs carry
53 landslides of one form or another and coastal instability and retreat is the dominant
54 natural process.
55
56
57
58
59
60
61
62
63
64
65

1 National Grid References quoted all fall within 100km grid square SZ. The
2 photographs used are part of the BGS collection (prefixed by a P number) and are
3 available on the BGS website (www.bgs.ac.uk) via GeoScenic.

4 **2 Landslide distribution and geology**

7 Each of the exposed geological units is susceptible to landsliding as the result of its
8 own failure or as the result of failure of the underlying unit(s). Lithology provides the
9 major control on the style and extent of landslide. The larger scale and most extensive
10 landslides occur on the slopes around the Southern Downs (Fig. 1). Away from these
11 chalk downs, inland landslides are rare. Smaller-scale landslides are widespread
12 around most of the coast, and are only absent in low-lying areas such as Sandown Bay
13 and along the northeast coastline. This may be a function of the greater incidence of
14 coastal woodland helping to stabilise the low cliffs or indeed disguising the true
15 degree of failure, and some notable examples of failure are known within wooded
16 areas (e.g. Seagrove Bay).
17
18
19

20 **2.1 Wealden Group**

22 The Wealden Group, consisting of interbedded mudstones and sandstones, can be
23 divided into the Wessex and Vectis formations (Table 1). The Wessex Formation is
24 composed of a non-marine sequence comprising multicoloured, predominantly red,
25 mudstones with subordinate sandstones. The Vectis Formation is principally
26 composed of dark grey siltstones and mudstones which are thought to have been
27 deposited in a shallow lagoonal environment. It is subdivided into three members:
28 Cowleaze Chine, Barnes High Sandstone and Shepherd's Chine Members (Daley &
29 Stewart, 1979). Both formations are highly susceptible to landslide failures,
30 particularly where they are exposed to storms on the southwest coast. The character of
31 the failure is determined by the presence and thickness of sandstone units within both
32 the Wessex and Vectis formations. This is illustrated in Fig. 2 using an example from
33 the Wessex Formation along the Brook Undercliff. Here three basic types of failure
34 can be identified (Zones A to C) associated with both formations. These are illustrated
35 in Figs. 2 to 5.
36
37
38
39
40

41 Zone A represents whole cliff activity, with thinner sandstones being incorporated
42 into landslides as rotated blocks and where the sandstones only act to limit failure
43 when outcropping at beach level and the cliff above is relatively low (Fig. 3). Zone B
44 represents cliffs where a thick competent sandstone acts as a natural revetment behind
45 the beach or as a reinforcement to the cliff when above the beach level. The degree
46 and inland extent of failure of the units above the competent sandstone is determined
47 by the remaining height of the cliff. Failure of the sandstone leads to debris cones of
48 material spilling onto the beach and abutting the sandstone unit (Fig. 4). In general,
49 clay-rich units, when exposed below the competent sandstone, retain a steeper profile
50 as they are frequently removed during storm events. In this scenario the sandstone
51 generally fails by undercutting and block-failure. Zone C is illustrative of the cliff
52 where no significant sandstone reinforcement is present and the whole cliff is active
53 even where cliffs are low (Fig. 5). General observations indicate that the retreat of the
54 cliffline is different within each of the zones; there is an increase of activity from
55 Zone B into Zone A and the most severe retreat is associated with Zone C.
56
57
58
59
60
61
62
63
64
65

1 Along the southwest coast, these zones can be most clearly seen between Roughland
2 Cliff [38578 83404] and Chilton Chine [40795 82100], where the Wessex Formation
3 is visible; similar features can be seen from Chilton Chine to Barnes High [43782
4 80666], from Barnes High along the coast toward Cowleaze Chine [44411 80020] and
5 Shepherds Chine [46450 79815]. Here the Barnes High Sandstone (Fig. 6) forms the
6 middle unit of three within the Vectis Formation. Landsliding is much more advanced
7 (i.e. well developed backscar, bench and lobe features) where the whole cliff is
8 composed of the Vectis Formation. Large scale rotational landslides and mudslides
9 have developed in the Shepherds Chine Member (Fig. 7) beneath the Atherfield Clay
10 Formation of the overlying Lower Greensand Group. These failures are mirrored at
11 Compton Bay [37650 84175] where the Wealden Group and the Lower Greensand
12 Group are eroding via a series of massive active landslides. The smallest of this group
13 of landslides is destroying the car park at Shippards Chine [37715 84000]. This
14 particular shallow, rotational failure, investigated by Bromhead et al. (2002), occurs
15 within mudstone of the Wessex Formation and has a basal slide surface controlled by
16 a bedding plane.
17
18
19
20

21 In Sandown Bay, on the southeast coast, landslides are present along the undefended
22 section of coast between the landing stage near Yaverland [61207 85016] and the
23 steep cliffs formed of higher strata at the eastern end of the bay [62726 85482]. The
24 landslides along this section form a continuous belt of shallow mud and debris slides
25 in mudstones of the Wessex Formation (Fig. 8). They pose a threat not only to users
26 of the beach but also to the buildings that back onto the cliffs, some at a distance of
27 only 60 m. By analogy with failures on the southwest coast, this section is
28 characterised by failures of the type in Zone C, as discussed above.
29
30

31 **2.2 Lower Greensand Group**

32
33

34 The Lower Greensand Group (Table 1) is dominated by mainly sand lithologies and
35 ranges from sandy mudstones of the Atherfield Clay Formation, to unconsolidated
36 sands, siltstones, mudstones and limestones of the Sandrock Formation. The Monks
37 Bay Sandstone Formation comprises coarse ferruginous sands and grits whilst the
38 Ferruginous Sands Formation is highly variable with sandy clays, glauconitic sands
39 pebble beds, siltstones and mudstones. Strata within the group are frequently subject
40 to landsliding where it forms outcrop at the coast. The Lower Greensand Group is
41 also involved with, but not necessarily the causal feature of, inland landsliding.
42
43
44

45 The largest landslide complex in Compton Bay [37132 84921] involves the Atherfield
46 Clay Formation and most of the outcrop of the Ferruginous Sands Formation.
47 Landsliding decreases in intensity as dips increase to the north-northwest into the
48 Brighstone Anticline. The failure is within the Atherfield Clay Formation and
49 probably in the upper part of the underlying Vectis Formation. The overlying
50 Sandrock and Monks Bay Sandstone formations only appear to be susceptible to
51 isolated rockfalls at this location. This probably results from the oblique orientation of
52 the bedding strike of the units with respect to the cliff-line and their steep dip into the
53 cliffs. The largest of the failures in the Compton Bay landslide complex is
54 approximately 750 m wide. Local degradation and vegetation (Fig. 9) indicate the
55 antiquity of parts of the landslide. However, rotated blocks, ponding and flows
56 indicate more recent activity.
57
58
59
60
61
62
63
64
65

1 The stratigraphical sequence in Compton Bay is repeated on the other side of the
2 island in Sandown Bay, but here the beds dip at approximately 10° to the north-
3 northeast. At Red Cliff [62360 85380] shallow translational landslides in the Wessex
4 and Vectis formations (see above) give way to a much larger landslide complex
5 developed in the Atherfield Clay Formation. The boundary of the Vectis and
6 Atherfield Clay formations forms a zone that is subject to large scale landsliding, as at
7 Cowleaze and Shepherd's Chine. Translational and rotational landsliding, and isolated
8 mudslides are present and large rotated blocks were observed at the time of the survey
9 in 2007 (Fig. 10). The landslide complex extends up to 150 m inland and is
10 approximately 115 m wide at beach level. It is known to have been almost
11 continuously active since at least 1912 (Hutchinson, 1965).
12
13

14 Considerably more complex failures occur within the Lower Greensand Group on the
15 coast at Chale Bay from Blackgang Chine [48400 76800] to Walpen Chine [46800
16 78200]. Here a bench has developed mid way up the cliff in the Ferruginous Sands
17 and Sandrock formations. Seepage erosion is taking place because of the presence of
18 clay aquitards within the sequence. Water movement along the top of the aquitards
19 leads to the undermining of material, leading to rock falls. Once on the bench the
20 landslide debris fails in a series of compound slides moving seaward. This process
21 produces a cliff top retreat rate that is more rapid than the marine erosion at the base
22 of the cliff, leading to the formation of the bench feature (Fig. 11) (Insole et al.,
23 1998).
24
25
26
27

28 **2.3 Selborne Group**

29

30 The Selborne Group consists of dark blue-grey silty clays of the Gault Clay
31 Formation passing transitionally upwards into glauconitic siltstone and fine-grained
32 sand and sandstone (with bands of calcareous and siliceous concretions) of the Upper
33 Greensand Formation. The Selborne Group is the controlling geological factor in the
34 landsliding of the Undercliff at Ventnor. This is an area that has become
35 internationally renowned as an example of good practice in landslide investigation
36 and remediation. The area has been the subject of two large conferences to
37 demonstrate best practice in dealing with landslide problems (McInnes and Jakeways,
38 2002, McInnes et al., 2007). There is a large volume of documentation relating to the
39 area (Hutchinson, 1991, Hutchinson et al., 1991a, Hutchinson et al., 1991b, Moore et
40 al., 2007 and Moore et al., 2010) and so the area is not discussed in detail here.
41 However the area is geologically similar to the Luccombe area discussed below.
42
43
44
45

46 The ancient landslide complex at Luccombe Chine [58090 79370] is very similar to
47 that at Ventnor. This landslide is known to have been reactivated in the 19th and 20th
48 Centuries and currently poses a potential threat to Luccombe Village. Lee and Moore
49 (1989) divided the landslide into 3 separate systems: ancient landslide units;
50 landsliding within a contemporary degradation zone; and erosion and landsliding of
51 modern sea cliffs. Ancient landsliding is still visible, represented by the degraded
52 multiple rotational blocks present above the village (Fig. 12). The back scars of these
53 features are within the Upper Greensand Formation, whilst it is postulated that the
54 shear surfaces lie within an upper clay-rich layer in the underlying Gault Formation
55 (Lee and Moore, 1989). Since the 1950s, reactivation of the contemporary
56 degradation zone has led to ground movements within Luccombe Village and the
57 destruction of several houses. Luccombe Village itself is built on a relict landslide
58
59
60
61
62
63
64
65

1 feature which has been reactivated due to cliff recession unloading the slope.
2 Construction of the houses, as well as drainage from them, is considered to have
3 contributed to this instability. Alteration of ground conditions, combined with intense
4 rainfall events, have led to periodic, major reactivations taking place. Demolition of at
5 least two houses was necessary after movements in 1987 and 1988 (Lee and Moore,
6 1989).

7
8 Large-scale inland landslides are also associated with the Selborne Group, particularly
9 on the south-western and northern margins of the Southern Downs. The hillsides
10 around the downs of St Catherine's [49360 78430], Appuldurcombe [53610 80450],
11 St Martin's [55570 80600], and Shanklin [57015 80165] all exhibit deep-seated
12 rotational landslides extending for several hundred metres across and down slope.
13 Large benches and blocks of Upper Greensand have failed and rotated on the
14 underlying Gault Formation. Prominent cliffs have formed in the upper portion of the
15 hillsides where backscars have formed in the Upper Greensand Formation (Fig.13).
16 Most of these features appear stable, but smaller scale failures within the broader
17 landslide zone display remarkably fresh landforms and they still occur depending on a
18 number of factors. Intense rainfall events are probably the most important, increasing
19 seepages within the chaotically deposited slip material and resulting in upslope
20 failures. The age of the larger landslide features is open to speculation but they
21 probably fall in the 'ancient landslide' category adopted by Lee and Moore (1989).
22 Very similar relict landslides occur in the same geological setting (e.g. Hopson, 1999)
23 near the Hampshire/Sussex border around the western closure of the Weald and here
24 they are considered to be Late- or immediately post-Devensian in age. They are
25 considered to have resulted from periglacial unlocking and remobilisation of
26 groundwater during the climatic amelioration at the end of the Devensian. This
27 resulted in high moisture content and pore pressures as well as spring-head erosion at
28 the base of the Upper Greensand. This is envisaged by the authors as the mechanism
29 of failure for these landslides on the Isle of Wight.
30
31
32
33
34
35

36 **2.4 White and Grey Chalk Subgroups**

37
38 The White and Grey Chalk Subgroups comprise parts of the previously named Upper,
39 Middle and Lower Chalk. The Grey Chalk Subgroup is a clayey chalk without flints
40 while the White Chalk Subgroup is a chalk with flints along with some discrete marl
41 seams. The chalk subgroups form steep, high cliffs between Compton Bay and the
42 Needles [28965 84840] in the west and at Culver Cliff [463710 85395] in the east
43 (Fig. 1). The cliffs in both areas are subject to falls of material. Some sliding is
44 controlled by the presence of well-developed bedding and joint planes and the attitude
45 of that bedding to the foreshore and erosion.
46
47
48

49
50 At Afton Down [36075 85580] in the west, failures in the chalk have required a major
51 realignment of the A3055 Military Road. This important route across the island runs
52 approximately 10-45 m from the edge of the 70 m high sea cliff. Fissuring in the
53 weathered chalk at the top of the cliff was observed to run roughly parallel to the cliff
54 edge during field work in 2007. At the top of the cliff, the beds dip steeply into the
55 cliff (65° towards the north), with widely spaced (1-2 m), low angled joints that are
56 parallel to the slope. Failure occurs by translational movement along the shoreward-
57 facing joint planes. Once the material has translated toward the edge of the cliff,
58 fissures develop in the soil and rock mass, leading to toppling failures (Anon, 2007).
59
60
61
62
63
64
65

1 Marine erosion at the site is also a significant factor leading to undercutting of the
2 cliff and failure of material by falling. Similar fissure-like features are found on the
3 southern margins of Culver Cliff where it abuts Sandown Bay [6350 8550], although
4 the fissuring does not appear to be as advanced. Little work has been conducted at this
5 site as it does not directly affect any infrastructure. Photographs collected during an
6 airborne geophysics survey by the BGS in September 2008 reveal the presence of
7 large tension cracks above Tennyson Down at the western end of the island,
8 suggesting that further failures along this section are also likely (Fig.14).
9

10 The chalk that forms the headland in the south of Alum Bay [30000 85000] has a
11 steep northerly dip into the bay with bedding planes being commonly undercut by
12 wave-action. This difference in orientation (e.g. compared to Afton Down) results in
13 failure on bedding planes rather than along joints. Typically, rock slides occur where
14 the dip of the beds is approximately parallel to the topographic slope. In Alum Bay
15 the chalk dips between 70-75° to the north, whilst the cliff slopes at about 60° to the
16 north. It is possible that the movement includes both translational movement and
17 falling (Fig.15).
18
19
20
21

22 **2.5 Lambeth, Thames, Bracklesham and Barton Groups**

23

24 The Reading Formation (Lambeth Group) is composed of predominantly mottled red
25 and purple clays with a basal conglomerate and sand. It outcrops in the west at Alum
26 Bay and in the east at Whitecliff Bay [463975 85830]. In both areas bedding is
27 vertical and the entire formation is subject to mudsliding (Fig. 16). The landslide in
28 Alum Bay is reported to have originally formed around the turn of the 20th Century
29 (Bromhead, 1979) and is most active between winter and early spring when the steep
30 slopes at its head become saturated due to higher rainfall levels. This causes the slope
31 to become unstable and fail as a mudslide, resulting in the formation of lobate toes
32 extending onto the beach (Bromhead, 1979). The toe area of the mudslide is subject to
33 marine erosion and is often eroded back to the cliff line during summer months, when
34 rainfall levels are generally lower and the slide is less active.
35
36
37
38

39 The Thames Group is represented by the London Clay Formation which consists of a
40 series of upward coarsening units composed of clays, silts and sands. The pattern of
41 landsliding in the London Clay Formation is similar to that in the Reading Formation.
42 Narrow gullies in both Alum Bay and Whitecliff Bay have formed between the
43 vertical strata as a result of the removal of material by mudsliding, although at both
44 localities the slides are not as active or advanced as those in the Reading Formation.
45
46

47 The Bracklesham Group is divided into six formations on the western side of the
48 island and four formations on the eastern side (Table 2). They generally consist of a
49 rhythmic sequence of glauconitic sands and clays at the base, succeeded by
50 interbedded clays, silts, and sands and also lignites. The Barton Group is divided into
51 four formations (Table 2) which are composed mainly of sands with subordinate clays
52 and silts. The Bracklesham Group and Barton Group in Alum Bay are not generally
53 susceptible to widespread landsliding, but are susceptible to shallow rotations,
54 mudslides and gully erosion. In Whitecliff Bay, a failure in the Wittering
55 Formation (Bracklesham Group) has occurred. This shallow mudslide is developed in
56 vertically-bedded laminated sandy muds and in 2007 had receded to within 20 m of a
57 caravan park.
58
59
60
61
62
63
64
65

2.6 Solent Group

The Solent Group is divided into three formations; the Headon Hill Formation is divided from the younger Bouldnor Formation by the Bembridge Limestone Formation. The Headon Hill Formation is separated into nine members (see Fig. 8 in Hopson, this issue) which are comprised of clays, sandy clays and limestones. The Bembridge Limestone Formation consists of limestone and marly limestone with shells, including peloidal limestone and calcrete with calcareous clay interbeds. The Bouldnor Formation comprises three members: the Bembridge Marls Member which is composed of grey-green clays and sandy clays, the Hamstead Member consisting of green clays and black organic sediments and the Cranmore Member is represented by shelly clays. The group contains interbedded units that may enhance or inhibit potential failure. For example where sand units are present they may carry groundwater or offer a conduit for surface run-off, or where thicker limestone units are present they may provide reinforcement to the cliff face. Examples of the types of landslide present are given below but there is a need for further study to provide a comprehensive appraisal of this group throughout the island.

Outcrops of the Headon Hill Formation are subject to active erosion and landsliding, especially along the coastline from Fort Albert [33010 89090] to Yarmouth [34715 89750]. Shallow mudslides and flows are currently eroding the cliffs at Fort Victoria [33615 89630], forming embayments and lobes. Larger landslides are also present on this stretch of coastline (Fig. 17). In the lower sections of the cliff, landsliding is occurring through failure at relatively shallow depth whilst higher in the cliff, deep-seated rotational landslides result in recession of the cliff line.

Within the wooded areas of Fort Victoria Country Park [33630 89580] there is a conspicuous landslide topography of degraded hummocks and ridges. These relict features indicate that there has previously been a more pervasive period of landsliding along this section of coastline and that it was more advanced than the shallow failures that are currently active within the cliff. It is likely that these ancient landslides formed when sea level was lower (Lambeck, 1997) than the present day, and their slip planes are thought to propagate beneath current beach level. Evidence for this larger-scale ancient activity is present in the beach in front of the cliff. The Fishbourne Beds and Osbourne Member of the Headon Hill Formation, seen as horizontally-bedded units in the mid-cliff section, are also found on the beach platform as steeply-dipping rotated, back-tilted blocks (Fig. 18) indicating a buried slip plane at depth. Hutchinson and Bromhead (2002) also described a raft of Headon Hill Formation with vertical strata near to this site at Bouldnor Cliff [38810 90955].

Rotational landslides and mudslides are a feature of the Bouldnor and Hamstead [39845 91630] cliffs on the NW coast. The cliffs along this section of coastline are composed entirely of the Bouldnor Formation, which is weak and susceptible to marine erosion. The mudslides are thought to have been initiated at the turn of the 20th Century and are seasonal (White, 1921). They appear to be controlled by high precipitation, high ground water levels and associated elevated pore water pressures (Bromhead, 1979). Denness (1970) suggested that the presence of a syncline behind Hamstead Cliffs plunging toward the sea provides a directed pathway for groundwater towards the cliff, resulting in increased activity at this site.

1 Another control on the type and activity of landsliding along this section of coastline
2 is the presence and height of the resistant Bembridge Limestone Formation beneath
3 the Bouldnor Formation. To the east of Bouldnor the Bembridge Limestone crops out
4 at beach level; the increased resistance provided by the limestone at the base of the
5 cliff reduces the rate of recession and decreases the activity and scale of mudsliding in
6 the overlying Bouldnor Formation (Hutchinson, 1983). To the west of Bouldnor, the
7 limestone is not exposed in the beach and this leads to increased marine erosion of the
8 soft muds. The rate of erosion at the toe is greater than the rate of mudsliding, leading
9 to an oversteepening of the cliffs. This, in turn, leads to the formation of deeper-
10 seated rotational landslides (Bromhead, 1979).
11

12
13 Landsliding of varying type, activity and age is present on the coast between Cowes
14 [49580 96380] and Gurnard [47240 95455]. Large, deep-seated, ancient, degraded
15 landslides form many of the coastal slopes in this area. Evidence for modern
16 landsliding is also apparent in both the developed and undeveloped areas. Houses
17 show evidence of displacement and settlement and cracks commonly appear in the
18 road. Heaving of the landslide toe is evident along the Esplanade. Further inland
19 along the coastal slope, benches and fresh scarps provide evidence for more recent
20 movement (Halcrow, 2000). Erosion rate at the toe exceeding the cliff-fall material
21 replenishment rate in this area has led to active recession of the backscar, as indicated
22 by the loss of part of a garden from a private residence on the cliff at Gurnard during
23 the winter of 2007/08.
24
25
26

27
28 Landslide activity at Seagrove Bay [63130 91070], on the northeast coast presents a
29 significant hazard to local infrastructure. There is a long history of property damage
30 due to ground movement (Winfield et al., 2007). The whole of the slope behind the
31 bay is critically stable, i.e. it is close to failure and has a factor of safety very close to
32 one. A change in the environmental conditions, such as an increase in the level of the
33 local water table, or removal of support from the toe of the landslide by beach
34 erosion, could result in failure of the slope.
35
36
37

38 **3 Discussion**

39

40 The majority of the geological strata that crop out on the Isle of Wight are susceptible
41 to landsliding, either directly or indirectly. The geological structure of the island also
42 influences landslide frequency, with the steeply-dipping strata around the central axis
43 (Fig. 1) facilitating a greater incidence of landsliding. However landslides are most
44 common around the coast. Ancient, advanced landslides are present on the slopes of
45 the Southern Downs. They formed during periglacial conditions but are currently
46 inactive. The stratigraphy here is similar to that of the active landslide complex at
47 Ventnor, but the slides are stable because no material is removed from the toe.
48 Therefore it is coastal erosion that is the primary causal factor of active landsliding on
49 the island.
50
51
52

53
54 Coastal aspect almost certainly has an influence on landslide frequency around the
55 island. Taking the example of the Wealden Group strata that crop out in the southern
56 half of the island, landslide frequency and thus coastal recession is likely to be highest
57 along the south west coast between The Needles [29000 84825] and St Catherine's
58 Point [49850 75265] where the full effect of storms and a wave fetch of several
59 thousand kilometres impact on the cliffs. Conversely in Sandown Bay, with a more
60
61
62
63
64
65

1 sheltered south-easterly aspect, the cliffs are more protected from storms and wave
2 energy is much lower. Therefore the frequency of landslides and coastal recession is
3 likely to be lower.

4
5 Cliff height influences landslide style due to the accommodation space, and therefore
6 mass of material, available for a landslide to occur (i.e. higher cliffs lead to larger,
7 more advanced landslides). The landsliding along the coastal section between
8 Compton Chine [36780 85180] and Hanover Point [37940 83770] provides a good
9 example of this. A large, relatively deep-seated, multiple, rotational landslide has
10 developed where the cliffs are highest at the western end of the bay. As the cliffs
11 become lower along the coast to the south east, landslides are less advanced and
12 consist of shallow rotations and mudflows due to less accommodation space being
13 available.
14
15

16
17 Extensive, active landslides are present along the northwest coast of the island. This
18 area of the island is currently underdeveloped and as a result there are very few
19 published studies of the landslides. With a likely future increase in population and the
20 associated increasing pressure on the landscape through residential and tourism
21 infrastructure, there is a need for more detailed understanding of the engineering
22 geological properties of the strata present along this stretch of coastline, and the
23 failure mechanisms of the landslides, to ensure sustainable development of the area.
24
25

26
27 Climate change forecasts such as an increase in the frequency and intensity of storm
28 events and sea level rise (Jenkins et al., 2009), will have serious implications for the
29 unstable coast around the Isle of Wight. The frequency and magnitude of landslides
30 are likely to increase leading to an increased rate of coastal recession. Higher rainfall
31 levels may also lead to the reactivation of the currently stable inland landslides around
32 the Southern Downs. Fresh scarps observed during the 2007 survey provide evidence
33 of more recent, small-scale movement and suggest that the larger, ancient landslides
34 could reactivate if rainfall event volumes and frequencies become greater in the
35 future.
36
37

38 39 **4 Conclusions**

40
41 Landslides on the Isle of Wight are highly variable. They include large deep-seated
42 multiple rotations associated with the Downs of Appuldurcombe, St Catherine's, St
43 Martin's and Shanklin, and shallower rotations and flows around the coast. The
44 majority of geological formations present on the Isle of Wight are susceptible to
45 landsliding. Many of the formations are composed of soft, poorly lithified sediments
46 making them highly susceptible to coastal erosion. Erosion removes support from the
47 base of the cliff and leads to subsequent failure. The widespread distribution of
48 landslides present potentially significant engineering and infrastructure problems;
49 notable large and active landslides occur at Ventnor, at Luccombe and Seagrove Bay
50 and along the Military Road at Afton Down.
51
52

53
54 The style and extent of landslides on the island is controlled by a complex interplay of
55 several factors: coastal position; the potential for unloading of the toe by marine
56 erosion; the degree of natural reinforcing within the lithological components of the
57 unit; the incidence of that reinforcement in respect of its position within the landform
58 concerned; geological structure; the incidence of significant climatic events (intense
59
60
61
62
63
64
65

1 storm activity and rising sea level). Landsliding is most frequent and active around the
2 coast due the effects of coastal erosion.

3
4 The northwest coastline of the island is a key target for future landslide research.
5 Focussed testing of engineering geological properties, and more detailed analysis of
6 the mechanisms of failure along this coast would ensure sustainable future
7 development.
8

9
10 If forecasts of climate change prove to be correct, then higher sea levels and increased
11 storm intensity and frequency are likely to exacerbate the frequency and scale of
12 landsliding on the Isle of Wight.
13

14 15 16 17 **Acknowledgements**

18
19 The authors would like to thank Anthony Cooper, Helen Reeves, Philip Wilby and
20 Chris Thomas for constructive advice and guidance during the drafting of this paper.
21 Gareth Jenkins, Claire Foster and Peter Hopson publish with the permission of the
22 Executive Director of the British Geological Survey (NERC).
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

References

- 1
2 Anon, 1993. Low-rise buildings on shrinkable clay soils: Part 1, Digest 240. Building
3 Research Establishment. Watford.
4
- 5 Anon 2007. International conference on Landslides and Climate Change- Challenges
6 and Solutions. Field Study Notes. Ventnor, Isle of Wight, 21-24th May 2007.
7
- 8 Bell, F. G. 2000. Engineering properties of soils and rocks. 4th Edition. Blackwell,
9 Oxford.
10
- 11 Booth, K. A. and Brayson, J. 2011. Geology, landscape and human interactions:
12 Examples from the Isle of Wight. Proceedings of the Geologists' Association. 122,
13 XXX-XXX.
14
- 15 Bromhead, E. N. (1979) Factors Affecting the Transition Between the Various Types
16 of Mass Movement in Coastal Cliffs Consisting Largely of Over-Consolidated Clay,
17 with Special Reference to Southern England, Quarterly Journal of Engineering
18 Geology. 12(4), 291-300.
19
- 20 Bromhead, E. N., Ibsen, M-L., Papanastassiou, X and Zemichael, A.A. 2002. Three-
21 dimensional stability analysis of a coastal landslide at Hanover Point, Isle of Wight.
22 Quarterly Journal of Engineering Geology & Hydrogeology. 35(1), 79-88.
23
- 24 British Geological Survey. 1976. Isle of Wight. Special Sheet. Part of sheets 330, 331.
25 Sheets 344, 345. Drift Edition. Ordnance Survey, Southampton.
26
- 27 Chadwick, A.1986. Extension tectonics in the Wessex Basin, southern England.
28 Journal of the Geological Society, London. 143, 465-488.
29
- 30 Daley, B. and Stewart, D. J. 1979. Week-end field meeting: the Wealden Group in the
31 Isle of Wight 17–19 June 1977. Proceedings of the Geologists' Association. 90, 51–
32 54.
33
- 34 Denness, B. 1970. A note on the landslip at Hampstead cliff, Isle of Wight.
35 Engineering Geology Group Report WN/EG/70/004. British Geological Survey.
36
- 37 Halcrow Group Ltd, 2000. Cowes to Gurnard coastal slope stability study: ground
38 behaviour assessment. Isle of Wight Council.
39
- 40 Hooker, J. J., Grimes, S. T., Matthey, D. P., Collinson, M. E. and Sheldon, N. D. 2009.
41 Refined correlation of the UK Late Eocene-Early Oligocene Solent Group and timing
42 of its climate history. *In* Koeberl, C and Montanari, A. (eds), The Late Eocene Earth –
43 Hothouse, Icehouse and Impacts: The Geological Society of America Special Paper
44 452, 179-195.
45
- 46 Hopson, P.M., 1999. Geology of the Fareham and Portsmouth district. Sheet
47 Description of the British Geological Survey, Sheet 316 and part of 331 (England and
48 Wales).
49
- 50 Hopson, P. M. 2011. The geological history of the Isle of Wight: an overview of the
51 'diamond in Britain's geological crown'. Proceedings of the Geologists' Association
52 122, XXX-XXX.
53
- 54 Hutchinson, J. N. 1965. A reconnaissance of coastal landslides in the Isle of Wight.
55 Building Research Station Note No. EN 11/65. Department of Scientific and
56 Industrial Research. Watford.
57
58
59
60
61
62
63
64
65

1 Hutchinson, J. N. 1983. A Pattern in the Incidence of Major Coastal Mudslides, Earth
2 Surface Processes and Landforms. 8, 391-397.

3 Hutchinson, J. N. 1991. The landslides forming the South Wight Undercliff. *In*
4 Chandler, R. J. (eds). Slope Stability Engineering: Developments and Applications.
5 Proceedings of the International Conference on Slope Stability Organized by the
6 Institution of Civil Engineers and held on the Isle of Wight on 15–18 April 1991 (ed.
7 R.J. Chandler), Thomas Telford, London. 157–68.

8
9 Hutchinson J. N., Brunsden D. and Lee E. M. 1991a. The geomorphology of the
10 landslide complexes at Ventnor, Isle of Wight. *In*: Chandler R.J. (ed.) Slope Stability
11 Engineering, Developments and applications. Proc. of the Int. Conf. on Slope Stability
12 organised by the ICE, Shanklin, Isle of Wight. 213-218.

13
14 Hutchinson, J. N., Bromhead, E. N. and Chandler, R. J. 1991b. Investigations of the
15 landslides at St Catherine’s Point, Isle of Wight. *In* Chandler, R. J. (ed). Slope
16 Stability Engineering: Developments and Applications. Proceedings of the
17 International Conference on Slope Stability Organized by the Institution of Civil
18 Engineers and held on the Isle of Wight on 15–18 April 1991, Thomas Telford,
19 London. 169-179.

20
21 Hutchinson, J. N. and Bromhead, E. N. 2002. Isle of Wight Landslides. *In* McInnes,
22 R. G. and Jakeways, J. (eds), Instability Planning and Management. Proceedings of
23 the International Conference. Ventnor, Isle of Wight, UK. 20-23rd May 2002.
24 Thomas Telford.

25
26 Insole, A., Daley, B. and Gale, A. 1998. The Isle of Wight. Geologists Association
27 Guide No.60. The Geologists Association.

28
29 Jenkins, G. J., Murphy, J. M., Sexton, D. M. H., Lowe, J. A., Jones, P. and Kilsby, C.
30 G. (2009). UK Climate Projections: Briefing report. Met Office Hadley Centre,
31 Exeter, UK.

32
33 Lambeck, K. 1997. Sea-level change along the French Atlantic and Channel coasts
34 since the time of the Last Glacial Maximum. *Palaeogeography, Palaeoclimatology,*
35 *Palaeoecology.* 129, 1-22.

36
37 Lee, E. M. and Moore, R. 1989. Report on the study of landsliding in and around
38 Luccombe Village. Department of the Environment.

39
40 McInnes, R. G. & Jakeways, J. 2002. Instability: Planning and Management.
41 Seeking sustainable solutions to ground movement problems. Proceedings of the
42 international conference organised by the Centre for the Coastal Environment, Isle of
43 Wight Council, Ventnor, Isle of Wight, UK, 20-23 May 2002.

44
45 McInnes, R. G., Jakeways, J., Fairbank, H. and Mathie, E. 2007. Landslides and
46 Climate Change: Challenges and Solutions. Proceedings of the International
47 Conference on Landslides and Climate Change, Ventnor, Isle of Wight, UK, 21-24
48 May 2007.

49
50 Moore, R., Turner, M. D., Palmer, M. J. and Carey, J. M. 2007. The Ventnor
51 Undercliff: Landslide model, mechanisms and causes, and the implications of climate
52 change induced ground behaviour and risk. *In* McInnes, R., Jakeways, J, Fairbank, H.
53 and Mathie, E. (eds), Landslides and Climate Change, Challenges and Solutions.
54 Proceedings of the International Conference on Landslides and Climate Change,
55 Ventnor, Isle of Wight, May 21-24 2007. 365-376. Thomas Telford.

56
57
58
59
60
61
62
63
64
65

1 Moore, R., Carey, J. M. and McInnes, R. G. 2010. Landslide behaviour and climate
2 change; predictable consequences for the Ventnor Undecliff, Isle of Wight. Quarterly
3 Journal of Engineering Geology & Hydrogeology. 43(4), 447-460.

4 Preece R. C., Scourse, J. D., Houghton, S .D., Knudsen, K. L. and Penney, D. N.,
5 1990. The Pleistocene sea-level and neotectonic history of the eastern Solent,
6 southern England. Philosophical Transactions of the Royal Society. B328, 425–477.
7

8 Radley, D. 2006. A Wealden guide II: the Wessex sub-basin. Geology Today. Vol 22
9 Issue 5. 187-193.
10

11 White H.J.O. 1921. A short account of the geology of the Isle of Wight. Memoir of
12 the Geological Survey of Great Britain (England and Wales). HMSO.
13

14 Winfield, P., Moses, E. and Woodruff, M. 2007. Combining slope stability and coast
15 protection at Seagrove Bay on the Isle of Wight. *In* McInnes, R., Jakeways, J,
16 Fairbank, H. and Mathie, E. (eds), Landslides and Climate Change, Challenges and
17 Solutions. Proceedings of the International Conference on Landslides and Climate
18 Change, Ventnor, Isle of Wight, May 21-24 2007. 365-376. Thomas Telford.
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Figure captions

1
2
3 Fig. 1. Simplified geological map of the Isle of Wight, with selected geological dip
4 readings. See Tables 1 and 2 for detailed stratigraphy. New landslide areas mapped
5 during the 2007 survey marked in orange. Geological data, British Geological Survey
6 © NERC. OS topography © Crown Copyright. All rights reserved. BGS
7 100017897/2011.
8

9
10 Fig. 2. Location of sandstones and their influence on landslides between Chilton
11 Chine and Roughlands (Modified from Insole *et al.*, 1998).
12

13 Fig. 3. Active landsliding near Chilton Chine (Fig. 2, Zone A). P760896.
14

15
16 Fig. 4. Presence of debris cones below a massive sandstone bed in the cliff at
17 Sudmoor Undercliff (Fig. 2, Zone B). P769944.
18

19 Fig. 5. Retreat of cliff line through active landsliding at Roughland Cliff, near Brook
20 (Fig. 2, Zone C). P769945.
21

22
23 Fig. 6. Failures in the Vectis Formation of the Wealden Group. P769946.
24

25 Fig. 7. Failures in the Shepherds Chine Member, Vectis Formation, east of Shepherds
26 Chine. P760858.
27

28
29 Fig. 8. Landsliding at Red Cliff, Sandown Bay. P769942.
30

31 Fig. 9. View across the large landslide complex in Compton Bay. P760901.
32

33
34 Fig. 10. Rotated block (Ferruginous Sands Formation) in the beach at Yaverland.
35 P769933.
36

37 Fig. 11. Bench feature between Walpen Chine and Blackgang Chine. P769966.
38

39
40 Fig. 12. View across Luccombe Coombe toward Knock Cliff. P769442.
41

42 Fig. 13. Backscars composed of the Upper Greensand Formation, St Catherine's
43 Down. P769458.
44

45
46 Fig. 14. Tension cracks above Tennyson Down which could lead to large scale
47 failures. P769938.
48

49
50 Fig. 15. Failure in steep chalk cliffs at White Cliffs, Alum Bay. P769937.
51

52 Fig. 16. Mudslide in the Reading Formation, Whitecliff Bay. P769939.
53

54
55 Fig. 17. Recent landslide at Fort Victoria. P769500.
56

57 Fig. 18. Steeply dipping bedded raft of Headon Hill Formation embedded in the beach
58 at Fort Victoria. P769941.
59
60
61
62
63
64
65

Table captions

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Table 1. The Cretaceous strata at outcrop on the Isle of Wight.

Table 2. The Palaeogene strata at outcrop on the Isle of Wight.

Table 1. The Cretaceous strata at outcrop on the Isle of Wight

Group or Subgroup		Formation
Chalk Group	White Chalk Subgroup	Portsdown Chalk Formation
		Culver Chalk Formation
		Newhaven Chalk Formation
		Seaford Chalk Formation
		Lewes Nodular Chalk Formation
		New Pit Chalk Formation
		Holywell Nodular Chalk Formation
	Grey Chalk Subgroup	Zig Zag Chalk Formation
		West Melbury Marly Chalk Formation
	Selborne Group	Upper Greensand Formation
		Gault Formation
Lower Greensand Group	Monk's Bay Sandstone Formation	
	Sandrock Formation	
	Ferruginous Sands Formation	
	Atherfield Clay Formation	
Wealden Group	Vectis Formation	
	Wessex Formation	

Table 2. The Palaeogene strata at outcrop on the Isle of Wight

Group	Formation	
Solent Group	Bouldnor Formation	
	Bembridge Limestone Formation	
	Headon Hill Formation	
Barton Group	WEST	EAST
	Becton Sand Formation	Becton Sand Formation
	Chama Sand Formation	Chama Sand Formation
	Barton Clay Formation	Barton Clay Formation
	Boscombe Sand Formation	
Bracklesham Group	WEST	EAST
	Branksome Sand Formation	Selsey Sand Formation
	Marsh Farm Formation	Marsh Farm Formation
	Poole Formation (Upper)	Earnley Sand Formation
	Wittering Formation (Upper)	Wittering Formation
	Poole Formation (Lower)	
Wittering Formation (Lower)		
Thames Group	London Clay Formation	
Lambeth Group	Reading Formation	

Figure 2
[Click here to download high resolution image](#)




Figure 3
[Click here to download high resolution image](#)



Figure 4
[Click here to download high resolution image](#)



Figure 5
[Click here to download high resolution image](#)



Figure 6
[Click here to download high resolution image](#)



Figure 7
[Click here to download high resolution image](#)




Figure 8
[Click here to download high resolution image](#)



Figure 9
[Click here to download high resolution image](#)



Figure 10
[Click here to download high resolution image](#)



Figure 11
[Click here to download high resolution image](#)




Figure 12
[Click here to download high resolution image](#)




Figure 13
[Click here to download high resolution image](#)




Figure 14

[Click here to download high resolution image](#)



Figure 15
[Click here to download high resolution image](#)



Figure 16
[Click here to download high resolution image](#)



Figure 17
[Click here to download high resolution image](#)




Figure 18

[Click here to download high resolution image](#)

