


BRITISH GEOLOGICAL SURVEY

Jim Carrigan Observatory Prudhoe Bay Monthly Magnetic Bulletin October 2008 08/10/JC



**British
Geological Survey**

NATIONAL ENVIRONMENT RESEARCH COUNCIL

1. JIM CARRIGAN OBSERVATORY MAGNETIC DATA

1. Introduction

Jim Carrigan Observatory is the fourth overseas geomagnetic observatory established by the British Geological Survey (BGS). The installation is a joint project between BGS and Sperry Drilling Services (SDS) in support of directional drilling programmes. SDS has operated a prototype station since 1997 which was upgraded by the BGS to a standard high-quality observatory in October 2003.

This bulletin is published to meet the needs of users of geomagnetic data. Magnetic observatory data is presented as a series of plots of one-minute, hourly and daily values, followed by a tabulation of monthly values. The operation of the observatory and presentation of data are described in the rest of this section.

Enquiries about the data should be addressed to:

National Geomagnetic Service
British Geological Survey
Murchison House, West Mains Road
Edinburgh EH9 3LA
Scotland, UK

Tel: +44 (0) 131 667 1000
Fax: +44 (0) 131 668 0265
E-mail: orba@bgs.ac.uk
Internet: www.geomag.bgs.ac.uk

2. Position

Jim Carrigan Observatory is situated at T-Pad, a man-made gravel bed close to the drilling sites at Prudhoe Bay, Alaska, USA.

The observatory co-ordinates are:-

Geographic: $70^{\circ} 21.4'N$ $211^{\circ} 12.1'E$
Geomagnetic: $70^{\circ} 04.5'N$ $254^{\circ} 01.7'E$
Height above mean sea level: *10m (approx)*

The geomagnetic co-ordinates are calculated using the 10th generation International Geomagnetic Reference Field (IGRF) at epoch 2008.5

3. The Observatory Operation

3.1 GDAS

The observatory operates under the control of the Geomagnetic Data Acquisition System (GDAS), which was developed by BGS and became fully operational from 27th October 2003. The system operates under the control of data acquisition software running on QNX computers, which controls the data logging and communications.

There are two sets of sensors used for making magnetic measurements. A triaxial linear-core fluxgate magnetometer, manufactured by the Danish Meteorological Institute, is used to measure the variations in the horizontal (H) and vertical (Z) components of the field. The third sensor is oriented perpendicular to these, and measures variations, which are proportional to the changes in declination (D). Measurements are made at a rate of 1 Hz.

In addition to the fluxgate sensors there is a proton precession magnetometer making measurements of the absolute total field intensity (F) at a rate of 0.1Hz.

The raw unfiltered data are retrieved automatically via internet connections to the BGS office in Edinburgh in near real-time. The fluxgate data are filtered to produce one-minute values using a 61-point cosine filter whilst the total field intensity samples are filtered using a 7-point cosine filter.

3.2 Absolute Observations

The GDAS fluxgate magnetometers accurately measure variations in the components of the geomagnetic field, but not the absolute magnitudes. Regular manual absolute measurements of the field are made throughout the year. A fluxgate sensor mounted on a theodolite is used to determine D and inclination (I); the GDAS PPM measurements, with a site difference correction applied, are used for F . The absolute observations are used in conjunction with the GDAS variometer measurements to produce a continuous record of the absolute values of the geomagnetic field elements as if they had been measured at the observatory reference pillar.

4. Data Presentation

The data presented in the bulletin are in the form of plots and tabulations described in the following sections.

4.1 Absolute Observations

The absolute observation measurements made during the month are tabulated. Also included are the corresponding baseline values, which are the differences between the absolute measurements and the variometer measurements of D , H and Z (in the sense absolute–variometer). These are also plotted (markers) along with the derived preliminary daily baseline values (line) throughout the year. Daily mean differences between the measured absolute F and the F computed from the baseline corrected H and Z values are plotted in the fourth panel (in the sense measured–derived). The bottom panel shows the daily mean temperature in the fluxgate chamber.

4.2 Summary magnetograms

Small-scale magnetograms are plotted which allow the month's data to be viewed at a glance. They are plotted 16 days a page and show the variations in D , H and Z . The scales are shown on the right-hand side of the page. On disturbed days the scales are multiplied by a factor, which is indicated above the panel for that day. The variations are centred on the monthly mean value, shown on the left side of the page.

4.3 Magnetograms

The daily magnetograms are plotted using one-minute values of D , H and Z from the fluxgate sensors, with any gaps filled using back-up data. The magnetograms are plotted to a variable scale; scale bars are shown to the right of each plot. The absolute level (the monthly mean value) is indicated on the left side of the plots.

4.4 Hourly Mean Value Plots

Hourly mean values of D , H and Z for the past 12 months are plotted in 27-day segments corresponding to the Bartels solar rotation number. Magnetic disturbances associated with active regions on the surface of the Sun may recur after 27 days: the same is true for geomagnetically quiet intervals. Plotting the data in this way highlights this recurrence, and also illustrates seasonal and diurnal variations throughout the year.

4.5 Daily and Monthly Mean Values

Daily mean values of D , H , Z and F are plotted throughout the year. In addition, a table of monthly mean values of all the geomagnetic elements is provided. These values depend on accurate specification of the fluxgate sensor baselines.


© NERC 2008

JIM CARRIGAN OBSERVATORY

ABSOLUTE OBSERVATIONS

		DECLINATION			INCLINATION							
Date	Day Number	Time (UT)	Absolute (°)	Baseline (°)	Time (UT)	Inclination (°)	Total Field Intensity (nT)	H Absolute (nT)	H Baseline (nT)	Z Absolute (nT)	Z Baseline (nT)	Observer
07-Oct-08	281	18:05	23.0269	24.3017	18:21	80.9704	57587.4	9038.2	9137.0	56873.7	56866.2	CWT
07-Oct-08	281	18:36	23.0175	24.2933	18:47	80.9691	57597.7	9041.0	9141.6	56883.7	56865.5	PS
07-Oct-08	281	19:02	23.0056	24.3017	19:17	80.9767	57601.9	9034.1	9137.5	56889.1	56865.9	CWT
11-Oct-08	285	21:19	22.8743	24.2883	21:35	80.9203	57585.1	9087.5	9140.4	56863.5	56866.3	PS
11-Oct-08	285	21:51	22.8113	24.2767	22:05	80.9367	57605.3	9074.3	9138.4	56886.1	56866.2	PS
11-Oct-08	285	22:18	22.8589	24.2867	22:27	80.9533	57615.0	9059.4	9140.3	56898.3	56866.3	PS
11-Oct-08	285	22:35	22.8409	24.2800	22:42	80.9456	57614.8	9067.1	9140.6	56896.9	56866.2	PS


Jim Carrigan Obs 2008




Jim Carrigan

October

2008






Date: 01-10-2008


Jim Carrigan

Day number: 275



Date: 02-10-2008


Day number: 276



Date: 03-10-2008


Jim Carrigan

Day number: 277



Date: 04-10-2008


Day number: 278



Date: 05-10-2008


Jim Carrigan

Day number: 279



Date: 06-10-2008


Day number: 280



Date: 07-10-2008


Jim Carrigan

Day number: 281



Date: 08-10-2008


Day number: 282



Date: 09-10-2008


Jim Carrigan

Day number: 283



Date: 10-10-2008


Day number: 284



Date: 11-10-2008


Jim Carrigan

Day number: 285



Date: 12-10-2008


Day number: 286



Date: 13-10-2008


Jim Carrigan

Day number: 287



Date: 14-10-2008


Day number: 288



Date: 15-10-2008


Jim Carrigan

Day number: 289



Date: 16-10-2008


Day number: 290



Date: 17-10-2008


Jim Carrigan

Day number: 291



Date: 18-10-2008


Day number: 292



Date: 19-10-2008


Jim Carrigan

Day number: 293



Date: 20-10-2008


Day number: 294



Date: 21-10-2008


Jim Carrigan

Day number: 295



Date: 22-10-2008


Day number: 296



Date: 23-10-2008


Jim Carrigan

Day number: 297



Date: 24-10-2008


Day number: 298



Date: 25-10-2008


Jim Carrigan

Day number: 299



Date: 26-10-2008


Day number: 300



Date: 27-10-2008


Jim Carrigan

Day number: 301



Date: 28-10-2008


Day number: 302



Date: 29-10-2008


Jim Carrigan

Day number: 303



Date: 30-10-2008


Day number: 304




Date: 31-10-2008

Jim Carrigan


Day number: 305




Jim Carrigan Observatory: Declination (degrees)




Jim Carrigan Observatory: Horizontal Intensity (nT)



Jim Carrigan Observatory: Vertical Intensity (nT)



Jim Carrigan Observatory 2008



Monthly Mean Values for Jim Carrigan Observatory 2008

Month	<i>D</i>	<i>H</i>	<i>I</i>	<i>X</i>	<i>Y</i>	<i>Z</i>	<i>F</i>
January	23° 14.8'	9030 nT	80° 58.8'	8297 nT	3564 nT	56887 nT	57599 nT
February	23° 12.1'	9011 nT	80° 60.0'	8282 nT	3550 nT	56893 nT	57602 nT
March	23° 9.1'	9013 nT	80° 59.9'	8287 nT	3544 nT	56897 nT	57607 nT
April	23° 6.4'	9019 nT	80° 59.6'	8295 nT	3540 nT	56903 nT	57613 nT
May	23° 4.0'	9040 nT	80° 58.3'	8318 nT	3542 nT	56892 nT	57606 nT
June	23° 1.7'	9041 nT	80° 58.2'	8321 nT	3537 nT	56889 nT	57603 nT
July	22° 59.2'	9038 nT	80° 58.4'	8320 nT	3529 nT	56890 nT	57604 nT
August	22° 58.2'	9038 nT	80° 58.4'	8321 nT	3527 nT	56891 nT	57605 nT
September	22° 55.0'	9028 nT	80° 59.0'	8316 nT	3515 nT	56896 nT	57608 nT
October	22° 53.4'	9017 nT	80° 59.7'	8307 nT	3507 nT	56901 nT	57611 nT

Note

- i. The values shown here are provisional.