

## Article (refereed) - postprint

---

von Zum Hof, Juergen; Schonrogge, Karsten; Cook, James M.; Gardner, Michael G. 2012 [Development of microsatellite markers using 454 sequencing for the rare socially parasitic hoverfly, \*Microdon mutabilis\*](#). *Australian Journal of Zoology*, 60 (2). 108-110. [10.1071/ZO12042](https://doi.org/10.1071/ZO12042)

This version available <http://nora.nerc.ac.uk/19884/>

NERC has developed NORA to enable users to access research outputs wholly or partially funded by NERC. Copyright and other rights for material on this site are retained by the rights owners. Users should read the terms and conditions of use of this material at <http://nora.nerc.ac.uk/policies.html#access>

**This document is the author's final manuscript version of the journal article, incorporating any revisions agreed during the peer review process. Some differences between this and the publisher's version remain. You are advised to consult the publisher's version if you wish to cite from this article.**

The definitive version is available at [www.publish.csiro.au/](http://www.publish.csiro.au/)

Contact CEH NORA team at  
[noraceh@ceh.ac.uk](mailto:noraceh@ceh.ac.uk)

**AUSTRALIAN JOURNAL OF ZOOLOGY - PRIMER NOTE: Development of microsatellite markers using 454 sequencing for the rare socially parasitic hoverfly *Microdon mutabilis*.**

Juergen von Zum Hof<sup>1</sup>, Karsten Schönrogge<sup>2</sup>, James M. Cook<sup>3</sup>, Michael G. Gardner<sup>1,4</sup>

<sup>1</sup>School of Biological Sciences, Flinders University GPO Box 2100, Adelaide, SA 5001, Australia.

<sup>2</sup>Centre for Ecology and Hydrology, CEH Wallingford,

Maclean Building, Benson Lane,

Wallingford, OX10 8BB, UK.

<sup>3</sup>School of Biological Sciences, University of Reading, Reading, Berkshire, RG6 6BX, UK.

<sup>4</sup>**Author for correspondence: Fax: +61 8 82013015; Email: [michael.gardner@flinders.edu.au](mailto:michael.gardner@flinders.edu.au)**

**Abstract**

To date there have only been limited fine-scale investigations into the molecular ecology of the European hoverfly *Microdon mutabilis* due to the paucity of available polymorphic markers. We describe the development of primers amplifying five novel using next generation sequencing (454) and three previously undescribed microsatellite *M. mutabilis* loci using enrichments. In hoverflies from a population in Ireland, the number of alleles per locus ranged from 2-16, and the observed heterozygosity ranged between 0.26 and 0.97.

**Keywords:** *Microdon mutabilis*, microsatellites, PCR, social parasite, genetic population structure, local adaptation

Running Head: *Microdon mutabilis* microsatellites

The rare European hoverfly *Microdon mutabilis* (Syrphidae: Microdontinae) is a cryptic myrmecophilous social parasite that displays extreme host specificity to *Formica lemani* ant colonies primarily within the British Isles and Scandinavia (Elmes *et al.* 1999, Schönrogge *et al.* 2006). Although cause and effect are unclear, the flies infest host ant nests that have high levels of genetic diversity (Gardner *et al.* 2007). The extreme local specificity is reflected in bioassays where geographic distance from natal nest of ovipositing females affects the survivorship of her eggs (for details see Elmes *et al.* 1999, Schönrogge *et al.* 2006) and we predict this could promote localised genetic structuring within the flies. To date there has only been limited fine scale investigations into the molecular ecology of *M. mutabilis* due to a lack of available polymorphic markers. An investigation at a fine scale will not only assist in understanding the measures that need to be implemented in the conservation of this rare fly, but also provide insights to the preservation of genetic diversity of its host ant species *F. lemani*. Here we detail the development of eight microsatellite loci, five novel loci and three of which were previously used by Schönrogge *et al.* (2006) but remain undescribed.

Two different methods of microsatellite development were used. Three loci were isolated using enrichments following Gardner *et al.* (1999) (Mmut9) or Hale *et al.* (2001) (CEH49, CEH52). All remaining loci were developed using next generation sequencing and subsequent bioinformatic screening following Gardner *et al.* (2011). Genomic DNA was extracted using the Gentra Puregene Tissue Kit (Qiagen, Doncaster, Australia) from the legs of 15 individuals of *M. mutabilis* collected from the Burren and from the Isle of Mull, and sequenced on a Titanium GS-FLX (454 Life Sciences/Roche FLX) at the Australian Genomic Research Facility (AGRF, Brisbane, Australia, [www.agrf.org.au](http://www.agrf.org.au)) following Gardner *et al.* (2011). A total of 55,312 reads were obtained with a mean read sequence length of 354bp from approximately 19.6 Mb of sequence. The program QDD ver 1.0 (Meglecz *et al.* 2010) was used on a windows system with the graphical user interface to

screen the 454 raw sequences for microsatellites (newer versions were not available at the time the study was conducted), check flanking regions to identify unique loci and design primers for perfect di-, tri- and tetranucleotides with a minimum of eight tandem repeats. Amplicon length was specified between 90-450bp and all other parameters in QDD were set to default. The installation of QDD ver 1.0 followed Meglec 2010 and also required the installation of ActivePerl ver 5.10 (<http://www.activestate.com/activeperl/>), BLAST ver 2.2.18 (<ftp://ftp.ncbi.nih.gov/blast/executables/>), CLUSTALw2 ver 1.83 (Larkin et al., 2007; <ftp://ftp.ebi.ac.uk/pub/software/clustalw2/>) and Primer3 (Rozen and Skaletsky, 2000; <http://primer3.sourceforge.net/>). This process yielded 94 putative microsatellite loci for *M. mutabilis*.

Following Gardner et al (2011), and using the primer pairs suggested as the “best” in QDD, 40 loci were tested for polymorphism on eight samples of *M. mutabilis*: four collected from each of The Burren (53°07'29"N, 9°05'19"W) and the Isle of Mull (56°32'44"N, 6°02'13"W). PCR amplifications for all loci except MiMut24 were performed in 15µl reaction volumes containing 1x ImmoBuffer, 0.2 mM each dNTP, 200nM each forward and reverse primer, 0.3U IMMOLASE DNA Polymerase (BIOLINE) and 20-40ng template gDNA. The reaction conditions for MiMut24 were the same except 2mM MgCl<sub>2</sub>, 1x PCR Gold Buffer and 0.3U AmpliTaq Gold DNA Polymerase (Applied Biosystems). The PCR cycling conditions were: initial denaturation step at 94°C (MiMut24) or 95°C (all other loci) for 9min, followed by 36 cycles at 94°C for 30 s, annealing temperature at 50°C for 30 s, and extension at 45°C for 45 s, followed by a single final extension step at 72°C for 30min. PCR products were inspected for approximate allele size after separation by electrophoresis in 3% Nu-sieve GTG agarose gel (Cambrex).

We selected eight putative loci which amplified products of the expected size from the 40 trialled, attached florescent labels (FAM, GeneWorks; and NED, PET or VIC, Applied Biosystems)

5' on each forward primer (Table 1) and amplified the loci using the PCR conditions described above. The PCR products were visually inspected on 6% polyacrylamide gels (Gel-scan 2000 Corbett Sydney, Australia), which revealed three loci did not produce clearly interpretable bands and these loci were subsequently removed. The five remaining loci and the three previously isolated CEH49, Mmut9 and CEH52 (Table 1) were amplified in 189 samples of *M. mutabilis* using the two single locus PCR reactions mentioned above. The PCR products were combined in two pools (Table 1) for genotyping on an ABI 3730 DNA Analyser (Applied Biosystems) with the size standard GS500 (-250) LIZ and scored with GENEMAPPER ver 3.7 (Applied Biosystems).

To assess the suitability of markers for analysis we genotyped individuals from the study site in The Burren, Ireland. Samples with more than three missing microsatellite loci were removed from the dataset. We used MICROCHECKER 2.2.3 (Van Oosterhout *et al.* 2004) to examine each locus for potential scoring errors due to genotyping, stuttering and large allele drop-out and found no evidence for errors due to these factors. For each locus (Table 1) we calculated: the polymorphic information content (PIC), the number of alleles, observed ( $H_O$ ), expected heterozygosity ( $H_E$ ), and null allele frequency using Cervus 3.0.3 (Kalinowski *et al.* 2007); and deviation from Hardy-Weinberg Equilibrium (HWE) and linkage disequilibrium between all pairs of loci using GENEPOP 4.1 (Raymond & Rousset 1995) (1,000 dememorizations, 100 batches, 10,000 iterations per batch). *P*-values from HWE tests were manually adjusted for multiple tests of significance using the sequential Bonferroni method (Hochberg 1988). No loci showed significant deviation from HWE expectation after Bonferroni adjustment. No loci pairs showed evidence for linkage disequilibrium after sequential Bonferroni adjustment. The number of alleles and observed heterozygosity ranged from 2-16 and 0.257-0.971 respectively. No loci exhibited large null allele frequencies (largest 0.135 for MiMut27). With a combined non-exclusion probability (sib identity)

of 0.00893, these loci are likely to be informative and valuable for future population-level genetic studies of *M. mutabilis*.

## Acknowledgements

We thank Alison Fitch, Kathy Saint and Terry Bertozzi at the South Australian Museum for technical support.

## References

- Elmes, G.W., Barr, B., Thomas, J.A., and Clarke, R.T. (1999) Extreme host specificity by *Microdon mutabilis* (Diptera : Syrphidae), a social parasite of ants. *Proceedings of the Royal Society of London Series B-Biological Sciences* **266**(1418), 447-453.
- Gardner, M.G., Cooper, S.J.B., Bull, C.M., and Grant, W.N. (1999) Isolation of microsatellite loci from a social lizard, *Egernia stokesii*, using a modified enrichment procedure. *Journal of Heredity* **90**, 301-304.
- Gardner, M.G., Fitch, A.J., Bertozzi, T., and Lowe, A.J. (2011) Rise of the machines – recommendations for ecologists when using next generation sequencing for microsatellite development. *Molecular Ecology Resources* **11**, 1093-1101.
- Gardner, M.G., Schonrogge, K., Elmes, G.W., and Thomas, J.A. (2007) Increased genetic diversity as a defence against parasites is undermined by social parasites: *Microdon mutabilis* hoverflies infesting *Formica lemni* ant colonies. *Proceedings of the Royal Society B-Biological Sciences* **274**, 103-110.
- Hale, M.L., Bevan, R., and Wolff, K. (2001) New polymorphic microsatellite markers for the red squirrel (*Sciurus vulgaris*) and their applicability to the grey squirrel (*S. carolinensis*). *Molecular Ecology Notes* **1**, 47-49.
- Hochberg, Y. (1988) A sharper Bonferroni procedure for multiple tests of significance. *Biometrika* **75**, 800-802.
- Kalinowski, S.T., Taper, M.L., and Marshall, T.C. (2007) Revising how the computer program CERVUS accommodates genotyping error increases success in paternity assignment. *Molecular Ecology* **16**, 1099-1106.
- Larkin M.A., Blackshields G., Brown N.P., Chenna R., McGettigan P.A., McWilliam H., Valentin F., Wallace I.M., Wilm A., Lopez R., Thompson J.D., Gibson T.J., and Higgins D.G. (2007) ClustalW and ClustalX version 2. *Bioinformatics* **23**, 2947–2948.
- Meglecz, E., Costedoat, C., Dubut, V., Gilles, A., Malausa, T., Pech, N., and Martin, J.-F. (2010) QDD: a user-friendly program to select microsatellite markers and design primers from large sequencing projects. *Bioinformatics* **26**, 403-404.
- Raymond, M., and Rousset, F. (1995) GENEPOP (version 1.2): Population genetics software for exact tests and ecumenicism. *Journal of Heredity* **86**, 248-249.
- Rozen, S. and Skaletsky, H. (2000) Primer3 on the WWW for general users and for biologist programmers. *Methods Molecular Biology* **132**, 365–386.

- Schönrogge, K., Gardner, M.G., Elmes, G.W., Napper, E.K.V., Simcox, D.J., Wardlaw, J.C., Breen, J., Barr, B., Knapp, J.J., Pickett, J.A., and Thomas, J.A. (2006) Host propagation permits extreme local adaptation in a social parasite of ants. *Ecology Letters* **9**, 1032-1040.
- Van Oosterhout, C., Hutchinson, W.F., Wills, D.P.M., Shipley, P. (2004) MICRO-CHECKER: software for identifying and correcting genotyping errors in microsatellite data. *Molecular Ecology Notes* **4**, 535-538.