BRITISH GEOLOGICAL SURVEY Port Stanley Observatory Monthly Magnetic **Bulletin** December 2012 PS







1. Introduction

Port Stanley observatory was installed by the British Geological Survey (BGS) with financial support from a consortium of oil companies and became operational in February 1994.

This bulletin is published to provide rapid access to the provisional geomagnetic observatory results. The information is freely available for personal, academic, educational and non-commercial research or use. Magnetic observatory data are presented as a series of plots of one-minute, hourly and daily values, followed by tabulations of monthly values. The operation of the observatory and presentation of data are described in the rest of this section.

Enquiries about the data should be addressed to:

Geomagnetism Team Earth Hazards and Systems British Geological Survey Murchison House, West Mains Road Edinburgh EH9 3LA Scotland, UK

 Tel:
 +44 (0) 131 667 1000

 Fax:
 +44 (0) 131 650 0265

 E-mail:
 enquiries@bgs.ac.uk

 Internet:
 www.geomag.bgs.ac.uk

2. Position

Port Stanley Observatory, one of the geomagnetic observatories maintained and operated by the British Geological Survey (BGS), is situated on a site at Sapper Hill near Port Stanley in the Falkland Islands. The observatory co-ordinates are:

Geographic:	51°42'15"S	302 <i>°06`25</i> "E
Geomagnetic:	42 <i>°07'26</i> "S	012°05'46"E
Height above m	ean sea level:	135 m

The geomagnetic co-ordinates are approximations, calculated using the 11th generation International Geomagnetic Reference Field (IGRF) at epoch 2012.5. On-line access to models (including IGRF), charts and navigational data are available at http://www.geomag.bgs.ac.uk/data_service/models_compass/home

3. The Observatory Operation

3.1 GDAS

The observatory operates under the control of the Geomagnetic Data Acquisition System (GDAS), which was developed by BGS staff, installed and became operational in August 2002. The data acquisition software, running on QNX operated computers, controls the data logging and the communications.

There are two sets of sensors used for making magnetic measurements. A tri-axial linear-core fluxgate magnetometer, manufactured by the Danish Meteorological Institute, is used to measure the variations in the horizontal (H) and vertical (Z) components of the field. The third sensor is oriented perpendicular to these, and measures variations, which are proportional to the changes in declination (D). Measurements are made at a rate of 1 Hz.

In addition to the fluxgate sensors there is a proton precession magnetometer (PPM) making measurements of the absolute total field intensity (F) at a rate of 0.1Hz.

The raw unfiltered data are retrieved automatically via Internet connections to the BGS office in Edinburgh in near real-time. The fluxgate data are filtered to produce one-minute values using a 61point cosine filter and the total field intensity samples are filtered using a 7-point cosine filter. The one-minute values provide input for various data products, available on-line at

www.geomag.bgs.ac.uk/data service/home

3.2 Absolute Observations

The GDAS fluxgate magnetometers accurately measure variations in the components of the geomagnetic field, but not the absolute magnitudes. Two sets of absolute measurements of the field are made manually twice per month. A fluxgate sensor mounted on a theodolite is used to determine D and inclination (1); the GDAS PPM measurements, with a site difference correction applied, are used for F. The absolute observations are used in conjunction with the **GDAS** variometer measurements to produce a continuous record of the absolute values of the geomagnetic field elements as if they had been measured at the observatory reference pillar.

4. Observatory Results

The data presented in the bulletin are in the form of plots and tabulations described in the following sections.

4.1 Absolute Observations

The absolute observation measurements made during the month are tabulated. Also included are the corresponding baseline values, which are the differences between the absolute measurements and the variometer measurements of D, H and Z (in the sense absolute–variometer). These are also plotted (markers) along with the derived preliminary daily baseline values (line) throughout the year. Daily mean differences between the measured absolute F and the F computed from the baseline corrected H and Z values are plotted in the fourth panel (in the sense measured–derived). The bottom panel shows the daily mean temperature in the fluxgate chamber.

4.2 Summary magnetograms

Small-scale magnetograms are plotted which allow the month's data to be viewed at a glance. They are plotted 16 days to a page and show the one-minute variations in D, H and Z. The scales are shown on the right-hand side of the page. On disturbed days the scales are multiplied by a factor, which is indicated above the panel for that day. The variations are centred on the monthly mean value, shown on the left side of the page.

4.3 Magnetograms

The daily magnetograms are plotted using oneminute values of D, H and Z from the fluxgate sensors, with any gaps filled using back-up data. The magnetograms are plotted to a variable scale; scale bars are shown to the right of each plot. The absolute level (the monthly mean value) is indicated on the left side of the plots.

4.4 Hourly Mean Value Plots

Hourly mean values of D, H and Z for the past 12 months are plotted in 27-day segments corresponding to the Bartels solar rotation number. Magnetic disturbances associated with active regions and/or coronal holes on the Sun may recur after 27 days: the same is true for geomagnetically quiet intervals. Plotting the data in this way highlights this recurrence. Diurnal variations are also clear in these plots and the amplitude changes throughout the year highlight the seasonal changes. Longer term secular variation is also illustrated.

4.5 Daily and Monthly Mean Values

Daily mean values of D, H, Z and F are plotted throughout the year. In addition, a table of monthly mean values of all the geomagnetic elements is provided. These values depend on accurate specification of the fluxgate sensor baselines. It is anticipated that these provisional values will not be altered by more than a few nT or tenths of arcminutes before being made definitive at the end of the year.

5. Conditions of Use

The data presented in this bulletin are provided for personal, academic, educational, non-commercial research or other non-commercial use and are not for sale or distribution to third parties without written permission from BGS.

Reproduction of any part of this bulletin should be accompanied by the statement: 'Reproduced with the permission of the British Geological Survey ©NERC. All rights Reserved'. Publications making use of the data should include an acknowledgment statement of the form: 'The results presented in this paper rely on the data collected at Port Stanley magnetic observatory, operated by the British Geological Survey.'

Commercial users can contact the geomagnetism team for information on the range of applications and services offered. Full contact details are available at www.geomag.bgs.ac.uk/contactus/staff


© NERC 2012. All rights reserved

Edinburgh


PORT STANLEY OBSERVATORY


ABSOLUTE OBSERVATIONS


	Declination			Inclination		Total Field		Horizontal Intensity		Vertical Intensity			
Date	Day Number	Time (UT)	Absolute (°)	Baseline (°)	Time (UT)	Absolute (°)	Site difference (nT)	Absolute corrected (nT)	Absolute (nT)	Baseline (nT)	Absolute (nT)	Baseline (nT)	Observer
10-Dec-12	345	15:09	3.2877	3.6717	15:15	-49.5352	-31.4	28548.7	18527.5	18847.2	-21720.0	-21941.1	NB
10-Dec-12	345	15:22	3.2927	3.6717	15:29	-49.5371	-31.4	28547.2	18525.9	18846.4	-21719.5	-21941.7	NB
20-Dec-12	355	16:41	3.3082	3.6783	16:37	-49.5450	-31.4	28558.3	18530.1	18848.6	-21730.5	-21940.4	NB
20-Dec-12	355	16:54	3.2969	3.6717	16:59	-49.5622	-31.4	28556.3	18522.3	18846.8	-21734.5	-21941.9	NB





Falkland Island 2012





















































Falkland Islands Observatory: Declination (degrees)




Falkland Islands Observatory: Horizontal Intensity InT/



Falkland Islands Observatory: Vertical Intensity (nT)





Monthly Mean Values for Port Stanley Observatory 2012

Month	D	Н	Ι	X	Y	Ζ	F
January February March April May June July August September October	3° 19.2′ 3° 18.8′ 3° 17.9′ 3° 17.2′ 3° 16.8′ 3° 16.5′ 3° 16.7′ 3° 16.3′ 3° 16.0′ 3° 15.2′	18587 nT 18571 nT 18544 nT 18544 nT 18542 nT 18534 nT 18522 nT 18528 nT 18521 nT 18521 nT 18510 nT	-49° 29.8′ -49° 31.3′ -49° 34.0′ -49° 34.4′ -49° 33.7′ -49° 34.2′ -49° 35.3′ -49° 34.3′ -49° 34.5′ -49° 35.3′	A 18556 nT 18540 nT 18514 nT 18509 nT 18512 nT 18504 nT 18492 nT 18497 nT 18491 nT 18480 nT	1077 nT 1074 nT 1067 nT 1063 nT 1061 nT 1059 nT 1059 nT 1057 nT 1056 nT 1050 nT	2 -21760 nT -21760 nT -21763 nT -21764 nT -21755 nT -21755 nT -21755 nT -21748 nT -21743 nT -21740 nT	28617 nT 28607 nT 28593 nT 28590 nT 28588 nT 28579 nT 28572 nT 28572 nT
November December	3° 14.8′ 3° 14.5′	18515 nT 18524 nT	-49° 34.5′ -49° 33.1′	18485 nT 18494 nT	1049 nT 1047 nT	-21740 nT -21735 nT -21728 nT	28552 nT

Note

i. The values shown here are provisional.