

Article (refereed) - postprint

Salmela, Matti J.; Cavers, Stephen; Cottrell, Joan E.; Iason, Glenn R.; Ennos, Richard A. 2013. **Spring phenology shows genetic variation among and within populations in seedlings of Scots pine (*Pinus sylvestris* L.) in the Scottish Highlands.**

© 2013 Botanical Society of Scotland and Taylor & Francis

This version available <http://nora.nerc.ac.uk/501122/>

NERC has developed NORA to enable users to access research outputs wholly or partially funded by NERC. Copyright and other rights for material on this site are retained by the rights owners. Users should read the terms and conditions of use of this material at <http://nora.nerc.ac.uk/policies.html#access>

This is an Author's Accepted Manuscript of an article published in *Plant Ecology & Diversity* (2013), 6(3-4). 523-536, copyright Botanical Society of Scotland and Taylor & Francis, available online at: <http://www.tandfonline.com/10.1080/17550874.2013.795627>

Contact CEH NORA team at
noraceh@ceh.ac.uk

1 **Spring phenology shows genetic variation among and within populations in seedlings of**
2 **Scots pine (*Pinus sylvestris* L.) in the Scottish Highlands**

3

4 Matti J. Salmela^{a,b,1}, Stephen Cavers^{b*}, Joan E. Cottrell^c, Glenn R. Iason^d, Richard A. Ennos^a

5

6 ^aInstitute of Evolutionary Biology, School of Biological Sciences, Ashworth Laboratories,
7 University of Edinburgh, Edinburgh EH9 3JT, UK

8 ^bNERC Centre for Ecology and Hydrology Edinburgh, Bush Estate, Penicuik, Midlothian
9 EH26 0QB, UK

10 ^cForest Research, Northern Research Station, Roslin, Midlothian EH25 9SY, UK

11 ^dJames Hutton Institute, Craigiebuckler, Aberdeen AB15 8QH, UK

12 ¹Present address: Department of Botany, 3165, 1000 E. University Ave., University of
13 Wyoming, Laramie, WY 82071, USA

14

15

16

17

18

19

20

21 Correspondence author. Email: scav@ceh.ac.uk, tel: +44 (0)131 445 4343, fax: +44 (0)131

22 445 3943

23

24

25

26

27

28 **Abstract**

29

30 Background: Genetic differentiation in phenotypic traits is often observed among forest tree
31 populations, but less is known about patterns of adaptive variation within populations. Such
32 variation is expected to enhance the survival likelihood of extant populations under climate
33 change.

34 Aims: Scots pine (*Pinus sylvestris*) occurs over a spatially and temporally heterogeneous
35 landscape in Scotland. Our goal was to examine whether populations had differentiated
36 genetically in timing of bud flush in response to spatial heterogeneity and whether variation
37 was also maintained within populations.

38 Methods: Two common-garden studies, involving maternal families of seedlings from 21
39 native pinewoods, were established and variation in the trait was measured at the beginning of
40 the second growing season.

41 Results: Populations showed genetic differences in the trait correlated with the length of
42 growing season at their site of origin, but the majority of variation was observed within
43 populations. Populations also differed in their levels of variation in the trait; a pattern that
44 may be influenced by spatial variation in the extent of temporal climate variability.

45 Conclusions: Our findings suggest that populations have adapted to their home environments
46 and that they also have substantial ability to adapt in situ to changes in growing season length.

47

48 Keywords: adaptation, adaptive potential, genetic differentiation, spatial heterogeneity,
49 temporal heterogeneity, variation within populations

50

51

52

53

54

55

56 **Introduction**

57

58 When a species is distributed across a spatially heterogeneous landscape, natural selection is
59 expected to favour different trait optima in divergent environments. For example, growth in
60 plants may continue furthest into the autumn in populations that experience the longest
61 growing seasons (e.g. Mikola 1982). This may lead to genetic differentiation in selected
62 phenotypic traits among populations and each population surviving and growing best at its
63 home site, i.e. local adaptation (Kawecki and Ebert 2004). Adaptations to local environments
64 have been described in many plant species (Linhart and Grant 1996), and in trees, numerous
65 common-garden studies have demonstrated that, for example, growth phenology and timing
66 of cold hardiness are optimised to local climates (Howe et al. 2003; Savolainen et al. 2007).
67 Local adaptation has also been demonstrated in transfer trials in which populations have been
68 grown at home and foreign sites (e.g. Persson and Ståhl 1990); a possible reason for poorer
69 survival and growth at foreign sites is a mismatch between annual climatic variation and
70 growth phenology (Eriksson et al. 1980).

71

72 Ongoing rapid climate change is affecting ecosystems globally, which is evident as range
73 shifts and changes in timing of growth and reproductive events in several species (Parmesan
74 2006). Due to the commercial importance of many species and conservation issues, predicting
75 evolutionary responses to a changing climate has become an active area of research (Hendry
76 et al. 2011; Hoffman and Sgrò 2011). For example, ecological modelling often considers
77 associations between species distributions and environmental factors: following
78 environmental change, ranges may expand into new areas that become suitable for a species,
79 while extinction may take place at sites that become unfavourable (Elith and Leathwick
80 2009). In trees, common-garden experiments replicated in multiple environments have been
81 used to estimate how environmental changes will affect contemporary populations and to
82 demonstrate that in the future, some populations might be exposed to non-optimal conditions
83 which might result in poorer growth (Rehfeldt et al. 2002; Reich and Oleksyn 2008).
84 However, such approaches can only be used to examine how existing populations will
85 respond to environmental changes as they neglect an important feature of natural populations:
86 their capacity to adapt to environmental changes in situ (Hoffman and Sgrò 2011). This

87 capability means that, as well as range shifts, responses to environmental change should also
88 involve genetic changes between generations exposed to different environments, allowing
89 populations to adapt to new conditions. Indeed, taking adaptation and phenotypic plasticity
90 into account in models can greatly enhance the survival likelihood of current populations
91 (Aitken et al. 2008; Benito Garzón et al. 2011). Adaptation to novel environments has
92 certainly taken place in many species when they expanded their ranges following the last ice
93 age (Davis and Shaw 2001), but it is possible that future changes will occur too rapidly for
94 populations to track them (Savolainen et al. 2004; Aitken et al. 2008) or that intensively
95 managed contemporary landscapes will be impermeable to migration.

96

97 In situ adaptive responses to a changing environment are possible when phenotypes vary
98 within populations and when such variation is due to genetic factors (Hoffman and Sgrò
99 2011). However, research on adaptation has largely focused on examining how populations
100 differ from each other in terms of trait means and how environmental differences among the
101 sites occupied by different populations contribute to such divergence (reviewed e.g. in
102 Savolainen et al. 2007). Thus, we currently have only a limited understanding of patterns of
103 adaptive trait variation within populations (Kramer and Havens 2009). When populations
104 adapt to their home environments, variation in traits affected by selection is expected to be
105 lost as individuals that differ too much from the local optimum have poorer chances of
106 survival (e.g. Falconer and Mackay 1996). Still, it is commonly found across different kinds
107 of organisms that significant genetic variation can be preserved even in traits under the
108 strongest type of selection (Houle 1992; Merilä and Sheldon 1999; Barton and Keightley
109 2002). Also in trees, within-population variation in traits under selection has been widely
110 found in common-garden studies (Howe et al. 2003). A similar pattern can also be seen in
111 trees observed in their natural habitats: in a stand of *Betula pendula* Roth growing in south-
112 eastern Finland, trees flushed at different times, among-tree differences being the smallest
113 during warm springs (Rousi and Heinonen 2007). The reasons for the persistence of such
114 variation in nature despite natural selection remain poorly understood, but a number of factors
115 might contribute to the maintenance of adaptive genetic diversity in forest trees. Due to their
116 longevity, trees are likely to experience a wide range of environmental conditions during their
117 lifespan (Petit and Hampe 2006), and mechanisms enabling individuals to modify their
118 phenotype according to the environment are expected to evolve under such conditions (Bull

119 1987). Indeed, phenotypic plasticity in initiation of growth allows trees to survive in
120 environments where temperature conditions in spring vary among years (Rousi and Heinonen
121 2007; Chmura et al. 2011). It is also possible that a population is found over a spatially
122 variable area so that selection pressure also varies across short distances (i.e., there is no
123 single trait optimum in a population; e.g. Campbell 1979), or that the environment varies
124 between years so that different age groups within populations may have experienced differing
125 selection pressure. This may maintain diversity if selection acts only on specific age groups
126 (e.g. young seedlings) while having less effect on others (Ellner and Hairston 1994). Variation
127 may also be introduced by gene flow via pollen from environmentally different sites (Yeaman
128 and Jarvis 2006). What the relative contributions of these factors in nature are remains largely
129 an unexplored topic.

130

131 Adaptive potential can be compromised especially in small and fragmented populations in
132 which random factors such as sudden population size changes may have shaped patterns of
133 genetic variation more than natural selection (e.g. Willi et al. 2006). Scots pine (*Pinus*
134 *sylvestris* L.) is the only pine native to northern Europe and has an extensive distribution
135 across Eurasia (Critchfield and Little 1966). The Scottish populations of the species are
136 geographically separated on the north-western edge of this range and have been subjected to
137 heavy human interference in the past. Currently, 84 discrete native pinewood sites of variable
138 size are recognised by the Forestry Commission of Great Britain which cover only about 1%
139 of their original postglacial maximum areal cover (Mason et al. 2004). Scots pine is a
140 foundation species upon which the persistence of many of the species in Scottish forests
141 depends. The native pinewoods are found over a geographically small but spatially highly
142 heterogeneous landscape, with steep gradients in temperatures and precipitation between the
143 oceanic west coast and the more continental east (Salmela et al. 2010). Despite a significant
144 decrease in abundance which has led to fragmentation, most of the populations are as diverse
145 at selectively neutral molecular markers as more continuous continental populations and show
146 very little differentiation for these neutral markers (Kinloch et al. 1986; Wachowiak et al.
147 2010). This suggests that at least historically, these populations have been connected by gene
148 flow. However, little is currently known about the patterns of adaptive trait variation in this
149 part of the species' distribution. In a recent experiment under natural climate conditions in
150 south-eastern Scotland, plants from eight populations were found to differ in their response to

151 winter and spring temperatures, which suggested environment-driven genetic differentiation
152 among some of the populations despite the small geographic scale (Salmela et al. 2011).
153 Similar differentiation was found for timing of growth initiation in spring, which was earlier
154 in populations from cooler, high-altitude locations.

155

156 In common with other parts of the world, increases in summer and winter temperatures and
157 changes in rainfall patterns are expected in Scotland in the coming decades (Ray 2008),
158 possibly leading to changes in selection pressures for traits related to timing of growth in
159 Scots pine. For current populations, these changes in climate have been predicted to be
160 detrimental (Ray 2008). However, the possibility of adaptation within populations has not
161 been considered in these predictions which might result in their conclusions being too
162 conservative and potentially in management actions detrimental to the genetic integrity of
163 current populations. Considering how allowing for adaptation influences model predictions on
164 the effects of climate change in tree populations (Aitken et al. 2008), it is important for the
165 conservation of the remaining native pinewood resources that the patterns of adaptive trait
166 variation among and within populations are investigated (Salmela et al. 2010).

167

168 Due to the highly variable climate conditions that they are found in, adaptive genetic
169 differentiation is expected to have taken place among pine populations from different parts of
170 Scotland. In addition to spatial environmental variation among and within populations, the
171 Scottish climate is also characterised by temporal (among-year) fluctuations, for instance in
172 the length of the growing season (Perry and Hollis 2005) and winter severity (Harrison 1997).
173 Such fluctuations probably account for phenomena such as the observed temporal variation in
174 timing of bud flush under natural climate conditions in two birch species, *Betula pubescens*
175 Ehrh. and *B. pendula* (Billington and Pelham 1991). The effects of climate fluctuations have
176 also been recognised in animals: sheep mortality on the island of St. Kilda has been found to
177 be higher in wet and warm winters which often coincide with positive phases of North
178 Atlantic Oscillation (Milner et al. 1999), a climatic phenomenon linked to the strength of
179 westerly winds across the northern Atlantic (Stenseth et al. 2002). Although temporally
180 variable selection has been suggested as one factor that may contribute to the maintenance of
181 adaptive diversity in long-lived trees (Howe et al. 2003; Westfall and Millar 2004; Yeaman

182 and Jarvis 2006), its potential role is yet to be studied in more detail. If populations are
183 exposed to different levels of spatial and temporal environmental heterogeneity, they might
184 also differ in the level of genetic diversity in adaptive traits and consequently their adaptive
185 capacity. In this study, we used Scots pine as a model system to examine how a phenological
186 trait varied among and within populations sampled across a spatially and temporally
187 heterogeneous landscape. More specifically, we collected families from 21 environmentally
188 diverse sites in Scotland and grew their progeny in two separate glasshouse experiments to
189 address the following questions:

190

- 191 1) When grown under common-garden conditions, are 21 native Scots pine populations
192 differentiated for timing of growth initiation in spring at the beginning of their second
193 growing season?
- 194 2) Are observed population differences associated with environmental variation among
195 their home sites?
- 196 3) Is there significant variation within populations? If so, what is the pattern of this
197 variation?
- 198 4) Could the relative amounts of within-population variation be accounted for by within-
199 site spatial and/or temporal variation in environment?

200

201

202 **Materials and methods**

203

204

205 *Measuring genetic diversity in timing of bud flush*

206

207 Phenotypes of adaptive traits are determined by both genetic factors and the environment
208 (Falconer and Mackay 1996). Therefore, to reveal differences due to the genetic component,
209 samples from different populations must be raised in a common-garden environment. To
210 estimate the levels of variation within populations, a family-structured design is needed so

211 that total variation in phenotype can be partitioned into among- and within-population
212 components. In tree populations, this can be accomplished by sampling multiple open-
213 pollinated seed from a number of mother trees in each population (e.g. White et al. 2007).
214 Due to high outcrossing rates (mother trees are generally pollinated by a large number of
215 pollen donors), such progeny are often assumed to consist mostly of half-siblings (i.e., family
216 members share only the maternal parent).

217

218

219 *Study populations*

220

221 A total of 21 native populations were sampled for this study, representing all parts of the
222 species' range in Scotland (Figure 1, Table 1). Cones were collected from 10 maternal trees in
223 each population in March 2007. Open-pollinated seed was extracted from cones and stored by
224 family.

225

226

227 *Provenance/progeny trials*

228

229 Sampled seed was used to establish two glasshouse-based common-garden trials located in
230 Edinburgh and Aberdeen (Figure 1) in late spring 2007. The two trials were set up by
231 independent investigators and they consequently had rather different germination conditions
232 and layout designs. In the Edinburgh trial located at the Centre for Ecology and Hydrology
233 (55.86° N, 3.21° W), seed were sampled from four mother trees per population (i.e. 84
234 families in total) and sown on trays (75:25 compost type John Innes 1: sand) in June 2007
235 under common-garden glasshouse conditions. After germination, seedlings were transferred to
236 pots of size 0.62 l (diameter 11 cm, depth 9.6 cm) and kept under natural light conditions
237 (glasshouse was shaded to avoid excess light) with watering applied two or three times per
238 week during the growing season. No heating was applied during winter. Each family
239 consisted of 40 progeny (~3,360 seedlings in total). The trial was divided into 40 blocks, each

240 having one member from each of 84 families, and the order of the families within blocks was
241 randomised.

242

243 The trial in Aberdeen was located at the James Hutton Institute (57.13° N, 2.16° W). Seed
244 were sampled from 10 mother trees per population (i.e. 210 families in total). Cones were
245 placed in a warm room (30 °C) for two weeks so that they opened and seed could be extracted
246 for germination. Seed from the individual trees were kept separate and were soaked in water
247 for 3 hours, then laid between sheets of damp paper towel placed in a cool room (3 °C) for
248 several weeks to break dormancy. Seed were taken out of the cool room and left (wrapped in
249 damp paper) in the laboratory at room temperature until they germinated. Germination took
250 approximately 7 days and seed from all the sampled families germinated at this time. On
251 germination they were transplanted into potting medium in the glasshouse into 8 × 8 × 9 cm
252 (0.4 l) pots. Each family consisted of eight progeny (~1,680 seedlings in total). The trial was
253 divided into 40 blocks with 42 plants per block, each block containing two plants from a
254 different mother from each population. The 84 mother trees sampled in the Edinburgh trial
255 were a subset of those included in the Aberdeen trial. Watering was applied automatically and
256 no artificial light was used.

257

258 In both trials, growth initiation at the beginning of the second growing season was considered
259 to have taken place when new green needle tips started to emerge from the apical bud (bud or
260 needle flush), and this was measured as the number of days since the first scoring date. Bud
261 flush was scored twice weekly in Edinburgh between March 23 and May 9 2008 and once
262 weekly in Aberdeen between March 31 and May 27 2008.

263

264

265 *Testing for genetic differentiation among populations and families*

266

267 Data from the Edinburgh trial were analysed using nested analysis of variance (ANOVA),
268 with populations considered as fixed and families within them and blocks as random factors.
269 Unbalanced nested ANOVA was applied to the data from the Aberdeen trial. To examine the

270 relative contributions of different factors to total variation in the trait, variance components
271 due to populations, families, and blocks were estimated using the restricted maximum
272 likelihood (REML) approach. Correlation analysis was used to test whether similar trends
273 were observed in the two trials. All statistical analyses in this study were carried out using
274 GenStat Ver. 13.1.0.4470.

275

276

277 *Measuring the level of variation within populations*

278

279 To examine the variability of the trait within each population in more detail, standard
280 deviations (SD_{POP}) were examined separately in the two trials. SD 's were calculated using raw
281 values because of their strong correlation ($r=0.94-0.99$) with values adjusted for the effects of
282 individual blocks. Because timing of bud flush does not have fixed means (i.e., means vary
283 depending on the date from which the timing is calculated), we used SD 's instead of
284 normalised measures of dispersion (see e.g. Garcia-Gonzalez et al. 2012). Correlation analysis
285 was used to test whether similar patterns of variation among SD_{POP} 's were observed in the
286 two trials.

287

288 In a study design consisting of families grouped within populations, within-population
289 variation can arise from two components: among families and within families (residual
290 variation). Variation among families is considered to reflect the level of heritable (additive)
291 genetic variation (Falconer and Mackay 1996), and populations with higher levels of heritable
292 variation are expected to have better adaptive potential (Houle 1992). Because the amount of
293 additive genetic variation is directly proportional to the amount of variation among families
294 (Falconer and Mackay 1996), we only used estimates of among-family variation in the
295 analyses presented. In order to examine whether the level of among and within-family
296 variation differed among populations, the REML approach was used to calculate variance
297 components due to families (V_{AF}) and individuals within families (V_{WF}) separately within each
298 trial and population (i.e., variation within each population was divided into components due to
299 families, blocks, and residual variation). Variation within families may include a component

300 due to the genetic diversity of pollen donors sampled by mother trees. SD 's were also
301 calculated for individual families (SD_{WF}).

302

303

304 *Climate data*

305

306

307 *Long-term means at the sampled sites*

308

309 To investigate how the sampled sites varied in terms of temperature conditions, UK
310 Meteorological Office 40-year mean (1961-2000) climate data (Perry and Hollis 2005) were
311 used to create climatic profiles of the populations' origins. Data were extracted for the length
312 of the growing season (GSL), number of growing degree days (GDD), and mean February
313 and July temperatures (FMT and JMT; these represent on average, the coldest and warmest
314 months, respectively). GSL is defined as the period bounded by daily mean temperature above
315 5 °C for more than five consecutive days and daily mean temperature below 5 °C for more
316 than five consecutive days (after 1 July), while GDD expresses the sum of daily heat sum
317 accumulation above 5.5 °C. Exact details on how the climate data were generated are given in
318 Perry and Hollis (2005). Climate data are available in 5 km × 5 km grids and are based on
319 interpolation of observations from the nearest weather stations. It is possible that due to
320 within-grid variation in the landscape, actual climate conditions experienced at our study sites
321 differ from the estimates, therefore, the climatic variables should only be considered as
322 proxies. The range of altitude sampled at each site was used as a proxy for fine-scale (within-
323 population) environmental variation.

324

325

326

327

328

329 *Temporal variation in the Scottish climate*

330

331 To investigate patterns of temporal variability in temperature in Scotland and at the sampled
332 sites, annual estimates of GSL, GDD, February and July temperatures (FT and JT) for the 5
333 km × 5 km grids in the period 1961-2000 were used. Using these 40-year data, mean absolute
334 deviations (*MAD*) of FT and JT, and coefficients of variation (*CV*) of GSL and GDD (CV_{GSL} ,
335 CV_{GDD}) were calculated separately for each site. A combined estimate of monthly temperature
336 variability was calculated as the average of the *MAD*'s of FT and JT. Annual means were also
337 calculated over all 21 sites. To test whether winter temperature in Scotland was associated
338 with the North Atlantic Oscillation (NAO), linear regression was also used to test for an
339 association between annual FTs and February NAO indices provided by the Climate Analysis
340 Section, NCAR, Boulder, USA, at NAO Index
341 <http://www.cgd.ucar.edu/cas/jhurrell/indices.html>. Linear regression was used to explore
342 whether the level of temporal variation in climate varied along latitudinal, longitudinal, or
343 altitudinal gradients.

344

345

346 *Associations between population means of timing of bud flush and climate at home site*

347

348 To investigate associations between variation in timing of bud flush, the locations of the
349 populations, and their climate, population means in the two trials were regressed against their
350 longitude, latitude, altitude, and long-term mean temperature estimates (GSL, GDD, FMT,
351 JMT) at origin.

352

353

354 *Associations between the level of variation within populations and latitude, longitude, and*
355 *altitude*

356

357 Linear regression was used to explore whether the level of variation within populations

358 (SD_{POP}) was linked to the range of altitude sampled at each site, or whether it varied along any
359 spatial gradients (longitude, latitude, altitude). We also tested for associations between the
360 locations of the sampled families and among and within-family trait variation (V_{AF} , V_{WF} ,
361 SD_{WF}) in each trial.

362

363

364 **Results**

365

366

367 *Genetic differentiation in timing of bud flush among populations and families*

368

369 In the Edinburgh trial, population means ranged between 11 days after March 23 for AC, BB,
370 GA, GD, and GT, and 18 days after March 23 for AB, BE, SD, and RD. ANOVA provided
371 some evidence of differences among populations ($P=0.058$), and differences among families
372 within populations, and among blocks were significant (Table 2a). The variance component
373 due to differences among families within populations (15.39; 22% of total variation) was
374 approximately five times larger than that among populations (2.98; 4% of total variation).

375

376 In the Aberdeen trial, the range of population means was from 16 days after March 31 for AC
377 and GL to 22 days after March 31 for AB. Significant differences were observed among
378 populations, families within populations, and blocks (Table 2b). The variance component due
379 to families (5.88; 11% of total variation) was approximately four times larger than that of
380 populations (1.45; 3% of total variation). Population means between the two trials were
381 significantly and moderately correlated ($r=0.48$, $P<0.05$). In both trials, the great majority of
382 the variation was residual, i.e. within families (Table 2a, b; 72% in the Edinburgh trial, 77% in
383 the Aberdeen trial).

384

385

386

387

388 *Associations between population means and climate at home site*

389

390 Temperature conditions varied greatly within Scotland: for instance, mean GSL ranged from
391 116 days in BB to 283 days in BE, and mean GDD from 446 to 1,329 dd at the same sites
392 (Table 1). Spatial variation was also found in timing of bud flush, and results from the two
393 trials showed similar trends although there were differences in the absolute values. In the
394 Edinburgh trial, when examining associations between population means and geographical
395 surrogates of environmental variation (latitude, longitude, altitude), means were best
396 associated with altitude at their site of origin. Low-altitude populations generally flushed later
397 than those from higher locations, and altitude explained 20% of the variation among
398 population means ($\beta_0=16.70$, $\beta_1=-0.010$, $P<0.05$). Altitude of the populations was negatively
399 correlated with mean GSL ($r=-0.80$, $P<0.001$) and GDD ($r=-0.65$, $P<0.01$) at origin, and
400 higher R^2 's were obtained when using these climate variables instead of altitude. Earlier bud
401 flush occurred in populations from areas with shorter GSL ($\beta_0=5.87$, $\beta_1=0.038$, $P<0.01$,
402 $R^2=31\%$), fewer GDD (Figure 2; $\beta_0=8.14$, $\beta_1=0.0066$, $P<0.01$, $R^2=35\%$), lower FMT
403 ($\beta_0=12.62$, $\beta_1=1.13$, $P<0.01$, $R^2=29\%$), and lower JMT ($\beta_0=0.35$, $\beta_1=1.13$, $P<0.010$, $R^2=27\%$).

404

405 A similar trend with altitude was also found in the Aberdeen trial ($\beta_0=20.25$, $\beta_1=-0.0035$), but
406 the association was not statistically significant ($P=0.22$, $R^2=3\%$). However, significant
407 associations were obtained when temperature estimates were used instead of altitude, and sites
408 with shorter GSL ($\beta_0=14.14$, $\beta_1=0.024$, $P<0.001$, $R^2=31\%$), fewer GDD (Figure 2; $\beta_0=15.70$,
409 $\beta_1=0.0040$, $P<0.01$, $R^2=32\%$), lower FMT ($\beta_0=18.38$, $\beta_1=0.69$, $P<0.01$, $R^2=28\%$), and lower
410 JMT ($\beta_0=9.68$, $\beta_1=0.79$, $P<0.01$, $R^2=35\%$) had earlier bud flush.

411

412

413 *Differences in the level of variation within populations*

414

415 Variation among SD_{POP} 's suggested that populations might have differed in the level of

416 variation in timing of bud flush (Table 3). In the Edinburgh trial, SD_{POP} 's varied between 6.25
 417 in GT and 9.53 in AB, while in the Aberdeen trial with a larger number of families within
 418 each population, SD_{POP} 's ranged from 4.29 in CG to 11.10 in GD. SD_{POP} 's across the two
 419 trials were not significantly correlated ($P=0.79$). In the Edinburgh trial, SD_{POP} 's were not
 420 associated with latitude, longitude, or altitude. However, in the Aberdeen trial, the pattern of
 421 variation among SD_{POP} 's was related to the geographic location of populations and individual
 422 mother trees: higher amounts of variation were observed at higher altitude sites (Figure 3;
 423 $\beta_0=5.14$, $\beta_1=0.0080$, $P<0.01$, $R^2=35\%$). The regression was strongly influenced by the three
 424 high-altitude sites with large SD_{POP} 's. When excluding these, the linear regression remained
 425 positive but became statistically non-significant ($\beta_0=5.59$, $\beta_1=0.0050$, $P=0.112$, $R^2=10\%$).

426

427 In the Aberdeen trial, differences among families were generally larger at higher altitudes, and
 428 altitude explained 16% of the variation among V_{AF} 's ($\beta_0=-0.65$, $\beta_1=0.024$, $P<0.05$); however,
 429 eight populations had V_{AF} estimates of 0. A positive correlation was observed between V_{AF} 's
 430 and the range of family means within each population ($r=0.77$, $P<0.001$). R^2 increased to 31%
 431 when the range of family means within each population was used instead of V_{AF} ($\beta_0=6.36$,
 432 $\beta_1=0.019$, $P<0.01$). In the Edinburgh trial, higher SD_{POP} 's were associated with higher V_{AF} 's
 433 ($r=0.69$, $P<0.001$). The association between altitude and V_{AF} 's was only suggestive of higher
 434 levels of among-family variation at higher altitudes ($\beta_0=6.074$, $\beta_1=0.040$, $P=0.11$, $R^2=9\%$)

435

436 Similarly to among-family differences, there was some evidence of larger V_{WF} 's at higher
 437 altitude sites in the Aberdeen trial ($\beta_0=27.39$, $\beta_1=0.070$, $R^2=13\%$, $P=0.062$). This pattern was
 438 also reflected in variation among SD_{WF} 's which ranged between 0 in nine families from seven
 439 populations and 17.92 in a family from AC. Altitude explained 4% of the variation among
 440 SD_{WF} 's ($\beta_0=4.76$, $\beta_1=0.0053$, $P<0.01$), but the association was non-significant ($P=0.53$) when
 441 the three highest-altitude sites were excluded. In the Edinburgh trial, there was no association
 442 between altitude and V_{WF} 's, but they were positively and significantly correlated with SD_{POP} 's
 443 ($r=0.78$, $P<0.001$). SD_{WF} 's varied between 4.16 in a family from GT and 11.04 in a family
 444 from CR, but they were not associated with altitude ($P=0.48$).

445

446

447

448 *Fine-scale environmental variation and the level of variation within populations*

449

450 Mother trees within populations were sampled at different altitudes, and consequently the
451 altitudinal range sampled at each site varied from 23 m at RM to 179 m at GC in the
452 Edinburgh trial, and from 34 at RM m to 199 at GC m in the Aberdeen trial. However,
453 increasing altitudinal range sampled within populations did not account for larger SD_{POP} 's
454 (the Edinburgh trial: $\beta_0=8.42$, $\beta_1=-0.0060$, $R^2=4\%$, $P=0.20$; the Aberdeen trial: $\beta_0=7.010$,
455 $\beta_1=0.001$, $R^2=0\%$, $P=0.95$).

456

457

458 *Temporal climate variation*

459

460 Climate differed markedly from year to year. For example, annual GSL and GDD of the sites
461 occupied by the 21 pinewoods showed extensive temporal fluctuation in the period 1961-2000
462 (Figure 4a). GSL varied between 174 days in 1968 and 271 days in 1989, while the lowest
463 GDD (756 dd) was reached in 1974 and the highest (1,167 dd) in 1995. Temporal variability
464 was also found in monthly winter and summer temperatures. The range of JTs was 9.90 °C in
465 1965 and 14.88 °C in 1983, and annual JTs were significantly correlated with GSL ($r=0.46$,
466 $P<0.01$) and GDD ($r=0.64$, $P<0.001$) in the same year. FTs varied between -2.38 °C in 1963
467 and 5.94 °C in 1998, and were found to be associated with the NAO, with colder temperatures
468 coinciding with lower NAO indices (Figure 4b; $\beta_0=1.29$, $\beta_1=0.58$, $P<0.0001$, $R^2=33\%$).

469

470 Populations from different parts of Scotland experienced different levels of temporal variation
471 in these climate features. The combined *MAD* of FT and JT increased with ascending altitudes
472 (Figure 4c; $\beta_0= 1.032$, $\beta_1=0.00072$, $P<0.0001$, $R^2=77\%$), while for GSL and GDD, temporal
473 variability increased very little from altitudes of 48 to 343 m, but was higher at the three sites
474 located above 450 m (Figure 4d).

475

476

477 **Discussion**

478

479 In this study, we combined phenotypic and climate data to examine the patterns of variation in
480 a phenological trait among and within native Scottish populations of Scots pine. Under
481 common-garden conditions, populations sampled across a spatially highly heterogeneous
482 landscape were found to differ in timing of bud flush at the beginning of the second growing
483 season which generally was earlier in populations from cooler locations. This suggests
484 environment-driven genetic differentiation. However, significant amounts of variation were
485 also found within populations. In addition, the data suggested that populations may differ in
486 their level of adaptive variation: in the Aberdeen trial, we found some evidence of higher
487 levels of such variation in populations from high-altitude sites that experience the most
488 among-year variation in temperature conditions.

489

490

491 *Populations are differentiated in timing of bud flush*

492

493 The annual cycle of temperate trees is divided into two phases: active growing period in
494 summer and winter dormancy (Howe et al. 2003). Due to differences in the length of the
495 frost-free period in the Northern hemisphere, phenological differences are common among
496 tree populations (Howe et al. 2003; Savolainen et al. 2007). However, such patterns have
497 mainly been examined across wide geographic areas, and less is known about genetic
498 differences among populations separated by shorter distances. Despite the small geographic
499 area (maximum distance between two native pinewoods is less than 200 km), spatial
500 heterogeneity in climate within Scotland is extensive (Salmela et al. 2010). Thus, conditions
501 are ideal for development of local adaptation. Indeed, population differences observed in our
502 study suggest that adaptive differentiation in response to environmental variation has
503 occurred. Under glasshouse conditions, bud flush generally took place earlier in populations
504 from the coolest high-altitude locations in the eastern Highlands and later in those from the
505 maritime west coast. However, possible home site advantage of these populations cannot be

506 inferred without reciprocal transplant experiments (Kawecki and Ebert 2004). Also note that
507 due to the format of the partially interpolated climate data (5 km × 5 km grids) and spatially
508 complex landscapes in Scotland, it is possible that the home site conditions of the populations
509 differ from those described in Table 1. Weather station coverage in the UK is especially sparse
510 in the Scottish Highlands which is likely to result in inaccuracies in the climate variables
511 (Perry and Holliss 2005).

512

513 In spring, growth is initiated from stem units formed in buds during the previous growing
514 season after genetically-determined chilling and heat sum requirements have been fulfilled
515 (Aitken and Hannerz 2001; Howe et al. 2003). The patterns observed in our study could
516 reflect longer chilling and higher heat sum requirements of populations from warmer
517 locations so that growth initiation is prevented under mild winter conditions (e.g. Leinonen
518 1996). Due to the strong dependence of the trait on temperature, these patterns of variation
519 may differ among years (e.g. Sagnard et al. 2002, but see Beuker 1994). However,
520 corresponding spatial patterns of variation in this trait have been found in provenance studies
521 on adult trees of the same species sampled across Eurasia (Steiner 1979) and along a
522 latitudinal gradient in North Europe (Beuker 1994). Also, in an outdoor trial consisting of a
523 small subset of seedlings from eight populations included in the Edinburgh trial, timing of bud
524 flush at the beginning of the fourth growing season was slightly earlier in populations from
525 cooler high-altitude locations (Salmela et al. 2011). Nonetheless, in accordance with earlier
526 findings also in other species (Aitken and Hannerz 2001), population differences in growth
527 initiation appeared small. Although similar overall trends were found in the two trials, there
528 were differences in absolute values. This may be due to differences in winter and spring
529 temperatures between the experimental sites.

530

531 It is possible that phenotypic variation is influenced not only by genetic variation due to
532 segregating genes among seedlings, but also by differences in seed maturation conditions
533 experienced by different mother trees in their home environments. These effects have been
534 shown to be strong for instance in *Picea abies* (L.) H.Karst (Johnsen et al. 2005). Although
535 we cannot exclude the possibility of such effects influencing the observed patterns of
536 variation also among Scottish pine populations, earlier studies suggest that in Scots pine, such

537 effects are not of the same magnitude as in *P. abies*. For example, Ruotsalainen et al. (1995)
538 found that seed maturing conditions did not have major effect on variation in another
539 phenological trait, timing of bud set, under common-garden conditions. Further, in addition to
540 young seedlings, evidence of adaptive differentiation among populations in timing of bud
541 flush has been observed when examining Scots pine trees aged over 10 (Steiner 1979) or
542 approximately 60 years (Beuker 1994). Thus, in further discussion we assume that the
543 differences observed in this study reflect mainly genetic variation among seedlings.

544

545

546 *Populations consist of genetically and phenotypically diverse individuals*

547

548 Despite the evidence of population differentiation, a much larger proportion of variation was
549 due to differences among families and individual seedlings within populations. This
550 observation is akin for instance to the one found by Sagnard et al. (2002) among *Abies alba*
551 Mill. populations from the south-western Alps and indicates that the trait is heritable and that
552 populations maintain considerable internal potential to adapt to changing conditions (e.g.
553 Kramer and Havens 2009; Hoffman and Sgrò 2011), such as variable growing season length.
554 Furthermore, our data suggested that populations differed in levels of internal variation:
555 family differences were larger and families more variable at high-altitude locations. This
556 pattern was found only in the Aberdeen trial which may be due to the larger number of
557 families sampled within each population. Evolutionary biology models predict the loss of
558 genetic variation in adaptive traits due to selection favouring only optimal phenotypes in each
559 population (Falconer and Mackay 1996), but significant levels of within-population variation
560 are often documented in adaptive traits that are differentiated among populations (Howe et al.
561 2003; Alberto et al. 2011; Savolainen et al. 2011). Still, population differences in the amount
562 of variation have been assessed only on few occasions (Savolainen et al. 2004; Notivol et al.
563 2007; Alberto et al. 2011).

564

565 Several factors could contribute to the population differences in the level of trait diversity
566 along an altitudinal gradient observed in our study. Although the areas covered by the 21

567 pinewoods vary in size (Mason et al. 2004), differences in population size are an unlikely
568 explanation, as earlier work using selectively neutral molecular markers in the nuclear
569 genome has shown no significant differences in molecular diversity across Scotland (Kinloch
570 et al. 1986; Wachowiak et al. 2010). Gene flow among heterogeneous sites may increase
571 variation within populations (Howe et al. 2003) and at least historically, Scottish populations
572 have been linked by gene flow (Kinloch et al. 1986). Whether population differences in the
573 extent of long-range gene flow contribute to the patterns observed here requires further
574 exploration. However, studies of pollen flow suggest that the great majority of the fertilising
575 pollen usually comes from local trees (Smouse and Sork 2004), suggesting that a large
576 proportion of within-population diversity can also arise from matings between local parents.
577 Thus, in a common-garden study design sampling open-pollinated seed from natural stands,
578 gene flow from other populations might contribute more to variation within than among
579 families because offspring from matings between parents located in different populations
580 might not be as well adapted to their home site as those with local parents and might not
581 become established in a population (Burczyk et al. 2004). The high levels of residual variation
582 found in both our trials might have resulted from effective outcrossing, while population
583 differences in the level of variation within families may reflect variation in the extent of long-
584 distance pollen flow and/or the genetic diversity of local pollen donors. However, the current
585 study design does not allow the separation of genetic effects from other possible causes of
586 residual variation.

587

588 Expression of phenotypic variation is strongly influenced by the environment and
589 consequently, artificial growing conditions in glasshouses may induce the expression of
590 variation that would normally be 'hidden' in nature (e.g. Hoffmann and Merilä 1999).
591 Although growth conditions are known to affect population means in trees (Oleksyn et al.
592 1998; Mimura and Aitken 2010), their effects on trait variances remain poorly characterised.
593 It is possible that seedlings have expressed different levels of their total potential if the
594 growing conditions in the glasshouses were more novel to some populations. This possibility
595 could be investigated further by experiments in additional growing environments. In addition
596 to differences in within-population sampling, discrepancies between the spatial patterns of
597 variation among SD_{POP} 's between the trials could be due to different growing environments or
598 scoring intervals.

599

600 Adaptive diversity may be increased in environments that are highly variable across space or
601 time (Yeaman and Jarvis 2006). Assuming that the altitudinal range sampled at each site also
602 reflects the level of fine-scale environmental variation within populations, there was no
603 evidence of increased spatial heterogeneity at high altitudes. Finer-scale climate data are
604 needed to explore how environmental conditions vary across short distances in complex
605 landscapes like the Scottish Highlands. However, there is significant temporal heterogeneity
606 in climate in Scotland, and for instance mid-winter temperatures were found to be associated
607 with the NAO phenomenon which is known to influence a variety of biological events in both
608 plants and animals (Stenseth et al. 2002). The effect of the NAO in Europe is known to be
609 particularly strong in winter, but positive phases of the NAO during spring (February-April)
610 have also been demonstrated to be associated with elevated temperatures and earlier
611 phenological events in plants (Chmielewski and Rötzer 2001). Such variation may also partly
612 explain the observed temporal variation in the length of the growing seasons and the number
613 of growing degree days in Scotland.

614

615 We also found that the extent of temporal variation varied spatially within Scotland, and
616 higher-altitude locations were characterised by more variable climates. Interestingly, our
617 common-garden data suggest that genetic diversity at least in timing of bud flush may be
618 higher in populations found at the environmentally most variable sites. The association of
619 temporal climate variability with altitude most likely arises from the fact that in our sampling
620 sites, the highest-altitude sites are located at the most continental sites in the eastern
621 Highlands, while the lowest-altitude sites are found on the maritime west coast (see Vasseur
622 and Yodzis 2004). Although the climate data were based on interpolation across a temporally
623 and spatially variable number of weather stations and the precision of estimates varies
624 depending on the variable being estimated (Perry and Hollis 2005), patterns of variation in
625 latewood density chronologies among five Scottish pinewoods provide indirect but
626 corresponding evidence of site differences in temporal variability at least in summer
627 temperatures (Hughes et al. 1984).

628

629 Temporal variation in temperatures across the whole year suggests that phenotypic optima in

630 phenological traits in populations vary from year to year. Furthermore, among-year variation
631 in trait optima within Scotland may be more pronounced at sites with more among-year
632 variation in climate. Large-scale climate fluctuations are known to have shaped the
633 distributions of many tree species (Westfall and Millar 2004) and among-year variation for
634 instance in summer temperatures has been documented to decrease the likelihood of good
635 seed years especially in harsh conditions at high latitudes (e.g. Hilli et al. 2008), but so far, the
636 role of temporal variation in factors likely to drive adaptation in plant populations has
637 received only little attention. The contribution of temporal heterogeneity in maintaining
638 adaptive diversity has often been considered weak (Hedrick 1986), but on the other hand,
639 variation in environmental factors has not been studied in such detail as variation in
640 phenotype. The role of temporally variable environment might be important especially in
641 long-lived trees with overlapping generations and low climate-related mortality in adults (see
642 Persson and Ståhl 1990; Ellner and Hairston 1994). Trees aged over 400 years have been
643 found in Scottish pinewoods (Fish et al. 2010), and considering the evidence for substantial
644 temporal heterogeneity in the European climate since 1500 (Luterbacher et al. 2004) and the
645 age structure of populations, it is possible that different age cohorts have experienced
646 differing selection pressure during their sensitive early life stages and that the current patterns
647 of diversity reflect adaptations to a range of past environments. Accordingly, a more variable
648 environment might also support higher levels of genetic variation and more diverse
649 phenotypes. This possibility could be explored further by examining genetic differences
650 among age groups from sites that differ in temporal heterogeneity: larger differences among
651 age groups in more variable environments would provide stronger evidence for a positive
652 association between the levels of environmental and genetic diversity. Phenotypic plasticity is
653 also expected to evolve in heterogenous environments (Valladares et al. 2007); whether
654 differences in plasticity contribute to our observations could be investigated in more detail by
655 examining the range of phenotypic variation expressed across multiple growing environments.

656

657

658

659

660

661 *Potential to adapt in situ is found in natural populations occupying heterogeneous*
662 *environments*

663

664 Genetic variation in adaptive traits has important biological implications. When assumed to
665 have no capacity to evolve, changing environmental conditions may result in reduced survival
666 or extinction of current populations in some areas. When the internal capacity of populations
667 to adapt is accounted for, the chance of survival is increased (Aitken et al. 2008). Thus,
668 natural populations inhabiting heterogeneous habitats should not be treated as fixed and
669 independent entities whose responses to changes will be determined solely by the
670 environment. Also, studies on adaption in tree and other plant populations would benefit from
671 more thorough examinations of environmental factors likely to drive adaptation processes
672 both within and among populations. Despite the evidence of temporal heterogeneity being a
673 general feature of natural environments (Vasseur and Yodzis 2004), local climates in
674 evolutionary studies have generally been considered to be rather static and non-overlapping,
675 and the potential role of environmental heterogeneity in maintaining genetic diversity has not
676 been extensively explored.

677

678 In this study, we have shown that the environmental variability natural populations are
679 exposed to may differ even across short geographic distances and that this may influence the
680 amount of adaptive diversity found within populations. Predicting the future of natural
681 populations is complicated by possible correlations between adaptive traits, complex effects
682 of the environment on the expression of phenotypes, and many non-genetic factors, but
683 clearly, the fluctuating behaviour of environmental factors and the ubiquitous finding of
684 adaptive potential at least in some key traits needs to be taken into account in further studies
685 which aim to predict the effects of global change on natural populations. More family-
686 structured trials grown across a range of sites are also needed to characterise within-
687 population variation and its causes in more detail. Thanks to the long-history of common-
688 garden studies with appropriate study designs in forest trees, existing data from a large
689 number of completed studies can be used to test for similar patterns in other species.

690

691

692 **Acknowledgements**

693

694 The authors wish to thank Scottish Forestry Trust for funding (MJS' Ph.D. studentship), Dave
695 Sim, Joan Beaton, Sheila Reid, and Ben Moore (James Hutton Institute) for making the seed
696 collections and experimental assistance, NERC, the Forestry Commission, the Scottish
697 Government's Rural and Environment Science and Analytical Services Division (RESAS),
698 and EU-funded Network of Excellence EVOLTREE for support, UK Meteorological Office
699 for the climate data, and anonymous reviewers for constructive comments on the manuscript.

700

701

702 **Notes on contributors**

703

704 Matti Salmela is an evolutionary biologist interested in local adaptation, its quantitative
705 genetic basis, and maintenance of adaptive genetic variation in natural populations.

706

707 Stephen Cavers is a senior scientist at CEH, studying genetic diversity, gene flow, and
708 adaptation in plants, in both tropical and temperate ecosystems.

709

710 Joan Cottrell is head of the molecular team at Forest Research. Her research interests include
711 the assessment of genetic diversity, adaptation, and gene flow in native British trees and their
712 associated woodland dwelling flora and fauna. She is particularly interested in translating
713 research results into informed conservation advice on policy and management of our
714 woodland resource.

715

716 Glenn Iason is an ecologist with interests in herbivore nutritional ecology and ecosystem
717 management. He is also studying the extent of genetic and environmental variation in plant
718 secondary metabolites and how these determine plant-herbivore interactions, associated
719 biodiversity and ecosystem function. He is head of chemical and molecular ecology in the
720 Ecological Sciences group of the James Hutton Institute.

721

722 Richard Ennos is reader in ecological genetics at the University of Edinburgh with particular
723 interests in evolution and adaptation of plants and fungi.

724

725

726 **References**

727

728 Aitken SN, Hannerz M. 2001. Genecology and gene resource management strategies for
729 conifer cold hardiness. In: *Conifer Cold Hardiness*. Dordrecht (Netherlands): Kluwer
730 Academic Publishers. p. 23-53.

731

732 Aitken SN, Yeaman S, Holliday JA, Wang T, Curtis-McLane S. 2008. Adaptation, migration
733 or extirpation: climate change outcomes for tree populations. *Evolutionary Applications* 1:95-
734 111.

735

736 Alberto F, Bouffier L, Louvet J-M, Lamy J-B, Delzon S, Kremer A. 2011. Adaptive responses
737 for seed and leaf phenology in natural populations of sessile oak along an altitudinal gradient.
738 *Journal of Evolutionary Biology* 24:1442-1454.

739

740 Barton NH, Keightley PD. 2002. Understanding quantitative genetic variation. *Nature*
741 *Reviews Genetics* 3:11-21.

742

743 Benito Garzón M, Alía R, Robson TM, Zavala MA. 2011. Intra-specific variability and
744 plasticity influence potential tree species distributions under climate change. *Global Ecology*
745 *and Biogeography* 20:766-778.

746

747 Beuker E. 1994. Adaptation to climatic changes of the timing of bud burst in populations of

748 *Pinus sylvestris* L. and *Picea abies* (L.) Karst. Tree Physiology 14:961-970.
749
750 Billington HL, Pelham J. 1991. Genetic variation in the date of budburst in Scottish birch
751 populations: implications for climate change. Functional Ecology 5:403-409.
752
753 Bull JJ. 1987. Evolution of phenotypic variance. Evolution 41:303-315.
754
755 Burczyk J, DiFazio SP, Adams WT. 2004. Gene flow in forest trees: how far do genes really
756 travel? Forest Genetics 11:179-192.
757
758 Campbell, R. K. (1979) Genecology of Douglas-fir in a watershed in the Oregon Cascades.
759 Ecology 60, 1036-1050.
760
761 Chmielewski F-M, Rötzer T. 2001. Response of tree phenology to climate change across
762 Europe. Agricultural and Forest Meteorology 108:101-112.
763
764 Chmura DJ, Anderson PD, Howe GT, Harrington CA, Halofsky JE, Peterson DL, Shaw DC,
765 St. Clair JB. 2011. Forest responses to climate change in the northwestern United States:
766 ecophysiological foundations for adaptive management. Forest Ecology and Management
767 261:1121-1142.
768
769 Critchfield WB, Little E. 1966. Geographic distribution of the pines of the world. U.S.
770 Department of Agriculture Forest Service Miscellaneous Publication, 991.
771
772 Davis MB, Shaw RG. 2001. Range shifts and adaptive responses to Quaternary climate
773 change. Science 292:673-679.
774

775 Elith J, Leathwick JR. 2009. Species distribution models: ecological explanation and
776 prediction across space and time. *Annual Review of Ecology, Evolution, and Systematics* 40:
777 677-697.

778

779 Ellner S, Hairston NG. Jr. 1994. Role of overlapping generations in maintaining genetic
780 variation in a fluctuating environment. *American Naturalist* 143:403-417.

781

782 Eriksson G, Andersson S, Eiche V, Ifver J, Persson A. 1980. Severity index and transfer effects
783 on survival and volume production of *Pinus sylvestris* in Northern Sweden. *Studia Forestalia*
784 *Suecica* 156:1-32.

785

786 Falconer DS, Mackay TFC. 1996. *Introduction to Quantitative Genetics*. Second edition.
787 Essex (UK): Pearson Education Limited.

788

789 Fish T, Wilson R, Edwards C, Mills C, Crone A, Kirchefer AJ, Linderholm AW, Loader NJ,
790 Woodley E. 2010. Exploring for senescence signals in native Scots pine (*Pinus sylvestris* L.)
791 in the Scottish Highlands. *Forest Ecology and Management* 260:321-330.

792

793 Garcia-Gonzalez F, Simmons LW, Tomkins JL, Kotiaho JS, Evans JP. 2012. Comparing
794 evolvabilities: common errors surrounding the calculation and use of coefficients of additive
795 genetic variation. *Evolution* 66:2341-2349.

796

797 Harrison SJ. 1997. Changes in the Scottish climate. *Botanical Journal of Scotland* 49:287-300.

798

799 Hedrick PW. 1986. Genetic polymorphism in heterogeneous environments - a decade later.
800 *Annual Review of Ecology and Systematics* 17:535-566.

801

802 Hendry AP, Kinnison MT, Heino M, Day T, Smith TB, Fitt G, Bergstrom CT, Oakeshott J,
803 Jørgensen PS, Zalucki MP et al. 2011. Evolutionary principles and their practical application.
804 Evolutionary Applications 4:159-183.

805

806 Hilli A, Hokkanen T, Hyvönen J, Sutinen M-L. 2008. Long-term variation in Scots pine seed
807 crop size and quality in northern Finland. Scandinavian Journal of Forest Research 23:395-
808 403.

809

810 Hoffmann AA, Merilä J. 1999. Heritable variation and evolution under favourable and
811 unfavourable conditions. Trends in Ecology and Evolution 14:96-101.

812

813 Hoffmann AA, Sgrò CM. 2011. Climate change and evolutionary adaptation. Nature 470:
814 479-485.

815

816 Houle D. 1992. Comparing evolvability and variability of quantitative traits. Genetics 130:
817 195-204.

818

819 Howe GT, Aitken SN, Neale DB, Jermstad KD, Wheeler NC, Chen THH. 2003. From
820 genotype to phenotype: unraveling the complexities of cold adaptation in forest trees.
821 Canadian Journal of Botany 81:1247-1266.

822

823 Hughes MK, Schweingruber FH, Cartwright D, Kelly PM. 1984. July-August temperature at
824 Edinburgh between 1721 and 1975 from tree-ring density and width data. Nature 308:341-
825 344.

826

827 Johnsen Ø, Dæhlen OG, Østregg G, Skråpaa T. 2005. Daylength and temperature during seed
828 production interactively affect adaptive performance of *Picea abies* progenies. New
829 Phytologist 168:589-596.

830

831 Kawecki TJ, Ebert D. 2004. Conceptual issues in local adaptation. *Ecology Letters* 7:1225-
832 1241.

833

834 Kinloch BB, Westfall RD, Forrest GI. 1986. Caledonian Scots pine - origins and genetic
835 structure. *New Phytologist* 104:703-729.

836

837 Kramer AT, Havens K. 2009. Plant conservation genetics in a changing world. *Trends in Plant*
838 *Science* 14:599-607.

839

840 Leinonen I. 1996. Dependence of dormancy release on temperature in different origins of
841 *Pinus sylvestris* and *Betula pendula* seedlings. *Scandinavian Journal of Forest Research* 11:
842 122-128.

843

844 Linhart YB, Grant MC. 1996. Evolutionary significance of local genetic differentiation in
845 plants. *Annual Review of Ecology and Systematics* 27:237-277.

846

847 Luterbacher J, Dietrich D, Xoplaki E, Grosjean M, Wanner H. 2004. European seasonal and
848 annual temperature variability, trends, and extremes since 1500. *Science* 303:1499-1503.

849

850 Mason WL, Hampson A, Edwards C. 2004. *Managing the Pinewoods of Scotland*. Edinburgh:
851 Forestry Commission.

852

853 Merilä J, Sheldon BC. 1999. Genetic architecture of fitness and nonfitness traits: empirical
854 patterns and development of ideas. *Heredity* 83:103-109.

855

856 Mikola J. 1982. Bud-set phenology as an indicator of climatic adaptation of Scots pine in

857 Finland. *Silva Fennica* 16:178-184.

858

859 Milner JM, Elston DA, Albon SD. 1999. Estimating the contributions of population density
860 and climatic fluctuations to interannual variation in survival of Soay sheep. *Journal of Animal*
861 *Ecology* 68:1235-1247.

862

863 Mimura M, Aitken SN. 2010. Local adaptation at the range peripheries of Sitka spruce.
864 *Journal of Evolution Biology* 23:249-258.

865

866 Notivol E, García-Gil MR, Alía R, Savolainen O. 2007. Genetic variation of growth rhythm in
867 the limits of a latitudinal cline in Scots pine. *Canadian Journal of Forest Research* 37:540-551.

868

869 Oleksyn J, Tjoelker MG, Reich PB. 1998. Adaptation to changing environment in Scots pine
870 populations across a latitudinal gradient. *Silva Fennica* 32:129-140.

871

872 Parmesan C. 2006. Ecological and evolutionary responses to recent climate change. *Annual*
873 *Review of Ecology, Evolution, and Systematics* 37:637-669.

874

875 Perry M, Hollis D. 2005. The generation of monthly gridded datasets for a range of climatic
876 variables over the UK. *International Journal of Climatology* 25:1041-1054.

877

878 Persson B, Ståhl EG. 1990. Survival and yield of *Pinus sylvestris* L. as related to provenance
879 transfer and spacing at high altitudes in northern Sweden. *Scandinavian Journal of Forest*
880 *Research* 5:381-395.

881

882 Petit RJ, Hampe A. 2006. Some evolutionary consequences of being a tree. *Annual Review of*
883 *Ecology, Evolution, and Systematics* 37:187-214.

884

885 Ray D. 2008. Forestry Commission Research Note 101: Impacts of climate change on forestry
886 in Scotland - a synopsis of spatial modelling research. Forestry Commission Scotland.

887

888 Rehfeldt GE, Tchebakova NM, Parfenova YI, Wykoff WR., Kuzmina NA, Milyutin LI. 2002.
889 Intraspecific responses to climate in *Pinus sylvestris*. Global Change Biology 8:912-929.

890

891 Reich PB, Oleksyn J. 2008. Climate warming will reduce growth and survival of Scots pine
892 except in the far north. Ecology Letters 11:588-597.

893

894 Rousi M, Heinonen J. 2007. Temperature sum accumulation effects on within-population
895 variation and long-term trends in date of bud burst of European white birch (*Betula pendula*).
896 Tree Physiology 27:1019-1025.

897

898 Ruotsalainen S, Nikkanen T, Haapanen M. 1995. Effect of seed-maturing conditions on the
899 growth and hardiness of one-year old *Pinus sylvestris* seedlings. Forest Genetics 2:189-198.

900

901 Sagnard F, Barberot C, Fady B. 2002. Structure of genetic diversity in *Abies alba* Mill. from
902 southwestern Alps: multivariate analysis of adaptive and non-adaptive traits for conservation
903 in France. Forest Ecology and Management 157:175-189.

904

905 Salmela MJ, Cavers S, Cottrell JE, Iason GR, Ennos RA. 2011. Seasonal patterns of
906 photochemical capacity and spring phenology reveal genetic differentiation among native
907 Scots pine (*Pinus sylvestris* L.) populations in Scotland. Forest Ecology and Management
908 262:1020-1029.

909

910 Salmela MJ, Cavers S, Cottrell JE, Wachowiak W, Iason GR, Ennos RA. 2010. Understanding
911 the evolution of native pinewoods in Scotland will benefit their future management and

912 conservation. *Forestry* 83:535-545.

913

914 Savolainen O, Bokma F, García-Gil R, Komulainen P, Repo T. 2004. Genetic variation in
915 cessation of growth and frost hardiness and consequences for adaptation of *Pinus sylvestris* to
916 climatic changes. *Forest Ecology and Management* 197:79-89.

917

918 Savolainen O, Kujala ST, Sokol C, Pyhäjärvi T, Avia K, Knürr T, Kärkkäinen K, Hicks S,
919 2011. Adaptive potential of northermost tree populations to climate change with emphasis on
920 Scots pine (*Pinus sylvestris* L.). *Journal of Heredity* 102:526-536.

921

922 Savolainen O, Pyhäjärvi T, Knürr T. 2007. Gene flow and local adaptation in trees. *Annual*
923 *Review of Ecology, Evolution, and Systematics* 38:595-619.

924

925 Smouse PE, Sork VL. 2004. Measuring pollen flow in forest trees: an exposition of alternative
926 approaches. *Forest Ecology and Management* 197:21-38.

927

928 Steiner KC. 1979. Patterns of variation in bud-burst timing among populations in several
929 *Pinus* species. *Silvae Genetica* 28:185-194.

930

931 Stenseth NC, Mysterud A, Ottersen G, Hurrell JW, Chan K-S, Lima M. 2002. Ecological
932 effects of climate fluctuations. *Science* 297:1292-1296.

933

934 Valladares F, Gianoli E, Gómez JM. 2007. Ecological limits to plant phenotypic plasticity.
935 *New Phytologist* 176:749-763.

936

937 Vasseur DA, Yodzis P. 2004. The color of environmental noise. *Ecology* 85:1146-1152.

938

939 Wachowiak W, Salmela MJ, Ennos RA, Iason G, Cavers S. 2010. High genetic diversity at the
940 extreme range edge: nucleotide variation at nuclear loci in Scots pine (*Pinus sylvestris* L.) in
941 Scotland. *Heredity* 106:775-787.

942

943 Westfall RD, Millar CI. 2004. Genetic consequences of forest population dynamics influenced
944 by historic climatic variability in the western USA. *Forest Ecology and Management*
945 197:159-170.

946

947 White TL, Adams WT, Neale DB. 2007. *Forest Genetics*. Wallingford (UK): CABI
948 Publishing.

949

950 Willi Y, Van Buskirk J, Hoffman AA. 2006. Limits to adaptive potential of small populations.
951 *Annual Review of Ecology, Evolution, and Systematics* 37:433-438.

952

953 Yeaman S, Jarvis A. 2006. Regional heterogeneity and gene flow maintain variance in a
954 quantitative trait within populations of lodgepole pine. *Proceedings of the Royal Society B:*
955 *Biological Sciences* 273:1587-1593.

956

957

958

959

960

961

962

963

964

965

966 **Tables**

967

968 Table 1 Populations of *Pinus sylvestris* in Scotland included in the study, their latitude (Lat.),
 969 longitude (Long.), altitudinal range sampled (Alt.), core pinewood area according to Mason et
 970 al. (2004), and mean (1961-2000) calculated climate features: growing season length (GSL;
 971 days), growing degree days (GDD: day degrees), and February and July mean temperatures
 972 (FMT and JMT).

973

Population	Lat.	Long.	Alt. (m)	Area (ha)	GSL	GDD	FMT (°C)	JMT (°C)
Abernethy (AB)	57.21	3.61	311-370	2452	211	990	1.15	12.73
Allt Cul (AC)	57.04	3.35	435-512	13	145	513	-1.01	10.41
Amat (AM)	57.87	4.60	39-201	181	214	892	1.22	12.29
Ballochbuie (BB)	56.98	3.30	421-531	775	116	446	-1.69	9.46
Beinn Eighe (BE)	57.63	5.40	17-91	182	283	1329	3.68	14.16
Black Wood of Rannoch (BW)	56.68	4.37	250-321	1011	254	1138	2.12	13.55
Coille Coire Chuilc (CCC)	56.42	4.71	222-311	67	226	928	1.64	12.32
Conaglen (CG)	56.79	5.33	89-193	189	246	887	2.20	11.73
Crannach (CR)	56.58	4.68	258-338	70	231	1019	1.81	12.62
Glen Affric (GA)	57.26	4.92	205-293	1532	210	769	0.88	11.62
Glen Cannich (GC)	57.35	4.95	182-381	301	212	778	0.96	11.71
Glen Derry (GD)	57.03	3.58	426-493	235	168	593	-0.46	11.34
Glen Einig (GE)	57.96	4.76	45-92	27	242	1089	2.19	13.15
Glen Loy (GL)	56.91	5.13	136-219	74	191	541	0.49	9.80
Glen Tanar (GT)	57.02	2.86	289-422	1564	235	1105	2.21	13.63
Loch Clair (LC)	57.56	5.36	98-166	126	277	1253	3.44	13.68
Meggernie (MG)	56.58	4.35	254-385	277	223	916	1.07	12.04
Rhidorroch (RD)	57.89	4.98	138-220	103	221	840	1.51	11.62
Rothiemurchus (RM)	57.15	3.77	295-329	1744	224	1087	1.39	13.15
Shieldaig (SD)	57.50	5.63	44-132	103	273	1093	3.21	12.83
Strath Oykel (SO)	57.98	4.61	35-160	14	257	1276	2.69	14.05

974

975

976

977

978

979

980

981 Table 2 Variation in timing of bud flush. Results of the nested ANOVA testing the effects of
982 population (fixed factor), families within populations (random factor), and blocks (random
983 factor) in the a) Edinburgh and b) Aberdeen trials.

984

a) Edinburgh trial

Source of variation	df	MS	F-ratio	P-value	Variance component
Population	20	1153.35	1.70	0.058	2.98
Families within populations	63	679.52	13.50	<0.001	15.39
Block	39	176.66	3.51	<0.001	1.50
Residual	3120	50.35			50.4

b) Aberdeen trial

Source of variation	df	MS	F-ratio	P-value	Variance component
Population	20	183.26	2.18	<0.01	1.45
Families within populations	188	83.96	2.01	<0.001	5.88
Block	39	221.37	5.29	<0.001	5.09
Residual	1216	41.87			42.26

985

Accepted Manuscript

986 Table 3 The amount of within-population variation in timing of bud flush. Standard deviations
 987 (SD_{POP}), variance components due to families (V_{AF}), residual variation (V_{WF}), and blocks
 988 (V_{Block}), and the range among family means in each population within the two trials.

Population	Edinburgh					Aberdeen				
	SD_{POP}	V_{AF}	Mean range	V_{WF}	V_{Block}	SD_{POP}	V_{AF}	Mean range	V_{WF}	V_{Block}
Abermethyl (AB)	9.53	43.95	15.63	70.12	0.00	8.48	0.00	14.09	76.92	0.00
Allt Cul (AC)	7.76	8.90	6.20	57.18	0.00	8.98	2.39	12.19	62.75	16.32
Amat (AM)	7.32	15.41	8.90	40.15	1.80	8.68	3.43	12.09	70.03	2.11
Ballochbuie (BB)	8.86	27.93	12.38	62.28	0.00	8.30	0.00	10.30	52.46	22.99
Beinn Eighe (BE)	7.62	9.56	6.72	48.00	2.84	6.29	0.00	6.09	34.88	6.64
Black Wood of Rannoch (BW)	8.00	11.79	8.24	55.55	0.00	7.92	0.00	9.62	27.07	35.22
Coille Coire Chuilc (CCC)	7.25	6.26	5.73	42.20	5.65	7.92	13.16	12.15	43.73	6.08
Conaglen (CG)	7.97	4.56	5.69	54.22	5.81	4.29	0.01	7.08	20.36	0.00
Crannach (CR)	9.19	55.18	16.34	51.34	0.80	5.52	11.40	10.02	23.81	0.00
Glen Affric (GA)	6.79	5.14	4.78	43.25	0.00	5.29	0.00	3.15	36.67	0.00
Glen Cannich (GC)	8.17	23.58	12.02	45.50	3.70	6.18	12.00	14.13	24.92	3.02
Glen Derry (GD)	7.95	30.24	11.85	39.48	1.07	11.10	29.05	24.03	94.57	5.24
Glen Einig (GE)	7.93	2.95	5.06	66.41	0.00	5.81	3.79	8.79	19.06	10.44
Glen Loy (GL)	6.78	7.31	6.75	37.61	2.75	6.71	7.03	10.19	41.43	0.00
Glen Tanar (GT)	6.25	8.09	6.22	32.43	0.44	8.17	11.87	16.91	53.83	2.84
Loch Clair (LC)	8.51	0.00	1.98	64.00	9.16	7.76	0.00	11.99	76.21	0.00
Meggernie (MG)	7.10	0.72	3.48	47.31	2.61	8.51	4.76	13.92	71.13	0.00
Rhidorroch (RD)	8.10	8.00	6.53	58.24	0.41	5.42	2.88	12.46	26.72	0.06
Rothiemurchus (RM)	9.02	22.73	10.55	61.92	2.35	6.17	11.53	12.84	18.53	6.97
Shieldaig (SD)	8.49	26.01	10.84	53.52	0.00	5.42	0.00	6.28	37.54	0.00
Strath Oykel (SO)	7.27	14.68	6.94	42.00	0.00	5.55	0.00	6.61	27.00	4.22

989

Accepted Manuscript

990 **Captions**

991

992 Figure 1 Map of the sampled native *Pinus sylvestris* populations and locations of the two trial
993 sites. Climatic features of the sites are given in Table 1. Inset: full distribution of *P. sylvestris*,
994 with study area highlighted by box.

995

996 Figure 2 Relationship between mean growing degree days (GDD) at origin and population
997 means of timing of bud flush in the two trials. In the Edinburgh trial: $\beta_0=8.14$, $\beta_1=0.0066$,
998 $P<0.01$, $R^2=35\%$; in the Aberdeen trial: $\beta_0=15.70$, $\beta_1=0.0040$, $P<0.001$, $R^2=32\%$. Error bars
999 indicate standard errors of the means.

1000

1001 Figure 3 Relationship between altitude at origin and the amount of variation (SD_{POP}) in timing
1002 of bud flush within populations in the Aberdeen trial ($\beta_0=5.14$, $\beta_1=0.0080$, $P<0.01$, $R^2=35\%$).


1003

1004 Figure 4 Among-year variation in the Scottish climate. a) Temporal variation in growing
1005 season length (GSL) and growing degree days (GDD); b) temporal variation in February
1006 temperature (FT) and February North Atlantic Oscillation (NAO) indices; c) relationship
1007 between the altitudes of the 21 native pinewood sites and among-year variability of winter
1008 and summer temperatures (combined MAD of FT and JT); d) temporal variation in GSL and
1009 GDD (coefficients of variation, CV) plotted against the altitudes of the 21 sampled sites. The
1010 climate data used cover the period 1961-2000. In a) and b), annual means were calculated
1011 over the $5\text{ km} \times 5\text{ km}$ grids within which the 21 pinewood sites are located.

1012

1013
1014 Figure 1.
1015


Accepted Manuscript



1017

1018 Figure 2.

1019




1020

Accepted Manuscript

1021 Figure 3.

1022

1023



1024

1025

1026

1027

1028


1029

1030

1031

Accepted Manuscript

1032 Figure 4. a)




1033

1034

1035

b)



1036

1037

1038

1039

1040

1041


1042

1043

1044

1045


c)



1046

1047

d)



1048

1049

1050