

## Article (refereed) - postprint

---

Gunn, Iain D.M.; Meis, Sebastian; Maberly, Stephen C.; Spears, Bryan M.  
2014. **Assessing the responses of aquatic macrophytes to the application of a lanthanum modified bentonite clay, at Loch Flemington, Scotland, UK.** *Hydrobiologia*, 737 (1). 309-320. [10.1007/s10750-013-1765-5](https://doi.org/10.1007/s10750-013-1765-5)

© Springer Science+Business Media Dordrecht 2013

This version available <http://nora.nerc.ac.uk/504207/>

NERC has developed NORA to enable users to access research outputs wholly or partially funded by NERC. Copyright and other rights for material on this site are retained by the rights owners. Users should read the terms and conditions of use of this material at <http://nora.nerc.ac.uk/policies.html#access>

**This document is the author's final manuscript version of the journal article, incorporating any revisions agreed during the peer review process. Some differences between this and the publisher's version remain. You are advised to consult the publisher's version if you wish to cite from this article.**

**The final publication is available at Springer via <http://dx.doi.org/10.1007/s10750-013-1765-5>**

Contact CEH NORA team at  
[noraceh@ceh.ac.uk](mailto:noraceh@ceh.ac.uk)

1 Assessing the responses of aquatic macrophytes to the application of a  
2 lanthanum modified bentonite clay, at Loch Flemington, Scotland, UK

3 IAIN D. M. GUNN\*

4 \*Corresponding author

5 Centre for Ecology & Hydrology, Bush Estate, Penicuik, Midlothian EH26 OQB, UK

6 E-mail: [idmg@ceh.ac.uk](mailto:idmg@ceh.ac.uk)

7

8 SEBASTIAN MEIS

9 Centre for Ecology & Hydrology, Bush Estate, Penicuik, Midlothian, EH26 OQB, UK

10 STEPHEN C. MABERLY

11 Centre for Ecology & Hydrology, Lancaster Environment Centre, Library Avenue, Bailrigg, Lancaster

12 LA1 4AP, UK

13 BRYAN M. SPEARS

14 Centre for Ecology & Hydrology, Bush Estate, Penicuik, Midlothian, EH26 OQB, UK

15

16 **ABSTRACT**

17

18 Loch Flemington is a shallow lake of international conservation and scientific importance. In recent decades its  
19 status has declined as a result of eutrophication and the establishment of non-native invasive aquatic  
20 macrophytes. As previous research had identified the lake bed sediments as an important source of phosphorus  
21 (P), the P-capping material Phoslock<sup>®</sup> was applied to improve water quality. This paper documents the  
22 responses of the aquatic macrophyte community by comparing data collected between 1988 and 2011. Summer  
23 water-column total P concentrations decreased significantly and water clarity increased following treatment.

24 Aquatic plant colonisation depth increased and plant coverage of the lake bed extended. However, the  
25 submerged vegetation remained dominated by the non-native *Elodea canadensis* Michx. Aquatic macrophyte  
26 community metrics indicated no significant change in trophic status. Species richness and the number of  
27 'natural' eutrophic characteristic species remained broadly similar with no records of rare species of  
28 conservation interest. Loch Flemington is still classified as being in 'unfavourable no change' condition based  
29 on its aquatic macrophytes despite the water quality improvements. The implications of these results are  
30 discussed in relation to the future management of Loch Flemington and in the wider context of trying to  
31 improve our understanding of lake restoration processes.

32 **KEYWORDS:** Eutrophication, Loch Flemington, lake restoration, P-capping material, aquatic macrophytes,  
33 lake management

34

## 35 **INTRODUCTION**

36

37 Eutrophication is regarded as being one of the most important factors causing degradation of lakes throughout  
38 Europe (Withers & Haygarth, 2007) and the rest of the world (Bennett *et al.*, 2001; Schindler, 2006). In shallow  
39 (< 3 m depth) freshwater lakes, elevated nutrient inputs from the catchment, resulting from agricultural, cultural  
40 and industrial activity, are the primary cause of this eutrophication (Scheffer, 2004). Increased loadings of  
41 phosphorus (P) commonly result in a shift from macrophyte to planktonic production, with associated  
42 phytoplankton blooms (particularly cyanobacterial), leading to deterioration in water quality and associated  
43 changes in the biological communities. European legislation has led in the UK to the setting of P targets for high  
44 alkalinity, shallow lakes to achieve 'favourable condition' (JNCC, 2005) under the Habitats Directive (Council  
45 of the European Communities, 1992) and 'good' and 'high' ecological status (UKTAG, 2008) under the Water  
46 Framework Directive (WFD) (Council of the European Communities, 2000). However, many shallow lakes in  
47 Europe, because of a legacy of eutrophication, already have nutrient concentrations which exceed these targets  
48 (EEA, 2012). For shallow lakes, designated to be of conservation importance, this deterioration in condition is  
49 also indicated by the decline or disappearance of rare and scarce vascular plants and/or assemblages of notable  
50 vascular plants and charophytes (JNCC, 2005).

51 On a global scale, research on eutrophication in relation to lake management has received considerable attention  
52 (Jeppesen *et al.*, 2003). Successful lake restoration depends on a good understanding of the site-specific drivers  
53 of eutrophication and using targeted management strategies which are based on good scientific evidence. In  
54 many lakes the main factor limiting phytoplankton abundance is P availability, so it is this nutrient which is  
55 usually targeted for control by most lake restoration strategies, although nitrogen (N) limitation in the summer  
56 months can also be important (Fisher *et al.*, 2009; May *et al.*, 2010). Reducing the external P load from a lake  
57 catchment is often the preferred remedial option where P is the limiting factor. However, its effectiveness is  
58 often reduced or delayed by internal recycling of P from the lake sediments (Søndergaard *et al.*, 2007). In such  
59 cases, alternative approaches to lake recovery must be used, coupled with reducing the external P load, such as  
60 bio-manipulation, biomass harvesting, lake flushing, sediment removal, vertical mixing, and P inactivation  
61 (Cooke *et al.*, 2005). The latter technique has in the past relied on adding iron, aluminium or similar salts to bind  
62 and precipitate available dissolved P from the water-column to the lake sediments. This prevents internal  
63 loading by maintaining, for example, a layer of iron over the sediments, but, often, after short-term success, pre-  
64 treatment levels of P release return within a few years (e.g. Foy, 1985). There have also been concerns that such  
65 techniques may have adverse effects on lake ecosystems (Randall *et al.*, 1999; Spears *et al.*, 2013a). A range of  
66 P-capping materials have been proposed for use in the UK (Spears *et al.*, 2013b). Phoslock<sup>®</sup> is a lanthanum (La)  
67 modified bentonite clay designed to enhance the capacity of lake sediments to adsorb dissolved (i.e. bio-  
68 available) P and significantly reduce its subsequent release (manufacturers reported binding capacity of 100g  
69 Phoslock<sup>®</sup>: 1 g of P). La<sup>3+</sup> ions that are strongly bound within the clay matrix react with dissolved P to form the  
70 mineral rhabdophane (LaPO<sub>4</sub>), a compound that is expected to be stable in the environment (i.e. beyond the pH  
71 range 4 – 9; Ross *et al.*, 2008).

72 The expected general responses of the aquatic macrophyte community, following eutrophication management,  
73 have recently been summarised in a literature review which incorporated long-term data from oligotrophication  
74 case studies (Jeppesen *et al.*, 2005; Verdonschot *et al.*, 2011). Typical macrophyte community responses  
75 included increases in colonisation depth, species richness (including relative characean abundance), the number  
76 of nutrient intolerant species and species distribution as TP concentrations decreased. Submerged macrophyte  
77 vegetation becomes increasingly restricted to shallower waters in response to decreasing light levels at the  
78 sediment surface as phytoplankton population levels increase. One simple measure of macrophyte abundance in  
79 lakes is maximum macrophyte growing depth (MMGD). MMGD is sensitive to eutrophication (Jupp & Spence,  
80 1977; May & Carvalho, 2010; Søndergaard *et al.*, 2005; Spears *et al.*, 2009) and can provide a direct measure of

81 the success of lake management measures to shift primary production in lakes from a phytoplankton-dominated  
82 to a more macrophyte-dominated state (Sas, 1989).

83 This paper examines how the aquatic macrophyte community in Loch Flemington responded following the  
84 application of the P-capping material Phoslock<sup>®</sup> to the lake bed sediments and, on the assumption that growing  
85 conditions will improve as a result of better water quality, tested the following hypotheses:

- 86 • There will be no negative impacts of the Phoslock<sup>®</sup> application on aquatic macrophyte community
- 87 • Macrophyte maximum growing depth and coverage of lake bed will increase
- 88 • Species diversity will increase and species of conservation interest will re-appear
- 89 • Aquatic macrophyte community attributes and metrics will indicate a change towards improvement in  
90 Loch Flemington's condition and trophic status
- 91 • Mechanical control of non-native aquatic species *Elodea canadensis* Michx. may encourage the  
92 recovery of native aquatic macrophyte species when used in combination with Phoslock<sup>®</sup>

93

## 94 **METHODS**

### 95 **Study site**

96 Loch Flemington (N 57° 32.570', W 3° 59.399') is a shallow (mean and maximum depths of 0.75 m and 2.9 m),  
97 lowland (altitude 46 m), high alkalinity (1240 µeqL<sup>-1</sup>), eutrophic lake, of glacial origin situated near Inverness in  
98 Scotland, UK (Figure 1). It has a surface area of 15 ha, a maximum fetch of 0.75 km and a perimeter of 2.7 km  
99 (UKLakes Database). The lake is situated in an area consisting of glaciofluvial sand and gravel within the  
100 Kildrummie Kames esker system (Gordon & Auton, 1993). Significant hydrological modifications were made to  
101 the lake in the 19<sup>th</sup> century by blocking the natural surface-water outflow and significantly increasing the surface  
102 area and depth of the loch (May *et al.*, 2001). Today, the lake has no natural surface water outflow and the Croy  
103 Burn is the only surface water inflow. Water leaves the lake by evaporation and by percolating through  
104 permeable gravels along the northwest shore, leading to an estimated water retention time of around 40 days  
105 (May *et al.*, 2001).

106 Loch Flemington lies within the Kildrummie Kames Site of Special Scientific Interest (SSSI) which was  
107 notified in 1974, in part, for Loch Flemington's eutrophic lake habitat and for supporting populations of aquatic  
108 macrophyte species of conservation interest: the Nationally Rare *Potamogeton rutilus* Wolfg. and the  
109 Nationally Scarce *Potamogeton filiformis* Pers. In addition, macrofossils of the Nationally Scarce and European  
110 Protected Species *Najas flexilis* (Willd.) Rostk. & Schmidt (listed in Annexes II and IV of the Habitats  
111 Directive (Council of the European Communities, 1992)) have recently been recorded in Loch Flemington  
112 (Bennion *et al.*, 2008). From the 1960s onwards Loch Flemington had become increasingly eutrophic due to  
113 nutrient-laden waste entering from the catchment resulting in a lake flora dominated by three submerged  
114 species *E. canadensis*, *Myriophyllum spicatum* L. and *Potamogeton crispus* L. (May *et al.*, 2001).  
115 Palaeolimnological records indicated that the aquatic macrophyte community had actually changed from a rich  
116 macrophyte flora characteristic of mesotrophic conditions before 1850 towards a more species poor community  
117 indicative of nutrient enrichment (Bennion *et al.*, 2008). This resulted in a build up of P within the lake which  
118 caused, by the mid 1970s, troublesome algal blooms to develop and fish kills to occur. Nuisance algal blooms,  
119 particularly those dominated by cyanobacteria, continued throughout the 1980s and 1990s with increasing  
120 frequency, threatening the conservation status of the lake. By 2003/04 the SSSI eutrophic lake habitat feature  
121 was found to be in 'unfavourable condition'. The main reasons for this negative classification were the lake's  
122 continuing poor water quality and the domination of the submerged vegetation by the non-native *E. canadensis*  
123 plus the disappearance of all the aquatic macrophyte species of national conservation interest. Restoration  
124 efforts at Loch Flemington up to this point had been focussed solely on improving water quality by reducing  
125 catchment point sources of nutrients. An assessment of total P (TP) loads to the lake indicated that, by 2001,  
126 diffuse (mainly agricultural) sources, represented ~ 80% and septic tanks ~ 18% of the external TP load,  
127 respectively (May *et al.*, 2001). However, the same study also found that the internal, sediment-driven load to  
128 the water-column (680 kg TP yr<sup>-1</sup>) was probably far greater than the external load from the catchment (120 kg  
129 TP yr<sup>-1</sup>).

### 130 **Phoslock® application**

131 Phoslock® was applied to Loch Flemington by Phoslock® Europe GmbH over a three day period, March 13<sup>th</sup> –  
132 15<sup>th</sup> 2010. A pilot study had estimated that the dosage of Phoslock® required to be applied to Loch Flemington  
133 was 25 tonnes in order to control a maximum of ~ 22 kg total phosphorus (TP) in the water column and ~ 210

134 kg TP in the upper 3 cm of lake sediment, on the basis that 100 g of Phoslock<sup>®</sup> binds 1 g of phosphorus (Meis *et*  
135 *al.*, 2012).

### 136 **Pre- and post-application water quality and ecological (non-aquatic macrophyte) monitoring**

137 A range of water quality and ecological (non-aquatic macrophyte) parameters were monitored at Loch  
138 Flemington on a monthly basis, pre- and post-application of Phoslock<sup>®</sup>, at five open water sampling sites  
139 between May 2009 and March 2011, and at three of these sites from April 2011 to November 2011, with the  
140 exception of July 2011 when all five sites were sampled. Field water quality measurements included surface  
141 water pH, conductivity, temperature and dissolved oxygen concentrations while water clarity was assessed using  
142 a Secchi disc. Water samples were collected for later analysis. Parameters measured included soluble reactive P  
143 (SRP), TP, total lanthanum (TLa), soluble lanthanum, chlorophyll *a* and phytoplankton abundance, community  
144 composition and bioassessment of biovolume. Crustacean zooplankton and macroinvertebrates were also  
145 collected for later analysis at the open water sample sites. For further details of sampling processing and  
146 analyses see Spears *et al.* (2012). In this paper only the mean annual surface water TP and chlorophyll *a*  
147 concentration results are reported on.

### 148 **Aquatic macrophyte community monitoring**

149 Assessment of the aquatic macrophyte communities at Loch Flemington, pre - and post - application of  
150 Phoslock<sup>®</sup>, involved three separate but related sampling monitoring programmes: Site Condition Monitoring  
151 (SCM) surveys, monthly qualitative assessments of the submerged vegetation and monitoring of the maximum  
152 macrophyte growing depths. Aquatic macrophyte data from a comprehensive 1988 survey of Loch Flemington,  
153 carried out by the Nature Conservancy Council Scottish Loch Survey team (NCC, 1988), were used to help  
154 draw historical comparisons.

### 155 ***Site Condition Monitoring (SCM) surveys***

156 The Site Condition Monitoring (SCM) survey method followed the Common Standards Monitoring (CSM)  
157 protocols produced by the UK Joint Nature Conservation Committee (JNCC) for assessing the aquatic  
158 macrophyte communities of standing waters of conservation importance (JNCC, 2005; JNCC, 2009). Pre-  
159 Phoslock<sup>®</sup> application SCM surveys were carried out in 2003/04 and 2009 and post-Phoslock<sup>®</sup> application  
160 aquatic macrophyte surveys were carried out in 2010 and 2011 (Table 1). Although the sampling effort differed  
161 slightly between these different SCM-style surveys, the resultant data are considered to be broadly comparable.

162 The SCM aquatic macrophyte survey techniques employed at Loch Flemington involved three main elements  
163 and are described in detail by (Gunn *et al.*, 2010; JNCC, 2005; JNCC, 2009). These were as follows: (1)  
164 perimeter strandline searches; (2) shore-wader depth transect surveys; and (3) boat-based depth transect surveys.

165 All three methods were based on representative 100 m sections or sectors selected around the perimeter of the  
166 lake in areas thought to be suitable for sustaining good aquatic macrophyte populations. Both the shore-wader  
167 and boat-based depth transects involved 20 sampling points per survey sector. These SCM methods were  
168 designed to produce data which are statistically robust (Gunn *et al.*, 2010).

#### 169 ***Qualitative assessments of aquatic macrophyte vegetation***

170 Seven monthly pre-application and thirteen monthly post-application qualitative assessments of the aquatic  
171 macrophyte community composition and abundance were carried out at Loch Flemington between May 2009  
172 and July 2011. These assessments involved random searches, by boat, for a fixed period of two hours. These  
173 searches were standardised by starting at a fixed point and going back and forth across the lake in straight  
174 transect lines sampling the submerged vegetation en route using a double-headed rake and/or using a  
175 bathyscope.

#### 176 ***Macrophyte maximum growing depth (MMGD) monitoring***

177 The MMGD of submerged macrophytes at Loch Flemington were assessed using a boat along five fixed  
178 transects that were evenly spaced around the deepest, north-eastern bay of the lake. Pre-application monitoring  
179 of the MMGD was carried out on a monthly basis from July 2009 to November 2009 and just before the  
180 Phoslock<sup>®</sup> application on the 11th March 2010. Post-application monitoring of MMGD was also carried out on a  
181 monthly basis from April 2010 through to March 2011 with an additional monitoring occasion in July 2011  
182 using methods as described by Jupp *et al.* (1974). Water depth and plant occurrence were measured at about 2 m  
183 horizontal intervals along each transect until a point was reached at which no plants were collected. MMGD was  
184 corrected for changes in water level by measuring water level at a common datum and subtracting water level  
185 height from the measured MMGD. The common datum was fixed at an average water depth of 0.38 m (range  
186 0.14 to 0.69 m). The methodology followed Spears *et al.* (2009). Data sets were arranged into seasons consisting  
187 of three months in line with the standard meteorological definition (Trenberth, 1983) resulting in four seasons:  
188 summer (June – August), autumn (September – November), winter (December – February) and spring (March –  
189 May). Statistical analyses were conducted using the software package Minitab 16 (Minitab<sup>®</sup> 16.1.1, Minitab

190 Ltd., Coventry, UK). Data sets of MMGD were not normally distributed (Anderson-Darling test,  $\alpha < 0.05$ ) even  
191 after a range of transformations (including  $x' = \log(x)$ ,  $x' = \ln(x)$ ,  $x' = \sqrt{x}$ ,  $x' = x^2$ ). Therefore a non-  
192 parametric Mann-Whitney U-test (MWU) was used to test for significant variation in MMGD between years for  
193 a given season, for seasons consisting of at least two out of three possible months as it is assumed that at least  
194 two months are required to represent a season. This procedure resulted in the following statistical comparisons:  
195 summer 2009 vs. summer 2010 and autumn 2009 vs. autumn 2010.

#### 196 **Clear-cutting experiment to mechanically control *E. canadensis***

197 To assess the response of native macrophytes species in the absence of the non-native *E. canadensis* in Loch  
198 Flemington, a replicated clear-cutting experiment was carried out (summer 2010) following the application of  
199 Phoslock<sup>®</sup>. A diver manually cleared five 4 m<sup>2</sup> plots of *E. canadensis* and marked a further five plots as  
200 controls. The percentage volume inhabited (PVI) and the aquatic plant community composition were tracked in  
201 each plot over the growing season (May till October) using visual underwater inspection. Statistical analyses  
202 were conducted using Minitab 16 (Minitab<sup>®</sup> 16.1.1, Minitab Ltd., Coventry, UK). Data sets were normally  
203 distributed (Anderson-Darling test,  $\alpha > 0.05$ ) but failed of a test for equal variance (Levene's test,  $\alpha < 0.05$ ) even  
204 after a range of transformations (including  $x' = \log(x)$ ,  $x' = \ln(x)$ ,  $x' = \sqrt{x}$ ,  $x' = x^2$ ). Therefore, a non-  
205 parametric MWU test was used to test for significant variation in PVI between clear cut plots and controls (both  
206  $n = 5$ ) in a given month.

#### 207 **Condition assessment of Loch Flemington based on aquatic macrophyte monitoring, pre- and post-** 208 **Phoslock<sup>®</sup> application**

209 Loch Flemington was judged against the targets set for each of the attributes listed for natural lakes with  
210 *Magnopotamion* or *Hydrocharition*-type vegetation in the UK CSM Guidance for Standing Waters (JNCC  
211 2005; JNCC, 2009). To be classified as being in 'favourable condition' Loch Flemington should have an aquatic  
212 macrophyte community with species characteristic of a natural eutrophic lake and comply with the specific  
213 aquatic macrophyte community composition and structure targets for that type of standing water (JNCC 2005;  
214 JNCC, 2009).

215 In order to evaluate if there has been any change in the trophic status in Loch Flemington, following the  
216 application of Phoslock<sup>®</sup>, the whole lake macrophyte assemblages recorded in the pre-and post-application  
217 surveys were assessed using two indices: the Trophic Ranking Score (TRS) (Palmer, 1992; Palmer *et al.*, 1992)

218 and the Plant Lake Ecotype Index (PLEX) (Duigan *et al.*, 2006; Duigan *et al.*, 2007). Both scoring systems are  
219 based around assigning a score to species based on their affiliation to particular trophic conditions. Scores of  
220 qualifying species can be summed for a site, and an average score per taxon can be calculated to give a site  
221 TRS/PLEX score.

222

## 223 **RESULTS**

### 224 **Summary of key water quality findings from pre- and post-Phoslock<sup>®</sup> application monitoring**

225 The application of Phoslock<sup>®</sup> led to a significant decrease in annual mean TP concentrations to levels below the  
226 WFD target for the lake (Table 2). This led, in turn, to a significant reduction in algal levels in Loch Flemington,  
227 as indicated by decreases in annual mean chlorophyll *a* concentrations (to below the WFD target (Table 2)) and  
228 by 77% ( $P < 0.001$ ) and 95% ( $P < 0.001$ ) reductions in summer chlorophyll *a* concentrations in 2010 and 2011,  
229 respectively. There was also an increase in Secchi disc water clarity (from  $< 0.5$  m in summer 2009 to *c.* 1.4 m in  
230 summer 2011) (Spears *et al.*, 2012).

231 Monthly estimates of MMGD between 2009 and 2011 showed that the aquatic macrophyte colonisation depth  
232 had significantly increased following the application of Phoslock<sup>®</sup>. An increase in MMGD was observed in  
233 summer (57% deeper;  $W = 55$ ;  $p < 0.001$ ;  $n_1 = 10$ ,  $n_2 = 15$ ) and autumn 2010 (15% deeper;  $W = 162$ ;  $p < 0.01$ ;  
234  $n_1 = n_2 = 15$ ) when compared to 2009, indicating an overall improvement in water quality corresponding with  
235 the increase in water clarity (Figure 2). With the lowering of the seasonal and annual mean chlorophyll *a*  
236 concentrations (Table 2), following the Phoslock<sup>®</sup> application, it is estimated that the depth limits of the  
237 dominant elodeid plant functional group (i.e. *E. canadensis*) increased (depending on the background light  
238 attenuation) from *c.* 1.4 - 1.6 m in 2009 to *c.* 2.3 - 2.9 m in 2011. Linked to this increase in MMGD, the  
239 coverage of aquatic macrophytes was estimated to also have increased by between 30 and 40% to *c.* 80% of the  
240 lake bed, after the application of Phoslock<sup>®</sup>.

### 241 **Condition assessment of Loch Flemington based on aquatic macrophyte monitoring, pre- and post-** 242 **Phoslock<sup>®</sup> application**

243 The number and abundance of 'natural eutrophic' characteristic species recorded in the post-application 2010  
244 and 2011 SCM surveys marginally increased. Although three characteristic eutrophic species were recorded in

245 these surveys, occurring in 28% and 36% of sampling points, respectively (Table 3), this was well below the  
246 target of six characteristic species occurring in at least 60% of sampling points. Nevertheless, this was an  
247 improvement on the situation in 1988 and 2003 (at a frequency of occurrence of 13% of sampling points), when  
248 only one characteristic species, *Potamogeton obtusifolius* Mert. & W. D. J. Koch was recorded and two in 2009  
249 (at a frequency of occurrence of 16% of sampling points) (Table 3). Two other characteristic species,  
250 *Potamogeton perfoliatus* L. and *P. x zizii* W. D. J. Koch ex Roth were recorded on the monthly qualitative  
251 assessments in 2009 and assessed as “rare” (Table 3). Overall, the aquatic macrophyte community, as indicated  
252 by the 2010 and 2011 post-Phoslock<sup>®</sup> application surveys, remained broadly similar in terms of species  
253 composition and richness to the 2009 pre-application surveys and improved compared with the 1988 and the  
254 2003/04 surveys (Table 3). There were no obvious negative impacts of the Phoslock<sup>®</sup> application on the aquatic  
255 vegetation. However, the submerged vegetation of Loch Flemington remained dominated by the non-native *E.*  
256 *canadensis*, which occurred at a frequency well above the 25% target threshold in the SCM survey sampling  
257 points (Table 3).

258 Analysis of the monthly qualitative assessments of the aquatic macrophyte vegetation indicated that the relative  
259 abundance of *E. canadensis* and another invasive non-native *Crassula helmsii* (Kirk) Cockayne had increased  
260 following the Phoslock<sup>®</sup> application, although this was not statistically significant. *C. helmsii* was first recorded  
261 as having colonised areas of shallows and margins along the northern shore Loch Flemington in 2009, prior to  
262 the Phoslock<sup>®</sup> application. In addition, none of the three species of national conservation value *N. flexilis*, *P.*  
263 *filiformis* and *P. rutilus* were re-recorded during any of the post-application aquatic macrophyte surveys  
264 including the extensive targeted searches carried out as part of the 2010 SCM survey. Overall, on the basis of  
265 the above assessments, measured against the set targets for a naturally eutrophic lake, Loch Flemington would  
266 still be classified as being in ‘unfavourable no change’ condition, post-Phoslock<sup>®</sup> application.

267 Table 3 summarises all the species and their abundances as recorded in the various different aquatic macrophyte  
268 surveys in Loch Flemington over the period of 1988-2011 plus their associated TRS and PLEX scores. Overall,  
269 there had been a decline in both the TRS and PLEX average score per taxon values between 1988 and 2011,  
270 reflected in the re-appearance of *Apium inundatum* (L.) Rchb. f., *Chara virgata* Kütz. and *Myriophyllum*  
271 *alterniflorum* DC., species characteristic of more mesotrophic conditions (Preston & Croft, 1997; Stewart, 2004)  
272 although *M. alterniflorum* was recorded in the lake before the Phoslock<sup>®</sup> application. Pre- and post application  
273 the mesotrophic characteristic species *Potamogeton gramineus* L. has also regularly been recorded. Recordings

274 of these more mesotrophic species are consistent with the palaeoecological results which indicated the lake  
275 supported mesotrophic species in the past (Bennion *et al.*, 2008). However, Loch Flemington on the basis of its  
276 current aquatic plant community would still be classified as a Type 10 eutrophic standing water body (Palmer,  
277 1992; Palmer *et al.*, 1992) and as a species-poor example of a Group G Central and Eastern, above neutral,  
278 lowland lake with *Lemna minor*, *Elodea canadensis*, *Potamogeton natans* and *Persicaria amphibia* (Duigan *et*  
279 *al.*, 2006; Duigan *et al.*, 2007).

#### 280 **Clear-cutting experiment to mechanically control *E. canadensis***

281 The result of the replicated clear-cutting experiment on *E. canadensis* indicated that although the PVI was  
282 reduced significantly ( $W = 5$ ;  $p < 0.05$ ;  $n_1 = n_2 = 5$ ) in the cleared plots the presence of *E. canadensis* was not  
283 reduced and no desirable macrophyte species began to colonise the cleared patches (Figure 3).

284

#### 285 **DISCUSSION**

286 The application of Phoslock<sup>®</sup> to Loch Flemington led to a reduction in water column P concentrations and  
287 summer algal blooms, resulting in increased water clarity which potentially improved the light climate for  
288 aquatic macrophytes. As expected, the treatment appeared to have had no negative effect on the aquatic  
289 macrophyte community. However, although the light climate had significantly improved the only evidence of  
290 changes in the diversity of the aquatic macrophyte community were several new records of mesotrophic species  
291 (reflected in a slight decline in the average TRS/PLEX metric scores). Analysis of plant macrofossils taken  
292 from a sediment core, collected from the littoral zone of Loch Flemington in 2006, indicated a greater diversity  
293 of species characteristic of meso-eutrophic conditions in the years post-1850 than in more recent times (Bennion  
294 *et al.*, 2008). These findings suggest that a recovering Loch Flemington should exhibit greater species diversity  
295 with more species present which are indicative of meso-eutrophic lakes, including *Chara* species, *Isoetes*  
296 *lacustris* L. *N. flexilis*, *Nitella* species and *Potamogeton praelongus* Wulfen. Overall, Loch Flemington, post-  
297 Phoslock<sup>®</sup> application, remains classified as a eutrophic lake in ‘unfavourable’ condition on the basis of its  
298 failure to achieve a number of its set conservation targets. These included the continued absence of species of  
299 conservation interest, *N. flexilis*, *P. filiformis* and *P. rutilus* and the continued dominance of the aquatic  
300 vegetation by the non-native *E. canadensis* and the recent colonisation by the invasive *C. helmsii*.

301 Although in the short-term the Loch Flemington aquatic macrophyte community has yet to show the response to  
302 the Phoslock<sup>®</sup> treatment that are consistent with community shifts reported in other re-oligotrophication studies  
303 (e.g. Jeppesen *et al.*, 2005), this is perhaps not surprising. Verdonschot *et al.* (2011), in their literature review on  
304 the ecological recovery of macrophyte communities from eutrophication, indicated that full recovery of species  
305 composition was rarely recorded, often as a result of physical barriers to distribution and/or the loss of nutrient  
306 intolerant seed banks in cases where eutrophic conditions had been prevalent for many years. At a structural  
307 level, macrophyte colonisation responses are generally observed relatively quickly (i.e. less than five years) after  
308 reductions in TP load, as was the case in this study. However, at a community composition level, the recovery  
309 timescales for macrophytes to shift from eutrophic to more mesotrophic conditions were generally reported to  
310 take place on a longer time-scale from 2 to 40+ years than reported here for Loch Flemington. Verdonschot *et*  
311 *al.* (2011) highlighted a number of factors which could explain such delayed responses in shallow lakes  
312 including the following: the suppression of native aquatic macrophyte species by the invasion of more  
313 competitive species; constant disturbance of lake sediments, e.g. by wind, linked to changes in climatic  
314 conditions (Spears & Jones, 2010), leading to a turbid state predominating (e.g. Lake Apopka, USA; Havens *et*  
315 *al.*, 2001); recovery of macrophyte community being, in part, dictated by the presence of individual species seed  
316 banks, growth traits, distribution networks and pathways with species characterised by being slower growing,  
317 having higher root: shoot ratios, and being longer lived becoming increasingly prevalent as nutrient  
318 concentrations reduce (Riis & Biggs, 2011).

319 In the case of Loch Flemington the recovery of the aquatic macrophyte community may have been hampered by  
320 the presence of the invasive species *E. canadensis* and *C. helmsii*. *E. canadensis* holds a competitive advantage  
321 over many native species because it can maintain some above-sediment plant growth in the winter (Simpson,  
322 1984) enabling it to outcompete other non-evergreen species that have to regenerate totally from turions or  
323 seeds. *E. canadensis* can also utilise bicarbonate rather than carbon dioxide for photosynthesis (Bowmer *et al.*,  
324 1995; Maberly, 1983), which may give it a competitive advantage over species such as *N. flexilis*, which is  
325 reliant on carbon dioxide (Wingfield, *et al.*, 2005). *N. flexilis* is an annual plant which spreads by underwater  
326 pollination and cannot reproduce vegetatively, relying instead on seed production to survive (Wingfield *et al.*,  
327 2004). Whilst it is possible that some seeds may survive in the seed bank for more than one year, their longevity  
328 is not well understood (Wingfield *et al.*, 2004). However, although no survey of contemporary aquatic  
329 macrophyte communities have ever recorded *N. flexilis* in Loch Flemington, a recent palaeoecological study  
330 found macrofossils of the species in the lake sediments indicating that it had been present for a minimum of 100

331 years (Bennion *et al.*, 2008). This suggests that there would be a readily available seed bank for *N. flexilis*,  
332 depending on seed longevity, to extend its coverage in the lake should suitable areas of the lake bed be opened  
333 up from *E. canadensis* dominance. Like *N. flexilis*, *P. rutilus* is typically found in unpolluted mesotrophic lakes  
334 in northern Scotland (Preston, 1995) and as such it might be expected to re-establish itself naturally in Loch  
335 Flemington if there were any local sources from which the plant could re-colonise. Partly because it is rare and  
336 partly because it grows often in deeper water where it is difficult to locate (Preston & Croft, 1997), little is  
337 known about the long-term survival of *P. rutilus* turions in the sediment. However, it is apparent from the clear-  
338 cutting experiment carried out at Loch Flemington, following the Phoslock<sup>®</sup> application, that local scale  
339 mechanical control methods did not help the process of re-establishing the native aquatic macrophyte  
340 community, at least in the short term.

341 While *E. canadensis* is well established at the site and is the dominant submerged macrophyte species, *C.*  
342 *helmsii* appears to be in the early stages of colonisation having first been recorded in Loch Flemington in 2009,  
343 prior to the Phoslock<sup>®</sup> application. Therefore, the options for management of these two species are very  
344 different. In the case of *E. canadensis*, although P concentrations in Loch Flemington have significantly dropped  
345 following the Phoslock<sup>®</sup> application, they are still too high to cause a decline in the population, as the species is  
346 known to exist in a wide range of nutrient concentrations in the UK (Preston & Croft, 1997). Although  
347 mechanical control methods such as clear-cutting may be used to contain the spread of *E. canadensis* it requires  
348 implementation on a regular basis, is often labour intensive and even then success in achieving long-term  
349 control is unlikely (Howard-Williams *et al.*, 1996). Wade (1990) showed that the recovery of pre-existing  
350 submerged vegetation is often rapid after being cut, particularly by species such as *E. canadensis*, which can re-  
351 colonise and spread quickly from undifferentiated plant fragments left behind and which is even less susceptible  
352 to control by cutting compared to other competitive disturbance-tolerant species such as *Myriophyllum spicatum*  
353 (Abernethy *et al.*, 1996). As well as clear cutting a number of other measures have been proposed for the control  
354 of dense stands of *E. canadensis*, including the use of jute mats (Caffrey *et al.*, 2010), and in extreme cases,  
355 draw-down followed by the application of a herbicide (Cooke *et al.*, 2005) but as far as the authors are aware,  
356 there have been no examples of the successful complete eradication of *E. canadensis* from lakes similar to Loch  
357 Flemington. In contrast to *E. canadensis*, an opportunity to eradicate *C. helmsii* from the lake exists as it is not  
358 abundant or widespread. In 2011, colonies of *C. helmsii* were restricted to shoreline areas and had not yet  
359 reached the open water areas of the lake. Control measures of *C. helmsii*, so far, have focused on the use of  
360 shading. This has involved identification and mapping existing colonies of *C. helmsii* prior to laying plastic

361 sheeting in an attempt to shade out some of the colonies but this work is extremely time consuming and labour  
362 intensive but important, particularly as it may aid the potential re-colonisation by the Nationally Scarce *P.*  
363 *filiformis*, which typically grows in shallow open water areas such as where the *C. helmsii* populations are  
364 currently concentrated. Future efforts to manage the *C. helmsii* population, during these early stages of  
365 colonisation, are likely to focus on the continued use of plastic sheeting supplemented by the possible use of  
366 selective herbicides.

367 The sediment treatment work in Loch Flemington has, so far, been shown to be successful in reducing P  
368 concentrations and summer algal blooms, and improving macrophytes growing conditions without any  
369 noticeable negative impacts on the ecology of the aquatic macrophyte community. However, it is also clear that  
370 the biological community, as exemplified by the aquatic macrophytes, will take longer than the water chemistry  
371 to recover. This study shows that reducing P loading is not sufficient in itself to rapidly restore the aquatic  
372 macrophyte species diversity of a lake, including the desirable re-colonisation of species of conservation value,  
373 if there is a pre-existing problem with invasive, non-native species or if there is a lack of plant propagules. In the  
374 case of Loch Flemington, while *C. helmsii* can be hopefully eradicated before becoming fully established,  
375 management of *E. canadensis* will need to focus on control and containment, with the aim of facilitating the  
376 recovery of some of the former aquatic macrophyte species diversity, for which the site was renowned. It is also  
377 important to emphasise that without reducing nutrient loads from the catchment, the water quality improvements  
378 so far brought about by the application of a P-capping agent such as Phoslock<sup>®</sup>, may not persist. Therefore, it is  
379 crucial that catchment nutrient sources continue to be monitored and managed where necessary.

380

## 381 **ACKNOWLEDGEMENTS**

382

383 The authors wish to thank the following for carrying out aquatic macrophyte fieldwork at Loch Flemington: Dr  
384 Laurence Carvalho, Alex Kirika and Kathy Dale for the 2003/04 SCM surveys; to Sue Bell and Sabrina Eckert  
385 for the 2010 SCM surveys; to Mike O'Malley, Amy Anderson and Myriam Kellou for their help with fieldwork  
386 over the 2009-2011 period. We are especially grateful to Dr Mattie O'Hare for his help and advice in preparing  
387 this paper. The input of Dr Linda May and Dr Rupert Perkins into the Loch Flemington project is gratefully  
388 acknowledged. We are also grateful to Mr. John Pottie and Dr. Alastair Noble for their continued support of the

389 monitoring programme. The authors also wish to thank Nigel Traill and Said Yasseri from Phoslock® Europe  
390 GmbH, Ian Milne from the Scottish Environment Protection Agency, and Tim Dawson, Dr Mary Hennessy, Ben  
391 Leyshon and Dr Iain Sime from Scottish Natural Heritage for their valuable support of this research. We also  
392 thank two anonymous reviewers for their constructive comments which improved the paper. Phoslock® Europe  
393 GmbH was not involved in the monitoring design, sample analysis or interpretation of the data. Sebastian Meis  
394 was jointly funded by CEH and by a DAAD Scholarship (agreement number D/08/42393).

395

## 396 REFERENCES

397

398 Abernethy, V. J., M. R. Sabbatini & K. J. Murphy, 1996. Response of *Elodea canadensis* Michx and  
399 *Myriophyllum spicatum* L to shade, cutting and competition in experimental culture. *Hydrobiologia* 340: 219-  
400 224.

401

402 Bennett, E. M., S. R. Carpenter & N. F. Caraco, 2001. Human impact on erodible phosphorus and  
403 eutrophication: a global perspective. *Bioscience* 51: 227-234.

404

405 Bennion, H., H. Clarke, T. Davidson, D. Morley, N. Rose, S. Turner & H. Yang, 2008. Palaeoecological study  
406 of seven mesotrophic lochs. Report to the Scottish Environment Protection Agency and Scottish Natural  
407 Heritage. Research Report Number: 121. ISSN: 1366-7300.

408

409 Bowmer, K. H., S. W. L. Jacobs & G. R. Sainty, 1995. Identification, biology and management of *Elodea*  
410 *canadensis*, Hydrocharitaceae. *Journal of Aquatic Plant Management* 17: 4-12.

411

412 Caffrey, J. M., M. Millane, S. Evers, H. Moran & M. Butler, 2010. A novel approach to aquatic weed control  
413 and habitat restoration using biodegradable jute matting. *Aquatic Invasions* 5: 123-129.

414

415 Cooke, G. D., E. B. Welch, S. A. Peterson & S. A. Nichols, 2005. Restoration and Management of Lakes and  
416 Reservoirs, 3<sup>rd</sup> ed. CRC Press, Taylor & Francis Group, Boca Raton, Florida.

417

418 Council of the European Communities, 1992. Council Directive 92/43/EEC of 21 May 1992 on the  
419 conservation of natural habitats and of wild fauna and flora. Official Journal of the European Communities  
420 L206.

421

422 Council of the European Communities, 2000. Directive 2000/60/EC of the European Parliament and of the  
423 Council of 23 October 2000 establishing a framework for Community action in the field of water policy.  
424 Official Journal of the European Communities L327: 1-73.

425

426 Duigan, C., W. Kovach & M. Palmer, 2006. Vegetation Communities of British Lakes: a revised classification.  
427 Joint Nature Conservation Committee, Peterborough.

428

429 Duigan, C., W. Kovach & M. Palmer, 2007. Vegetation Communities of British Lakes: a revised classification  
430 for conservation. Aquatic Conservation: Marine and Freshwater Ecosystems 17:147-173.

431

432 EEA, 2012. European waters – assessment of status and pressures. EEA report No 8/2012. European  
433 Environment Agency, Copenhagen, Denmark.

434

435 Fisher, J., T. Barker, C. James & S. Clarke, 2009. Water quality in chronically nutrient-rich lakes: the example  
436 of the Shropshire-Cheshire Meres. Freshwater Reviews 2: 79-99.

437

438 Foy, R. H., 1985. Phosphorus inactivation in a eutrophic lake by the direct addition of ferric aluminium  
439 sulphate: impact of iron and phosphorus. *Freshwater Biology* 15: 613-629.

440

441 Gordon, J. E. & C. A. Auton, 1993. The Kildrummie Kames. In J. E. Gordon & D. G. Sutherland (eds),  
442 Quaternary of Scotland. Chapman and Hall, London: 176-181.

443

444 Gunn, I. D. M., M. O'Hare, L. Carvalho, D. B. Roy, P. Rothery & A. M. Darwell, 2010. Assessing the condition  
445 of lake habitats: a test of methods for surveying aquatic macrophyte communities. *Hydrobiologia* 656: 87-97.

446

447 Havens, K. E., K. R. Jin, A. J. Rodusky, B. Sharfstein, M. A. Brady, T. L. East, N. Iricanin, R. T. James, M. C.  
448 Harwell & A. D. Steinman, 2001. Hurricane effects on a shallow lake ecosystem and its response to a controlled  
449 manipulation of water level. *Scientific World Journal* 1: 44-70.

450

451 Howard-Williams, C., A. M., Schwarz & V. Reid, 1996. Patterns of aquatic weed regrowth following  
452 mechanical harvesting in New Zealand hydro-lakes. *Hydrobiologia* 340: 229-234.

453

454 Jeppesen, E., M. Søndergaard, J. P. Jensen & T. L. Lauridsen, 2003. Restoration of eutrophic lakes: A global  
455 perspective. In Kumagai, M. & W. F. Vincent (eds), *Freshwater Management: Global Versus Local*  
456 *Perspectives*. Springer-Verlag, Tokyo:135-151.

457

458 Jeppesen, E., M. Søndergaard, J. P. Jensen, K. E. Havens, O. Anneville, L. Carvalho, M. F. Coveney, R.  
459 Deneke, M. T. Dokulil, B. Foy, D. Gerdeaux, S. E. Hampton, S. Hilt, K. Kangur, J. Kohler, E. Lammens, T. L.  
460 Lauridsen, M. Manca, M. R. Miracle, B. Moss, P. Noges, G. Persson, G. Phillips, R. Portielje, C. L. Schelske,  
461 D. Straile, I. Tatrai, E. Willen & M. Winder, 2005. Lake responses to reduced nutrient loading - an analysis of  
462 contemporary long-term data from 35 case studies. *Freshwater Biology* 50: 1747-1771.

463

464 JNCC, 2005. Common Standards Monitoring Guidance for Standing Waters. Joint Nature Conservation  
465 Committee, Peterborough.

466

467 JNCC, 2009. Common Standards Monitoring Guidance for Standing Waters. Revised 10 November 2009, Joint  
468 Nature Conservation Committee, Peterborough.

469

470 Jupp, B. P., D. H. N. Spence & R. H. Britton, 1974. The distribution and production of submerged macrophytes  
471 in Loch Leven, Kinross. Proceedings of the Royal Society of Edinburgh – B 74: 195-208.

472

473 Jupp, B. P. & D. H. N. Spence, 1977. Limitations on macrophytes in a eutrophic lake, Loch Leven. 1. Effects of  
474 phytoplankton. Journal of Ecology 65: 175-186.

475

476 Maberly, S. C., 1983. The interdependence of photon irradiance and free carbon-dioxide or bicarbonate  
477 concentration on the photosynthetic compensation points of fresh-water plants. New Phytologist 93: 1-12.

478

479 May, L. & L. Carvalho, 2010. Maximum growing depth of macrophytes in Loch Leven, Scotland, United  
480 Kingdom, in relation to historical changes in estimated phosphorus loading. Hydrobiologia 646: 123-131.

481

482 May, L., I. Gunn & A. Kirika, 2001. Phosphorus study at Loch Flemington. Unpublished report to Scottish  
483 Natural Heritage. Centre for Ecology & Hydrology.

484

485 May, L., B. M. Spears, B. J. Dudley & T. W. Hatton-Ellis, 2010. The importance of nitrogen limitation in the  
486 restoration of Llangorse Lake, Wales, UK. Journal of Environmental Monitoring 12: 338-346.

487

488 Meis, S., B. M. Spears, S. C. Maberly, M. B. O'Malley & R. G. Perkins, 2012. Sediment amendment with  
489 Phoslock<sup>®</sup> in Clatto Reservoir (Dundee, UK): Investigating changes in sediment elemental composition and  
490 phosphorus fractionation. *Journal of Environmental Management* 93: 185-193.

491

492 NCC, 1988. Loch Flemington: Nature Conservancy Council Loch Survey aquatic macrophyte field data  
493 (19/07/1988). Joint Nature Conservation Committee, Peterborough.

494

495 Palmer, M. A., 1992. A botanical classification of standing waters in Great Britain and a method for the use of  
496 macrophyte flora in assessing changes in water quality incorporating a reworking of data 1992. *Research and  
497 Survey in Nature Conservation*, No. 19. Joint Nature Conservation Committee, Peterborough.

498

499 Palmer, M. A., S. L. Bell & I. Butterfield, 1992. A botanical classification of standing waters in Britain:  
500 applications for conservation and monitoring. *Aquatic Conservation: Marine and Freshwater Ecosystems* 2: 125-  
501 143.

502

503 Preston, C. D., 1995. *Pondweeds of Great Britain and Ireland*, BSBI Handbook No. 8. Botanical Society of the  
504 British Isles. London.

505

506 Preston, C. D. & J. M. Croft, 1997. *Aquatic Plants in Britain and Ireland*. Harley Books, Colchester.

507

508 Randall, S., Harper, D. & B. Brierley, 1999. Ecological and ecophysiological impacts of ferric dosing in  
509 reservoirs. *Hydrobiologia* 395/396: 355-364.

510

511 Riis, R. & B. J. F. Biggs, 2001. Distribution of macrophytes in New Zealand streams and lakes in relation to  
512 disturbance frequency and resource supply - a synthesis and conceptual model. New Zealand Journal of Marine  
513 and Freshwater Research 35: 255-267.

514

515 Ross, G., F. Haghseresht & T. E. Cloete, 2008. The effect of pH and anoxia on the performance of Phoslock<sup>(R)</sup>,  
516 a phosphorus binding clay. Harmful Algae 7: 545-550.

517

518 Sas, H., 1989. Lake restoration by reduction of nutrient loading: expectations, experiences, Extrapolations. Acad  
519 Verlag Richarz.

520

521 Scheffer, M., 2004. Ecology of shallow lakes. Kluwer Academic Publishers, Dordrecht, The Netherlands

522

523 Schindler, D. W., 2006. Recent advances in the understanding and management of eutrophication. Limnology  
524 and Oceanography 51: 356-363.

525

526 Simpson, D. A., 1984. A short history of the introduction and spread of *Elodea* Michx. in the British Isles.  
527 *Watsonia* 15: 1-9.

528

529 Søndergaard, M., E. Jeppesen, J. P. Jensen & S. L. Amsinck, 2005. Water Framework Directive: Ecological  
530 classification of Danish lakes. *Journal of Applied Ecology* 42: 616-629

531

532 Søndergaard, M., E. Jeppesen, T. L. Lauridsen, C. Skov, E. H. Van Nes, R. Roijackers, E. Lammens & R.  
533 Portielje, 2007. Lake restoration: successes, failures and long-term effects. *Journal of Applied Ecology* 44:  
534 1095-1105.

535

536 Spears, B. M., I. D. M. Gunn, L. Carvalho, I. J. Winfield, B. Dudley, K. Murphy & L. May, 2009. An evaluation  
537 of methods for sampling macrophyte maximum colonisation depth in Loch Leven, Scotland. *Aquatic Botany* 91:  
538 75–81.

539

540 Spears, B. M. & I. Jones, 2010. The long-term (1979-2005) effects of the north Atlantic oscillation on wind  
541 induced wave mixing in Loch Leven (Scotland). *Hydrobiologia* 646: 49-59.

542

543 Spears, B. M., S. Meis, S. C. Maberly, I. D. M. Gunn & L. May, 2012. Assessing recovery from eutrophication in  
544 Loch Flemington following the control of sediment phosphorus release using Phoslock<sup>®</sup>. Unpublished report to  
545 Scottish Natural Heritage. Centre for Ecology & Hydrology.

546

547 Spears, B.M., B. Dudley, K. Reitzel & E. Rydin, 2013a. Geo-engineering in lakes – a call for consensus.  
548 *Environmental Science & Technology*, 47: 3953-3954.

549

550 Spears, B. M., S. Meis, A. Anderson & M. Kellou, 2013b. Comparison of phosphorus (P) removal properties of  
551 materials proposed for the control of sediment P release in UK lakes. *Science of the Total Environment* 442:  
552 103-110.

553

554 Stewart, N. F., 2004. Important Stonewort Areas. An assessment of the best areas for stoneworts in the United  
555 Kingdom. Plantlife International, Salisbury, UK.

556

557 Trenberth, K. W., 1983. What are the seasons? *Bulletin of the American Meteorological Society* 64: 1276-1277.

558

559 UKLakes Database: <http://www.UKLakes.net/>

560

561 UK TAG, 2008. UK Environmental Standards and Conditions (Phase 2). UK Technical Advisory Group on the  
562 Water Framework Directive.

563

564 Verdonschot, P., D. Angler, À. Boria, S. Brucet, C. K. Field, I. Gunn, R. Johnson, J. Kail, M. Kernan, A.  
565 Marzin, L. May, S. Meis & B. Spears, 2011. WISER Deliverable D6.4-2: Report on the differences between  
566 cause-effect-recovery chains of different drivers within the water categories. European Commission.

567

568 Wade, P. M., 1990. Physical control of aquatic weeds. In Pieterse, A. H. & K. J. Murphy (eds), Aquatic Weeds  
569 The Ecology and Management of Nuisance Aquatic Vegetation. Oxford Science Publications: 93-135.

570

571 Wingfield, R. A., K. Murphy, P. Hollingworth & M. Gaywood, 2004. The Ecology of *Najas flexilis*. Scottish  
572 Natural Heritage Commissioned Report No. 017. Scottish Natural Heritage.


573

574 Wingfield, R., K. Murphy & M. Gaywood, 2005. Lake habitat suitability for the rare European macrophyte  
575 *Najas flexilis* (Willd.) Rostk & Schmidt. Aquatic Conservation: Marine and Freshwater Ecosystems 15: 227-  
576 241.

577

578 Withers, P. J. A. & P. M. Haygarth, 2007. Agriculture, phosphorus and eutrophication: a European perspective.  
579 Soil Use and Management 23: 1-4.

580




582

583

584

585


586 **Fig. 1.** Bathymetric Map of Loch Flemington



587

588 **Fig. 2.** Seasonal variation in macrophyte maximum growing depth (MMGD) in Loch Flemington. Dashed  
 589 vertical line indicates timing of the Phoslock® application, error bars represent standard error of the mean (n =  
 590 variable) and seasons in which no sampling occurred are marked with 'X'. Significant differences (\*\*, p < 0.01;  
 591 \*\*\*, p < 0.001) in a season between pre- and post-application years and percent change are indicated

592



593

594

595 **Fig. 3.** Seasonal variation in percentage volume inhabited (PVI) of *Elodea canadensis* following clear-cutting in  
 596 five 4 m<sup>2</sup> plots compared with five uncut 4 m<sup>2</sup> plots in Loch Flemington in 2010. Error bars represent the  
 597 standard error of the mean.

598

599 **Table 1.** Summary of SCM aquatic macrophyte surveys carried out at Loch Flemington, pre- and post-  
 600 Phoslock<sup>®</sup> application.

	SCM surveys pre-Phoslock <sup>®</sup> application		SCM surveys post-Phoslock <sup>®</sup> application	
	3 July 2003 + 5 August 2004	12 August 2009	26/27 July + 2/3 August 2010	18/19 <sup>th</sup> July 2011
Number of perimeter strandline searches	4	4	5	4
Number of shore-wader depth transects	4	4	3	3
Number of boat-based depth transects	-	1	2	2
Other	Targeted search for rare species in open-water	-	Targeted search for rare species in open water	-

601

602 **Table 2.** Summary table of annual mean surface water TP and chlorophyll *a* concentrations, pre - and post-  
603 Phoslock<sup>®</sup> application

<b>Year</b>	<b>Mean TP concentrations (<math>\mu\text{g P l}^{-1}</math>) (n = variable)</b>	<b>Mean chlorophyll <i>a</i> concentrations (<math>\mu\text{g l}^{-1}</math>) (n = variable)</b>
WFD target	32	16
2009 (Pre-application )	60	51
2010 (Post-application)	31	25
2011 (post-application)	27	12

604

605

606

607

608

609

610

611

612

613

614

615

616

617

618

619

620

621

622

623

624

625

626 **Table 3.** Summary of aquatic macrophyte species occurrence, species abundance (% of SCM survey sampling  
627 points), Trophic Ranking Scores (TRS) and Plant Lake Ecotype Complex (PLEX) scores for Loch Flemington  
628 from 1988 - 2011.

Species	Pre - Phoslock® application					Post – Phoslock® application	
	TRS	PLEX	1988 <sup>a</sup>	2003/04	2009 <sup>c</sup>	2010 <sup>c</sup>	2011 <sup>c</sup>
<i>Apium inundatum</i>	7.00	7.50	-	-	-	1%	present
<i>Chara virgata</i> <sup>d</sup>	7.30	7.69	-	-	-	present	-
<i>Crassula helmsii</i>	X	X	-	-	present	present	present
<i>Elodea canadensis</i>	8.50	7.95	LD	56%	62%	52%	51%
<i>Lemna minor</i> <sup>d</sup>	9.00	8.85	-	-	3%	5%	28%
<i>Littorella uniflora</i>	6.70	4.23	O	6%	3%	3%	present
<i>Myriophyllum alterniflorum</i>	5.50	4.23	-	-	14%	9%	7%
<i>Myriophyllum spicatum</i>	10.00	8.85	-	1%	1%	present	-
<i>Persicaria amphibia</i>	9.00	7.95	LD	28%	33%	16%	22%
<i>Potamogeton gramineus</i>	7.30	7.31	-	present	3%	present	1%
<i>Potamogeton natans</i>	6.70	4.23	-	7%	17%	8%	6%
<i>Potamogeton obtusifolius</i> <sup>d</sup>	7.30	6.54	O-F	13%	13%	27%	9%
<i>Potamogeton perfoliatus</i> <sup>d</sup>	7.30	7.69	-	-	present	-	present
<i>Potamogeton x zizii</i> <sup>d</sup>	X <sup>b</sup>	7.69	-	-	present	-	-
<i>Ranunculus aquatilis</i>	8.50	7.95	R	-	-	-	-
<b>Total number of species</b>			5	7	12	12	11
<b>Mean number of species per SCM survey transect (wader &amp; boat)</b>			N/A	3.75	4.00	4.17	4.20
<b>Total number of characteristic<sup>d</sup> species</b>			1	1	4	3	3
<b>% frequency of sampling points with characteristic<sup>d</sup> species</b>			N/A	13	16	28	36
<b>Total TRS</b>			40	55.50	77.3	84.3	74.3
<b>Total PLEX</b>			34.62	47.07	75.52	75.33	66.48
<b>Mean Score Per Taxon - TRS</b>			8.00	7.93	7.73	7.66	7.43
<b>Mean Score Per Taxon - PLEX</b>			6.92	6.72	6.87	6.85	6.65

629

630 <sup>a</sup> in 1988 DAFOR (Dominance, Abundant, Frequent, Occasional, Rare) abundance scale used; LD = Locally  
631 Dominant

632 <sup>b</sup> X indicates no scores allocated to species

633 <sup>c</sup> species list includes data from monthly qualitative assessments in addition to SCM survey data – only SCM  
634 survey data used for abundances

635 <sup>d</sup> characteristic species of natural eutrophic lakes with *Magnopotamion* or *Hydrocharition*-type vegetation  
636 (JNCC 2005; JNCC, 2009)

637

638