

British **Geological Survey**

Friday 10th January 2014 RAS Meeting: 'Geomagnetic field dynamics and structure on timescales from minutes to decades'

NATURAL ENVIRONMENT RESEARCH COUNCIL

THE ROLE OF GEOMAGNETIC OBSERVATORY DATA DURING THE SWARM MISSION

VICTORIA RIDLEY (vicley@bgs.ac.uk), SUSAN MACMILLAN (smac@bgs.ac.uk) and BRIAN HAMILTON British Geological Survey, West Mains Road, Edinburgh EH9 3LA, United Kingdom

SUMMARY	2. QUALITY OF OBSERVATORY DATA
The ESA Swarm mission will measure magnetic signals from all sources of the geomagnetic field with unprecedented accuracy. The scientific use of Swarm data is greatly enhanced when used in combination with observatory data and indices and this has increased interest in ground based measurements. As part of the Swarm Level-2 data activities, plans are in place to distribute such data along with the satellite data [1]. Here, we also discuss how observatory data can be used for the Calibration/Validation of Swarm and how observatory data may better constrain the time parameterisation of global field models.	 To aid quality control of global observatory data prior to joint analyses with Swarm data, misfits of spherical harmonic models can be inspected in the temporal & spatial domains [1]. This pre-processing and modelling removes all signals that can be modelled, except at high latitudes, and the misfits represent measurement artefacts on the 0-10 nT scale. Tests using definitive hourly mean data contemporary with the Ørsted and CHAMP missions showed that the method was useful for detecting these small artefacts. The plot shows a subset of the misfits: in the geomagnetic south component for observatories between geomagnetic latitudes 58° (top) and 46° (bottom). The data used were a combination of definitive and quasi-definitive hourly mean data 2009-2013 spanning the gap between CHAMP and the launch of Swarm. High quality data is reflected by values ³⁰/₄₀

1. TIMELINESS OF OBSERVATORY DATA

- Every 3 months during the whole of the Swarm mission we aim to update files of observatory hourly mean values.
- During the first 6 months of the mission we aim to update files of observatory minute mean values to provide independent data for ground-truthing purposes.
- Advantage is taken of INTERMAGNET and other efforts in Norway, France & UK to improve the timeliness of quasi-definitive (QD) observatory data.
- QD data aims to be within 5 nT of definitive data (averaging on a monthly basis) and available within 3 months of measurement. The map shows the observatories producing such data as of December 2013.



- being:
- a) Close to zero (more so the lower the latitude),
- b) No discernible discontinuities present
- c) Coherent with geomagnetic latitude if large (indicating a natural storm signal).
- To achieve this, poor quality data are excluded and time series are split to account for unmeasured jumps
- In addition to being available from the ESA data centre once the Swarm Level 2 processing starts, the cleaned-up hourly mean observatory data will also be regularly updated at ftp://ftp.nmh.ac.uk/geomag/smac/AU X OBS 2/



2010

2005

2005

Time (vear

2005 2010

Time (years)

2005

Time (vears)

0.65yr

Time (years)

3. GROUND-TRUTHING SWARM DATA FOR CAL/VAL

- The Calibration/Validation (Cal/Val) period, forming the first 3 months of the Swarm mission, will be used to confirm the instruments are operating as expected.
- Can Swarm measurements be ground-truthed with QD observatory data to aid the Cal/Val effort?
- The following method will be applied to each satellite during Cal/Val. We will look at how the global results a) vary **between satellites**
 - b) compare with the results obtained when a similar approach has been applied to CHAMP data.

0

3a. Approach

3b. Number of passes

4. OBSERVATORY DATA & MODEL TIME PARAMETERISATION

- Global geomagnetic field models typically fit temporal changes to the Earth's core field using order 6 B-splines Spline knot spacings and regularisation varies: CHAOS-3 [2] & GRIMM-2 [3] use 0.5 year spacing and smooth the 2nd/3rd time derivative; C³FM2 [4] use 1.2 year knot spacing and smooth by minimising changes to core surface flow
- Using different knot spacings we fit order 6 B-splines to data from 53 observatories in an attempt to ascertain the optimal knot-spacings and identify whether a minimum spacing is required to fit the data well
- Prior to fitting splines, the data are detrended to remove annual and semi-annual signals

- LER0

MEA0

SIT2

UPS0

NUR0

STJ0

VAL0

BFE0

OTT0

NEW0

SHU0

VIC0

HAD0

SBL0

BOX0

HLP0

YAK0

MAB0

2008





$$\left|\Delta \sigma_{n}\right| = \left(\frac{B_{sat} - \overline{X}_{sat}}{\sigma_{sat}}\right) - \left(\frac{B_{obs} - \overline{X}_{obs}}{\sigma_{obs}}\right) - \frac{1}{2} \left(\frac{B_{obs} - \overline{X$$

- Preliminary study carried out for CHAMP and definitive minute-mean data from >100 observatories between 2001-2011
- Example here shows ESK
- $|\Delta \sigma_{\rm N}|$ generally <0.5
- Seen more clearly when comparing individual passes (lower plots)

Good correlation between B_{obs} and B_{sat} for CHAMP at: all local times; different disturbance levels and all latitudes. Similar findings expected for Swarm, but difficult to predict how QD data might compare with the minute means used here for CHAMP



ESK $ \Delta \sigma_N $	<0.5	<1.0	<2.0
% of 2001-2011 passes	94.8	98.9	99.6

Sum of squared spinle in an $X 10_5$	x 10 ⁴	x 10 ^{4t}	\times 10 ⁵ x 10 ⁵	Sum of squared spinle misin $x 10^4$	Sum of squared spiller mit
--------------------------------------	-------------------	--------------------	--	--------------------------------------	----------------------------

Examples for three observatories are shown here

- As expected: with low knot spacing the fit to the data is improved, but the spline dB/dt is unrealistically complex; with higher knot spacing, the fit to the data is worse and dB/dt is much smoother
- The trade-off curve knees signify where further increasing the knot-spacing does not greatly alter the amount of structure in dB/dt, but does increase the data misfit
- Before the knee, a significant increase in spline complexity is seen for lower knot spacings, signifying rapid, unrealistic rates of change being modelled
- There are well defined trade-off curve knees for many observatories (e.g. PPT) but also some more ambiguous results (e.g. ESA); results also vary between Bx, By and Bz
- In some cases, just half a year difference in knot spacing significantly changes the fit and complexity (e.g. VIC)

Though results vary between observatories (reflecting different secular variation rates by location), knot spacings of <1.0 year are preferred, even without spline smoothing. For most observatories, spacings of at least 0.55-0.70 years are required to adequately fit the field variations.

REFERENCES

[1] Macmillan, S.; Olsen, N.. 2013 Observatory data and the Swarm mission [in special issue: Swarm] Earth Planets Space, 65. 1355-1362. 10.5047/eps.2013.07.011 [2] Lesur, V., Wardinski, I., Hamoudi, M. & Rother, M., 2010. The second generation of the GFZ reference internal magnetic model: GRIMM-2, Earth Planets Space, 62, 765–773 [3] Olsen, N., Mandea, M., Sabaka, T.J. & Tøffner-Clausen, L., 2010. The CHAOS-3 geomagnetic field model and candidates for the 11th generation IGRF, Earth Planets Space, **62**, 719–727.

[4] Lesur, V., Wardinski, I., Asari, S., Minchev, B. & Mandea, M., 2010. Modelling the Earth's core magnetic field under flow constraints, *Earth Planets Space*, **62**, 503–516