

1989

NSORURE

CEH Lancaster  
Archive Copy  
Please do NOT remove  
Not for loan

SOIL SEED BANKS AND WEED GERMINATION:  
A SELECTIVE BIBLIOGRAPHY

Trevor R Harwood

Land Use Research Group  
Institute of Terrestrial Ecology  
Merlewood Research Station  
Grange-over-Sands  
Cumbria LA11 6JU

December 1989

SOIL SEED BANKS AND WEED GERMINATION:  
A SELECTIVE BIBLIOGRAPHY

Trevor R Harwood

Land Use Research Group  
Institute of Terrestrial Ecology  
Merlewood Research Station  
Grange-over-Sands  
Cumbria LA11 6JU

December 1989

## INTRODUCTION

This work does not claim to be complete: the majority of entries are of recent and "classical" works on British agricultural and semi-natural habitats.

Additional references include frequently-quoted examples from North America and elsewhere; plus those on habitat types poorly covered in UK literature, such as marshes and other waterlogged sites.

Papers on the seed behaviour of single species or groups of similar species have generally been omitted: these and many further references may be found in the larger references herein, such as Grime, Hodgson and Hunt (1988). Also not included are references to seed behaviour in arid areas such as those of Australia and North America.

- AMEN, R.D. 1968. A model of seed dormancy. Botanical Review, 34, 1-31.
- ARCHIBOLD, O.W. 1981. Buried viable propagules in native prairie and adjacent agricultural sites in central Saskatchewan. Canadian Journal of Botany, 59, 701-706.
- ARTIGAS, F.J. & BOERNER, R.E.J. 1989. Advance regeneration and seed banking of woody plants in Ohio pine plantations: implications for landscape change. Landscape Ecology, 2, 139-150.
- AYERS, A.D. 1952. Seed germination as affected by moisture and soil salinity. Agronomy Journal, 44, 82-84.
- BAKER, H.G. 1972. Seed weight in relation to environmental conditions in California. Ecology, 53, 997-1010.
- BAKKER, J.P. 1983. Seedbank and species diversity in moist grassland communities. Acta Botanica Neerlandica, 32, 246.
- BAKKER, J.P. 1985. The impact of grazing on plant communities, plant populations and soil conditions on salt marshes. Vegetatio, 62, 391-398.
- BASKIN, J.M. & BASKIN, C.C. 1985. The annual dormancy cycle in buried weed seeds: a continuum. Bioscience, 35, 492-498.
- BERTNESS, M.D., WISE, C. & ELLISON, A.M. 1987. Consumer pressure and seed set in a salt marsh perennial plant community. Oecologia (Berlin), 71, 190-200.
- BOORMAN, L.A. 1968. Some aspects of the reproductive biology of *Limonium vulgare* Mill., and *Limonium humile* Mill. Annals of Botany, 32, 803-824.
- BOSTOCK, S.J. 1978. Seed germination strategies of five perennial weeds. Oecologia (Berlin) 36, 113-126.
- BRENCHLEY, W.E. 1918. Buried weed seeds. Journal of Agricultural Science, 9, 1-31.
- BRENCHLEY, W.E. & WARINGTON, K. 1930. The weed seed population of arable soil. I. Numerical estimation of viable seeds and observations on their natural dormancy. Journal of Ecology, 18, 235-272.
- BRENCHLEY, W.E. & WARINGTON, K. 1933. The weed seed population of an arable soil. II. Influence of crop, soil and methods of cultivation upon the relative abundance of viable seeds. Journal of Ecology, 21, 103-127.
- BRENCHLEY, W.E. & WARINGTON, K. 1936. The weed seed population of an arable soil. III. The re-establishment of weed species after reduction by fallowing. Journal of Ecology, 24, 479-501.
- BRENCHLEY, W.E. & WARRINGTON, K. 1945. The influence of periodic fallowing on the prevalence of viable weed seeds in arable soils. Annals of Applied Biology, 32, 285-296.
- BROWN, A.H.F. 1979. Management effects in lowland coppice woods. Viable seed content of soils of conifer-planted areas in ancient coppice woods. Natural Environment Research Council contract report to the Nature Conservancy Council. Grange-over-Sands: Institute of Terrestrial Ecology.

- BROWN, A.H.F. 1981. The recovery of ground vegetation in coppice wood: the significance of buried seed. In: Forest and Woodland Ecology, edited by F T Last & A S Gardiner, 41-44. Cambridge: Institute of Terrestrial Ecology.
- BROWN, A.H.F. & OOSTERHUIS, L. 1981. The role of buried seed in coppice woods. Biological Conservation, 21, 19-38.
- CAVERS, P.B. 1983. Seed demography. Canadian Journal of Botany, 61, 3578-3590.
- CHAMPNESS, S.S. 1949. Note on the technique of sampling soil to determine the content of buried viable seeds. Journal of the British Grassland Society, 4, 115-118. X
- CHAMPNESS, S.S. & MORRIS, K. 1948. The population of buried viable seeds in relation to contrasting pasture and soil types. Journal of Ecology, 36, 149-173. X
- CHANCELLOR, R.J. 1964. Emergence of weed seedlings in the field and the effects of different frequencies of cultivation. Proceedings of the Seventh British Weed Control Conference, 2, 599-606.
- CHANCELLOR, R.J. 1966. The identification of weed seedlings of farm and garden. Oxford: Blackwell Scientific.
- CHANCELLOR, R.J. 1978. Grass seeds beneath pastures. In: Changes in sward composition and productivity, Occasional Symposium of the British Grassland Society, 10, 147-150. X
- CHANCELLOR, R.J. 1984. Dormancy and germination of grass seeds. Proceedings, 7th International Symposium on Weed Biology, Ecology and Systematics, Versailles, 21-27.
- CHANCELLOR, R.J. 1985. Changes in the weed flora of an arable field cultivated for 20 years. Journal of Applied Ecology, 22, 491-501. v
- CHANCELLOR, R.J. 1986. Decline of arable weed seeds during 20 years in soil under grass and the periodicity of seedling emergence after cultivation. Journal of Applied Ecology, 23, 631-637. v
- CHANCELLOR, R.J. & FROUD-WILLIAMS, R.J. 1983. Weeds of cereals in central southern England. Weed Research Organisation Tenth Report, 1982-3, 27-32.
- CHANCELLOR, R.J. & FROUD-WILLIAMS, R.J. 1984. A second survey of cereal weeds in central southern England. Weed Research, 24, 29-36.
- CHANCELLOR, R.J., FRYER, J.D. & CUSSANS, G.W. 1984. The effects of agricultural practices on weeds in arable land. In: Agriculture and the Environment, edited by D. Jenkins, 89-94. Cambridge: Institute of Terrestrial Ecology.
- CHIPPINDALE, H.G. & MILTON, W.E.J. 1934. On the viable seeds present in the soil beneath pastures. Journal of Ecology, 22, 508-531. v
- CRESWELL, E. & GRIME, J.P. 1981. Induction of a light requirement during seed development and its ecological consequences. Nature, 291, 583-585.

- CROCKER, W. & BARTON, L.V. 1953. Physiology of seeds. Waltham: Chronica Botanica.
- DEANE, R.J.L. 1989. Expanded field margins - their costs to the farmer and benefits to wildlife; a report to the Nature Conservancy Council. Tewkesbury: Kemerton Court.
- DICKIE, J.B., GAJJAR, K.H., BIRCH, P. & HARRIS, J.A. 1988. The survival of viable seeds in stored topsoil from opencast coal workings and its implications for site restoration. Biological Conservation, 43, 257-265.
- DONELAN, M. & THOMPSON, K. 1980. Distribution of buried viable seeds along a successional series. Biological Conservation, 17, 297-311.
- DOUGLAS, G. 1965. The weed flora of chemically-renewed lowland swards. Journal of the British Grassland Society, 20, 91-100.
- DURING, H.J. & TER HORST, B. 1983. The diaspore bank of bryophytes and ferns in chalk grassland. Lindbergia, 9, 57-64.
- FEAST, P.M. & ROBERTS, H.A. 1973. Note on the estimation of viable weeds in soil samples. Weed Research, 13, 110-113.
- FENNER, M. 1978a. A comparison of the abilities of colonizers and closed-turf species to establish from seed in artificial swards. Journal of Ecology, 66, 953-963.
- FENNER, M. 1978b. Susceptibility to shade in seedlings of colonising and closed turf species. New Phytologist, 81, 739-744.
- FENNER, M. 1985. Seed ecology. London: Chapman & Hall.
- FOWLER, N.L. 1988. The effects of environmental heterogeneity in space and time on the regulation of populations and communities. In: Plant population ecology, edited by M.J. Hutchings, A.J. Davy & A.R. Watkinson, 249-269. Oxford: Blackwell Scientific.
- FOX, J.F. 1983. Germinable seed banks of interior Alaskan tundra. Arctic and Alpine Research, 15, 405-411.
- FREEDMAN, B., HILL, N., SVOBODA, J. & HENRY, G. 1982. Seed banks and seedling occurrence in a high Arctic oasis at Alexandra Fjord, Ellesmere Island, Canada. Canadian Journal of Botany, 60, 2112-2118.
- FROUD-WILLIAMS, R.J. & CHANCELLOR, R.J. 1982. A survey of grass weeds in cereals in central southern England. Weed Research, 22, 163-171.
- FROUD-WILLIAMS, R.J., DRENNAN, D.S.H. & CHANCELLOR, R.J. 1983a. Influence of cultivation regime on weed floras of arable cropping systems. Journal of Applied Ecology, 20, 187-197.
- FROUD-WILLIAMS, R.J., CHANCELLOR, R.J. & DRENNAN, D.S.H. 1983b. Influence of cultivation regime on buried weed seeds in arable cropping systems. Journal of Applied Ecology, 20, 199-298.
- FROUD-WILLIAMS, R.J., CHANCELLOR, R.J. & DRENNAN, D.S.H. 1984. The effects of seed burial and soil disturbance on emergence and survival of arable weeds in relation to minimal cultivation. Journal of Applied Ecology, 21, 629-641.

- FRYER, J.D. & CHANCELLOR, R.J. 1970. Herbicides and our changing weeds. In: The Flora of a Changing Britain, edited by F. Perring, 105-118. Farringdon: Classey.
- GARTNER, B.L., CHAPIN, F.S., III & SHAVER, G.R. 1983. Demographic patterns of seedling establishment and growth of native graminoids an an Alaskan tundra disturbance. Journal of Applied Ecology, 20, 965-980.
- GASHWILLER, J.W. 1967. Conifer seed survival in a western Oregon clear cut. Ecology, 48, 431-438.
- GRAHAM, D.J. & HUTCHINGS, M.J. 1988a. Estimation of the seed bank of a chalk grassland ley established on former arable land. Journal of Applied Ecology, 25, 241-252.
- GRAHAM, D.J. & HUTCHINGS, M.J. 1988b. A field investigation of germination from the seed bank of a chalk grassland ley on former arable land. Journal of Applied Ecology, 25, 253-263.
- GRANSTROM, A. 1982. Seed banks in five boreal forest stands originating between 1810 and 1963. Canadian Journal of Botany, 60, 1815-1821.
- GRANSTROM, A. 1986. Seed banks in forest soils and their role in vegetation succession after disturbance. Umea: Swedish University of Agricultural Sciences.
- GRANSTROM, A. 1987. Seed viability of fourteen species during five years of storage in a forest soil. Journal of Ecology, 75, 321-331.
- GRANSTROM, A. 1988. Seed banks at six open and afforested heathland sites in southern Sweden. Journal of Applied Ecology, 25, 297-306.
- GRIME, J.P. 1966. Shade avoidance and shade tolerance in flowering plants. In: Light as an ecological factor, edited by R. Bainbridge, G.C. Evans & O. Rackham, 187-207. Oxford: Blackwell Scientific.
- GRIME, J.P. 1979. Plant strategies and vegetation processes. Chichester: John Wiley & Sons.
- GRIME, J.P. & JARVIS, B.C. 1975. Shade avoidance and shade tolerance in flowering plants II. Effects of light on the germination of species of contrasted ecology. In: Light as an ecological factor II, edited by G.C. Evans, R. Bainbridge & O. Rackham, 525-532. Oxford: Blackwell Scientific.
- GRIME, J.P. & JEFFREY, D.W. 1965. Seedling establishment in vertical gradients of sunlight. Journal of Ecology, 53, 621-642.
- GRIME, J.P., HODGSON, J.G. & HUNT, R. 1988. Comparative Plant Ecology: A functional approach to common British plants and communities. London: Unwin Hyman.
- GRIME, J.P., MASON, G., CURTIS, A.V., RODMAN, J., BAND, S.R., MOWFORTH, M.A.G., NEAL, A.M. & SHAW, S. 1981. A comparative study of the germination characteristics in a local flora. Journal of Ecology, 69, 1017-1059.

- GRUBB, P.J. 1976. A theoretical background to the conservation of ecologically distinct groups of annuals and biennials in the chalk grassland ecosystem. Biological Conservation, 10, 53-76.
- GRUBB, P.J. 1977. The maintenance of species-richness in plant communities: the importance of the regeneration niche. Biological Reviews, 52, 107-145. x
- GRUBB, P.J. 1988. The uncoupling of disturbance and recruitment, two kinds of seed bank, and persistence of plant populations at the regional and local scales. Annales Zoologici Fennici, 25, 23-36.
- HANF, M. 1976. Weeds and their seedlings. Ipswich: BASF United Kingdom Limited.
- HARPER, J.L. (ed). 1960. The biology of weeds. Oxford: Blackwell Scientific.
- HARPER, J.L. 1977. The population biology of plants. London: Academic Press.
- HARPER, J.L. & BENTON, R.A. 1966. The behaviour of seeds in soil. II. The germination of seeds on the surface of a water supplying substrate. Journal of Ecology, 54, 151-166.
- HARPER, J.L., LOVELL, P.H. & MOORE, K.G. 1970. The shapes and sizes of seeds. Annual Review of Ecology and Systematics, 1, 327-356.
- HARPER, J.L., WILLIAMS, J.T. & SAGAR, G.R. 1965. The behaviour of seeds in soil. I. The heterogeneity of soil surfaces and its role in determining the establishment of plants from seed. Journal of Ecology, 53, 273-286.
- HEYDECKER, W. (Ed) 1973. Seed ecology. London: Butterworths.
- HILL, M.O. 1980. The changing flora of aging plantations of sitka spruce and other conifers. Proceedings of the Conference on Weed Control in Forestry, 1-2 April, 1980, University of Nottingham, England, edited by D Atkinson, W J McCavish, D H Spencer-Jones, M G O'Keefe, M G Allen & R J Makepeace, 45-53. Wellesbourne: Association of Applied Biologists.
- HILL, M.O. & STEVENS, P.A. 1981. The density of viable seed in soils of forest plantations in upland Britain. Journal of Ecology, 69, 693-709.
- HILL, N.M., PATRIQUIN, D.G. & VANDER KLOET, S.P. 1989. Weed seed bank and vegetation at the beginning and end of the first cycle of a 4-course crop rotation with minimal weed control. Journal of Applied Ecology, 26, 233-246.
- HOBBS, R.J. & GIMINGHAM, C.H. 1984. Studies on fire in the Scottish heathland communities. II. Post-fire vegetation development. Journal of Ecology, 72, 585-610.
- HOBBS, R.J., MALLIK, A.U. & GIMINGHAM, C.H. 1984. Studies on fire in Scottish heathland communities. III. Vital attributes of the species. Journal of Ecology, 72, 963-976.
- HOPKINS, D.R. & PARKER, V.T. 1984. A study of the seed bank of a salt marsh in northern San Francisco Bay. American Journal of Botany, 71, 348-355.



- HOUGHTON, A., BANNISTER, N., McLAREN, D. & WRIGHT, S. 1988. A review of the opportunities presented by extended field margins for conservation, recreation and agriculture. Ashford: CEAS Consultants (Wye) Ltd.
- HOWE, C.D. & CHANCELLOR, R.J. 1983. Factors affecting the viable seed content of soils beneath lowland pastures. Journal of Applied Ecology, 20, 915-922. X
- HUTCHINGS, M.J. & RUSSELL, P.J. 1989. The seed regeneration dynamics of an emergent salt marsh. Journal of Ecology, 77, 615-637.
- JALLOQ, M. 1975. The invasion of molehills by weeds as a possible factor in the degeneration of reseeded pasture. I. The buried viable seed population of molehills from four reseeded pastures in west Wales. Journal of Applied Ecology, 12, 643-657. y
- JENSEN, H.A. 1969. Content of buried seeds in arable soil in Denmark and its relation to the weed population. Dansk Botanisk Arkiv, 27 (2), 1-56.
- JERLING, L. 1983. Composition and viability of the seed bank along a successional gradient on a Baltic sea shore meadow. Holarctic Ecology, 6, 150-156.
- JOENJE, W. 1979. Plant succession and nature conservation of newly embanked tidal flats in the Lauwerzeepolder. In Ecological Processes in Coastal Environments, edited by R.L. Jefferies & A.J. Davy, 617-634. Oxford: Blackwell Scientific.
- JOHNSON, E.A. 1975. Buried seed populations in the subarctic forest east of Great Slave Lake, Northwest Territories. Canadian Journal of Botany, 53, 2933-2941.
- JOHNSTON, A., SMOLIAK, S. & STRINGER, P.W. 1969. Viable seed populations in Alberta prairie topsoils. Canadian Journal of Plant Science, 49, 75-82.
- KEENLYSIDE, C. & MORRIS, D. 1989. Management options for expanded field margins. Peterborough: Nature Conservancy Council. o
- KEIZER, P.J., VAN TOOREN, B.F. & DURING, H.J. 1985. Effects of bryophytes on seedling emergence and establishment of short-lived forbs in chalk grassland. Journal of Ecology, 73, 493-504.
- KELLMAN, M.C. 1974a. The viable weed seed content of some tropical agricultural soils. Journal of Applied Ecology, 11, 669-677.
- KELLMAN, M.C. 1974b. Preliminary seed budgets for two plant communities in Coastal British Columbia. Journal of Biogeography, 1, 123-133.
- KELLMAN, M.C. 1978. Microdistribution of viable weed seed in two tropical soils. Journal of Biogeography, 5, 291-300.
- KELLY, D. 1989. Demography of short-lived plants in chalk grassland, I-III. Journal of Ecology, 77, 747-798.
- KING, T.J. 1975. Inhibition of seed germination under leaf canopies in *Arenaria serpyllifolia*, *Veronica arvensis* and *Cerastium holosteoides*. New Phytologist, 75, 87-90.

- KING, T.J. 1976. The viable seed contents of ant-hill and pasture soil. New Phytologist, 77, 143-147. x
- KING, T.J. 1977. The plant ecology of ant-hills in calcareous grasslands. III. Factors affecting the population size of selected species. Journal of Ecology, 65, 279-315.
- KROPAC, Z. 1966. Estimation of weed seeds in arable soil. Pedobiologia, 6, 105-128.
- LEAKEY, R.R.B. 1981. Adaptive biology of vegetatively regenerating weeds. Advances in Applied Biology, 6, 57-90.
- LECK, M.A. 1980. Germination in Barrow, Alaska, tundra soil cores. Arctic and Alpine Research, 12, 343-349.
- LECK, M.A. & GRAVELINE, K.K. 1979. The seed bank of a freshwater tidal marsh. American Journal of Botany, 66, 1006-1015.
- LECK, M.A. & SIMPSON, R.A. 1987. Seed bank of a freshwater tidal wetland: turnover and relationship to vegetation change. American Journal of Botany, 74, 360-370.
- LECK, M.A., PARKER, V.T. & SIMPSON, R.L. 1989. Ecology of soil seed banks. London: Academic Press. c213224
- LEWIS, J. 1973. Longevity of crop and weed seeds: survival after 20 years in the soil. Weed Research, 13, 179-191.
- LIVINGSTON, R.B. & ALESSIO, M.L. 1968. Buried viable seed in successional field and forest stands, Harvard Forest, Massachusetts. Bulletin of the Torrey Botanical Club, 95, 58-69.
- MACDONALD, N. & WATKINSON, A.R. 1981. Models of an annual plant population with a seed bank. Journal of Theoretical Biology, 93, 643-653.
- MCGRAW, J.B. 1980. Seed bank size and distribution of seeds in cottongrass tussock tundra, Eagle Creek, Alaska. Canadian Journal of Botany, 58, 1607-1611.
- MCGRAW, J.B. 1987. Seed bank properties of an Appalachian sphagnum bog and a model of the depth distribution of viable seeds. Canadian Journal of Botany, 65, 2028-2035.
- MCRILL, M. 1979. The ingestion of weed seed by earthworms. Proceedings 12th British Weed Control Conference (1974), 2, 519-525.
- MAHMOUD, A. & GRIME, J.P. 1974. A comparison of negative relative growth rates in shaded seedlings. New Phytologist, 73, 1215-1219.
- MAJOR, J. & PYOTT, W.T. 1966. Buried, viable seeds in two California bunchgrass sites and their bearing on the definition of a flora. Vegetatio, 13, 253-282.
- MALLIK, A.U. & GIMINGHAM, C.H. 1985. Ecological effects of heather burning II. Effects on seed germination and vegetative regeneration. Journal of Ecology, 73, 633-644.

- MALLIK, A.U., HOBBS, R.J. & LEGG, C.J. 1984. Seed dynamics in *Calluna-Arctostaphylos* heath in north-eastern Scotland. Journal of Ecology, 72, 855-871.
- MANN, H.H. 1957. Weed herbage of slightly acid arable soils as affected by manuring. Journal of Ecology, 45, 149-156.
- MARLETTE, G.M. & ANDERSON, J.E. 1986. Seed banks and propagule dispersal in crested wheatgrass stands. Journal of Applied Ecology, 23, 161-175.
- MARQUIS, D.A. 1975. Seed storage and germination under northern hardwood forests. Canadian Journal of Forest Research, 5, 478-484.
- MARSHALL, E.J.P. 1985. Weed distributions associated with cereal field edges - some preliminary observations. Aspects of Applied Biology, 9, 49-58.
- MARSHALL, E.J.P. 1986. Studies in the flora in arable field margins. Long Ashton Research Station Technical Report, 96.
- MARSHALL, E.J.P. 1989. Distribution patterns of plants associated with arable field edges. Journal of Applied Ecology, 26, 247-258.
- MAYER, A.M. & POLJAKOFF-MAYBER, A. 1963. The germination of seeds. Oxford: Pergamon.
- MILES, J. 1972. Experimental establishment of seedlings on a southern English heath. Journal of Ecology, 60, 225-234.
- MILES, J. 1973a. Early mortality and survival of self-sown seedlings in Glenfeshie, Inverness-shire. Journal of Ecology, 61, 93-98.
- MILES, J. 1973b. Natural recolonization of experimentally bared soil in Callunetum in north-east Scotland. Journal of Ecology, 61, 399-412.
- MILES, J. 1974a. Experimental establishment of new species from seed in Callunetum in north-east Scotland. Journal of Ecology, 62, 527-551.
- MILES, J. 1974b. Effects of experimental interference with stand structure on establishment of seedlings in Callunetum. Journal of Ecology, 62, 675-687.
- MILES, J. 1975. Performance after six growing seasons of new species established from seed in Callunetum in north-east Scotland. Journal of Ecology, 63, 891-902.
- MILLER, G.R. & CUMMINS, R.P. 1976. Seed production on mountains. Annual Report Institute of Terrestrial Ecology 1975, 35-36.
- MILLER, G.R. & CUMMINS, R.P. 1981. Population dynamics of buried seeds on mountains. Annual Report Institute Terrestrial Ecology 1980, 75-76.
- MILLER, G.R. & CUMMINS, R.P. 1985. How important are banks of buried seeds in the Cairngorm mountains? Annual Report Institute of Terrestrial Ecology 1984, 101-104.
- MILLER, G.R. & CUMMINS, R.P. 1987. Role of buried viable seeds in the recolonization of disturbed ground by heather (*Calluna vulgaris* [L] Hull) in the Cairngorm Mountains, Scotland, UK. Arctic and Alpine Research, 19, 396-401.

- MILTON, W.E.J. 1939. The occurrence of buried viable seeds in soils at different elevations and on a saltmarsh. Journal of Ecology, 27, 149-159.
- MILTON, W.E.J. 1948. The buried viable-seed content of upland soils in Montgomeryshire. Empire Journal of Experimental Agriculture, 16, 163-177.
- MOORE, J.M. & WEIN, R.W. 1977. Viable seed populations by soil depth and potential site recolonization after disturbance. Canadian Journal of Botany, 55, 2408-2412.
- MOORE, P.D. 1980. Soil seed banks. Nature, 284, 123-124.
- MULLER, F.M. 1978. Seedlings of the North-western European lowland. The Hague: Junk.
- NICHOLSON, A. & KEDDY, P.A. 1983., The depth profile of a shoreline seed bank in Matchedash Lake, Ontario. Canadian Journal of Botany, 61, 3293-3296.
- NOBLE, I.R. & SLATYER, R.O. 1980. The use of vital attributes to predict successional changes in plant communities subject to recurrent disturbances. Vegetatio, 43, 5-21.
- ODUM, S. 1978. Dormant seeds in Danish ruderal soils. Horsholm: Royal Veterinary and Agricultural University, Horsholm Arboretium.
- OLMSTED, N.W. & CURTIS, J.D. 1947. Seeds of the forest floor. Ecology, 28, 49-52.
- OOSTING, H.J. & HUMPHREYS, M.E. 1940. Buried viable seed in a successional series of old field and forest soils. Bulletin of the Torrey Botanical Club, 67, 253-273.
- PARKER, V.T. & LECK, M.A. 1985. Relationships of seed banks to plant distribution patterns in a freshwater tidal marsh. American Journal of Botany, 72, 161-174.
- POLLARD, F. & CUSSANS, G.W. 1981. The influence of tillage on the weed flora in a succession of winter cereal crops on a sandy loam soil. Weed Research, 21, 185-190.
- RABINOWITZ, D. 1981. Buried viable seeds in a North American tall grass prairies: the resemblance of their abundance and composition to dispersing seeds. Oikos, 36, 191-195.
- RABINOWITZ, D. & RAPP, J.K. 1980. Seed rain in a North American tall grass prairie. Journal of Applied Ecology, 17, 793-802.
- RATCLIFFE, D. 1961. Adaptation to habitat in a group of annual plants. Journal of Ecology, 49, 187-203.
- ROBERTS, H.A. 1958. Studies on the weeds of vegetable crops. I. Initial effects of cropping on the weed seeds in the soil. Journal of Ecology, 46, 759-768.
- ROBERTS, H.A. 1962. Studies on the weeds of vegetable crops. II. Effect of

- six years of cropping on the weed seeds in the soil. Journal of Ecology, 50, 803-813.
- ROBERTS, H.A. 1963. Studies on the weeds of vegetable crops. III. Effect of different primary cultivations on the weed seeds in the soil. Journal of Ecology, 51, 83-95.
- ROBERTS, H.A. 1963. Studies on the weeds of vegetable crops. IV. Further observations on the effects of different primary cultivations. Journal of Ecology, 51, 323-332.
- ROBERTS, H.A. 1964. Emergence and longevity in cultivated soil of seeds of some annual weeds. Weed Research, 4, 296-307.
- ROBERTS, H.A. 1968. The changing population of viable weed seeds in an arable soil. Weed Research, 8, 253-256.
- ROBERTS, H.A. 1970. Viable weed seeds in cultivated soils. Report National Vegetable Research Station 1969, 25-28.
- ROBERTS, H.A. (ed.) 1972. Viability of seeds. London: Chapman & Hall.
- ROBERTS, H.A. 1979. Periodicity of seedling emergence and seed survival in some Umbelliferae. Journal of Applied Ecology, 16, 195-201.
- ROBERTS, H.A. 1981. Seed banks in soils. Advances in Applied Biology, 6, x 1-55.
- ROBERTS, H.A. 1986. Persistence of seeds of some grass species in cultivated soil. Grass and Forage Science, 41, 273-276. x
- ROBERTS, H.A. 1986. Seed persistence in soil and seasonal emergence in plant species from different habitats. Journal of Applied Ecology, 23, x 639-656.
- ROBERTS, H.A. & BODRELL, J.E. 1983. Seed survival and periodicity of seedling emergence in ten species of annual weeds. Annals of Applied Biology, 102, 523-532.
- ROBERTS, H.A. & BODDRELL, J.E. 1984. Seed survival and seasonal emergence of seedlings of some ruderal plants. Journal of Applied Ecology, 21, 617-628.
- ROBERTS, H.A. & DAWKINS, P.A. 1967. Effect of cultivation on the numbers of viable weed seeds in soil. Weed Research, 7, 290-301.
- ROBERTS, H.A. & FEAST, P.M. 1972. Fate of seeds of some annual weeds in different depths of cultivated and undisturbed soil. Weed Research, 12, 316-324.
- ROBERTS, H.A. & FEAST, P.M. 1973a. Emergence and longevity of seeds of annual weeds in cultivated and undisturbed soils. Journal of Applied Ecology, 10, 133-143.
- ROBERTS, H.A. & FEAST, P.M. 1973b. Changes in the numbers of viable weed seeds in soil under different regimes. Weed Research, 13, 298-303.

- ROBERTS, H.A. & NEILSON, J.E. 1981a. Seed survival and periodicity of seedling emergence in twelve weedy species of Compositae. Annals of Applied Biology, 97, 325-334.
- ROBERTS, H.A. & NEILSON, J.E. 1981b. Changes in the soil seed bank of four long-term crop/herbicide experiments. Journal of Applied Ecology, 18, 661-668.
- ROBERTS, H.A. & NEILSON, J.E. 1982. Seed banks of soils under vegetable cropping in England. Weed Research, 22, 13-16.
- ROBERTS, H.A. & POTTER, M.E. 1980. Emergence patterns of weed seedlings in relation to cultivation and rainfall. Weed Research, 20, 377-386.
- ROBERTS, H.A. & RICKETTS, M.E. 1979. Quantitative relationships between the weed flora after cultivation and the seed population in the soil. Weed Research, 19, 269-275.
- ROBERTS, H.A. & STOKES, F.G. 1965. Studies on the weeds of vegetable crops. V. Final observations on an experiment with different primary cultivations. Journal of Applied Ecology, 2, 307-315.
- ROBERTS, H.A. & STOKES, G. 1966. Studies on the weeds of vegetable crops. VI. Seed populations of soil under commercial cropping. Journal of Applied Ecology, 3, 181-190.
- ROSS, M.A. & HARPER, J.L. 1972. Occupation of biological space during seedling establishment. Journal of Ecology, 60, 77-88.
- SAGAR, G.R. & MORTIMER, A.M. 1976. An approach to the study of the population dynamics of plants with special reference to weeds. Applied Biology, 1, 1-47.
- SALISBURY, E.J. 1942. The reproductive capacity of plants. London: Bell.
- SALONEN, V. 1989. Weed communities of cereal crops grown on differently revegetated cut-over peatland sites. Journal of Applied Ecology, 26, 563-569.
- SCHENKEVELD, A.J. & VERKAAR, H.J. 1984. The ecology of short-lived forbs in chalk grasslands: distribution of germinative seeds and its significance for seedling emergence. Journal of Biogeography, 11, 251-260.
- SCHERING AGRICULTURE 1986. Weed Guide. Nottingham: Schering Agriculture.
- SHELDON, J.C. 1974. The behaviour of seeds in soil. Journal of Ecology, 62, 47-66.
- SILVERTOWN, J.W. 1980. Leaf-canopy induced seed dormancy in a grassland flora. New Phytologist, 85, 109-118.
- SILVERTOWN, J.W. 1981a. Seed size, life span, and germination date as coadapted features of plant life history. The American Naturalist, 118, 860-864.
- SILVERTOWN, J.W. 1981b. Micro-spatial heterogeneity and seedling demography in species-rich grassland. New Phytologist, 88, 117-129.

- SILVERTOWN, J.W. 1982. Introduction to plant population ecology. London: Longman.
- SILVERTOWN, J.W. 1988. The demographic and evolutionary consequences of seed dormancy. In: Plant population ecology, edited by M J Hutchings, A J Davy & A R Watkinson, 205-219. Oxford: Blackwell Scientific.
- SILVERTOWN, J.W. & DICKIE, J.B. 1981. Seedling survivorship in natural populations of nine perennial chalk grassland plants. New Phytologist, 88, 555-558.
- SMITH, L.M. & KADLEC, J.A. 1983. Seed banks and their role during drawdown of a North American marsh. Journal of Applied Ecology, 20, 673-684.
- SMITH, L.M. & KADLEC, J.A. 1985. The effects of disturbance on marsh seed banks. Canadian Journal of Botany, 63, 2133-2137.
- STEBBINS, G.L. 1971. Adaptive radiation of reproductive characteristics of angiosperms. II. Seeds and seedlings. Annual Review of Ecology and Systematics, 2, 237-260.
- STRICKLER, G.S. & EDGERTON, P.J. 1976. Emergent seedlings from coniferous litter and soil in eastern Oregon. Ecology, 57, 801-807.
- THOMPSON, K. 1978. The occurrence of buried viable seeds in relation to environmental gradients. Journal of Biogeography, 5, 425-430.
- THOMPSON, K. 1985. Buried seed banks as indicators of seed output along an altitudinal gradient. Journal of Biological Education, 19, 137-140.
- THOMPSON, K. 1986. Small-scale heterogeneity in the seed bank of an acidic grassland. Journal of Ecology, 74, 733-738.
- THOMPSON, K. 1987. Seeds and seed banks. New Phytologist, 106 (Suppl), 23-34.
- THOMPSON, K. & GRIME, J.P. 1979. Seasonal variation in the seed banks of herbaceous species in ten contrasting habitats. Journal of Ecology, 67, 893-921.
- THOMPSON, K. & GRIME, J.P. 1983. A comparative study of germination responses to diurnally-fluctuating temperatures. Journal of Applied Ecology, 20, 141-156.
- THOMPSON, K., GRIME, J.P. & MASON, G. 1977. Seed germination in response to diurnal fluctuations of temperature. Nature, 267, 147-149.
- UNGAR, I.A. 1987. Population ecology of halophyte seeds. Botanical Review, 53, 301-334.
- UNGAR, I.A. & RIEHL, T.E. 1980. The effect of seed reserves on species composition in zonal halophyte communities. Botanical Gazette, 141, 447-452.
- VAN DER VALK, A.G. & DAVIS, C.B. 1976. The seed banks of prairie glacial marshes. Canadian Journal of Botany, 54, 1832-1838.
- VAN DER VALK, A.G. & DAVIS, C.B. 1978. The role of seed banks in the vegetation dynamics of prairie glacial marshes. Ecology, 59, 322-335.

- VAN DER VALK, A.G. & DAVIS, C.R. 1979. A reconstruction of the recent vegetational history of a prairie marsh, Eagle Lake, Iowa, from its seed bank. Aquatic Botany, 6, 29-51.
- VERKAAR, H.J. & SCHENKEVELD, A.J. 1984a. On the ecology of short-lived forbs in chalk grasslands: seedling development under low photon flux density conditions. Flora, 175, 135-141.
- VERKAAR, H.J. & SCHENKEVELD, A.J. 1984b. On the ecology of short-lived forbs in chalk grassland: life history characteristics. New Phytologist, 98, 659-672.
- VERKAAR, H.J. & SCHENKEVELD, A.J. 1984c. On the ecology of short-lived forbs in chalk grassland: semelparity and seed output of some species in relation to various levels of nutrient supply. New Phytologist, 98, 673-682.
- VERKAAR, H.J., SCHENKEVELD, A.J. & KLASHORST, M.P., VAN DE 1983a. The ecology of short-lived forbs in chalk grasslands: dispersal of seeds. New Phytologist, 95, 335-344.
- VERKAAR, H.J., SCHENKEVELD, A.J. & BRAND, J.M. 1983b. On the ecology of short-lived forbs in chalk grassland: micro-site tolerance in relation to vegetation structure. Vegetatio, 52, 91-102.
- WARINGTON, K. 1958. Changes in the weed flora on Broadbalk permanent wheat field during the period 1930-55. Journal of Ecology, 46, 101-113.
- WATKINSON, A.R. 1981. The population ecology of winter annuals. In: The Biological Aspects of Rare Plant Conservation, edited by H. Synge, 253-264. Chichester: Wiley.
- WAY, J.M. (ed) 1969. Road Verges. Their function and management. Huntingdon: Institute of Terrestrial Ecology.
- WAY, J.M. & GREIG-SMITH, P.W. (eds) 1987. Field Margins. Monograph no. 35. Thornton Heath: British Crop Protection Council.
- WEINS, D. 1984. Ovule survivorship, broad size, life history, breeding systems, and reproductive success in plants. Oecologia, 64, 47-53.
- WELCH, D. 1985. Studies in the grazing of heather moorland in north-east Scotland. IV. Seed dispersal and plant establishment in dung. Journal of Applied Ecology, 22, 461-472.
- WESSON, G. & WAREING, P.F. 1967. Light requirements of buried seeds. Nature, 213, 600-601.
- WESSON, G. & WAREING, P.F. 1969a. The role of light in the germination of naturally occurring populations of buried weed seeds. Journal of Experimental Botany, 20, 402-413.
- WESSON, G. & WAREING, P.F. 1969b. The induction of light sensitivity in weed seeds by burial. Journal of Experimental Botany, 20, 414-425.
- WESTOBY, M. 1981. How diversified seed germination is selected. American Naturalist, 118, 882-885.



- WHIPPLE, S.A. 1978. The relationship of buried, germinating seeds to vegetation in an old-growth Colorado subalpine forest. Canadian Journal of Botany, 56, 1505-1509.
- WILLIAMS, E.D. 1983a. Germinability and enforced dormancy in seeds of species of indigenous grassland. Annals of Applied Biology, 102, 557-566.
- WILLIAMS, E.D. 1983b. Effects of temperature fluctuation, red and far-red light and nitrate on seed germination of five grasses. Journal of Applied Ecology, 20, 923-935.
- WILLIAMS, E.D. 1984. Changes during 3 years in the size and composition of the seed bank beneath a long-term pasture as influenced by defoliation and fertilizer regime. Journal of Applied Ecology, 21, 603-615. x

