

Article (refereed) - postprint

Arroyo, Beatriz; Leckie, Fiona; Amar, Arjun; McCluskie, Aly; Redpath, Steve. 2014. **Ranging behaviour of hen harriers breeding in Special Protection Areas in Scotland.** *Bird Study*, 61 (1). 48-55.

[10.1080/00063657.2013.874976](https://doi.org/10.1080/00063657.2013.874976)

Copyright © 2014 British Trust for Ornithology

This version available <http://nora.nerc.ac.uk/509535/>

NERC has developed NORA to enable users to access research outputs wholly or partially funded by NERC. Copyright and other rights for material on this site are retained by the rights owners. Users should read the terms and conditions of use of this material at

<http://nora.nerc.ac.uk/policies.html#access>

This is a pre-copy-editing, author-produced PDF of an article accepted for publication in Bird Study following peer review. The definitive publisher-authenticated version is available online at Bird Study, 61 (1). 48-55.:

<http://dx.doi.org/10.1080/00063657.2013.874976>

Contact CEH NORA team at
noraceh@ceh.ac.uk

1 **Ranging behaviour of Hen Harriers breeding in Special Protection**
2 **Areas in Scotland**

3

4 BEATRIZ ARROYO^{1,2}, FIONA LECKIE^{1,3}, ARJUN AMAR^{4,5}, ALY
5 MCCLUSKIE^{1,7} & STEVE REDPATH^{1,6}

6

7 *1-Centre for Ecology and Hydrology (CEH), Hill of Brathens, Banchory,*
8 *Aberdeenshire, AB31 4BW UK*

9 *2-Instituto de Investigación en Recursos Cinegéticos (IREC) (CSIC-UCLM-JCCM),*
10 *Ronda de Toledo s/n, 13005 Ciudad Real, Spain*

11 *3-Natural Research Ltd, Hill of Brathens, Banchory, Aberdeenshire, AB31 4BW UK*

12 *4-Game Conservancy Trust, c/o CEH - Banchory, Hill of Brathens, Banchory,*
13 *Aberdeenshire, AB31 4BW, UK*

14 *5-Percy FitzPatrick Institute of African Ornithology, University of Cape Town,*
15 *DST/NRF Centre of Excellence, 7701, Rondebosch, South Africa*

16 *6-Aberdeen Centre for Environmental Sustainability, University of Aberdeen,*
17 *Aberdeen AB24 2TZ, UK.*

18 *7-RSPB, 2 Lochside View, Edinburgh Park, Edinburgh, EH12 9DH.*

19

20 Running head: Hen Harrier ranging distances

21

22 Keywords:

23 Hunting distances, protected areas, management

24

25

26 Correspondence author (email):

27 Beatriz Arroyo (beatriz.arroyo@uclm.es)

28

29

30 **Capsule** Breeding female Hen Harriers hunted mostly within 1 km from the nest and
31 males mostly within 2 km.

32 **Aims** To quantify temporal and spatial variation in home range sizes and hunting
33 distances of breeding male and female Hen Harriers.

34 **Methods** We radio-tracked ten breeding harriers (five males and five females) in
35 three Special Protection Areas (SPAs) in Scotland between 2002-2004.

36 **Results** Male Hen Harriers travelled up to 9 km from nests but had a home range size
37 that averaged only 8 km² (90% kernel); average home range size for females was 4.5
38 km² . Hunting distances did not vary throughout the season. No significant differences
39 were found among study areas, but there was large individual variability.

40 **Conclusions** Our results provide information on foraging harriers to support
41 management: actions within 1 km of nesting sites will favour both sexes, and within
42 2km will mostly favour males. Our data also suggest overlap between foraging areas
43 of neighbouring birds. Thus, there is the potential for good foraging areas to be
44 utilised by multiple breeding pairs.

45

46 Habitat loss and land use change are recognised as major threats to many bird
47 populations, including raptors (Newton 1979). Populations of raptors have been
48 shown to decline due to loss of their preferred habitats (Donazar et al. 1993, Amar &
49 Redpath 2005, Thiollay 2006). Legislative protection of habitats is thus a major
50 conservation tool used all over the world. In Europe the two most influential pieces of
51 protective legislation relating to nature conservation are the Habitats (92/42/EEC) and
52 Birds Directives (2009/147/EC). These Directives give EU member states the power
53 and responsibility to create Special Protection Areas (SPAs) to protect birds which are
54 rare or vulnerable in Europe, forming the European network of protected areas known
55 as Natura 2000. SPAs are intended to safeguard the habitats of the species for which
56 they are designated and to protect the birds from significant disturbance. There may
57 be financial incentives for sustainable management of the land, in ways that have been
58 recognised as beneficial to the species either directly, for example by providing
59 nesting habitat, or indirectly, for example by providing habitats for their prey species.

60 A number of studies have highlighted that effective management of areas for
61 vulnerable species must consider their foraging needs in addition to their nesting
62 needs (Donazar et al. 1993, Martin & Possingham 2005, García et al. 2006). Studies
63 have shown that availability of good foraging areas around nest sites can influence
64 breeding success (e.g. Tella et al. 1998, Rodriguez et al. 2006, Amar et al. 2008,
65 Hinan & Clair 2008). Furthermore, some birds may regularly forage far away from
66 their nests, so protected areas based only on distribution of nests may be insufficient
67 to contain all resources needed for a given species (Martinez et al. 2007, Guixé &
68 Arroyo 2011). Information on ranging behaviour may thus provide critical
69 information for management of protected species in protected areas.

70 The Hen Harrier *Circus cyaneus* is a medium-size raptor which is listed on
71 Annex 1 of the EU Birds Directive. In the UK, it breeds predominantly in heather
72 moorland (including grouse moors, Redpath et al. 1998, Sim et al. 2007, Hayhow et
73 al. 2014), where it preys mainly on small passerines and small mammals, although
74 they also sometimes take larger prey like grouse, waders and young rabbits (Redpath
75 et al. 2002, Amar et al. 2003). When breeding in moorland, the best foraging habitats
76 for the species include areas of heather *Calluna vulgaris* mixed with rough grass
77 habitats (Amar & Redpath 2005, Arroyo et al. 2009), where prey abundance is highest
78 (e.g. Smith et al. 2001, Vanhinsbergh & Chamberlain 2001, Amar & Redpath 2005).

79 National surveys for this species over recent decades have shown that there have been
80 marked declines in some regions and the population is currently well below its
81 potential population size and range (Sim et al. 2007, Anderson et al. 2009, Fielding et
82 al.2011, Hayhow et al. 2014). The conservation status of the species in the UK is
83 threatened because Hen Harriers can, in certain circumstances, reduce the numbers of
84 red grouse available for recreational shooting (Thirgood et al. 2000), and as a result
85 they are illegally killed on certain grouse moors (Etheridge et al. 1997). There is,
86 therefore, a strong conservation concern for this species, with UK government listing
87 the species as a conservation priority, and a series of SPAs have been identified in the
88 UK for this species (<http://jncc.defra.gov.uk/pdf/UKSPA/UKSPA-A6-47A.pdf>).

89 Accurate information on home range size of Hen Harriers is important to
90 understand whether all the needs for the species are likely to be covered within these
91 SPAs. Evaluation of hunting distances will also provide information on the ideal
92 locations to deploy conservation measures in support of the SPA, such as agro-
93 environmental support schemes (Amar et al. 2011). This information will also be
94 useful for development issues such as placement of windfarms (Madders & Whitfield
95 2006, Whitfield & Madders 2006), or in the context of the conflict with grouse
96 shooting (Redpath & Thirgood 2009, Thompson et al. 2009, Sotherton et al. 2009).
97 For example, management of SPAs may include measures to reduce the impact of
98 predation on grouse (e.g. Langholm Moor Demonstration Project,
99 <http://www.langholmproject.com/index.html>), if part of the area is used for
100 commercial shooting.

101 Published information on the home range sizes for this species is limited.
102 Picozzi (1978) estimated foraging range of male harriers in NE Scotland as 14 km²
103 based on observations of hunting birds. Radio-tracking studies of the closely related
104 Northern Harrier *Circus hudsonius* in Idaho, USA, produced an estimated average
105 breeding male range size of 16 km² (Martin 1987). Both of these estimates were
106 however based on Minimum Convex Polygons, which may overestimate ranging
107 areas if there are outlying locations (Kenward 2001). Beyond these studies, there
108 exists only a limited amount of indirect information about maximum hunting
109 distances based on observations of hunting birds in continental Europe (Schipper
110 1977, García & Arroyo 2005).

111 This paper aims to investigate the ranging behaviour of breeding Hen Harriers.
112 Specifically, we aim to evaluate the average home range size and maximum hunting

113 distances of breeding Hen Harriers, and test whether home ranges varied between
114 sexes or study areas and whether there was any temporal variation in ranging
115 distances over the course of the nestling period.

116

117 **METHODS**

118

119 **Study areas and radio-tracking data**

120

121 The study was carried out on three Scottish SPAs over three years. Harrier nests were
122 located in each area early in the breeding season. Breeding adults were trapped, under
123 the appropriate licences, during the nestling period (using dho-ghaza collapsible nets
124 set close to the nest with a nest predator decoy, or mono-filament noose bonnets on a
125 plastic eagle owl) and fitted with 8g tail mounted radio telemetry tags (Biotrack Ltd,
126 Dorset). In total twelve adults were tagged: three birds (one male and two females) in
127 Langholm in 2002, three birds (two males and one female) in Orkney in 2003 and six
128 birds (two males and four females) in Galloway in 2004.

129 Locations of birds were evaluated through bi or tri-angulations from multiple
130 vantage points distributed throughout the study areas: observers stationed at elevated
131 fixed points conducted scans for each tagged individual using a 3 bar Yagi antennae
132 and radio-receiver. When a signal was located, observers communicated using two-
133 way radios, and simultaneously took a compass bearing for that signal. Positions were
134 then calculated by plotting compass bearings on 1:25 000 maps.

135 We calculated the error in the estimation of the locations derived with this
136 method using tags attached to poles located in certain (immobile) positions unknown
137 to observers, which were asked to provide a fix for them ($n = 133$ crossings on 20
138 dummy tags in Langholm; $n = 142$ crossings on 25 dummy tags in Orkney; $n = 31$
139 crossings on 4 dummy tags in Galloway). Locations of these fixed tags based on bi- or
140 triangulations were associated with an error of x meters (range 501-728 m). Accuracy
141 depended mainly on the angle between the bearings: error was greater when bearings
142 crossed at angles higher than 135° or lower than 45° . When eliminating these fixes,
143 the error made with bi- or triangulations was not significantly different ($P > 0.3$), and
144 averaged 308 ± 172 m (mean \pm sd, $n = 6$) in Langholm, 65 ± 220 m ($n = 28$) in
145 Orkney, and 206 ± 125 m ($n = 19$) in Galloway. This figure may not necessarily be

146 comparable to the error in fixing moving birds, because there is probably less time for
147 observers to obtain a locational fix, however the signal from transmitters in the air is
148 better than that of transmitters closer to the ground (which was the case for those used
149 to estimate errors).

150 In Galloway and Orkney, fix locations were taken every ten to fifteen minutes
151 from the same vantage point for a period of several hours, and repeated every few days.
152 In Langholm, the monitoring was less intensive, with one or two bearings being taken
153 per day per bird, repeated every few days. Locations were obtained throughout the
154 nestling period, until the chicks had left the nest. A total of 1146 fixes were obtained
155 (all birds combined). We carried out an initial selection of these fixes, eliminating
156 those ($n = 523$) based on bearings crossing at angles lower than 45 or higher than 135
157 degrees. After that selection, the average time between successive fixes on the same
158 bird in 2003-2004 was 33 ± 33 min (2-198). As some bearings were taken at short
159 intervals, some fixes may not have been independent (Kenward 2001), therefore we ran
160 autocorrelation analyses with Ranges VI, and calculated Shoenener's (1981) test of
161 Time to Independence between fixes (Kenward 2001) for each bird. This analysis
162 indicated that locations were independent for all birds but one (a female, tag 658, in
163 2004), for which time to independence was 1100 minutes, a figure much larger than our
164 recording sessions. That particular female moved little around the nest (see results). We
165 therefore included all fixes for this female in further analyses, while noting its spatially
166 restricted behaviour. In contrast, we eliminated data from two females (one in
167 Langholm and one in Galloway), for which only 3 and 6 fixes (respectively) were
168 available after selection, because this sample size was insufficient to calculate home
169 range size. The average number of fixes for the other tracked birds was 61 ± 33 ($n = 10$,
170 range 11-116).

171

172 **Analyses**

173

174 Home range size was estimated with ArcView 3.2, using Kernel Contours least
175 squares cross validation (LSCV) method to provide 50, 70 and 90% kernels. Kernel-
176 based LSCV home-range estimators are generally favoured with respect to space use
177 patterns (Worton 1989, Boitani & Fuller 2000). Kernel estimators provide an
178 indication of the relative frequency of use of different areas within the home range,

179 thus providing biologically meaningful information, and can give stable area estimates
180 with only 15-20 fixes (Kenward 2001). Minimum Convex Polygons (MCP) from
181 fixes were also calculated to allow comparisons with other studies.

182 We examined the relationship between hunting distance (distance from the
183 nest to tracking fix, calculated with ArcView) and the phase of the nestling cycle
184 using General Linear Mixed Models, with a normal distribution and an identity link
185 function, using “individual” and “area” as random variables to account for the lack of
186 independence of observations of the same bird and fixes within the same study area.
187 We defined a “relative date” with day 1 being the hatching date of a tracked bird’s
188 brood. In two cases in Orkney, monitored males were bigamous. In those cases, we
189 considered the hatching date of the earliest female, and distance to the nest from each
190 fix was evaluated as the distance to the nearest nest.

191 Differences in home range size among areas or among sexes were tested with
192 General Linear Models, fitting the response variables (home range size in km²) with a
193 normal distribution and an identity link function.

194 Statistical analyses were carried out using SAS 9.2 (SAS Institute Inc. 2004)

195
196

197 **RESULTS**

198

199 Most female fixes (67%, $n = 272$) were within 1 km of the nest (Fig. 1). In contrast,
200 only 44% ($n = 343$) of male fixes were within that distance. The maximum distance
201 from the nest at which a male was recorded was 8.5 km (Fig. 1). The average
202 proportion of male fixes beyond 2 km was $24 \pm 16\%$ ($n = 5$, range 9-45).

203 Distance from the nest did not vary in relation to relative date (days from
204 hatching), but varied in relation to sex (relative date: $F_{1,603} = 0.001$, $P = 0.95$; sex:
205 $F_{1,603} = 5.18$, $P = 0.02$, LS Means for males: 1.52 ± 0.23 ; for females 0.85 ± 0.22 ; Fig.
206 1).

207 There was large variability in home range size between individuals, for both
208 sexes (Table 1, Figs. 2-4). However, average male home range size was almost twice
209 the size of females, irrespective of which method of estimation was used (Tables 1 &
210 2). Differences between sexes were statistically significant, whereas differences in

211 home range sizes between study areas were not, although sample size was small
212 (Table 3).

213

214

215 **DISCUSSION**

216

217 Our results showed that male Hen Harriers in Scotland mostly hunted within 2 km of
218 their nest and the estimated 90% kernel of their home ranges averaged 8 km². Female
219 harriers mostly hunted within 1 km of their nest and average home range estimates
220 were half the size of that of males. These figures did not vary significantly among the
221 three study areas, although there was large individual variability.

222 Geographical variations in home range are expected as a result of differences
223 in habitat and food (Tella et al. 1998, Jedrzejewski et al. 2007, Schmidt 2008). The
224 fact that we did not find statistical differences among study areas may be a
225 consequence of the large individual variation and our small sample size: our data may
226 thus lack power for between-region comparisons. However, our results suggest that, at
227 least within the study areas, these differences are not extremely marked. The two
228 previous studies calculating estimates of home range size for this species or the
229 closely related Northern Harrier in the US were larger, at 14 km² (Picozzi 1978) and
230 16 km² (Martin 1987). Both studies used minimum convex polygons to estimate
231 ranges, and those values are similar to the 17 km² we estimated in our study using that
232 method. The lack of important differences in average home range sizes among areas
233 (both in this study and in relation to the two other previous ones) may reflect similar
234 prey abundances in all studies, or that there is a maximum distance from the nest
235 beyond which it is unprofitable for this species to regularly forage.

236 Sexual differences in ranging behaviour such as those found in this study were
237 not unexpected. Martin's (1987) study of radio-tracked breeding northern harriers
238 found that female harriers never ranged further than 2 km from their nest sites,
239 whereas males spent 26% of their time ranging over 2 km from the nest, which is,
240 again, very similar to our findings from this current study. Other previous studies have
241 also suggested that males hunt further away from their nests than females, both in the
242 UK (Picozzi 1978, Thirgood et al. 2003) and in Spain (García & Arroyo 2005). This
243 may also explain why habitat around the nest affected prey delivery to the nest by
244 females, but not males, at Langholm (Amar et al. 2004). Hunting closer to the nest

245 may enable females to quickly return to brood the young if weather conditions change
246 (Redpath et al. 2002) or to observe their nesting area and protect the nestlings from
247 predation (Amar & Burthe 2001).

248

249 Knowledge about the degree of overlap in home ranges of neighbouring
250 individuals provides important information on whether good quality foraging patches
251 can benefit more than one breeding pair. In our study, it was not possible to quantify
252 the degree of overlap between neighbouring ranges because not all birds nested
253 adjacent to each other. However, home ranges of the two neighbouring males in
254 Galloway did overlap extensively, as did those of two females, to a certain extent
255 (Fig. 2), although the smaller size of female home ranges and the tendency for the
256 range to be centred around the nest implied that the overlap for females in general
257 might be less extensive. In Langholm and Orkney, it was not possible to evaluate
258 overlap, because trapped birds were from non-neighbouring nests (Orkney), or data
259 came from different sexes (Langholm). However, the home ranges of all three males
260 included the nest sites of other birds (Arroyo et al. 2006, and Fig. 2), suggesting that
261 they must have overlapped with the ranges of at least some of the neighbouring birds.
262 These results also support Redpath (1992), who noted that the hunting ranges of birds
263 in Highland Scotland overlapped considerably. These results have implications for
264 conservation management, because they suggest that when creating good foraging
265 areas there is the potential for them to be utilised by multiple breeding pairs, and
266 therefore their benefit as a conservation measure can be maximised if they are located
267 within close enough proximity to multiple nesting territories.

268 SPA management should consider as a priority the creation or maintenance of
269 favoured foraging habitats for harriers (Arroyo et al. 2009). Our results provide
270 information about where to implement management to favour foraging harriers: any
271 action within 2 km of existing nesting sites will favour males, but management within
272 1 km will be needed to favour foraging females.

273

274 **ACKNOWLEDGEMENTS**

275

276 This work was a partnership project between Royal Society for the Protection of Birds
277 (RSPB), Scottish National Heritage (SNH), Game Conservancy Trust (GCT) and
278 Centre for Ecology and Hydrology (CEH). We are very grateful to everyone that

279 helped during fieldwork, particularly to Paula Keane, Louise Ross, Chris Rhodes,
280 Jude Hamilton, Liz Wickens, Rebecca Hughes, Genevieve Jones, Winks Emmerson,
281 Charlie Parks, Ricky Gladwell and Geoff Sheppard. We are very grateful to farmers
282 and landowners of all other study sites for allowing access, and also to Buccleuch
283 Estates Ltd for providing accommodation during the field season in Langholm. We
284 are also thankful to Alan Fielding, Phil Whitfield and Will Cresswell for constructive
285 comments on the MS.

286

287

288

289 **REFERENCES**

290

291 **Amar, A. & Burthe, S.** 2001. Observations of predation of Hen Harrier nestlings by
292 Hooded Crows in Orkney. *Scottish Birds* **22**: 65-66.

293 **Amar, A. & Redpath, S.** 2005. Habitat use by hen harriers *Circus cyaneus* on
294 Orkney: implications of land use change on this declining population. *Ibis* **147**:
295 37-47.

296 **Amar, A., Arroyo, B., Redpath, S. & Thirgood S.** 2004. Habitat predicts losses
297 of red grouse to individual hen harriers. *Journal of Applied Ecology* **41**: 305-
298 314.

299 **Amar, A., Arroyo, B., Meek E., Redpath, S. & Riley H.** 2008. Influence of
300 habitat on breeding performance of Hen Harriers in Orkney. *Ibis* **150**: 400-
301 404.

302 **Amar, A., Redpath, S. & Thirgood, S.** 2003. Evidence for food limitation in a
303 declining raptor population. *Biological Conservation* **111**: 377-384.

304 **Amar, A., Grant, M., Buchanan G., Sim, I., Wilson, J., Pearce-Higgins, J.W. &**
305 **Redpath, S.** 2011. Exploring the relationships between wader declines and
306 current land-use in the British uplands. *Bird Study* **58**: 13-26

307 **Anderson, B.J., Arroyo, B., Collingham, Y.C., Etheridge, B., Fernandez-de-**
308 **Simon, J., Gillings, S., Gregory, R., Leckie, F., Thomas, C. D., Travis, J.**
309 **& Redpath, S.M.** 2009. Using distribution models to test alternative
310 hypotheses about a species' environmental limits and recovery prospects.
311 *Biological Conservation* **142**: 488-499.

312 **Arroyo, B., Leckie, F. & Redpath, S.** 2006. Habitat use and range management
313 on priority areas for hen harriers: final report. *Report to Scottish Natural*
314 *Heritage, Edinburgh, UK.* 57 pp.

315 **Arroyo, B., Amar, A., Leckie, F., Buchannan, G., Wilson, J. & Redpath, S.**
316 2009. Hunting habitat selection by hen harriers on moorland: implications
317 for conservation. *Biological Conservation* **142**: 586-596.

318 **Boitani, L. & Fuller, T.K.** 2000. *Research techniques in animal ecology:*
319 *controversies and consequences.* New York: Columbia University Press.

- 320 **Donazar, J.A., Negro, J.J. & Hiraldo, F.** 1993. Foraging habitat selection, land-use
 321 changes and population decline in the lesser kestrel *Falco naumanni*. *Journal of*
 322 *Applied Ecology* **30**: 515-522.
- 323 **Etheridge, B., Summers, R.W. & Green, R.E.** 1997. The effects of illegal killing
 324 and destruction of nests by humans on the population dynamics of the hen
 325 harrier *Circus cyaneus* in Scotland. *Journal of Applied Ecology* **34**: 1081-105.
- 326 **Fielding, A., Haworth, P., Whitfield, P., McLeod, D. & Riley, H.** 2011. A
 327 *Conservation Framework for Hen Harriers in the United Kingdom*. JNCC
 328 Report 441. Joint Nature Conservation Committee, Peterborough.
- 329 **García, J.T. & Arroyo, B.E.** 2005. Food-niche differentiation in sympatric Hen
 330 and Montagu's harriers. *Ibis* **147**: 144-154.
- 331 **García, J.T., Morales, M.B., Martínez, J., Iglesias L., De-la-Morena, E.G.,**
 332 **Suarez, F. & Vinuela, J.** 2006. Foraging activity and use of space by Lesser
 333 Kestrel *Falco naumanni* in relation to agrarian management in central Spain.
 334 *Bird Conservation International* **16**: 83-95
- 335 **Guixé D., & Arroyo, B.** 2011. Appropriateness of Special Protection areas for
 336 wide ranging species: the importance of scale and protecting foraging, not
 337 just nesting habitats. *Animal Conservation* **14**: 391-399
- 338 **Hayhow, D.B., Eaton, M.A., Bladwell, S., Etheridge, B., Ewing, S.R., Ruddock,**
 339 **M., Saunders, R., Sharpe, C., Sim, I.M.W. & Stevenson, A.** 2014. The status
 340 of the Hen Harrier, *Circus cyaneus*, in the UK and the Isle of Man in 2010. *Bird*
 341 *Study* **60**: 446-458.
- 342 **Hinam, H.L. & Clair, C.C.S.** 2008. High levels of habitat loss and fragmentation
 343 limit reproductive success by reducing home range size and provisioning rates
 344 of Northern saw-whet owls. *Biological Conservation* **141**: 524-535.
- 345 **Jedrzejewski, W., Schmidt, K., Theuerkauf, J., Jedrzejewska, B. & Kowalczyk,**
 346 **R.** 2007. Territory size of wolves *Canis lupus*: linking local (Bialowieza
 347 Primeval Forest, Poland) and Holarctic-scale patterns. *Ecography* **30**: 66-76
- 348 **Kenward, R.E.** 2001. *A manual for wildlife radio tagging*. Academic Press, San
 349 Diego, California.
- 350 **Madders, M. & Whitfield, D.P.** (2006). Upland raptors and the assessment of wind
 351 farm impacts. *Ibis* **148** (Suppl. 1), 43-56.
- 352 **Martin, T.G. & Possingham, H.P.** 2005. Predicting the impact of livestock grazing
 353 on birds using foraging height data. *Journal of Applied Ecology* **42**: 400-408.

- 354 **Martin, J.W.** 1987. Behaviour and habitat use of breeding Northern harriers in
355 southwestern Idaho. *Journal of Raptor Research* **21**: 57-66.
- 356 **Martínez, J.E., Pagan, I., Palazón, J.A. & Calvo, J.F.** 2007. Habitat use of Booted
357 Eagles (*Hieraaetus pennatus*) in a Special Protection Area: implications for
358 conservation. *Biodiversity and Conservation* **16**: 3481-3488.
- 359 **Newton, I.** 1979. *Population Ecology of Raptors*. Calton: T & AD Poyser.
- 360 **Picozzi, N.** 1978. Dispersion, breeding and prey of the hen harrier *Circus cyaneus* in
361 Glen Dye, Kincardineshire. *Ibis* **120**:498-509.
- 362 **Redpath, S. & Thirgood, S.** 2009. Hen harriers and red grouse: moving towards
363 consensus? *Journal of Applied Ecology* **46**: 961-963.
- 364 **Redpath, S.M.** 1992 Behavioural interactions between hen harriers and their
365 moorland prey. *Ornis Scandinavica* **23**: 73-80.
- 366 **Redpath, S.M., Madders, M., Donnelly, E., Anderson, B., Thirgood, S., Martin,**
367 **A. & McLeod, D.** 1998. Nest site selection by Hen Harriers in Scotland. *Bird*
368 *Study* **45**: 51-61.
- 369 **Redpath, S.M., Arroyo, B.E., Etheridge, B., Leckie, F, Bouwman, K. &**
370 **Thirgood, S.J.** 2002. Temperature and hen harrier productivity: from local
371 mechanisms to geographical patterns. *Ecography* **25**: 533-540.
- 372 **Rodriguez, C., Johst, K. & Bustamante, J.** 2006. How do crop types influence
373 breeding success in lesser kestrels through prey quality and availability? A
374 modelling approach. *Journal of Applied Ecology* **43**: 587-597.
- 375 **SAS Institute Inc.** 2004. SAS/STAT 9.1 User's Guide. SAS Institute Inc., Cary, NC.
- 376 **Schipper, W.J.A.** 1977. Hunting in three European harriers (*Circus*) during the
377 breeding season. *Ardea* **65**: 53-72
- 378 **Schmidt, K.** 2008. Behavioural and spatial adaptation of the Eurasian lynx to a
379 decline in prey availability. *Acta Theriologica* **53**: 1-16.
- 380 **Schoenener, T.W.** 1981. An empirically based estimate of home range. *Theoretical*
381 *Population Biology* **20**: 281-325
- 382 **Sim, I.M.W., Dillion, I.A., Eaton, M.A., Etheridge, B., Lindley, P., Riley, H.,**
383 **Saunders, R., Sharpe, C. & Tickner, M.** 2007. Status of the Hen Harrier
384 *Circus cyaneus* in th UK and the Isle of Man in 2004, and a comparison with the
385 1988/89 and 1998 surveys. *Bird Study* **54**: 256-67.

- 386 **Smith, A.A., Redpath, S.M., Campbell, S.T. & Thirgood, S.J.** 2001. Meadow
387 pipits, red grouse and the habitat characteristics of managed grouse moors.
388 *Journal of Applied Ecology* **38**: 390-400.
- 389 **Sotherton, N., Tapper, S., & Smith, A.** 2009. Hen harriers and red grouse: economic
390 aspects of red grouse shooting and the implications for moorland conservation.
391 *Journal of Applied Ecology* **46**: 955-960.
- 392 **Tella, J.L., Forero, M.G., Hiraldo, F. & Donazar, J.A.** 1998. Conflicts between
393 lesser kestrel conservation and European agricultural policies as identified by
394 habitat use analyses. *Conservation Biology* **12**: 593-604.
- 395 **Thiollay, J.M.** 2006. The decline of raptors in west Africa: long term assessment and
396 the role of protected areas. *Ibis* **148**: 240-254.
- 397 **Thirgood, S., Redpath, S., Newton, I. & Hudson, P.** 2000. Raptors and Red Grouse:
398 Conservation conflicts and management solutions. *Conservation Biology* **14**:
399 95-104.
- 400 **Thirgood, S., Redpath, S. & Graham, I.** 2003. What determines the foraging
401 distribution of raptors in heather moorland? *Oikos* **100**: 15-24.
- 402 **Thompson, P. S., Amar, A., Hoccom, D.G., Knott, J. & Wilson, J.D.** 2009.
403 Resolving the conflict between driven-grouse shooting and conservation of hen
404 harriers. *Journal of Applied Ecology* **46**: 950-954.
- 405 **Vanhinsbergh, D.P. & Chamberlain, D.E.** 2001. Habitat associations of breeding
406 Meadow Pipits *Anthus pratensis* in the British uplands. *Bird Study* **48**: 159-172.
- 407 **Whitfield, D.P. & Madders, M.** (2006). *A review of the impacts of wind farms on*
408 *hen harriers *Circus cyaneus* and an estimation of collision avoidance rates.*
409 Natural Research Information Note 1 (revised). Natural Research Ltd,
410 Banchory.
- 411 **Worton, B.J.** 1989. Kernel methods for estimating the utilization distribution in
412 home-range studies. *Ecology* **70**: 164-168.
- 413
414
415
416

417 Table 1. Home range size of the ten radio tracked hen harriers according to different
 418 methods of calculation, areas shown in km². *n* = sample size (number of fixes). MCP
 419 = Minimum Convex Polygon
 420

ID	<i>n</i>	MCP	Kernel home range estimations		
			50%	70%	90%
Langholm					
Female 257	13	3.38	0.58	1.50	5.33
Male 279	11	5.90	0.95	2.41	8.26
Orkney					
Female 115	89	11.22	1.00	1.58	4.92
Male 286	80	11.92	0.92	1.59	3.96
Male 296	59	12.70	1.71	3.24	7.59
Galloway					
Female 35	61	6.25	0.46	0.97	3.37
Female 155	34	9.50	1.09	2.46	8.23
Female 658	69	4.02	0.19	0.34	0.80
Male 233	77	36.57	2.44	4.92	13.39
Male 543	116	22.04	1.70	4.48	8.38
Average Males	Mean	17.53	1.54	3.33	8.31
	sd	12.14	0.63	1.39	3.36
Average Females	Mean	6.87	0.66	1.37	4.53
	sd	3.41	0.38	0.78	2.73


421
 422
 423

424 Table 2. Results from a General Linear Model testing for both site and sex differences
 425 in three different home range size estimators from the 10 hen harriers radio tracked in
 426 the three Scottish SPAs. Results are for the Type III (partial) tests with both sex and
 427 site fitted in each model.
 428

	df	Chi-square	<i>P</i>	Parameter estimate (mean ± se)
50% Kernel				Intercept: 1.73 ± 0.25
Sex	1	7.5	0.006	Female -0.92 ± 0.27
Site	2	1.9	0.34	Langholm 0.5 ± 0.35; Orkney -0.21 ± 0.31
70% Kernel				Intercept: 3.93 ± 0.52
Sex	1	8.9	0.003	Female -2.17 ± 0.57
Site	2	2.7	0.25	Langholm -0.89 ± 0.73; Orkney -1.08 ± 0.66
90% Kernel				Intercept: 9.40 ± 1.49
Sex	1	5.2	0.023	Female -4.28 ± 1.64
Site	2	1.6	0.44	Langholm -0.47 ± 2.11; Orkney -2.48 ± 1.89

429
 430

431 Figure 1. Frequency distribution of the distances to the nest for each fix of the radio-
432 tracked hen harrier females ($n = 272$) and males ($n = 340$) within three study areas in
433 Scotland.



434
435

Figure 2. Home ranges of the monitored birds female (left) and male (right) in relation to nest site (star) and other nests (white circles) and the limits of the SPAs (in thick lines) in Langholm.




Figure 3. Home ranges of the monitored birds in relation to nest site (star) and other nests (white circles) and the limits of the SPAs (in thick lines) in Orkney. The bottom right range corresponds to a female, the two others to males.



Figure 4. Home ranges of the monitored females (top panels) and males (bottom panels) in relation to nest site (star) and other nests (white circles) and the limits of the SPAs (in thick lines) in Galloway.

