

## Article (refereed) - postprint

---

Øverli, Øyvind; Winberg, Svante; Pottinger, Tom G. **2005 Behavioral and neuroendocrine correlates of selection for stress responsiveness in rainbow trout - a review.** *Integrative and Comparative Biology*, 45 (3). 463-474. [10.1093/icb/45.3.463](https://doi.org/10.1093/icb/45.3.463)

Copyright © 2005 [Oxford University Press](https://www.oxforduniversitypress.com/)

This version available <http://nora.nerc.ac.uk/510675/>

NERC has developed NORA to enable users to access research outputs wholly or partially funded by NERC. Copyright and other rights for material on this site are retained by the rights owners. Users should read the terms and conditions of use of this material at <http://nora.nerc.ac.uk/policies.html#access>

This is a pre-copy-editing, author-produced PDF of an article accepted for publication in *Integrative and Comparative Biology* following peer review. The definitive publisher-authenticated version Øverli, Øyvind; Winberg, Svante; Pottinger, Tom G. **2005 Behavioral and neuroendocrine correlates of selection for stress responsiveness in rainbow trout - a review.** *Integrative and Comparative Biology*, 45 (3). 463-474. [10.1093/icb/45.3.463](https://doi.org/10.1093/icb/45.3.463) is available online at: [www.icb.oxfordjournals.org](http://www.icb.oxfordjournals.org)

Contact CEH NORA team at  
[noraceh@ceh.ac.uk](mailto:noraceh@ceh.ac.uk)

1 **Behavioral and Neuroendocrine Correlates of Selection for Stress Responsiveness in**  
2 **Rainbow Trout - A Review<sup>1</sup>**

3  
4 RUNNING TITLE: SELECTION FOR STRESS RESPONSIVENESS IN TROUT

5  
6 ØYVIND ØVERLI<sup>2\*</sup>, SVANTE WINBERG<sup>†</sup>, AND TOM G. POTTINGER<sup>‡</sup>

7  
8 \* Norwegian University of Life Sciences, Dept. of Animal and Aquacultural Sciences.  
9 P.O. Box 5003, NO-1432 Ås, Norway.

10  
11 †Department of Comparative Physiology, Uppsala University, SE-75236, Uppsala,  
12 Sweden

13  
14 ‡ NERC Centre For Ecology and Hydrology, Windermere Laboratory, Far Sawrey,  
15 Ambleside, Cumbria LA22 0LP, United Kingdom

16  
17 <sup>1</sup>From the Symposium *Selection Experiments as a Tool in Evolutionary and Comparative*  
18 *Physiology: Insights into Complex Traits*

19  
20 <sup>2</sup>Corresponding author. E-mail: oyvind.overli@umb.no, phone ++ 47 64965100, fax  
21 ++47 64947960

22

23 **SYNOPSIS.**

24 In rainbow trout the magnitude of the cortisol response to stress shows both consistency  
25 over time and a moderate to high degree of heritability, and high responding (HR) and  
26 low responding (LR) lines of rainbow trout have been generated by individual selection  
27 for consistently high or low post-stress cortisol values. Using 2nd and 3rd generation fish,  
28 we tested the hypothesis that differential stress responsiveness is associated with  
29 behavioral alterations in the HR-LR trout model. LR fish showed a tendency to become  
30 socially dominant, a rapid recovery of food intake after transfer to a novel environment,  
31 and a reduced locomotor response in a territorial intrusion test. Furthermore, stress  
32 induced elevation of brain stem and optic tectum concentrations of the monoamine  
33 neurotransmitters serotonin, dopamine, and norepinephrine and their metabolites suggests  
34 that both synthesis and metabolism of these transmitters were elevated after stress to a  
35 larger degree in HR than in LR trout. A divergent pattern was seen in the hypothalamus,  
36 where LR fish displayed elevated levels of 5-hydroxyindoleacetic acid (a serotonin  
37 metabolite) and 3-methoxy-4-hydroxyphenylglycol (a norepinephrine metabolite). Thus,  
38 selection for a single trait, cortisol responsiveness, in rainbow trout is associated with  
39 concurrent changes in both behavior and central signaling systems. The apparent parallel  
40 to genetically determined stress coping styles in mammals, and the existence of similar  
41 trait associations in unselected populations of rainbow trout, suggests an evolutionarily  
42 conserved correlation between multiple traits. Continuing studies on the HR and LR trout  
43 lines are aimed at providing the physiological and genetic basis for new marker-assisted  
44 selection strategies in the rapidly developing finfish aquaculture industry, as well as  
45 increased knowledge of the function and evolution of central neuroendocrine signaling  
46 systems.

47

48 INTRODUCTION

49 The rainbow trout *Oncorhynchus mykiss* (Walbaum, 1792) is native to the Pacific coast  
50 of North America and Russia, but has been widely cultivated and introduced as a food  
51 and game fish in temperate regions around the world. Wide occurrence, availability and  
52 ease of culture have made the rainbow trout one of the most intensively studied fish  
53 species in biological research - this species has been used in fields ranging from  
54 evolutionary ecology to behavior, physiology, genetics, toxicology and cancer research  
55 (see Thorgaard et al., 2002). Rainbow trout also attracts interest as a model species for  
56 the commercially important family salmonidae, with focus on nutrition, reproduction, and  
57 stress physiology.

58 Intensive aquaculture of rainbow trout and other salmonids is a relatively new  
59 innovation, compared to the cultivation of most other animals. The process of  
60 domestication has only just begun in salmonids, and considerable effort is devoted to  
61 selecting for and defining the heritability of economically important traits in these fish,  
62 such as growth rate, age at maturity, fillet composition, and disease resistance (see e.g.  
63 Gall and Huang 1988a,b; Gjerde, 1993; Gjøen and Bentsen, 1997; Gjedrem, 2000;  
64 Midtlyng et al., 2002). Natural populations of salmonids are characteristic in displaying  
65 an immense variability in genetic composition, life history, and a range of phenotypic  
66 traits (Ryman, 1983; Taylor, 1991; Hershberger, 1992; Carlsson et al., 1999; Waples et  
67 al., 2001; Hansen et al., 2002), which makes these fish make excellent raw material for  
68 genetic studies and selection experiments.

69 Fish under intensive culture conditions are exposed to a regime of acute and  
70 chronic stressors, which have adverse effects on growth, reproduction,

71 immunocompetence, and flesh quality, among other things (Barton et al., 1987; Maule et  
72 al., 1989; Barton and Iwama, 1991; Lowe et al., 1993; Pickering, 1993; Balm, 1997;  
73 Pankhurst and Van der Kraak, 1997; Sigholt et al., 1997; Schreck et al., 2001). As in  
74 other vertebrates, the blood concentration of corticosteroid hormones is a major index of  
75 stress in fish, and elevated levels of these hormones arise from activation of the  
76 hypothalamus-pituitary-interrenal (HPI) axis (Wendelaar-Bonga, 1997). The main  
77 corticosteroid in teleost fish is cortisol (Kime, 1987), and this steroid is a causal factor in  
78 many of the deleterious effects of stress (Barton et al., 1987; Barton and Iwama, 1991;  
79 Harris and Bird, 2000; Pankhurst and Van der Kraak, 2000; Schreck et al., 2001; Consten  
80 et al., 2002; Bernier et al., 2004). In rainbow trout the magnitude of the cortisol response  
81 to stress is an individual characteristic which is stable over time, with a moderate to high  
82 degree of heritability (Pottinger et al., 1992, 1994; Fevolden et al., 1999). Thus, post-  
83 stress cortisol levels provide a trait of functional and economical significance upon which  
84 selection pressure can be directed, and lines of high- (HR) and low-responsive (LR) trout  
85 have been established at the Windermere laboratory of the UK Centre for Ecology and  
86 Hydrology (Pottinger and Carrick, 1999, 2001a).

87         In association with the physiological response, stress may lead to drastic changes  
88 in behavior, which has been classified and interpreted in various ways (see e.g. Koolhaas  
89 et al., 1999; Wingfield, 2003). A distinction is often made between proactive (active  
90 coping, or ‘fight or flight’) and reactive (passive coping, or ‘conservation-withdrawal’)  
91 responses (Engel and Schmale, 1972; Henry and Stephens, 1977; Benus et al., 1991;  
92 Henry, 1993; Koolhaas et al., 1999). In both mammals and fish, a transition from  
93 behavioral activation to an inhibiting effect is often seen with increasing duration or

94 severity of the challenge (Haller et al., 1998; Øverli et al., 2004a). Seminal studies in  
95 mammals indicated that the threshold at which the shift from an active to a passive  
96 behavioral response occurs is subject to great individual variation (Engel and Schmale,  
97 1972; Henry and Stephens, 1977).

98         More recently, it has been established that individual differences in the  
99 physiological stress response are associated with differences in behavior. For instance, a  
100 pro-active stress coping style is behaviorally characterized by a high level of active  
101 avoidance, aggression, and other behavioral patterns indicating an active attempt to  
102 counteract the stressful stimulus (e.g. defensive burying in rodents) (Bohus et al., 1987;  
103 Benus et al., 1989, 1991a; Sluyter et al., 1996). Reactive (passive) coping, on the other  
104 hand, involves immobility and low levels of aggression. Physiologically, the pro-active  
105 strategy is associated with low hypothalamus-pituitary-adrenal axis (HPA axis, the  
106 mammalian equivalent of the teleost HPI axis) responsiveness, but high sympathetic  
107 reactivity, while the opposite is true for reactive coping (de Boer et al., 1990; Korte et al.,  
108 1992; Fokkema et al., 1995). A genetic basis for the expression of behavioral and  
109 physiological components of individual coping styles has repeatedly been demonstrated  
110 (e.g. Driscoll et al., 1998; Ellenbroek and Cools, 2002; de Boer et al., 2003; Veenema et  
111 al., 2003). Limited information, however, is available on whether different behavioral-  
112 physiological stress coping styles are present in teleost fish (but see Francis, 1990; Van  
113 Raaij et al., 1996; Budaev et al., 1999).

114         In fish, like in other vertebrates, behavioral and physiological stress responses are  
115 to a large degree linked by common control mechanisms in the brain, and the monoamine  
116 neurotransmitters serotonin (5-hydroxytryptamine, 5-HT), dopamine (DA), and

117 norepinephrine (NE) play a vital role in this co-ordination (Winberg and Nilsson, 1993;  
118 Winberg et al., 1997, 2001; Øverli et al., 1998, 1999; Höglund et al., 2001, 2002a,b;  
119 Lepage et al., 2002, 2003; Clements et al., 2003; Larson et al., 2003; Perreault et al.,  
120 2003). Genetically determined variation in behavior and stress responsiveness has been  
121 associated with differences in brain monoaminergic function in both fish and mammals  
122 (e.g. Benus et al., 1991b; Popova et al., 1991a,b; Nikulina et al., 1992; Rots et al.,  
123 1996a,b,c; de Kloet et al., 1996; Sallinen et al., 1999; Lepage et al., 2000; Giorgi et al.,  
124 2003). In this paper we review the effects of selection for post stress cortisol production  
125 in rainbow trout, with particular reference to a series of studies investigating the  
126 association between altered plasma cortisol dynamics, behavior, and brain monoamine  
127 neurotransmitters. In other words, we examine the hypothesis that behavioral and  
128 physiological traits are linked in such a way that altering one trait, post-stress plasma  
129 cortisol concentrations, incurs differences in other putative components of individual  
130 stress coping styles. It should be noted that several additional physiological differences  
131 between HR and LR lines, such as metabolic changes and interrenal sensitivity to  
132 adenocorticotrophic hormone (ACTH) also has been reported, but for details of this  
133 experimental work we refer to the original papers (Pottinger and Carrick, 2001b,  
134 Trenzado et al., 2003).

135

136

137

## MATERIAL AND METHODS

138 *Generation of HR and LR trout lines*

139 The selection procedure and the effect of the breeding program on the cortisol response  
140 has been thoroughly described elsewhere (Pottinger and Carrick, 1999, 2001a), and will  
141 only be briefly reviewed here. The parental generation of the HR and LR trout lines was  
142 established in 1996 by repeated stress testing (3h confinement in 50 L water in groups of  
143 6-7 individuals once monthly) of passive integrated transponder (PIT) tagged 2-year-old  
144 rainbow trout. Following confinement, blood samples (0.5 ml) and PIT-tag readings were  
145 obtained from anaesthetized (2-phenoxyethanol, 1:2000) fish, and plasma was later  
146 analyzed for cortisol content by a previously validated radioimmunoassay (RIA)  
147 procedure (Pickering et al., 1987). The mean post-stress plasma cortisol content across  
148 five episodes of confinement was then calculated for each fish, and individuals were  
149 ranked accordingly. Between testing fish had been kept in groups of 25 in 1500 L holding  
150 tanks, and the four most high-responding (HR) and the four most low-responding (LR)  
151 fish in each tank were removed from their home tank, segregated by sex, and kept  
152 separate in 4 tanks based on group and sex until maturation.

153         Confinement stress testing of the 1st generation offspring (F1), consisting of 15  
154 HR and 14 LR families each resulting from a unique male-female crossing, were carried  
155 out on five different occasions between September 1997 and September 1998, and on five  
156 occasions in 1999. A highly significant regression of mid-parent cortisol response on  
157 progeny response was seen (estimated  $r^2$  [h<sup>2</sup>] value = 0.41) and the six LR families with  
158 the lowest mean cortisol response and the six HR families with the highest mean cortisol  
159 response were identified and used for further work (Pottinger and Carrick, 1999). An  
160 unselected (US) population resulting from random breeding of fish not designated as HR  
161 or LR in the parental group (6 families) was tested on one occasion, and showed a



162 cortisol response intermediate to the mean of these most divergent HR and LR groups.  
163 Adult female F1 generation HR and LR fish were later used in a study investigating the  
164 effect of selection for stress responsiveness on behavior and brain monoamine  
165 neurotransmitters (Øverli et al., 2001, 2002a).

166 The 2nd generation of offspring consisted of 11 HR, 11 LR and 3 US families.  
167 The regression of midparent [(male + female)/2] cortisol response on progeny cortisol  
168 response provided an estimated  $h^2$  of 0.6 in this generation, while male and female  
169 parent–progeny regressions provided estimates for  $h^2$  of 0.73 and 0.44, respectively  
170 (Pottinger and Carrick, 2001a).

171

## 172 *Behavioral experiments*

### 173 Locomotor activity, feed intake, and brain monoaminergic activity in HR and LR trout:

174 These experiments are described in detail in Øverli et al. (2001, 2002a). Recovery of feed  
175 intake after transfer to a new and potentially stressful environment was used as an index  
176 of adaptiveness. Only adult female F1 fish were available for these studies. Locomotor  
177 activity when in isolation and in response to a territorial intruder was also analyzed, along  
178 with basal and stress-induced brain 5-HT, DA and NE activity. During April 2000 adult  
179 F1 female HR (n=18) and LR (n=18) rainbow trout weighing  $987.5 \pm 39.5$  g (mean  $\pm$  SE)  
180 were transferred individually from communal holding tanks to rearing in isolation in  
181 white 250 L polypropylene observation tanks. From day 1 after transfer to rearing in  
182 isolation, fish were fed daily by hand (1.5% of body weight), and fish were observed for  
183 3 min after distribution of food to register food intake. Behavioral observations and blood  
184 sampling (see below) were carried out between 10.00 and 14.00.

185           After being held for 6 days in isolation, locomotor activity was quantified by  
186 observing time spent moving during 20 min for each fish, starting 1 h after feeding.  
187 Locomotor activity was quantified again the next day, this time for 20 min immediately  
188 following the introduction of an intruder fish in the observation tank. Intruder fish were  
189 smaller (< 50% body weight of the resident fish) group reared rainbow trout from a  
190 hatchery population, and previously unfamiliar to the test fish.

191           On the day after the intruder test, 50% of the fish from each line (HR, LR) were  
192 randomly selected for individual stress testing in 50 L confinement tanks. After 1h in the  
193 confinement tanks, fish were netted, anaesthetized in 0.5 ml/L 2-phenoxyethanol, and a  
194 blood sample was obtained from the caudal sinus into a heparinized syringe. The  
195 remaining 50% of the fish were sampled directly from observation tanks to serve as  
196 undisturbed controls. Immediately following blood sampling fish were killed by  
197 decapitation and dissected, and the presence or absence of food in the stomach and / or  
198 intestines was registered. Brains were removed and dissected into telencephalon  
199 (excluding the olfactory bulbs), hypothalamus (excluding the pituitary), optic tectum, and  
200 brain stem (excluding the cerebellum). Brain samples were immediately wrapped in  
201 aluminum foil and snap frozen in liquid nitrogen, where after concentrations of  
202 monoamines and monoamine metabolites in brain samples were analyzed by HPLC with  
203 electrochemical detection (Øverli et al., 1999). Plasma cortisol levels were determined  
204 using the RIA procedure described by Pickering et al. (1987).

205           Tests for social dominance in HR and LR trout: Tests to investigate the relative  
206 tendency toward dominant or subordinate social status of the HR and LR lines were  
207 carried out in September and October 2000 using juvenile F2 progeny (Pottinger and

208 Carrick, 2001a). Mixed-sex juvenile fish were used in these tests since they are generally  
209 more territorial than adults, and there is no effect of sex on the ability to gain dominance  
210 at this life stage (Ø. Øverli and S. Winberg, unpublished results). Dyadic contests were  
211 performed following broadly the protocol of Øverli et al. (1999). In all, 47 size matched  
212 pairs consisting of 1 HR and 1 LR individual were transferred from holding tanks to  
213 rearing in glass observation aquaria (63 L volume) covered with black plastic on three  
214 sides. In each aquarium two fish were kept separate by a removable opaque plastic  
215 barrier.

216 Fish were offered food (five crumbs, Trouw Fry 18) twice each day after being  
217 transferred to the aquaria. Previous experiments have shown that holding juvenile  
218 salmonid fish in isolation for approximately one week with sufficient access to food is  
219 highly effective in inducing territoriality and motivation to express aggressive behavior  
220 (Winberg et al., 1991; Winberg and Lepage, 1998; Øverli et al., 1999). Barriers  
221 separating HR/LR pairs were removed after 5 days of acclimation, resulting in escalated  
222 contests for social dominance within each pair. Behavioral observations started at 3 h  
223 after the onset of interaction, by which time the conflict was resolved and the identity of  
224 the dominant and subordinate individuals within each pair was clear. At this time  
225 dominant fish were characterized by holding a midwater position, displaying extensive  
226 movement around the tank, nipping and/or chasing the subordinate, and intercepting  
227 food. Subordinate fish were typically located on the base of the aquarium, usually  
228 immediately adjacent to the aquarium wall, exhibited little swimming activity, and  
229 directed no aggressive acts toward the dominant individual. After 5 h of social interaction

230 fish were killed and, a blood sample was obtained, and blood plasma was analyzed for  
231 cortisol content using a RIA with ethyl acetate extraction (Pottinger and Carrick, 2001a).

232

## 233 RESULTS AND DISCUSSION

### 234 *Cortisol responsiveness in HR and LR rainbow trout*

235 Post-stress blood plasma cortisol levels of HR and LR trout for every occasion on which  
236 they were tested, from the establishment of the parental generation (F0) up to present  
237 (F3), is shown in Figure 1. There is considerable variation in the magnitude of the cortisol  
238 response over time, and between generations, but a difference between the lines  
239 (HR>LR) is always evident. The exception to this is the final sample for the F0 fish in  
240 which reproductive status may have confounded the normal previously observed  
241 divergence. Some of the variation in overall magnitude of the stress response can no  
242 doubt be accounted for by seasonal changes in water temperature (Sumpter et al., 1985;  
243 Barton and Schreck, 1987; Pickering and Pottinger, 1987) and in reproductive status of  
244 the fish (Pottinger et al., 1995; Pottinger and Carrick, 2000). However, it must be borne  
245 in mind that these tests were carried out for a variety of purposes and did not employ  
246 exactly the same protocol on each occasion. Therefore, while it is appropriate to conclude  
247 that divergence in stress responsiveness has been sustained across three generations, it is  
248 not possible to directly compare successive time points and evaluate temporal trends.

249 Notwithstanding this caveat, the type of test that were employed for the first  
250 occasion on which each generation was assessed were similar and the results of these  
251 tests may be directly compared. The magnitude of the difference in cortisol  
252 responsiveness between lines has remained similar over 3 successive generations for

253 juvenile fish subjected to their first stress test (age 6-7 months, tested in September 1997,  
254 September 2000 and October 2003). The post-stress plasma cortisol levels in HR  
255 juveniles represents 135%, 224%, and 220% that of the LR fish for the F1-F3 generations  
256 respectively. The similarity in the magnitude of divergence of these generations might  
257 indicate that there are physiological constraints on the range of plasma cortisol  
258 responsiveness to stress in trout. Such constraints may be imposed upon low-responders  
259 by the need to retain a functional cortisol response and upon high-responders by problems  
260 associated with hypercortisolism.

261 Finally, it should be noted that strain differences in plasma cortisol levels was  
262 never observed in unstressed fish. Thus, it seems likely that the effect of the selection  
263 program on post-stress levels of cortisol represents divergent responses to stressors,  
264 rather than differences in basal HPI-axis activity, which in turn could affect  
265 responsiveness.

266

#### 267 *Behavioral effects of selection for stress responsiveness*

268 Pottinger and Carrick (2001a) reported a tendency of LR fish to establish social  
269 dominance over HR fish when held in pairs (Figure 2), and Øverli et al. (2001) observed  
270 that only LR fish had regained feed intake within one week after transfer to a new  
271 environment (Figure 3). Interestingly, an association between rapidly regaining feed  
272 intake after environmental change and the ability to win fights for social dominance was  
273 also observed in an unselected population of rainbow trout, using juvenile fish of both  
274 sexes (Øverli et al., 2004b). These observations indicate that the HR and LR lines are  
275 based on multi-trait phenotypes that display a significant amount of variation also in

276 unselected populations, but it has not yet been studied whether the divergence in behavior  
277 has increased with successive generations.

278           Locomotor activity in isolated HR and LR female rainbow trout, when held alone  
279 and when challenged with a conspecific intruder, is depicted in Figure 4. HR rainbow  
280 trout displayed higher locomotor activity than LR trout in the presence of an intruder, but  
281 there was no significant difference between the two groups in the isolated condition.  
282 However, both HR and LR rainbow trout increased their activity level when the intruder  
283 was present. Thus, it appears that some behavioral differences between HR and LR lines  
284 occur only under an acute challenge, which may indicate that they depend on control  
285 mechanisms activated in synchronization with the physiological stress response.

286

#### 287 *Brain monoaminergic activity in HR and LR trout*

288 Concentrations of the three monoamine neurotransmitters 5-HT, DA, and NE and their  
289 respective metabolites 5-hydroxyindoleacetic acid (5-HIAA), 3,4-dihydroxyphenylacetic  
290 acid (DOPAC), and 3-methoxy-4-hydroxyphenylglycol (MHPG), and corresponding  
291 metabolite/monoamine ratios in four different brain regions of stressed and control HR  
292 and LR trout are shown in Table I. Since monoamine neurotransmitters are not exposed  
293 to monoamine oxidase (MAO) while stored in vesicles, increased concentrations of their  
294 deaminated metabolites are thought to indicate increased release and turnover of the  
295 neurotransmitter (Fillenz, 1993).

296           In particular, metabolite/monoamine ratios are frequently used as an index of  
297 neural activity. However, attention should be paid to the fact that altered metabolite /  
298 monoamine ratios may be caused by changes in the concentrations of monoamine

299 neurotransmitters as well as metabolites, or by combinations of such changes. Thus,  
300 differential patterns of activation may be assumed depending on whether concentrations  
301 or ratios are studied. For instance, brain stem 5-HIAA concentrations were significantly  
302 affected by confinement stress only in HR fish (Table I). Therefore, it could be concluded  
303 that the brain 5-HT system was activated by confinement stress to a larger extent in HR  
304 than in LR fish. On the other hand, brain stem 5-HIAA/5-HT ratios were higher in LR  
305 than HR fish both in the control condition and after stress. This apparent contradiction is  
306 probably caused by the fact that HR fish, but not LR fish, responded to stress by an  
307 increase in 5-HT concentrations in the brain stem. Apart from that, three possible  
308 explanations can be given to the observation that 5-HIAA/5-HT ratios were elevated in  
309 LR fish relative to HR fish: 1. The proportion of 5-HT that was actually released, and  
310 thereby exposed to the action of MAO after re-uptake from the intercellular space, was  
311 greater in LR fish. 2. MAO enzyme activity was decreased in HR fish. 3. Re-uptake of 5-  
312 HT was more effective in LR fish.

313         Nonetheless, one of the most evident findings of this study was that HR trout  
314 reacted to stress by an increase in the tissue concentrations of serotonin (brain stem),  
315 dopamine (brain stem), and norepinephrine (optic tectum, telencephalon), whereas low-  
316 responsive fish did not (Table I). Brain stem and optic tectum concentrations of  
317 monoamine metabolites were also elevated after stress in HR, but not in LR fish. A  
318 divergent pattern was seen in the hypothalamus, where LR fish displayed elevated levels  
319 of 5-HIAA and MHPG. Both populations had elevated telencephalic concentrations of  
320 these metabolites after stress. Some differences were also seen in fish sampled directly  
321 from rearing in isolation, suggesting that the rearing environment was not entirely

322 optimal, and the experimental fish may have experienced a mild stress even in the  
323 undisturbed condition. This notion is also in line with the lack of aggressive behavior and  
324 low level of feed intake in these fish (Øverli et al., 2002a). Alternatively, differential  
325 stress responsiveness in HR and LR fish is associated with permanent differences in brain  
326 monoaminergic systems that are expressed even in unstressed individuals.

327

328 *Correlated physiological and behavioral stress responses in HR and LR trout lines*

329 The creation of HR and LR lines of rainbow trout confirm that the magnitude of the  
330 cortisol response to a standardized stressor is an individual, heritable characteristic in this  
331 species (Pottinger et al., 1992, 1994; Fevolden et al., 1999). Furthermore, it appears that  
332 the magnitude of stress-induced elevation of blood cortisol is part of a complex trait  
333 incorporating several correlated physiological and behavioral responses (Pottinger and  
334 Carrick 2001a,b; Øverli et al., 2001, 2002; Trenzado et al., 2003). Taken together, these  
335 observations suggest that the HR and LR rainbow trout may represent selection for  
336 different physiological/behavioral stress-coping styles, as defined by Koolhaas et al.  
337 (1999). These authors stated: “*A coping style can be defined as a coherent set of*  
338 *behavioral and physiological stress responses which is consistent over time and which is*  
339 *characteristic to a certain group of individuals.*”

340 As noted previously, the plasma cortisol response is an individual trait which is  
341 consistent over time in rainbow trout (Pottinger et al., 1992; Pottinger and Carrick, 1999).  
342 The degree to which behavioral traits are consistent in individual HR and LR fish has not  
343 been determined, but comparisons of behavior between the two lines have revealed  
344 several striking differences (*c.f.* figure 2-4). Behavioral and physiological stress responses



345 are to a large degree controlled by common neuroendocrine signaling systems. Thus, if  
346 the cortisol response is a consistent individual trait, it seems likely permanent differences  
347 exist with respect to the behavioral components of different stress coping styles.

348         Most likely, some degree of trait associations exist in all animals with respect to a  
349 fundamental response such as stress coping, but the evolutionary success of different  
350 profiles may vary between species. It is not clear whether the behavioral and endocrine  
351 traits of LR and HR rainbow correspond exactly to the proactive (active) and reactive  
352 (passive) coping styles described in mammals. The proactive stress coping style in  
353 rodents involves low glucocorticoid production (de Boer et al., 1990; Korte et al., 1992;  
354 Fokkema et al., 1995), while in HR trout high cortisol responsiveness was associated with  
355 increased swimming activity during an intruder test (figure 4). The impression of the  
356 observers was that the HR trout showed anxiety-like erratic behavior, rather than targeted  
357 attempts to actively cope with the experimental situation. If that is the case, the  
358 behavioral strategy of the LR fish (to remain passive and ignore the presence of an  
359 intruder in a low-quality territory) may be more equivalent to the active coping style that  
360 is normally associated with low HPA axis responsiveness in rats (Koolhaas et al., 1999).  
361 Furthermore, the observation that LR trout tend to win encounters for social dominance  
362 (figure 2) is in accordance with an active coping style (see e.g. Verbeek et al., 1996;  
363 Klomberg et al., 2002; Zhukov and Vinogradova, 2002).

364         Several rodent models consisting of two contrasting lines/strains that respond  
365 differently to stressful environments have also been developed through bidirectional  
366 selection. Examples are the Roman high (RHA) and low (RLA) avoidance rats (Driscoll  
367 et al., 1998; Steimer and Driscoll, 2003), the Wistar high (HAB) and low (LAB) anxiety-

368 related behavior lines (Liebsch et al., 1998; Yilmazer-Hanke et al., 2004), the Maudsley  
369 reactive and nonreactive strains (Blizard and Adams, 2002), the short (SAL) and long  
370 attack latency (LAL) house mice (Veenema et al., 2003), apomorphine susceptible and  
371 unsusceptible rats (Ellenbroek and Cools, 2002), and several others (e.g. Klomberg et al.,  
372 2002; Viggiano et al., 2002, 2003; Brush, 2003; Ramos et al., 2003).

373         Like the HR and LR rainbow trout, these models were created by selective mating  
374 of animals with the highest and lowest scores for a given behavioral or physiological trait  
375 over several generations. In general, it is assumed that maximizing differences in this way  
376 produces one line with more genes that affect the selected trait positively, and one line  
377 carrying more genes with negative effects on the same trait (Ramos and Mormède, 1998;  
378 Crabbe, 1999). Ideally, correlated responses should be consistent across several replicate  
379 lines to indicate the presence of genetic correlations (Henderson, 1997). Thus, in the case  
380 of the HR and LR trout lines, founder effects, unique mutations and random genetic drift  
381 can not be ruled out as contributing factors in the simultaneous divergence of multiple  
382 traits. However, the apparent parallel to genetically determined stress coping styles in  
383 mammals, and the existence of similar trait associations in unselected populations (Øverli  
384 et al., 2004b), suggest an evolutionarily conserved correlation between multiple traits.

385

### 386 *Neuroendocrine mechanisms integrating physiological and behavioral stress responses*

387 The mechanisms integrating the behavioral and physiological characteristics of HR and  
388 LR trout lines remain largely unknown. The behavioral differences between HR and LR  
389 trout are consistent with some reported effects of corticosteroid hormones in  
390 poikilotherms (decreased appetite: Gregory and Wood, 1999; increased locomotor

391 activity: Cash and Holberton, 1999; Øverli et al., 2002b). In fish, like in mammals, these  
392 steroids typically have time-, context-, and dose-dependent effects (Øverli et al., 2002b;  
393 Bernier et al., 2004). Thus, altered competitive ability in a stressful situation such as a  
394 fight for social dominance (Pottinger and Carrick, 2001a) may also be directly caused by  
395 differences in circulating hormone levels.

396         The behavioral and physiological characteristics of HR and LR rainbow trout may  
397 also be functionally linked through a number of factors which influence both endocrine  
398 and behavioral responses. For instance, it seems likely that corticotrophin releasing  
399 hormone (CRH) is involved in the increase in locomotor activity observed in HR trout. In  
400 juvenile chinook salmon (*Oncorhynchus tshawytscha*), intracerebroventricular injections  
401 of CRH induced hyperactivity, an effect that was shown to depend on concurrent (i.e.  
402 CRH induced) 5-HT activation (Clements et al., 2003).

403         Interestingly, CRH administration also increases DA concentrations in dorsal  
404 medial hypothalamus of newts (*Taricha granulosa*) (Lowry et al., 2001), and one of the  
405 main neurochemical differences between HR and LR rainbow trout was that HR fish  
406 responded to stress by increased DA concentrations and turnover in several brain areas,  
407 while LR fish did not (Table I). Genetically determined differences in DA systems has  
408 been reported in several other models (Rots et al., 1996a,b,c; Lecca et al., 2004).  
409 However, increased DA synthesis and release may also be an effect of acute elevations in  
410 glucocorticoid concentrations (Dunn et al., 1978; Piazza et al., 1996a,b; Barrot et al.,  
411 2000, 2001). Effects of glucocorticoids on dopaminergic activity are also strongly context  
412 (Piazza et al., 1996b) and regionally dependent (Lucas et al., 1998; Barrot et al., 2000,  
413 2001). Thus, at present it is not known whether differences in DA systems between HR

414 and LR trout are a cause or a consequence of hormone dynamics, but there is an  
415 interesting parallel to mammalian models which suggest the presence of evolutionary  
416 conserved trait correlations. Notably, a similar negative relationship between DA  
417 reactivity, stressor or novelty-induced locomotor activity, and social competitive ability  
418 has also been demonstrated in cynomolgus monkeys (*Macaca fascicularis*) (Morgan et  
419 al., 2000). Individually housed monkeys with high levels of locomotion in an open-field  
420 test after a low dose of cocaine (which increases CNS dopamine) were more likely to  
421 become subordinate in subsequent group housing (Morgan et al., 2000).

422         Like with DA, corticosteroids may affect brain 5-HT signaling directly and  
423 through interaction with other neurotransmitter systems (Chaouloff, 2000). Inheritable  
424 properties of the 5-HT system have also been associated with HPA-axis activity,  
425 personality and mood alterations in human and other animals. Examples are differences  
426 in MAO and polymorphisms in the 5-HT transporter gene or promoter region (Lesch et  
427 al., 1996; Shih et al., 1999; Fernandez et al., 2003).

428         The interaction between different signaling systems involved in the stress  
429 response is, in fact, so complex that on occasions it appears futile to disentangle causes  
430 and consequences. Most research on neuroendocrine control of behavioral and  
431 physiological stress responses has been carried out on mammals. However, the  
432 complexity of the interaction between a single neurotransmitter, 5-HT, and stress is well  
433 illustrated by a series of studies with salmonid fish: Stress affects 5-HT metabolism and  
434 most likely also functional release (Winberg and Nilsson, 1993; Øverli et al., 1998, 1999,  
435 2001; Lepage et al., 2002) and 5-HT in turn affects behavior (Winberg et al., 1993, 2001)  
436 and cortisol release (Winberg et al., 1997; Höglund et al., 2002b; Lepage et al.,

437 2002,2003). Cortisol also affects behavior (Øverli et al., 2002), while both stressful and  
438 rewarding behavior influence 5-HT as well as stress hormones (Winberg and Lepage  
439 1997; Øverli et al., 1999, 2004a). Finally, there is preliminary evidence that circulating  
440 cortisol act on brain 5-HT neurons and can modify neurotransmission also in fish (Øverli  
441 et al., 2003).

442 Thus, it is not surprising that selection for stress responsiveness is also associated  
443 with alterations in brain 5-HT activity (Øverli et al., 2001, Table I). However, at present  
444 it is not known to what degree the physiological and behavioral profiles of HR and LR  
445 trout are a result of innate differences in central signaling systems, of glucocorticoid  
446 influence on brain function, or a combination of these factors. Finally, it should be kept  
447 in mind that the differences in stress induced cortisol concentrations between HR and LR  
448 fish might arise from differences in interrenal function, rather than in central HPI-axis  
449 control (Pottinger and Carrick 2001b). Similarly, seasonal variations in stress-induced  
450 plasma corticosteroid levels was correlated to alterations in adrenocortical cell  
451 steroidogenic function in lizards (*Sceloporus undulatus*) (Carsia and John-Alder, 2003).  
452 Thus, the possibility should be considered that differences in neurochemistry and  
453 behavior between HR and LR trout are a result of variation in interrenal influence on  
454 brain functions, rather than the opposite. However, differences between unstressed HR  
455 and LR lines also have been observed, and in these cases interrenal function are unlikely  
456 to be directly responsible.

457

458 *Further experiments with HR and LR trout lines*

459 In summary, experiments carried out on the HR and LR trout lines demonstrate a tight  
460 coupling of physiological and behavioral stress responses. The neuroendocrine control  
461 mechanisms behind this connection appear to be evolutionarily conserved, and are well  
462 illustrated by genetically selected strains of animals that display simultaneous differences  
463 in physiology and behavior. In many ways, selection models are superior to targeted gene  
464 knock-outs, since an entire suite of neuroendocrine mechanisms are selected for, instead  
465 of altering just one specific gene product. The systems that produce behavioral and  
466 neuroendocrine stress responses work in an integrated fashion, and selection models keep  
467 that integration intact.

468 Continuing studies on the 3rd generation HR and LR trout lines are providing  
469 evidence that the range of behavioral traits in which there are pronounced differences  
470 between the two lines far exceeds what has so far been published. For instance, a recent  
471 study has demonstrated that the extinction of a conditioned response occurs more rapidly  
472 among fish from the HR line than fish from the LR line, suggesting that the two lines  
473 differ in cognitive function as well as in behavioral characteristics (Moreira et al., 2004).  
474 It is, however, not known whether this result depends on differences in memory  
475 formation, consolidation, or retrieval. This latter point is of particular interest, since some  
476 studies suggest dual effects of glucocorticoids on specific memory phases (Rooszendaal,  
477 2002). Furthermore, the results of Moreira et al. (2004) suggest that densities and  
478 composition of *N*-methyl-D-aspartate (NMDA) are also likely to differ between the lines,  
479 in view of the important role of these receptors in excitatory neurotransmission, synaptic  
480 plasticity, and cognition in other vertebrates.

481           One neurotransmitter system that is also likely to play a central role in shaping  
482 behavioral profiles, but that has not been studied in the HR/LR trout model, is CRH. It is,  
483 for instance, not known whether basal or stress-induced CRH activity differs between the  
484 lines, although their behavioral and physiological profile strongly suggests such a  
485 difference. As suggested by Clements et al. (2003), it would also be interesting to  
486 compare the responses of these two lines to CRH treatment.

487           In a different venue of research, recently developed genomic tools, such as  
488 microarray technology, are being utilized in an EU program (STRESSGENES) to  
489 identify candidate genes associated with resistance to stress. Knowledge of the genes  
490 responsible for trait variability will further point out the pathways responsible for the  
491 phenotypical differences between HR and LR lines. This should provide the  
492 physiological and genetic basis for new marker-assisted selection strategies in the rapidly  
493 developing finfish aquaculture industry, as well as increased knowledge of the function  
494 and evolution of central neuroendocrine signaling systems.

495

496

#### ACKNOWLEDGEMENTS

497 TGP was supported by the Natural Environment Research Council (U.K.) and the  
498 European Community (Contract FAIR-CT95-0152). ØØ and SW received support from  
499 Norwegian (NFR) and Swedish (SJFR/FORMAS) research councils.

500

501

502 REFERENCES

- 503 Balm, P.H.M. 1997. Immune-endocrine interactions. *In* G. K. Iwama, A. D. Pickering,  
504 J. P. Sumpter, and C. B. Schreck (eds.), *Fish stress and health in aquaculture*, pp.  
505 195-221. Cambridge University Press, Cambridge.
- 506 Barrot, M., M. Marinelli, D. N. Abrous, F. Rougé-Pont, M. Le Moal, and P. V. Piazza.  
507 2000. The dopaminergic hyper-responsiveness of the shell of the nucleus  
508 accumbens is hormone-dependent. *Eur. J. Neurosci.* 12:973-979.
- 509 Barrot, M., D. N. Abrous, M. Marinelli, F. Rougé-Pont, M. Le Moal, and P. V. Piazza.  
510 2001. Influence of glucocorticoids on dopaminergic transmission in the rat  
511 dorsolateral striatum. *Eur. J. Neurosci.* 13:812-818.
- 512 Barton, B.A. and C. B. Schreck. 1987. Influence of acclimation temperature on interrenal  
513 and carbohydrate stress responses in juvenile Chinook salmon (*Oncorhynchus*  
514 *tshawytscha*). *Aquaculture* 62:299-310.
- 515 Barton, B. A., C. B. Schreck, and L. D. Barton. 1987. Effects of chronic cortisol  
516 administration and daily acute stress on growth, physiological conditions, and  
517 stress responses in juvenile rainbow-trout. *Dis. Aquat. Organ.* 2:173-185.
- 518 Barton, B. A., and G. K. Iwama. 1991. Physiological changes in fish from stress in  
519 aquaculture with emphasis on the response and effects of corticosteroids. *Ann.*  
520 *Rev. Fish Diseases* 1:3-26.
- 521 Benus, R. F., B. Bohus, J. M. Koolhaas, and G. A. van Oortmerssen. 1989. Behavioral  
522 strategies of aggressive and nonaggressive male-mice in active shock avoidance.  
523 *Behav. Process.* 20:1-12.



524 Benus, R. F., B. Bohus, J. M. Koolhaas, and G. A. van Oortmerssen. 1991a. Heritable  
525 variation for aggression as a reflection of individual coping strategies. *Experientia*  
526 47:1008-1019.

527 Benus, R. F., B. Bohus, J. M. Koolhaas, and G. A. van Oortmerssen. 1991b. Behavioral-  
528 differences between artificially selected aggressive and nonaggressive mice:  
529 response to apomorphine. *Behav. Brain Res.* 43:203-208.

530 Bernier, N. J., N. Bedard, and R. E. Peter. 2004. Effects of cortisol on food intake,  
531 growth, and forebrain neuropeptide Y and corticotropin-releasing factor gene  
532 expression in goldfish. *Gen. Comp. Endocrinol.* 135:230-240.

533 Blizard, D. A. and N. Adams. 2002. The maudslay reactive and nonreactive strains: A  
534 new perspective. *Behav. Genet.* 32:277-299.

535 Bohus, B., R. F. Benus, D.S. Fokkema, J. M. Koolhaas, C. Nyakas, G. A. Van  
536 Oortmerssen, A. J. A. Prins, A. J. H. de Ruitter, A. J. W. Scheurink, and A. B.  
537 Steffens. 1987. Neuroendocrine states and behavioral and physiological stress  
538 responses. *Prog. Brain Res.* 72: 57-70.

539 Brush, F. R. 2003. The syracuse strains, selectively bred for differences in active  
540 avoidance learning, may be models of genetic differences in trait and state  
541 anxiety. *Stress.* 6:77-85.

542 Budaev, S. V., D. D. Zworykin, and A. D. Mochek. 1999. Consistency of individual  
543 differences in behaviour of the lion-headed cichlid, *Steatocranus casuarius*.  
544 *Behav. Process.* 48:49-55.

545 Carlsson, J., K. H. Olsén, J. Nilsson, Ø. Øverli, and O. B. Stabell. 1999. Microsatellites  
546 reveal fine-scale genetic structure in stream-living brown trout. *J. Fish Biol.*  
547 55:1290-1303.

548 Carsia, R. V. and H. John-Alder. 2003. Seasonal alterations in adrenocortical cell  
549 function associated with stress responsiveness and sex in the eastern fence lizard  
550 (*Sceloporus undulatus*). *Horm. Behav.* 43:408-420.

551 Cash, W. B. and R. L. Holberton. 1999. Effects of exogenous corticosterone on  
552 locomotor activity in the red-eared slider turtle, *Trachemys scripta elegans*.  
553 *J. Exp. Zool.* 284:637-644.

554 Chaouloff, F. 2000. Serotonin, stress and corticoids. *J. Psychopharmacol.* 14:139-151.

555 Clements, S., F. L. Moore, and C. B. Schreck. 2003. Evidence that acute serotonergic  
556 activation potentiates the locomotor-stimulating effects of corticotropin-releasing  
557 hormone in juvenile chinook salmon (*Oncorhynchus tshawytscha*). *Horm. Behav.*  
558 43:214-221.

559 Crabbe, J. C. 1999. Animal models in neurobehavioral genetics: Methods for estimating  
560 genetic correlation. In B. C. Jones and P. Mormède (eds.), *Neurobehavioral*  
561 *genetics: Methods and applications*, pp. 121–138. CRC Press, Boca Raton.

562 Gonsten, D., J. G. D. Lambert, H. Komen and H. J. T. Goos. 2002. Corticosteroids affect  
563 the testicular androgen production in male common carp (*Cyprinus carpio* L.)  
564 *Biol. Reprod.* 66:106-111.

565 de Boer, S. F., R. de Beun, J. L. Slangen, and J. Van der Gugten. 1990. Dynamics of  
566 plasma-catecholamine and corticosterone concentrations during reinforced and  
567 extinguished operant-behavior in rats. *Physiol. Behav.* 47:691-698.

568 de Boer, S. F., B. J. Van der Vegt, and J. M. Koolhaas. 2003. Individual variation in  
569 aggression of feral rodent strains: A standard for the genetics of aggression and  
570 violence? *Behav. Genet.* 33:485-501.

571 de Kloet, E. R., S. M. Korte, N. Y. Rots, and M. R. Kruk. 1996. Stress hormones,  
572 genotype, and brain organization - implications for aggression. *Ann. N.Y. Acad.*  
573 *Sci.* 794: 179-191

574 Driscoll, P., R. M. Escorihuela, A. Fernandez-Teruel, O. Giorgi, H. Schwegler, T.  
575 Steimer, A. Wiersma, M. G. Corda, J. Flint, J. M. Koolhaas, W. Langhans, P. E.  
576 Schulz, J. Siegel, and A. Tobeña. 1998. Genetic selection and differential stress  
577 responses - the Roman lines/strains of rats. *Ann. N.Y. Acad. Sci.* 851: 501-510

578 Dunn, A. J., N. B. Gildersleeve, and H. E. Gray. 1978. Mouse brain tyrosine hydroxylase  
579 and glutamic acid decarboxylase following treatment with adrenocorticotrophic  
580 hormone, vasopressin or corticosterone. *J. Neurochem.* 31:977-982.

581 Ellenbroek, B. A. and A. R. Cools. 2002. Apomorphine susceptibility and animal models  
582 for psychopathology: genes and environment. *Behav. Genet.* 32:349-361.

583 Engel, B. T., and A. H. Schmale. 1972. Conservation-withdrawal: A primary  
584 regulator process for organic homeostasis. *In* R. Porter and J. Knight (eds.),  
585 *Physiology, emotions and psychosomatic illness* vol. 8, pp. 57-76. CIBA  
586 Foundation Symposium. Elsevier Science Publishers, New York.

587 Fernandez, F., S. Sarre, J. M. Launay, S. Aguerre, V. Guyonnet-Dupérat, M. P. Moisan,  
588 G. Ebinger, Y. Michotte, P. Morméde, and F. Chaouloff. 2003. Rat strain  
589 differences in peripheral and central serotonin transporter protein expression and  
590 function. *Eur. J. of Neurosci.* 17:494-506.

591 Fevolden, S. E., K. H. Røed, K. T. Fjalestad, and J. Stien. 1999. Poststress levels of  
592 lysozyme and cortisol in adult rainbow trout: heritabilities and genetic  
593 correlations. *J. Fish Biol.* 54:900-910.

594 Fillenz, M. 1993. Neurochemistry of stress: introduction to techniques. *In* S. C. Stanford  
595 and P. Salmon (eds.), *Stress from synapse to syndrome*, pp. 247-279. Academic  
596 Press, London.

597 Fokkema, D. S., J. M. Koolhaas, and J. Van der Gugten. 1995. Individual characteristics  
598 of behavior, blood-pressure, and adrenal hormones in colony rats. *Physiol.*  
599 *Behav.* 57:857-862.

600 Francis, R. C. 1990. Temperament in a fish - a longitudinal-study of the development of  
601 individual-differences in aggression and social rank in the midas cichlid. *Ethology*  
602 86:311-325.

603 Gall, G. A. E. and N. Huang. 1988a. Heritability and selection schemes for rainbow trout:  
604 body-weight. *Aquaculture* 73:43-56.

605 Gall, G. A. E. and N. Huang. 1988b. Heritability and selection schemes for rainbow trout:  
606 female reproductive-performance. *Aquaculture* 73:57-66.

607 Giorgi, O., D. Lecca, G. Piras, P. Driscoll, and M. G. Corda. 2003. Dissociation between  
608 mesocortical dopamine release and fear-related behaviours in two

609 psychogenetically selected lines of rats that differ in coping strategies to aversive  
610 conditions. *Eur. J. Neurosci.* 17:2716-2726.

611 Gjedrem, T. 2000. Genetic improvement of cold-water fish species. *Aquacult. Res.*  
612 31:25-33.

613 Gjerde, B. 1993. Breeding and selection. *In* K. Heen, R. Monahan, and F. Utter (eds.),  
614 *Salmon aquaculture*, pp. 187–208. Blackwell, Oxford.

615 GjØen, H. M. and Bentsen, H. B. 1997. Past, present, and future of genetic improvement  
616 in salmon aquaculture. *ICES J. Mar. Sci.* 54:1009-1014.

617 Gregory, T. R. and C. M. Wood. 1999. The effects of chronic plasma cortisol elevation  
618 on the feeding behaviour, growth, competitive ability, and swimming  
619 performance of juvenile rainbow trout. *Physiol. Biochem. Zool.* 72:286-295.

620 Haller, J., J. Halasz, G. B. Makara, and M. R. Kruk. 1998. Acute effects of  
621 glucocorticoids: behavioral and pharmacological perspectives. *Neurosci.*  
622 *Biobehav. Rev.* 23:337-344.

623 Hansen, M. M., D. E. Ruzzante, E. E. Nielsen, D. Bekkevold, and K. L. D. Mensberg.  
624 2002. Long-term effective population sizes, temporal stability of genetic  
625 composition and potential for local adaptation in anadromous brown trout (*Salmo*  
626 *trutta*) populations. *Mol. Ecol.* 11:2523-2535.

627 Harris, J. and D. J. Bird. 2000. Modulation of the fish immune system by hormones. *Vet.*  
628 *Immunol. Immunopathol.* 77:163-176.

629 Henderson, N. D. 1997. Spurious associations in unreplicated selected lines. *Behav.*  
630 *Genet.* 27:145-154.

- 631 Henry, J. P. and P. M. Stephens. 1977. *Stress, health, and the social environment. A*  
632 *sociobiologic approach to medicine*. Springer, New York.
- 633 Henry, J. P. 1993. Biological basis of the stress response. *News Physiol. Sci.* 8:69-73.
- 634 Hershberger, W. K. 1992. Genetic variability in rainbow trout populations.  
635 *Aquaculture* 100:51-71.
- 636 Höglund, E., N. Kolm, and S. Winberg. 2001. Stress-induced changes in brain  
637 serotonergic activity, plasma cortisol and aggressive behavior in Arctic charr  
638 (*Salvelinus alpinus*) is counteracted by L-DOPA. *Physiol. Behav.* 74:381-389.
- 639 Höglund, E., P. H. M. Balm, and S. Winberg. 2002a. Behavioural and neuroendocrine  
640 effects of environmental background colour and social interaction in Arctic charr  
641 (*Salvelinus alpinus*). *J. Exp. Biol.* 205:2535-2543.
- 642 Höglund, E., P. H. M. Balm, and S. Winberg. 2002b. Stimulatory and inhibitory effects  
643 of 5-HT<sub>1A</sub> receptors on adrenocorticotrophic hormone and cortisol secretion in a  
644 teleost fish, the Arctic charr (*Salvelinus alpinus*). *Neurosci. Lett.* 324:193-196.
- 645 Kime, D.E., 1987. The steroids. *In* I. Chester-Jones, P. Ingleton, J.G. Phillips (eds.),  
646 *Fundamentals of comparative vertebrate endocrinology*, pp. 3–56. Plenum Press,  
647 New York.
- 648 Klomberg, K. F., T. Garland, J. G. Swallow, and P. A. Carter. 2002. Dominance, plasma  
649 testosterone levels, and testis size in house mice artificially selected for high  
650 activity levels. *Physiol. Behav.* 77:27-38.
- 651 Koolhaas, J. M., S. M. Korte, S. F. De Boer, B. J. Van der Vegt, C. G. Van Reenen, H.  
652 Hopster, I. C. De Jong, M. A. W. Ruis, and H. J. Blokhuis. 1999. Coping styles in

653 animals: current status in behavior and stress-physiology. *Neurosci. Biobehav.*  
654 *Rev.* 23:925-935.

655 Korte, S. M., B. Buwalda, G. A. H. Bouws, J. M. Koolhaas, F. W. Maes, and B. Bohus.  
656 1992. Conditioned neuroendocrine and cardiovascular stress responsiveness  
657 accompanying behavioral passivity and activity in aged and in young rats.  
658 *Physiol. Behav.* 51:815-822.

659 Larson, E. T., D. O. Norris, and C. H. Summers. 2003. Monoaminergic changes  
660 associated with socially induced sex reversal in the saddleback wrasse.  
661 *Neuroscience* 119:251-263.

662 Lecca D., G. Piras, P. Driscoll, O. Giorgi, and M. G. Corda. 2004. A differential  
663 activation of dopamine output in the shell and core of the nucleus accumbens is  
664 associated with the motor responses to addictive drugs: a brain dialysis study in  
665 Roman high- and low-avoidance rats. *Neuropharmacology* 46: 688-699.

666 Lepage, O., Ø. Øverli, E. Petersson, T. Jarvi, and S. Winberg. 2000. Differential stress  
667 coping in wild and domesticated sea trout. *Brain. Behav. Evol.* 56:259-268.

668 Lepage, O., O. Tottmar, and S. Winberg. 2002. Elevated dietary intake of L-tryptophan  
669 counteracts the stress-induced elevation of plasma cortisol in rainbow trout  
670 (*Oncorhynchus mykiss*). *J. Exp. Biol.* 205:3679-3687.

671 Lepage, O., I. M. Vilchez, T. G. Pottinger, and S. Winberg. 2003. Time-course of the  
672 effect of dietary L-tryptophan on plasma cortisol levels in rainbow trout  
673 *Oncorhynchus mykiss*. *J. Exp. Biol.* 206:3589-3599.

674 Lesch, K. P., D. Bengel, A. Heils, S. Z. Sabol, B. D. Greenberg, S. Petri, J. Benjamin, C.  
675 R. Muller, D. H. Hamer, and D. L. Murphy. 1996. Association of anxiety-related  
676 traits with a polymorphism in the serotonin transporter gene regulatory region.  
677 *Science* 274:1527-1531.

678 Liebsch, G., A. Montkowski, F. Holsboer, and R. Landgraf. 1998. Behavioural profiles of  
679 two Wistar rat lines selectively bred for high or low anxiety-related behaviour.  
680 *Behav. Brain Res.* 94:301-310.

681 Lowe, T. E., J. M. Ryder, J. F. Carragher, and R. M. G. Wells. 1993. Flesh quality in  
682 snapper, *Pagrus auratus*, affected by capture stress. *J. Food Sci.* 58:770-773.

683 Lowry, C. A., K. A. Burke, K. J. Renner, F. L. Moore, and M. Orchinik. 2001. Rapid  
684 changes in monoamine levels following administration of corticotropin-releasing  
685 factor or corticosterone are localized in the dorsomedial hypothalamus. *Horm.*  
686 *Behav.* 39:195-205.

687 Lucas, L. R., P. Pompei, J. Ono, and B. S. McEwen. 1998. Effects of adrenal steroids on  
688 basal ganglia neuropeptide mRNA and tyrosine hydroxylase radioimmunoreactive  
689 levels in the adrenalectomized rat. *J. of Neurochem.* 71:833-843.

690 Maule, R.A., R.A. Tripp, S.L. Kaattari, and C.B. Schreck. 1989. Stress alters immune  
691 function and disease resistance in chinook salmon (*Oncorhynchus tshawytscha*). *J.*  
692 *Endocrinol.* 120:135–142.

693 Midtlyng, P. J., A. Storset, C. Michel, W. J. Slierendrecht, and N. Okamoto. 2002.  
694 Breeding for disease resistance in fish. *Bull. Eur. Assn Fish Pathol.* 22:166-172.



695 Moreira, P. S. A., K. G. T. Pulman, and T. G. Pottinger. 2004. Extinction of a conditioned  
696 response in rainbow trout selected for high or low responsiveness to stress. *Horm.*  
697 *Behav.* 46: 450-457.

698 Morgan, D., K. A. Grant, O. A. Prioleau, S. H. Nader, J. R. Kaplan, and M. A. Nader.  
699 2000. Predictors of social status in cynomolgus monkeys (*Macaca fascicularis*)  
700 after group formation. *Am. J. Primatol.* 52:115-131.

701 Nikulina, E. M., D. F. Avgustinovich, and N. K. Popova. 1992. Selection for reduced  
702 aggressiveness towards man and dopaminergic activity in Norway rats. *Aggress.*  
703 *Behav.* 18:65-72.

704 Øverli, Ø., S. Winberg, B. Damsgård, and M. Jobling. 1998. Food intake and spontaneous  
705 swimming activity in Arctic char (*Salvelinus alpinus*): role of brain serotonergic  
706 activity and social interactions. *Can. J. Zool.* 76:1366-1370.

707 Øverli, Ø., C. A. Harris, and S. Winberg. 1999. Short-term effects of fights for social  
708 dominance and the establishment of dominant-subordinate relationships on brain  
709 monoamines and cortisol in rainbow trout. *Brain. Behav. Evol.* 54:263-275.

710 Øverli, Ø., T. G. Pottinger, T. R. Carrick, E. Øverli, and S. Winberg. 2001. Brain  
711 monoaminergic activity in rainbow trout selected for high and low stress  
712 responsiveness. *Brain. Behav. Evol.* 57:214-224.

713 Øverli, Ø., T. G. Pottinger, T. R. Carrick, E. Øverli, and S. Winberg. 2002a. Differences  
714 in behaviour between rainbow trout selected for high- and low-stress  
715 responsiveness. *J. Exp. Biol.* 205:391-395.

716 Øverli, Ø., S. Kotzian, and S. Winberg. 2002b. Effects of cortisol on aggression and  
717 locomotor activity in rainbow trout. *Horm. Behav.* 42:53-61.

718 Øverli, Ø., S. Winberg, E. T. Larson, W. J. Korzan, O. Lepage, T. R. Summers, K. J.  
719 Renner, and C. H. Summers. (2003). Opposing effects of stress and exogenous  
720 cortisol on hindbrain serotonin. *Soc. Neurosci. Abstract.* 713.15.

721 Øverli, Ø., W. J. Korzan, E. T. Larson, S. Winberg, O. Lepage, T. G. Pottinger, K. J.  
722 Renner, and C. H. Summers. 2004a. Behavioral and neuroendocrine correlates of  
723 displaced aggression in trout. *Horm. Behav.* 45: 324-329.

724 Øverli, Ø., W. J. Korzan, E. Höglund, S. Winberg, H. Bollig, M. Watt, G. L. Forster,  
725 B. A. Barton, E. Øverli, K. J. Renner, and C. H. Summers. 2004b. Stress coping  
726 style predicts aggression and social dominance in rainbow trout. *Horm. Behav.*  
727 45: 235-241.

728 Pankhurst, N.W. and G. Van der Kraak. 1997. Effects of stress on reproduction and  
729 growth of fish. *In* G.K. Iwama, A.D. Pickering, J.P. Sumpter, and C.B. Schreck  
730 (eds.), *Fish stress and health in aquaculture*, pp. 73-93. Cambridge University  
731 Press, Cambridge.

732 Pankhurst, N.W. and G. van der Kraak. 2000. Evidence that acute stress inhibits ovarian  
733 steroidogenesis in rainbow trout in vivo, through the action of cortisol. *Gen.*  
734 *Comp. Endocrinol.* 117:225-237.

735 Perreault, H. A. N., K. Semsar, and J. Godwin. 2003. Fluoxetine treatment decreases  
736 territorial aggression in a coral reef fish. *Physiol. Behav.* 79:719-724.

737 Piazza, P. V., M. Barrot, F. Rougé-Pont, M. Marinelli, S. Maccari, D. N. Abrous, H.  
738 Simon, and M. Le Moal. 1996a. Suppression of glucocorticoid secretion and  
739 antipsychotic drugs have similar effects on the mesolimbic dopaminergic  
740 transmission. Proc. Natl. Acad. Sci. USA 93:15445-15450.

741 Piazza, P. V., F. Rougé-Pont, V. Deroche, S. Maccari, H. Simon, and M. Le Moal. 1996b.  
742 Glucocorticoids have state-dependent stimulant effects on the mesencephalic  
743 dopaminergic transmission. Proc. Natl. Acad. Sci. USA 93:8716-8720.

744 Pickering, A. D. and T. G. Pottinger. 1987. Poor water quality suppresses the cortisol  
745 response of salmonid fish to handling and confinement. J. Fish Biol. 30:363-374.

746 Pickering, A. D., T. G. Pottinger, J. Carragher, and J. P. Sumpter. 1987. The effects of  
747 acute and chronic stress on the levels of reproductive hormones in the plasma of  
748 mature male brown trout, *Salmo trutta* L. Gen. Comp. Endocrinol. 68:249-259.

749 Pickering, A. D. 1993. Growth and stress in fish production. Aquaculture 111:51-63.

750 Popova, N. K., N. N. Voitenko, A. V. Kulikov, and D. F. Avgustinovich. 1991a.  
751 Evidence for the involvement of central serotonin in mechanism of domestication  
752 of silver foxes. Pharmacol. Biochem. Behav. 40:751-756.

753 Popova, N. K., A. V. Kulikov, E. M. Nikulina, E. Y. Kozlachkova, and G. B. Maslova.  
754 1991b. Serotonin metabolism and serotonergic receptors in Norway rats selected  
755 for low aggressiveness to man. Aggress. Behav. 17:207-213.

756 Pottinger, T. G., A. D. Pickering, and M. A. Hurley. 1992. Consistency in the stress  
757 response of individuals of two strains of rainbow trout, *Oncorhynchus mykiss*.  
758 Aquaculture 103:275-289.

759 Pottinger, T. G., T. A. Moran, and J. A. W. Morgan. 1994. Primary and secondary  
760 indexes of stress in the progeny of rainbow-trout (*Oncorhynchus mykiss*) selected  
761 for high and low responsiveness to stress. J. Fish Biol. 44:149-163.

762 Pottinger, T. G., P. H. M. Balm, and A. D. Pickering. 1995. Sexual maturity modifies the  
763 responsiveness of the pituitary-interrenal axis to stress in male rainbow trout. Gen.  
764 Comp. Endocrinol. 98:311-320.

765 Pottinger, T. G. and T. R. Carrick. 1999. Modification of the plasma cortisol response to  
766 stress in rainbow trout by selective breeding. Gen. Comp. Endocrinol. 116:122-  
767 132.

768 Pottinger, T. G., and T. R. Carrick. 2000. Contrasting seasonal modulation of the stress  
769 response in male and female rainbow trout. J. Fish Biol. 56:667-675.

770 Pottinger, T. G. and T. R. Carrick. 2001a. Stress responsiveness affects dominant-  
771 subordinate relationships in rainbow trout. Horm. Behav. 40:419-427.

772 Pottinger, T. G. and T. R. Carrick. 2001b. ACTH does not mediate divergent stress  
773 responsiveness in rainbow trout. Comp. Biochem. Physiol. A 129:399-404.

774 Ramos, A. and P. Mormède. 1998. Stress and emotionality: a multidimensional and  
775 genetic approach. Neurosci. Biobehav. Rev. 22:33-57.

776 Ramos, A., E. C. Correia, G. S. Izídio, and G. R. Brüske. 2003. Genetic selection of two  
777 new rat lines displaying different levels of anxiety-related behaviors. Behav.  
778 Genet. 33:657-668.

779 Rots, N. Y., J. Workel, M. S. Oitzl, A. Béro, W. Rostène, A. R. Cools, and E. R. de  
780 Kloet. 1996a. Development of divergence in dopamine responsiveness in

781 genetically selected rat lines is preceded by changes in pituitary-adrenal activity.  
782 Dev. Brain Res. 92:164-171.

783 Rots, N. Y., A. R. Cools, M. S. Oitzl, J. de Jong, W. Sutanto, and E. R. de Kloet. 1996b.  
784 Divergent prolactin and pituitary-adrenal activity in rats selectively bred for  
785 different dopamine responsiveness. Endocrinology 137:1678-1686.

786 Rots, N. Y., A. R. Cools, A. Béroud, P. Voorn, W. Rostène, and E. R. de Kloet. 1996c.  
787 Rats bred for enhanced apomorphine susceptibility have elevated tyrosine  
788 hydroxylase mRNA and dopamine D<sub>2</sub>-receptor binding sites in nigrostriatal and  
789 tuberoinfundibular dopamine systems. Brain Res. 710:189-196.

790 Roozendaal, B., 2002. Stress and memory: opposing effects of glucocorticoids on  
791 memory consolidation and memory retrieval. Neurobiol. Learn. Mem: 78, 578-  
792 595.

793 Ryman, N. 1983. Patterns of distribution of biochemical genetic variation in salmonids:  
794 Differences between species. Aquaculture 33:1-21.

795 Sallinen, J., A. Haapalinna, E. MacDonald, T. Viitamaa, J. Lähdesmäki, E. Rybnikova,  
796 M. Peltö-Huikko, B. K. Kobilka, and M. Scheinin. 1999. Genetic alteration of the  
797  $\alpha_2$ -adrenoceptor subtype C in mice affects the development of behavioral despair  
798 and stress-induced increases in plasma corticosterone levels. Mol. Psychiatry 4:  
799 443-452.

800 Schreck, C. B., W. Contreras-Sanchez, and M. S. Fitzpatrick. 2001. Effects of stress on  
801 fish reproduction, gamete quality, and progeny. Aquaculture 197:3-24.

802 Shih, J. C., K. Chen, and M. J. Ridd. 1999. Monoamine oxidase: From genes to behavior.  
803 Ann. Rev. Neurosci. 22:197-217.

804 Sigholt, T., U. Erikson, T. Rustad, S. Johansen, T. S. Nordtvedt, and A. Seland. 1997.  
805 Handling stress and storage temperature affect meat quality of farm-raised  
806 Atlantic salmon (*Salmo salar*). J. Food Sci. 62:898-905.

807 Sluyter, F., S. M. Korte, B. Bohus, and G. A. Van Oortmerssen. 1996. Behavioral stress  
808 response of genetically selected aggressive and nonaggressive wild house mice in  
809 the shock-probe/defensive burying test. Pharmacol. Biochem. Behav. 54:113-116.

810 Steimer, T. and P. Driscoll. 2003. Divergent stress responses and coping styles in  
811 psychogenetically selected roman high-(RHA) and low-(RLA) avoidance rats:  
812 Behavioural, neuroendocrine and developmental aspects. Stress 6:87-100.

813 Sumpter, J.P., A.D. Pickering, and T. G. Pottinger. 1985. Stress induced elevation of  
814 plasma  $\alpha$ -MSH and endorphin in brown trout, *Salmo trutta* L. Gen. Comp.  
815 Endocrinol. 59:257-265.

816 Taylor, E. B. 1991. A review of local adaptation in salmonidae, with particular reference  
817 to Pacific and Atlantic salmon. Aquaculture 98:185-207.

818 Thorgaard, G. H., G. S. Bailey, D. Williams, D. R. Buhler, S. L. Kaattari, S. S. Ristow, J.  
819 D. Hansen, J. R. Winton, J. L. Bartholomew, J. J. Nagler, P. J. Walsh, M. M.  
820 Vijayan, R. H. Devlin, R. W. Hardy, K. E. Overturf, W. P. Young, B. D. Robison,  
821 C. Rexroad, and Y. Palti. 2002. Status and opportunities for genomics research  
822 with rainbow trout. Comp. Biochem. Physiol. B. 133:609-646.

823 Trenzado, C. E., T. R. Carrick, and T. G. Pottinger. 2003. Divergence of endocrine and  
824 metabolic responses to stress in two rainbow trout lines selected for differing  
825 cortisol responsiveness to stress. *Gen. Comp. Endocrinol.* 133:332-340.

826 Van Raaij, M. T. M., D. S. S. Pit, P. H. M. Balm, A. B. Steffens, and G. Van denThillart.  
827 1996. Behavioral strategy and the physiological stress response in rainbow trout  
828 exposed to severe hypoxia. *Horm. Behav.* 30:85-92.

829 Veenema, A. H., O. C. Meijer, E. R. de Kloet, and J. M. Koolhaas. 2003. Genetic  
830 selection for coping style predicts stressor susceptibility. *J. Neuroendocrinol.*  
831 15:256-267.

832 Verbeek, M. E. M., A. Boon, and P. J. Drent. 1996. Exploration, aggressive behavior and  
833 dominance in pair-wise confrontations of juvenile male great tits. *Behaviour*  
834 133:945-963.

835 Viggiano, D., D. Vallone, H. Welzl, and A. G. Sadile. 2002. The naples high- and low-  
836 excitability rats: Selective breeding, behavioral profile, morphometry, and  
837 molecular biology of the mesocortical dopamine system. *Behav. Genet.* 32:315-  
838 333.

839 Viggiano, D., D. Vallone, L. A. Ruocco, and A. G. Sadile. 2003. Behavioural,  
840 pharmacological, morpho-functional molecular studies reveal a hyperfunctioning  
841 mesocortical dopamine system in an animal model of attention deficit and  
842 hyperactivity disorder. *Neurosci. Biobehav. Rev.* 27:683-689.

843 Waples, R. S., R. G. Gustafson, L. A. Weitkamp, J. M. Myers, O. W. Johnson, P. J.  
844 Busby, J. J. Hard, G. J. Bryant, F. W. Waknitz, K. Neely, D. Teel, W. S. Grant, G.

845 A. Winans, S. Phelps, A. Marshall, and B. M. Baker. 2001. Characterizing  
846 diversity in salmon from the Pacific Northwest. *J. Fish Biol.* 59:1-41.

847 Wendelaar-Bonga, S.E. 1997. The stress response in fish. *Physiol. Rev.* 77:591-625.

848 Winberg, S., G. E. Nilsson, and K. H. Olsén. 1991. Social rank and brain  
849 levels of monoamines and monoamine metabolites in Arctic charr, *Salvelinus*  
850 *alpinus* (L.). *J. Comp. Physiol. A* 168: 241-246.

851 Winberg, S. and G. E. Nilsson. 1993. Roles of brain monoamine neurotransmitters in  
852 agonistic behavior and stress reactions, with particular reference to fish. *Comp.*  
853 *Biochem. Physiol. C.* 106:597-614.

854 Winberg, S., G. E. Nilsson, B. M. Spruijt, and U. Hoglund. 1993. Spontaneous  
855 locomotor-activity in Arctic charr measured by a computerized imaging  
856 technique: role of brain serotonergic activity. *J. Exp. Biol.* 179:213-232.

857 Winberg, S., A. Nilsson, P. Hylland, V. Söderström, and G. E. Nilsson. 1997. Serotonin  
858 as a regulator of hypothalamic-pituitary-interrenal activity in teleost fish.  
859 *Neurosci. Lett.* 230:113-116.

860 Winberg, S. and O. Lepage. 1998. Elevation of brain 5-HT activity, POMC expression,  
861 and plasma cortisol in socially subordinate rainbow trout. *Am. J. Physiol. R.*  
862 43:645-R654.

863 Winberg, S., Ø. Øverli, and O. Lepage. 2001. Suppression of aggression in rainbow trout  
864 (*Oncorhynchus mykiss*) by dietary L-tryptophan. *J. Exp. Biol.* 204:3867-3876.

865 Wingfield, J. C. 2003. Control of behavioural strategies for capricious environments.  
866 *Anim. Behav.* 66:807-815.



867 Yilmazer-Hanke, D. M., A. Wigger, R. Linke, R. Landgraf, and H. Schwegler. 2004.  
868 Two Wistar rat lines selectively bred for anxiety-related behavior show opposite  
869 reactions in elevated plus maze and fear-sensitized acoustic startle tests. *Behav.*  
870 *Genet.* 34:309-318.

871 Zhukov, D. A. and K. P. Vinogradova. 2002. Learned helplessness or learned inactivity  
872 after inescapable stress? Interpretation depends on coping styles. *Integr. Physiol.*  
873 *Behav. Sci.* 37:35-43.

874

875 Figure legends

876 Figure 1: Post-stress blood plasma cortisol levels (mean  $\pm$  SE) of LR and HR rainbow  
877 trout (mixed sex samples) for all test occasions from the establishment of the parental  
878 generation (F0) up to present (F3).

879 Figure 2: The number of LR and HR rainbow trout identified as either dominant or  
880 subordinate during paired contests. HR subordinates > LR subordinates,  $p < 0.001$ ,  $\chi^2$  test  
881 (data from Pottinger and Carrick, 2001a).

882 Figure 3. The number of LR and HR rainbow trout regaining feed intake within 1 week  
883 of transfer to a new environment. LR feeding > HR feeding,  $p = 0.003$ ,  $\chi^2$  test (data from  
884 Øverli et al., 2002a).

885 Figure 4. Locomotor activity in HR and LR rainbow trout quantified as time spent  
886 moving during a 20 min observation period, with or without the presence of a conspecific  
887 intruder. Asterisks indicate an effect of the intruder, asterisks in brackets [\*] indicates a  
888 difference between HR and LR fish, \* =  $p < 0.01$ , \*\* =  $p < 0.001$ , \*\*\* =  $p < 0.0001$ , Kruskal–  
889 Wallis analysis of variance followed by groupwise comparisons by the Mann–Whitney  
890 U-test (data from Øverli et al., 2002a).

891 Table I. Tissue concentrations of monoamines and monoamine metabolites (ng/g), and  
892 corresponding metabolite/monoamine ratios (mean  $\pm$  SE) in different brain regions of LR  
893 and HR rainbow trout when reared in isolation (controls) or following 1 h confinement  
894 stress. Ratios are in italics. Values that are significantly higher in one selection line (LR  
895 vs HR) or in stressed than control fish are in bold font. Asterisks indicate an effect of  
896 stress, asterisks in brackets [\*] indicates a difference between HR and LR fish, \* =  
897  $p < 0.01$ , \*\* =  $p < 0.001$ , \*\*\* =  $p < 0.0001$ , analysis of variance followed Tukey HSD post-  
898 hoc test (data from Øverli et al., 2001).

899

901 Table I.

	LR control	HR control	LR stressed	HR stressed
<i>Telencephalon</i>				
DOPAC	17.1±4.2	15.0±3.9	23.6±5.9	22.4±2.8
DA	218.3±30.7	188.8±39.2	268.6±57.5	229.9±14.5
DOPAC/DA	0.074±0.009	0.076±0.005	0.086±0.008	0.095±0.008
MHPG	13.5±1.0	11.6±0.6	<b>19.8±1.2</b> ***	<b>21.0±1.2</b> ***
NE	1264±78	1282±50	1347±93	<b>1674±76</b> *
MHPG/NE	0.010±0.001	0.009±0.001	<b>0.015±0.001</b> ***	0.012±0.001
5-HIAA	752.8±39.3	595.1±31.6	<b>914.7±51.3</b> *	<b>838.6±48.0</b> ***
5-HT	1516±75	1490±34	1528±64	1588±77
5-HIAA/5-HT	<b>0.50±0.02</b> [*]	0.40±0.02	<b>0.60±0.03</b> *	<b>0.53±0.03</b> **
<i>Hypothalamus</i>				
DOPAC	3.62±0.41	4.80±0.89	4.87±0.66	4.93±0.86
DA	754.8±15.6	734.8±37.8	696.2±23.6	758.6±36.2
DOPAC/DA	0.0049±0.0006	0.0065±0.0012	0.0072±0.0011	0.0065±0.0012
MHPG	8.08±0.84	7.27±0.88	<b>11.3±0.89</b> *	10.2±0.58
NE	403.5±29.7	379.2±23.7	418.9±31.4	418.4±34.3
MHPG/NE	0.020±0.002	0.020±0.003	0.028±0.003	0.025±0.002
5-HIAA	<b>362.6±19.3</b> [*]	283.0±21.6	<b>420.0±18.9</b> [*]	345.2±14.5
5-HT	1883±243	2080±306	1923±263	2058±303
5-HIAA/5-HT	0.21±0.03	0.16±0.03	0.23±0.03	0.20±0.03
<i>Optic tectum</i>				
DOPAC	6.00±0.25	6.90±0.49	6.31±0.25	<b>8.46±0.41</b> [*]
DA	61.0±3.6	59.3±3.2	75.2±8.5	73.7±5.8
DOPAC/DA	0.099±0.003	0.117±0.007	0.090±0.007	<b>0.117±0.004</b> [*]
MHPG	6.13±0.38	6.21±0.67	7.43±0.26	<b>8.54±0.31</b> **
NE	189.9±11.7	205.2±5.9	200.4±7.5	<b>262.8±15.6</b> [*]
MHPG/NE	0.032±0.001	0.030±0.003	0.037±0.001	0.033±0.002
5-HIAA	114.7±7.4	121.8±11.8	145.1±8.0	154.1±9.8
5-HT	496.2±19.9	534.7±44.1	557.1±39.8	588.8±49.3
5-HIAA/5-HT	0.23±0.01	0.23±0.01	0.26±0.01	0.27±0.01
<i>Brain stem</i>				
DOPAC	4.03 ± 0.13	4.51±0.19	4.26±0.17	<b>5.40±0.21</b> [*]
DA	140.3±6.4	139.3±5.3	127.0±5.2	<b>149.9±5.7</b> [*]
DOPAC/DA	0.029±0.002	0.033±0.002	0.034±0.002	0.036±0.001
MHPG	5.74±0.7	5.8±0.4	7.51±0.5	<b>7.95±0.48</b> *
NE	248.6±15.1	240.7±7.8	234.6±8	253.6±10.6
MHPG/NE	0.023±0.003	0.024±0.002	<b>0.034±0.003</b> *	0.031±0.001
5-HIAA	125.6±6.3	103.6±4.9	144.2±6.9	<b>140.6±5.7</b> ***
5-HT	595.8±31.1	610.6±31.6	532.8±24.3	<b>646.1±28.8</b> [*]
5-HIAA/5-HT	<b>0.21±0.006</b> [*]	0.17±0.005	<b>0.27±0.016</b> ***	<b>0.21±0.005</b> [*]

902


903


904 Figure 1

905

906

907






909 Figure 3

910

911

912



913 Figure 4

914

