

1 Demersal fish communities of the shelf and slope of South Georgia and Shag Rocks (Southern Ocean)

2

3 Susan Gregory <sup>1</sup>, Martin A. Collins <sup>2</sup> and Mark Belchier <sup>1</sup>

4

5 1: British Antarctic Survey (NERC), High Cross, Madingley Road, Cambridge, CB3 0ET, UK

6 2: Government of South Georgia & the South Sandwich Islands, Government House, Stanley, Falkland Islands, FIQQ 1ZZ

7

8 Corresponding author: Susan Gregory. Email: [sgreg@bas.ac.uk](mailto:sgreg@bas.ac.uk); Tel: +44 (0)1223 221386; Fax: +44 (0)1223 362616

9

## Abstract

This research is the first to investigate deepwater demersal fish distribution and community structure around South Georgia and Shag Rocks. Analysis of catch data from a trawl survey conducted in 2003 indicated that depth and location have a marked influence over demersal fish community structure in the region. Three distinct, depth stratified fish assemblages were observed. The demersal fish assemblage found on the shelf to depths of around 400m was dominated by nototheniids and was comprised largely of species endemic to the Southern Ocean and Scotia Sea. At the greatest depths sampled (>600m) the demersal fish community was dominated by gadiform fishes including members of the Macrouridae and Moridae, many of which are not endemic to the Southern Ocean. From 400-600m there was a transitional zone with demersal fish representatives of both the shelf and deeper slope communities. Clear geographic differences in the shelf community were apparent with differences observed in community structure between South Georgia and Shag Rocks to depths of around 400m. These data provide valuable baseline information to aid environmental management decisions and assess potential impacts of rapid ocean warming around South Georgia.

**Keywords:** South Georgia • fish • demersal • slope • community • distribution

## Introduction

Fish assemblages are comprised of all the species in a particular area, regardless of whether they interact with each other or not (Wootton 1990) and there are many factors that will influence both the size and diversity of those communities. Depth is well recognised as a key factor in structuring a community (Merrett & Haedrich 1997), however abiotic factors such as a physical lack of space (for example on small plateaux), substrates/habitats that are unsuitable as nursery grounds, isolation caused by oceanic current patterns, and both current and historical fishing pressures, can all impact the community structure with potential effects on both the size structure and diversity of an assemblage (Bianchi et al. 2000; Neat & Campbell 2011; Sundblad et al. 2013, Young et al. 2015).

Deepwater fisheries are generally deemed to be those that target species which routinely live at depths greater than 400-500m (Thiel 2003; Gordon 2005). Patterns of species richness in deep-sea demersal fish assemblages can be determined by local phenomena, and a combination of physical factors and prey availability can regulate the diversity of the fish community (e.g. Priede et al. 2010). It is recognised however that our knowledge of deep sea fish assemblages may be biased by the types of sampling gear used (Gordon 2005) and that the use of different fishing methodologies and baited cameras can enhance knowledge of community structures (Priede et al. 1994, 2010).

South Georgia and Shag Rocks are outcrops of the Scotia Arc, a predominantly undersea ridge that forms the boundary of the Scotia Sea in the Atlantic sector of the Southern Ocean. The continental shelf that surrounds the islands has an average depth of 200m and extends 30-150km in width before dropping away to the abyssal plain. The waters around the islands are highly productive, lying south of the Polar Front and in the path of the Antarctic Circumpolar Current which transports krill to the region from the colder waters of the Antarctic Peninsula and Weddell Sea. Since 1987 twenty one groundfish surveys have been carried out at the sub-Antarctic islands of South Georgia and Shag Rocks (CCAMLR sub-area 48.3). These random stratified bottom trawl surveys have principally been designed to obtain standing stock estimates to assess stock status of the commercial finfish species *Champscephalus gunnari* (mackerel icefish) and also to provide an index of abundance of pre-recruit Patagonian toothfish (*Dissostichus eleginoides*). In addition they have provided information on the relative abundance and ecology of species taken as by-catch in the *C. gunnari* fishery whilst providing estimates of abundance and demographic data on many of the demersal fish species found on the shelf region of South Georgia, including previously over-exploited species. These surveys have largely been confined to the shelf region, covering relatively shallow areas up to maximum depths of ~500m but with the majority sampling at depths less than 350m. Data from these surveys has contributed substantially to our understanding of the ecology of the demersal fish assemblage of the South Georgia and Shag Rocks shelves (Belchier 2013) and has provided data on many of the individual species which are found in this habitat (e.g. Everson et al. 1999; Reid et al. 2007; Belchier & Collins 2008; Main & Collins 2011; Gregory et al. 2014). In contrast, research into the ecology of species found at greater depths on the continental slope around South Georgia has been much more limited.

61 Ecosystem based fisheries management (EBFM), such as that practised by the Commission for the Conservation of Antarctic  
62 Marine Living Resources (CCAMLR) (Constable et al. 2000), requires a thorough understanding of the biology and ecology of  
63 the target species, the dependent prey and by-catch and sympatric species found within the ecosystem. The relatively recent  
64 development (since 1988) of a deepwater demersal long line fishery for Patagonian toothfish (*Dissostichus eleginoides*) has  
65 greatly increased our understanding of the demersal fish assemblage found on the South Georgia slope. However sampling has  
66 been limited to the target species and those caught as by-catch (e.g. Marlow et al. 2003; Morley et al. 2004; Endicott 2010). The  
67 rough bathymetry and deepwater of the slope has meant that fisheries independent surveys have been very limited, restricted to  
68 a handful of studies of the scavenging fauna using baited camera systems (Yau et al. 2000 & 2002; Collins et al. 2002 & 2003)  
69 and cephalopods (Collins et al. 2004). There has, to date, been no research undertaken which has specifically focussed on the  
70 demersal fish assemblages found at greater depths of the continental slope at South Georgia.

71  
72 In addition to informing fisheries management decisions, baseline information on biodiversity and species interactions in the  
73 region is increasingly important at a time when South Georgia is facing rapid ocean warming (Whitehouse et al. 2008) and new  
74 marine protected areas are being introduced to protect the ecosystem (Trathan et al. 2014). In January 2003 a deepwater bottom  
75 trawl survey was undertaken to investigate the biota found on areas of the shelf and continental slope at depths down to 1000m.  
76 This is the only deepwater demersal trawl survey that has been successfully carried out to date at South Georgia and this paper  
77 reviews the data collected during that survey and examines the distribution and community structure of demersal fish species in  
78 the region.

## 79 **Materials and Methods**

### 80 *2003 Deepwater Survey sample collection*

81  
82 The survey was carried out on the research vessel *Dorada* between 11<sup>th</sup> and 31<sup>st</sup> January 2003 (Collins et al. 2004). A radial  
83 transect design was used, with trawls undertaken at pre-determined locations along transects aligned onshore/offshore at Shag  
84 Rocks and South Georgia. Historically the study area has been divided into five sectors when carrying out research surveys –  
85 Shag Rocks and four distinct sectors around South Georgia (North East, North West, South East and South West). We have used  
86 the same geographic boundaries to divide the survey area in our analyses to allow comparisons with previously published work.  
87 Each transect comprised five hauls, one within each pre-designated depth strata (0-199m, 200-399m, 400-599m, 600-799m and  
88 800-1000m). Hauls were carried out using a commercial sized FP120 trawl net with a rubber ‘Rockhopper’ bobbin rig (see  
89 Pilling & Parkes 1995) with tows lasting 30 minutes (bottom time) at a speed of approximately four knots, giving a trawled  
90 distance of approximately two nautical miles at each station. A total of 45 trawls were carried out in nine transects (three transects  
91 at Shag Rocks and six at South Georgia) (Fig 1; Appendix 1, Table 3). Sensor failure during trawl 11 (NW) and trawl 60 (SE)  
92 meant no depth data were recorded so these hauls have been excluded from all analyses. Trawls 20 and 36 (both at Shag Rocks)  
93 were aborted after <12 minutes bottom time due to the net coming fast and being badly damaged, although the catches from  
94 these hauls have been included in species presence/absence analysis.

### 95 *Catch processing*

96  
97 Catches were sorted into individual species and a total wet weight for each species within a trawl was measured using motion  
98 compensated scales. Any unidentified fish were frozen and returned to the King Edward Point Research Station, South Georgia,  
99 for identification using published guides (e.g. Gon & Heemstra 1990). Detailed biological measurements (length, weight, sex,  
100 maturity) and samples (stomach contents, otoliths, muscle and liver) were taken from a range of species including nototheniids,  
101 channichthyids, macrourids, myctophids and rajids for additional research.

### 102 *Data analysis*

103  
104 In the context of this study, the term ‘demersal fish’ refers to both benthic and benthopelagic ichthyofauna (Gordon, 2005).  
105 Known meso- and bathypelagic fish (e.g. bathylagids, gempylids, nemichthyids, paralepidids, myctophids, scopelarchids and  
106 stomiids) which would most likely have been caught in the water column during net retrieval were excluded from community  
107 structure analyses, as were all invertebrates such as crustaceans, cephalopods and medusae. Following recent genetic studies  
108  
109  
110  
111

(Fitzcharles et al. 2012; McMillan et al. 2012) grenadiers identified as *Macrourus whitsoni* on the survey have been reclassified as *Macrourus caml*.

The locations of all hauls were plotted using ArcGIS. The mean species richness (number of fish taxa present) was calculated for the different geographic sectors and depth strata sampled in each area, then ANOVA were used to compare geographic and depth variations. The depth range where each species was caught was also illustrated. The influence of depth and haul location on community structure was analysed using PRIMER-E statistical software, version 6 (Primer-E Ltd, Plymouth, UK). Using non-metric multi-dimensional scaling (NMDS), a resemblance matrix was applied to presence/absence data from each identified fish species with a zero-adjusted Bray-Curtis similarity coefficient and visualised in a two dimensional ordination plot (minimum stress 0.001, 250 restarts). PERMANOVA was used to check for significant differences between the depth zones before hierarchical cluster analysis of presence/absence data was used to identify species assemblages. The contribution of individual species to the assemblages was determined using similarity percentage (SIMPER) analysis. Within assemblages the consolidating species were classed as those that cumulatively contributed >70% to the similarity, and between assemblages species contributing >70% of dissimilarity were classed as discriminating.

## Results

The total catch weight of fish species for all trawls was 6,715.8kg (6,565.9kg excluding pelagic species). A total of 69 fish taxa were recorded from the 42 trawls, and of these, 51 were identified to species level, 7 to genus and 11 to family (see Appendix 1, Table 4). Of the fish identified to species level, the highest number were from the suborder Notothenioidei (10 species), with 7 species of Gadiformes, 7 species of Aulopiformes, 3 species of Zoarcidei, 2 species of Rajiformes, and 2 species of Stomiiformes. Other orders were represented by a single species. Eight of the identified species, four of the taxa classified to genus level, and eight of the taxa classified to family level were pelagic which have been discounted from subsequent analyses.

Demersal species richness (number of taxa) for individual trawls ranged from 3 taxa (trawls 24 and 35, both at Shag Rocks, 400-599m) to 13 taxa (trawl 46 - South East, 600-799m). Analysis by sector indicated the largest number of fish species were caught in trawls in the North West sector (30 taxa) with a mean of 9.75 taxa/trawl (Fig 2). The mean number of taxa differed significantly between sectors (ANOVA:  $F(4, 38) = 8.05, p = <0.0001$ ). A post-hoc Tukey test demonstrated that species richness at Shag Rocks was significantly lower than other sectors. Point estimates indicated the highest species richness occurred in trawls from the depth zone from 600-799m (25 taxa) with a mean of 9.00 taxa/trawl (Fig 3) however this was not found to be significantly greater than mean species number/tow found within the other depth strata. (ANOVA:  $F(4, 38) = 1.49, p = 0.225$ ).

### Community structure

Multivariate analysis based on the presence/absence of each fish species from trawls at South Georgia (all 4 sectors combined) and Shag Rocks showed some degree of community structure differentiation between the sectors (Fig 4), however a greater structuring influence was exerted by the depth zone. PERMANOVA analysis found significant differences in the community structure between all depth zones except 600-799m and 800-1000m ( $p < 0.05$ ) so for further analysis the two deepest zones were pooled into one group, 600-1000m (Fig 5).

Hierarchical cluster analysis showed three major species assemblages at the resemblance level of *c.* 30% (Fig 6). Assemblage A was comprised entirely of shallow trawls (12 in total) – eleven of the twelve trawls in this cluster were <400m at South Georgia plus one Shag Rocks trawls <200m. At a similarity level of ~35% the geographic separation within the assemblage is evident, and at ~50% the division between 0-199m and 200-399m trawls at South Georgia occurs. Assemblage C comprised 14 trawls, and contained the deepest trawls – all the trawls in the 600-1000m range from both Shag Rocks and South Georgia, plus one 400-599m trawl at South Georgia. Assemblage B was comprised of the highest number of trawls (17) which included all remaining trawls <600m at Shag Rocks, and all except one trawl from 400-599m at South Georgia. Again at ~50% similarity a geographic separation within Assemblage B can be seen (Fig 6).

SIMPER analysis showed moderate average within-group similarity ranging from 49.6% to 55.2% (Table 1). The consolidating species differed between assemblages with no single species contributing to all three clusters. The only overlap occurred with *Dissostichus eleginoides* (contributing 28.1% to Assemblage B and 17.7% to Assemblage C) (Table 1).

164

165 In Assemblage A *Gobionotothen gibberifrons* contributed the highest percentage (25.6%) to the consolidating species (those  
166 cumulatively contributing >70% to the similarity), while the three species of Channichthyidae (*Pseudochaenichthys georgianus*,  
167 *Chaenocephalus aceratus* and *Champocephalus gunnari*) cumulatively contributed another 46.2%. There were only three  
168 consolidating species in Assemblage B – *Dissostichus eleginoides* (28.1%), *Mancopsetta maculata antarctica* (a bothid flatfish,  
169 23.7%) and *Lepidonotothen squamifrons* (18.2%). Assemblage C had the greatest number of consolidating species with  
170 gadiformes making the highest overall contribution (*Antimora rostrata*: 26.0%; *Macrourus caml*: 10.9%; *Muraenolepis microps*:  
171 9.0%; and *Macrourus carinatus*: 7.7%). *D. eleginoides* was the only other consolidating species in Assemblage C (17.7%).  
172 (Table 1).

173

174 The dissimilarity levels between the assemblages were much higher than the within-assemblage similarities, with the highest  
175 dissimilarity (91.5%) between A (the shallowest trawls) and C (the deepest trawls). Assemblage B was mainly comprised of  
176 mid-depth trawls and the dissimilarities between A & B and B & C were lower, 74.3% and 73.3% respectively. Discriminating  
177 species (those cumulatively contributing >70% of the dissimilarity) between assemblages were more numerous than the  
178 consolidating species within assemblages, ranging from 10 to 16 species. Between A & C the dissimilarity was primarily a result  
179 of species that were absent from one or other of the assemblages, however between A & B and B & C the dissimilarity was  
180 driven more by differences in average abundance rather than presence/absence (Table 2).

181

### 182 *Bathymetric distribution*

183

184 Nototheniodei were found at all trawled depths, however individual species ranges were more localised (Fig 7). *Dissostichus*  
185 *eleginoides* was the only nototheniodei found at all depths, with species such as *Pseudochaenichthys georgianus* and *Artedidraco*  
186 *mirus* restricted to shallow hauls, *Patagonotothen guntheri* and *Lepidonotothen squamifrons* in shallow to mid-range trawls,  
187 *Aethotaxis mitopteryx* and *Bathydraco joannae* in mid-range to deep trawls and *Pleuragramma antarcticum* only in deep trawls.  
188 Rajids also showed species range differences with *Raja georgiana* being caught in a much wider range of depths compared to  
189 *Bathyraja meridionalis*. All species of Aulopiformes and Stomiiformes were found in mid-range trawls, whilst Zoarceidei and  
190 Gadiformes were primarily found in the mid-range to deep trawls (with the exception of *Muraenolepis microps* which was also  
191 found in shallow trawls) (Fig 7).

192

### 193 **Discussion**

194

195 At the sub Antarctic island of South Georgia three decades of research trawl surveys of the shelf area have provided a  
196 considerable body of information on the ecology of the demersal fish assemblage down to depths of approx. 350- 400m (e.g.  
197 North 2005; Stowasser et al. 2012). In contrast, our understanding of the demersal fish assemblage found at depths greater than  
198 400m in the shelf-break and slope region is far more limited.

199

200 It is widely documented that the composition of fish assemblages changes with depth (Haedrich 1997). Our data demonstrate  
201 that at South Georgia water depth plays a clear role in driving the observed patterns in community structure as indicated by the  
202 presence of three putative depth-separated demersal fish assemblages. These are represented by a shelf assemblage at depths  
203 between 100 and 400m, a 'shallow' slope assemblage located approximately between 400 and 600m and a deeper slope  
204 assemblage found at depths > 600m. The bathymetric distribution of demersal fishes has been very poorly studied in the Southern  
205 Ocean but our results are in broad agreement with the study of Pakhomov et al. (2006) from the sub-Antarctic Prince Edward  
206 archipelago.

207

208 In common with all shelf areas of the Southern Ocean, including the seasonal pack ice zone and the High Antarctic Zone  
209 (Mintenbeck et al. 2012), the South Georgia and Shag Rocks shelves are dominated by nototheniids and channichthyids (which  
210 comprise in excess of 96% of the demersal fish biomass) at depths < 400m. The relatively clear differences seen between the  
211 South Georgia and Shag Rocks shelf demersal fish communities (depths < 400m) seen in this study are in broad agreement with  
212 the multi-year analyses detailed in Belchier (2013). The lower species diversity at Shag Rocks has been attributed to the reduced  
213 number of habitats and reduced shelf area (Belchier 2013) whilst the presence of more temperate, Patagonian species such as  
214 *Patagonotothen guntheri* and absence of bathydraconids such as *Psilodraco breviceps* distinguish the Shag Rocks shelf demersal

215 fish assemblage from that found at South Georgia. The reduction in the number of Antarctic species at Shag Rocks has also been  
216 observed in other sub-Antarctic seamounts and other islands located on the periphery of the Southern Ocean (Kock 1992).

217  
218 Many of the species of nototheniid and channichthyid found at South Georgia and Shag Rocks have a broad geographical  
219 distribution throughout the seasonal pack ice zone. This includes *Champscephalus gunnari* and the nototheniids  
220 *Lepidonotothen squamifrons*, *Lepidonotothen larseni* and *Notothenia rossii* which are found south of the Antarctic Polar Frontal  
221 Zone (APFZ) across all sectors of the Southern Ocean, whilst other species have a distribution that is restricted to the Atlantic  
222 region (*Lepidonotothen nudifrons*, *Gobionotothen gibberifrons* and *Chaenocephalus aceratus*) (Knox 2007) with some endemic  
223 to the South Georgia region (*Psilodraco breviceps*) (Gon & Heemstra 1990). Very few of the South Georgia shelf species are  
224 found north of the APFZ, the exceptions being *D. eleginoides* which is found at depths < 300m in regions both north and south  
225 of the APFZ (Collins et al. 2010) and *P. guntheri* which is found both at Shag Rocks and on the shelf regions of the North Scotia  
226 Ridge and on the Patagonian shelf. *D. eleginoides* populations found north and south of the polar front are known to be  
227 genetically distinct with the Polar Front acting as a major barrier to gene flow (Shaw et al. 2004; Rogers et al. 2006)

228  
229 The abundant shelf-dwelling species of nototheniid such as *L. larseni*, *G. gibberifrons* and *Trematomus hansonii* and all of the  
230 channichthyids (icefishes) became very scarce or absent in trawls conducted deeper than 400m. The catches within these depths  
231 were dominated by two species of nototheniid, *L. squamifrons* and *D. eleginoides*, but was also characterised by the presence of  
232 the achiropsettoid flatfish *Mancopsetta maculata* (Assemblage B). These results are in very close agreement with the Pakhomov  
233 et al. (2006) study from the Prince Edward Islands that recorded *D. eleginoides* and *L. squamifrons* being the dominant species  
234 within the shallow slope assemblage. At the Prince Edward Island (Pakhomov et al. 2006) and in the high Antarctic region  
235 (Eastman 1993) the highest diversity of the Nototheniidae has been recorded between 300-600m. Whilst overall fish diversity  
236 was greatest at these depths at South Georgia, the greatest Nototheniidae diversity was actually recorded over the shelf  
237 (Assemblage A). The high diversity within Assemblage B results from it being a transitional region between the shelf and deeper  
238 slope communities and therefore having representatives of both.

239  
240 Assemblage C, found largely at depths >600m at both South Georgia and Shag Rocks is characterised by the presence of gadiform  
241 species including representatives of the Macrouridae, Moridae and Muraenolepididae. These families, which comprise a large  
242 part of the by-catch of the longline fishery for *D. eleginoides*, are not only distributed throughout the Southern Ocean but also in  
243 continental slope regions north of the APFZ (Smith et al. 2011). *Macrourus holotrachys*, *M. caml* and *M. carinatus* caught  
244 during our study are widely found in the Ross Sea (Hanchet et al. 2013) and the sub Antarctic islands of the Indian ocean sector  
245 of the Southern Ocean such as Heard Island (van Wijk et al. 2000), Prince Edward Island, Crozet and Kerguelen (De Broyer et  
246 al. 2014) and many of the seamounts of the sub Antarctic region (Wiff et al. 2012). *M. holotrachys* is caught as by catch in  
247 demersal longline fisheries operating in the Patagonian shelf region of the SW Atlantic (Laptikhovskiy 2005) and has been  
248 recorded as far north as the Juan Fernandez Archipelago at 33°S (Niklitschek et al. 2010). Many of the species of deepwater  
249 gadiforms found between 600-1000m at South Georgia have broad geographic distributions, for example the morids *Halargyreus*  
250 *johnsonii* and *Antimora rostrata* are known to have a global distribution and are ubiquitous in the north Atlantic (Mauchline &  
251 Gordon 1984).

252  
253 In 2002 Yau et al. described scavenging fauna observed using baited cameras around South Georgia at depths from 625-1520m.  
254 Only eight fish species were observed during the camera deployments, a fraction of the species caught during the groundfish  
255 survey. The most commonly observed fish species during camera deployments was a benthopelagic zoarcid (tentatively  
256 identified as *Lycodapus antarcticus*) which were seen at depths from 625m, however catches of this species were low in the  
257 groundfish survey and it made no contribution to assemblage similarities and <4% to the dissimilarity in community structure.  
258 Differences in trawl catches compared to camera observations are not unexpected as the sampling methodologies can target  
259 different components of the fish community in terms of both species and size composition (Priede et al. 1994). Camera derived  
260 data can be biased by factors such as bait predominately attracting scavengers and predators that use olfactory cues (King et al.  
261 2006; Bailey et al. 2007), oceanographic conditions affecting fish response to bait (Zinten et al. 2012), and an inability to  
262 discriminate between rare species (Priede et al. 2010). Underwater imagery can however complement extractive studies by  
263 allowing observations of species and age classes not targeted by nets or hooks (Priede et al. 1994, 2010) and offers a non-  
264 destructive method of sampling vulnerable species (McLean et al. 2014).

266 We are aware that there were limitations in the sampling design and the spatial extent covered by the survey. Consequently,  
267 whilst this study represents the most extensive fisheries independent sampling of the deepwater demersal fish assemblages of  
268 South Georgia to date, it does not provide a complete inventory of all the demersal fish species of South Georgia. During the  
269 survey only one haul occurred within 12nm of the coast and consequently a number of shallow water coastal species such as  
270 members of the Harpagiferidae and the rock cod *Notothenia coriiceps* were not sampled effectively. Similarly at the greatest  
271 depths sampled, concerns over gear damage or loss meant that the rougher areas of seabed were avoided. Limited numbers of  
272 hauls conducted within each of the depth strata coupled with uncertainty over bottom contact time and net damage in deep tows  
273 (> 600m) meant that a fully quantitative assessment of species density (i.e. through calculation of CPUE by species) was not  
274 considered to be appropriate. It is highly likely that increased sampling effort within the deeper depth strata will yield a greater  
275 number of fish species than were seen in this study.

276  
277 We can conclude that the deepwater demersal fish assemblage of the slope of South Georgia is characterised by the presence of  
278 gadiform species that have wide ranging distributions that extend far beyond the northern limits of the APFZ. The composition  
279 of this assemblage at South Georgia appears very similar to those found on the slopes of other archipelagos and continental shelf  
280 margins both within the APFZ and to the north. The composition of the deepwater slope assemblage contrasts with that found  
281 on the shelf which is comprised predominantly of nototheniid species almost all of which are endemic to the Southern Ocean  
282 and some of which are endemic to the South Georgia shelf.

283  
284 The APF forms a very strong barrier to the migration and dispersal of marine organisms including fish between the Southern  
285 Ocean and the more temperate waters located to the north of South Georgia in the South Atlantic (Strugnall & Allcock 2013).  
286 Consequently there are very few demersal species that are found on continental shelves on either side of the APF. *Patagonotothen*  
287 *guntheri* is one of the only species that is found either side the APF and its distribution to the south is restricted to the  
288 comparatively warmer waters of the Shag Rocks shelf. However, the Polar Front does not act as a barrier to the migration and  
289 dispersal of slope species in deeper habitats. The eurybathic nature of many slope species such as the Macrouridae (i.e.  
290 Laptikhovsky et al. 2008) coupled with water temperatures on the slope consistently well above freezing provides a more  
291 homogeneous environment over which the range of these species can extend. Similar contrast between the heterogeneous shelf  
292 assemblages and the more homogenous slope fish assemblage has recently been reported in the Antarctic Peninsula region  
293 (Amsler et al. 2016). The North Scotia Ridge which extends as a series of sub surface seamounts from the Patagonian shelf to  
294 South Georgia may also provide an additional mechanism which links the slope habitats of the Patagonian shelf with that at  
295 South Georgia.

296  
297 Further fisheries independent surveys are required to provide greater detail and quantitative data in order to better understand  
298 the observed patterns of demersal fish distribution in this highly productive but isolated environment.

## 299 **Acknowledgments**

300  
301 We would like to extend thanks to the captain and crew of the Fisheries Patrol/Research Vessel *Dorada*, and scientists from the  
302 British Antarctic Survey, Marine Resources Assessment Group and Aberdeen University who undertook the survey, and to Dr  
303 Mark Jessopp for advice on statistical tests. Sample processing was carried out at the King Edward Point Applied Fisheries  
304 Laboratory, South Georgia, and the research was funded by the Government of South Georgia and the South Sandwich Islands.  
305 We are grateful to three anonymous reviewers for their constructive comments on earlier versions of this manuscript.  
306

## 307 **References**

- 308  
309  
310 Amsler MO, Eastman JT, Smith KE, McClintock JB, Hanumant S, Thatje S, Aronson RB (2016) Zonation of demersal fishes  
311 off Anvers Island, western Antarctic Peninsula. *Ant Sci* 28:44-50  
312  
313 Belchier M (2013) Decadal trends in the South Georgia demersal fish assemblage. *CCAMLR WG-FSA-13/26*  
314  
315 Belchier M, Collins MA (2008) Recruitment and body size in relation to temperature in juvenile Patagonian toothfish  
316 (*Dissostichus eleginoides*) at South Georgia. *Mar Biol* 155:493-503  
317

318 Bailey DM, King NJ, Priede IG (2007) Cameras and carcasses: historical and current methods for using artificial food falls to  
319 study deep water animals. *Mar Ecol Prog Ser* 350:179-191  
320  
321 Bianchi G, Gislason H, Graham K, Hill L, Jin X, Koranteng K, Manickchand-Heileman S, Payá I, Sainsbury K, Sanchez F,  
322 Zwanenburg K (2000) Impact of fishing on size composition and diversity of demersal fish communities. *ICES J Mar Sci* 57:558-  
323 571  
324  
325 Collins MA, Yau C, Guilfoyle F, Bagley P, Everson I, Priede IG, Agnew D (2002) Assessment of stone crab (Lithodidae) density  
326 on the South Georgia slope using baited video cameras. *ICES J Mar Sci* 59(2):370-379  
327  
328 Collins MA, Everson I, Patterson R, Bagley PM, Yau C, Belchier M, Hawkins S (2003) *In situ* observations of the scavenging  
329 fauna of the South Georgia slope. In: Shotton R (ed.) *Deep Sea 2003: conference on the governance and management of deep-*  
330 *sea fisheries. Part 2: conference poster papers and workshop papers.* FAO, Rome, pp 487  
331  
332 Collins MA, Allcock AL, Belchier M (2004) Cephalopods of the South Georgia Slope. *J Mar Biol Ass UK.* 84:415-419  
333  
334 Collins MA, Brickle P, Brown J, Belchier M (2010) The Patagonian Toothfish: biology, ecology and fishery. *Adv Mar Bio*  
335 58:227-300  
336  
337 Constable, A.J., de la Mare, W.K., Agnew, D.J., Everson, I. & Miller, D. (2000) Managing fisheries to conserve the Antarctic  
338 marine ecosystem: practical implementation of the Convention on the Conservation of Antarctic Marine Living Resources  
339 (CCAMLR). *ICES J Mar Sci* 57(3):778-791  
340  
341 De Broyer C, Koubbi P, Griffiths HJ, Raymond B, d'Udekem d'Acoz C, Van de Putte AP, Danis B, David B, Grant S, Gutt J,  
342 Held C, Hosie G, Huettmann, F, Post A, Ropert-Coudert Y (eds.) (2014) *Biogeographic Atlas of the Southern Ocean.* SCAR,  
343 Cambridge  
344  
345 Eastman JT (1993) *Antarctic fish biology: evolution in a unique environment.* Academic Press, San Diego  
346  
347 Endicott MR (2010) The impact of the toothfish longline fishery on skate populations around South Georgia. Thesis (PhD),  
348 Imperial College London  
349  
350 Everson I, Parkes G, Kock KH, Boyd IL (1999) Variation in the standing stock of the mackerel icefish *Champsocephalus gunnari*  
351 at South Georgia. *J App Ecol* 4:591-603  
352  
353 Fitzcharles E, Brigden K, Gregory S, Belchier M, Brown J (2012) Molecular and morphological identification of *Macrourus*  
354 species caught as by-catch in the toothfish longline fisheries in CCAMLR Subareas 48.3 and 48.4. CCAMLR WG-FSA-12/35.  
355  
356 Gon O, Heemstra PC (eds.) (1990) *Fishes of the Southern Ocean.* JLB Smith Institute of Ichthyology, Grahamstown  
357  
358 Gordon JDM (2005) Environmental and biological aspects of deepwater demersal fishes. In: Shotton R (ed.), *Deep Sea 2003,*  
359 *Proceedings of the Conference on the Governance and Management of Deep-Sea Fisheries. Part 1: Conference reports.* FAO  
360 *Fisheries Proceedings, No. 3/1.* FAO, Rome, pp. 70–88  
361  
362 Gregory S, Brown J, Belchier M (2014) Ecology and distribution of the grey notothen, *Lepidonotothen squamifrons*, around  
363 South Georgia and Shag Rocks, Southern Ocean. *Antarct Sci* 26(3):239-249  
364  
365 Griffiths HJ, Whittle RJ, Roberts SJ, Belchier M, Linse K (2013) Antarctic Crabs: Invasion or Endurance? *PLoS One* 8(7):1-17  
366  
367 Haedrich R L (1997) Distribution and population ecology. In: Randall DJ & Farrell AP (ed.) *Deep-sea Fishes.* Academic Press,  
368 San Diego, pp. 79–114  
369



370 Hanchet SM, Stewart AL, McMillan PJ, Clark MR, O’Driscoll RL, Stevenson ML (2013) Diversity, relative abundance, new  
371 locality records, and updated fish fauna of the Ross Sea region. *Antarct Sci* 25(5):619-636  
372  
373 King NJ, Bagley PM, Priede IG (2006) Depth zonation and latitudinal distribution of deep-sea scavenging demersal fishes of the  
374 Mid-Atlantic Ridge, 42 to 53°N. *Mar Ecol Prog Ser* 319:263-274  
375  
376 Knox GA (2007) *Biology of the Southern Ocean*. Cambridge University Press, Cambridge  
377  
378 Kock KH (1992) *Antarctic fish and fisheries*. Cambridge University Press, Cambridge  
379  
380 Laptikhovsky VV (2005) A trophic ecology of two grenadier species (Macrouridae, Pisces) in deep waters of the Southwest  
381 Atlantic. *Deep Sea Res* 50:1502–1504  
382  
383 Laptikhovsky V, Arkhipkin A, Brickle P (2008) Biology and distribution of grenadiers of the family Macrouridae around the  
384 Falkland Islands. *Am Fish Soc Symp* 63:261–284  
385  
386 Main, C.E. & Collins, M.A. (2011) Diet of the Antarctic starry skate *Amblyraja georgiana* (Rajidae, Chondrichthyes) at South  
387 Georgia (Southern Ocean). *Polar Biol* 34:389-396  
388  
389 Marlow TR, Agnew DJ, Purves MG (2003) Movement and growth of tagged *Dissostichus eleginoides* around South Georgia  
390 and Shag Rocks (subarea 48.3). *CCAMLR Sci* 10:101-111  
391  
392 Mauchline J, Gordon JDM (1984) Diets and bathymetric distributions of the macrourid fish of the Rockall Trough, northeastern  
393 Atlantic Ocean. *Mar Biol* 81:107–121  
394  
395 McLean DL, Green M, Harvey ES, Williams A, Daley R, Graham KJ (2014) Comparison of baited longlines and baited  
396 underwater cameras for assessing the composition of continental slope deepwater fish assemblages off southeast Australia.  
397 *Deep-Sea Res Pt 1* 98:10-20  
398  
399 McMillan P, Iwamoto T, Stewart A, Smith PJ (2012) A New Species of Grenadier, Genus *Macrourus* (Teleostei, Gadiformes,  
400 Macrouridae) from the Southern Hemisphere and a Revision of the Genus. *Zootaxa* 3165:1–24  
401  
402 Merrett NR, Haedrich RL (1997) *Deep-Sea Demersal Fish and Fisheries*. Chapman and Hall, London  
403  
404 Mintenbeck K, Barrera-Oro ER, Brey T, Jacob U, Knust R, Mark FC, Moreira E, Strobel A, Arntz WE (2012) Impact of Climate  
405 Change on Fishes in Complex Antarctic Ecosystem. *Adv Ecol Res* 46:351-426  
406  
407 Morley SA, Mulvey T, Dickson J, Belchier M (2004) The biology of the bigeye grenadier at South Georgia. *J Fish Biol*  
408 64(6):1514-1529  
409  
410 Neat F, Campbell N (2011) Demersal fish diversity of the isolated Rockall plateau compared with the adjacent west coast shelf  
411 of Scotland. *Bio J Linnean Soc* 104:138-147  
412  
413 Niklitschek EJ, Cornejo-Donoso J, Oyarzun C, Hernandez E, Toledo P (2010) Developing seamount fishery produces localized  
414 reductions in abundance and changes in species composition of by catch. *Mar Ecol* 31(Suppl.1):168–182  
415  
416 North AW (2005) Mackerel icefish size and age differences and long-term change at South Georgia and Shag Rocks. *J Fish Biol*  
417 67(6):1666-1685  
418  
419 Pakhomov EA, Bushula T, Kaehler S, Watkins BP, Leslie RW (2006) Structure and distribution of the slope fish community in  
420 the vicinity of the sub-Antarctic Prince Edward Archipelago. *J Fish Biol* 68:1834-1866  
421  
422 Pilling G, Parkes G (1995) Performance and geometry of the FP-120 trawl used during UK fish stock assessment surveys around  
423 South Georgia, Subarea 48.3. *CCAMLR Sci* 2:51-69

424

425 Priede IG, Bagley PM, Smith A, Creasey S, Merrett NR (1994) Scavenging deep demersal fishes of the Porcupine Seabight,  
 426 North-East Atlantic: observations by baited camera, trap and trawl. *J Mar Biol Ass UK* 74:481-498

427

428 Priede IG, Godbold JA, King NJ, Collins MA, Bailey DM, Gordon JDM (2010). Deep-sea demersal fish species richness in the  
 429 Porcupine Seabight, NE Atlantic Ocean: global and regional patterns. *Mar Ecol* 31:247-260

430

431 Reid WDK, Clarke S, Collins MA, Belchier M (2007) Distribution and ecology of *Chaenocephalus aceratus* (Channichthyidae)  
 432 around South Georgia and Shag Rocks (Southern Ocean). *Polar Biol* 30:1523-1533

433

434 Rogers AD, Morley S, Fitzcharles E, Jarvis K, Belchier M (2006) Genetic structure of Patagonian toothfish (*Dissostichus*  
 435 *eleginoides*) populations on the Patagonian Shelf and Atlantic and western Indian Ocean Sectors of the Southern Ocean. *Mar*  
 436 *Biol* 149(4):915-924

437

438 Shaw PW, Arkhipkin AI, Al-Khairulla H (2004). Genetic structuring of Patagonian toothfish populations in the Southwest  
 439 Atlantic Ocean: the effect of the Antarctic Polar Front and deep-water troughs as barriers to genetic exchange. *Molecular Ecol*  
 440 13(11):3293-3303

441

442 Smith PJ, Steinke D, McMillan PJ, Stewart AL, McVeagh SM, Diaz de Astarloa JM, Welsford D, Ward RD (2011) DNA  
 443 barcoding highlights a cryptic species of grenadier *Macrourus* in the Southern Ocean. *J Fish Biol* 78:355–365

444

445 Stowasser G, Pond DW, Collins MA (2012) Fatty acid trophic markers elucidate resource partitioning within the demersal fish  
 446 community of South Georgia and Shag Rocks (Southern Ocean). *Mar Biol* 159:2299-2310

447

448 Strugnell JM, Allcock AL (2013) Southern Ocean evolution in a global context: A molecular viewpoint. In: Verde C, di  
 449 Prisco G (eds.) *Adaptation and Evolution in Marine Environments, Vol 2, From Pole to Pole*. Springer-Verlag Berlin  
 450 Heidelberg, pp 35-53

451

452 Sundblad G, Bergström U, Sandström A, Eklöv P (2013) Nursery habitat availability limits adult stock sizes of predatory coastal  
 453 fish. *ICES J Mar Sci* 71(3):672-680

454

455 Thiel H (2003) Anthropogenic impacts on the deep sea. In: Tyler, P.A. (Ed.) *Ecosystems of the Deep Oceans*. Elsevier Science  
 456 BV, Amsterdam, The Netherlands, pp. 427-472

457

458 Trathan PN, Collins MA, Grant S, Belchier M, Barnes DKA, Brown J, Staniland IJ (2014) The South Georgia and the South  
 459 Sandwich Islands MPA: protecting a biodiverse oceanic island chain situated in the flow of the Antarctic Circumpolar  
 460 Current. *Adv Mar Bio* 69:15–78

461

462 van Wijk EM, Constable AJ, Williams R, Lamb T (2000) Distribution and abundance of *Macrourus carinatus* on Banzare Bank  
 463 in the southern Indian Ocean. *CCAMLR Sci* 7:171-178

464

465 Whitehouse MJ, Meredith MP, Rothery P, Atkinson A, Ward P, Korb RE (2008) Rapid warming of the ocean around South  
 466 Georgia, Southern Ocean, during the 20<sup>th</sup> century: Forcings, characteristics and implications for lower trophic levels. *Deep Sea*  
 467 *Res I* 55:1218-1228

468

469 Wiff R, Belchier M, Quiroz JC, Arata J (2012) A characterisation of the toothfish fishery in Subarea 48.6 from 2003/04 to  
 470 2011/12. *CCAMLR WG-FSA-12/38*

471

472 Wootton RJ (1990) *Ecology of Teleost Fish*, 2<sup>nd</sup> Edition. Chapman & Hall Ltd, London

473

474 Yau C, Collins MA, Everson I (2000) Commensalism between a liparid fish (*Careproctus* sp.) and stone crabs (Lithodidae)  
 475 photographed *in situ* using a baited camera. *J Mar Biol Ass UK* 80:379-380

476  
477  
478  
479  
480  
481  
482  
483  
484  
485  
486  
487  
488  
489  
490  
491  
492  
493  
494  
495  
496  
497  
498  
499  
500  
501  
502  
503  
504  
505  
506  
507  
508  
509  
510  
511  
512  
513  
514  
515  
516

Yau C, Collins MA, Bagley P, Everson I, Priede IG (2002) Scavenging by megabenthos and demersal fish in the South Georgia slope. *Antarct Sci* 14:16-24

Young EF, Belchier M, Hauser L, Horsburgh GJ, Meredith MP, Murphy EJ, Pascoal S, Rock J, Tysklind N, Carvalho G (2015) Oceanography and life history predict contrasting genetic population structure in two Antarctic fish species. *Evolutionary Applications* 8(5):486-509

Zinten V, Anderson MJ, Roberts CD, Harvey ES, Stewart AL, Struthers CD (2012) Diversity and Composition of Demersal Fishes along a Depth Gradient Assessed by Baited Remote Underwater Stereo-Video. *PLoS One* 7: DOI: 10.1371/journal.pone.0048522

**Fig 1** Location of individual trawls grouped into the nine transects carried out at Shag Rocks and the four sectors around South Georgia. The 200m and 1000m isobaths are shown

**Fig 2** Mean number of demersal fish taxa (excluding pelagics) caught in the four sectors around South Georgia and at Shag Rocks during the 2003 groundfish survey (all depths). Error bars show standard deviation

**Fig 3** Mean number of demersal fish taxa (excluding pelagics) caught around South Georgia and Shag Rocks (all sectors combined) during the 2003 groundfish survey based on depth zone. Error bars show standard deviation

**Fig 4** Two dimensional ordination plot illustrating community structure differences between South Georgia (all four sectors combined) and Shag Rocks [South Georgia: triangle; Shag Rocks: circle. Numbers indicate the trawl ID]

**Fig 5** Two dimensional ordination plot illustrating community structure differences based on species presence/absence and the influence of depth (South Georgia and Shag Rocks combined) [triangle: 0-199m; circle: 200-399m; cross 400-599m; square 600-1000m]

**Fig 6** (a) Dendrogram and (b) ordination in two-dimensions using multidimensional scaling (MDS) showing the three fish community assemblages (Assemblages A, B and C) at Shag Rocks and South Georgia in 2003 based on species presence/absence (30% similarity level)

**Fig 7** Depth ranges of fish species caught during the 2003 groundfish survey [Black bars: demersal species; grey bars: pelagic species (not included in analyses). A: nototheniidae; B: channichthyidae; C: bathydraconidae; D: rajidae; E: zoarcidae; F: macrouridae; G: moridae; H: other demersal fish species; I: myctophidae; J: paralepididae; K: stomiidae; L: scopelarchidae; M: other pelagic fish species]

**Table 1** SIMPER results of demersal fish species contributing >70% of similarity for the three community assemblages (A, B & C) identified using cluster analysis

<b>Assemblage A</b>						
Average similarity: 55.22						
<b>Species</b>	<b>Av.Abund</b>	<b>Av.Sim</b>	<b>Sim/SD</b>	<b>Contrib%</b>	<b>Cum.%</b>	
<i>Gobionotothen gibberifrons</i>	1	14.15	4.84	25.62	25.62	
<i>Chaenocephalus aceratus</i>	0.92	11.21	2.05	20.3	45.91	
<i>Pseudochaenichthys georgianus</i>	0.83	8.65	1.42	15.66	61.57	
<i>Champscephalus gunnari</i>	0.67	5.64	0.83	10.21	71.78	
<b>Assemblage B</b>						
Average similarity: 51.53						
<b>Species</b>	<b>Av.Abund</b>	<b>Av.Sim</b>	<b>Sim/SD</b>	<b>Contrib%</b>	<b>Cum.%</b>	
<i>Dissostichus eleginoides</i>	0.88	14.5	1.68	28.14	28.14	
<i>Mancopsetta maculata antarctica</i>	0.82	12.23	1.35	23.74	51.89	
<i>Lepidonotothen squamifrons</i>	0.71	9.39	0.92	18.23	70.12	
<b>Assemblage C</b>						
Average similarity: 49.63						
<b>Species</b>	<b>Av.Abund</b>	<b>Av.Sim</b>	<b>Sim/SD</b>	<b>Contrib%</b>	<b>Cum.%</b>	
<i>Antimora rostrata</i>	1	12.91	4	26.01	26.01	
<i>Dissostichus eleginoides</i>	0.86	8.78	1.5	17.7	43.71	
<i>Macrourus caml</i>	0.71	5.39	0.97	10.85	54.56	
<i>Muraenolepis microps</i>	0.64	4.48	0.8	9.03	63.59	
<i>Macrourus carinatus</i>	0.57	3.82	0.64	7.7	71.28	

**Table 2** SIMPER results of demersal fish species contributing >70% of dissimilarity between the three community assemblages (A, B & C) at South Georgia and Shag Rocks in 2003 identified using cluster analysis

<b>Assemblages A &amp; C</b>						
Average dissimilarity = 91.51						
Species	Assemblage A		Assemblage C		Contrib %	Cum. %
	Av.Abund	Av.Abund	Av.Diss	Diss/SD		
<i>Antimora rostrata</i>	0	1	6.79	3.98	7.42	7.42
<i>Gobionotothen gibberifrons</i>	1	0	6.79	3.98	7.42	14.84
<i>Chaenocephalus aceratus</i>	0.92	0	6.07	2.57	6.63	21.47
<i>Pseudochaenichthys georgianus</i>	0.83	0	5.35	2	5.84	27.32
<i>Dissostichus eleginoides</i>	0.08	0.86	5.28	1.74	5.77	33.08
<i>Macrourus caml</i>	0	0.71	4.4	1.51	4.81	37.89
<i>Champscephalus gunnari</i>	0.67	0.07	4.26	1.24	4.66	42.55
<i>Macrourus carinatus</i>	0	0.57	3.8	1.09	4.16	46.71
<i>Bathyraco joannae</i>	0	0.57	3.55	1.11	3.88	50.58
<i>Muraenolepis microps</i>	0.58	0.64	3.3	0.91	3.6	54.18
<i>Macrourus holotrachys</i>	0	0.5	3.22	0.94	3.51	57.7
<i>Lepidonotothen larseni</i>	0.5	0	3.06	0.97	3.35	61.04
<i>Lepidonotothen nudifrons</i>	0.42	0	2.77	0.79	3.02	64.07
<i>Raja georgiana</i>	0.25	0.29	2.73	0.76	2.99	67.05
<i>Lepidonotothen squamifrons</i>	0.33	0.14	2.67	0.74	2.92	69.98
<i>Mancopsetta maculata antarctica</i>	0.33	0.14	2.66	0.75	2.9	72.88

<b>Assemblages A &amp; B</b>						
Average dissimilarity = 74.30						
Species	Assemblage A		Assemblage B		Contrib %	Cum. %
	Av.Abund	Av.Abund	Av.Diss	Diss/SD		
<i>Chaenocephalus aceratus</i>	0.92	0.06	6.92	2.23	9.32	9.32
<i>Gobionotothen gibberifrons</i>	1	0.18	6.79	1.9	9.14	18.45
<i>Dissostichus eleginoides</i>	0.08	0.88	6.68	1.87	9	27.45
<i>Pseudochaenichthys georgianus</i>	0.83	0	6.37	2.08	8.57	36.02
<i>Mancopsetta maculata antarctica</i>	0.33	0.82	4.85	1.17	6.53	42.55
<i>Champscephalus gunnari</i>	0.67	0.24	4.75	1.13	6.39	48.94
<i>Lepidonotothen squamifrons</i>	0.33	0.71	4.56	1.09	6.14	55.08
<i>Raja georgiana</i>	0.25	0.53	4.18	0.98	5.62	60.7
<i>Muraenolepis microps</i>	0.58	0.59	4.03	0.93	5.42	66.12
<i>Lepidonotothen nudifrons</i>	0.42	0.29	3.76	0.89	5.06	71.18

<b>Assemblages C &amp; B</b>						
Average dissimilarity = 73.33						
Species	Assemblage C		Assemblage B		Contrib %	Cum. %
	Av.Abund	Av.Abund	Av.Diss	Diss/SD		

<i>Antimora rostrata</i>	1	0.06	7.32	2.71	9.98	9.98
<i>Mancopsetta maculata antarctica</i>	0.14	0.82	5.59	1.47	7.63	17.61
<i>Lepidonotothen squamifrons</i>	0.14	0.71	5.13	1.23	6.99	24.6
<i>Macrourus caml</i>	0.71	0.12	4.72	1.35	6.44	31.04
<i>Macrourus carinatus</i>	0.57	0.12	4.24	1.05	5.79	36.83
<i>Bathyraco joannae</i>	0.57	0	4.01	1.12	5.46	42.29
<i>Raja georgiana</i>	0.29	0.53	3.95	0.96	5.39	47.68
<i>Muraenolepis microps</i>	0.64	0.59	3.78	0.91	5.15	52.84
<i>Macrourus holotrachys</i>	0.5	0.06	3.69	0.94	5.03	57.86
<i>Lycodapus antarcticus</i>	0.43	0	2.95	0.84	4.03	61.89
<i>Lepidion spp</i>	0.43	0	2.95	0.85	4.02	65.91
<i>Halargyreus johnsonii</i>	0.36	0	2.87	0.72	3.91	69.82
<i>Patagonotothen guntheri</i>	0	0.35	2.67	0.71	3.64	73.45

## Appendix One

**Table 3** 2003 deepwater survey haul locations at South Georgia and Shag Rocks

Event	Date	Start Time (UTC)	Duration (mins)	Start Latitude		Start Longitude		End Latitude		End Longitude		Heading (deg)	Minimum Depth (m)	Maximum Depth (m)	Mean Depth (m)
				Degs	Mins	Degs	Mins	Degs	Mins	Degs	Mins				
1	11/01/2003	19:57	30	-54	1.67	36	35.05	54	2.36	36	31.79	108	160	177.5	170
3	12/01/2003	09:02	33	-53	49.08	36	9.32	53	48.84	36	6.23	77	810	928	884
4	12/01/2003	13:43	30	-53	49.27	35	49.25	53	47.39	35	49.66	0	653	765	721
5	12/01/2003	17:12	31	-53	52.9	35	47.11	53	51.27	35	48.53	333	507	564	533
7	13/01/2003	08:09	32	-53	24.68	37	1.21	53	24.96	36	57.79	97	858	964	917
10	14/01/2003	10:33	30	-53	32.18	37	2.23	53	31.56	36	59.11	68	697	706.9	700
11 <sup>a</sup>	14/01/2003			-53	34.69	37	18.03	53	34.69	37	18.03				
12	14/01/2003	16:38	30	-53	34.69	37	18.03	53	34.87	37	21.37	271	487	519	508
13	14/01/2003	19:45	30	-53	42.28	37	30.39	53	42.08	37	26.83	84	297	330	317
14	15/01/2003	14:16	30	-53	54.92	39	1.35	53	54.69	38	57.85	163.2	150	179	
15	15/01/2003	17:19	30	-53	53.52	39	11.81	53	51.97	39	8.93	41	275	290	281.9
16	15/01/2003	21:09	30	-53	47.22	39	18.6	53	49.1	39	19.07	200	452	464	457.6
18	16/01/2003	09:24	31	-53	48.93	39	24.37	53	49.72	39	27.2	250	699	709	703
19	17/01/2003	07:34	30	-53	45.66	40	30.12	53	45.51	40	30.97	284	840	845	843
20 <sup>b</sup>	17/01/2003	11:49	2	-53	44.38	40	35.04	53	44.39	40	34.84	92	881	881	881
23	18/01/2003	10:35	30	-53	48.61	40	33.78	53	48.8	40	30.65	96	579.2	607.3	597.3
24	18/01/2003	17:34	23	-53	41.33	40	42.49	53	40.31	40	40.07	53	439	445	442
26	20/01/2003	12:00	30	-53	39.52	40	47.63	53	41.39	40	46.65	162	309	312	310
27	20/01/2003	15:25	30	-53	37.16	41	0.39	53	37.37	40	57.29	98	156	162	159
28	20/01/2003	17:22	30	-53	38.27	41	2.7	53	38.26	40	59.29	92	129.6	144.5	137
29	21/01/2003	09:58	30	-53	53.87	41	22.88	53	53.77	41	19.11	88	218	225	221
30	21/01/2003	18:03	30	-53	55.32	41	42.26	53	56.44	41	38.71	135	383	408	393
32	22/01/2003	11:53	30	-53	56.71	41	47.16	53	56.63	41	43.47	83	569	605	591
33	23/01/2003	10:37	30	-53	36.04	42	8.1	53	36.46	42	4.57	97	158	180	171
34	23/01/2003	14:45	30	-53	29.38	42	27.41	53	30.63	42	24.61	131	302.3	318	309.1

35	23/01/2003	18:31	30	-53	18.06	42	40.51	53	17.69	42	37.27	79	530.2	553.8	541.3
36 <sup>b</sup>	24/01/2003	07:12	12	-53	16.7	42	40	53	16.26	42	38.86	59	679.1	702.1	685.8
37	24/01/2003	19:13	30	-53	27.5	41	50.01	53	28.29	41	46.7	112	145.6	160.3	155.6
38	25/01/2003	12:02	30	-54	41.91	38	10.29	54	42.59	38	13.6	250	148.6	165.1	159.7
39	25/01/2003	14:35	30	-54	44.99	38	31.33	54	45.81	38	27.87	112	266.1	273.5	270.3
40	25/01/2003	18:38	29	-54	48.27	38	10.69	54	49.12	38	7.27	113	410	515.8	466.5
42	26/01/2003	11:43	27	-54	53.99	38	17.43	54	53.96	38	14.43	89	655.2	689.9	670.2
44	26/01/2003	17:26	26	-54	54.23	38	18.12	54	54.51	38	17.42	143	750.6	925	841.5
46	27/01/2003	11:02	30	-55	25.21	36	13.16	55	23.3	36	14.5	338	651	698	669.4
47	27/01/2003	14:07	30	-55	22.14	36	12.21	55	24.22	36	11.71	173	550.4	567.1	563.5
48	27/01/2003	19:25	30	-55	20.87	36	4.33	55	23	36	4.34	180	434.7	441.6	437.3
50	28/01/2003	10:25	30	-55	30.52	35	36.13	55	29.37	35	39.43	302	295.1	218.2	309.3
51	28/01/2003	14:26	30	-55	14.43	35	37.94	55	14.73	35	41.74	262	150.6	160	154.3
52	29/01/2003	07:09	30	-54	35.7	35	27.85	54	37.27	35	25.42	138	147.5	157.8	153.2
53	29/01/2003	10:18	30	-54	44.7	35	9.47	54	44.77	35	13.17	268	396.6	311.1	308.7
55	29/01/2003	17:26	30	-54	53.46	34	21.29	54	52.37	34	24.55	303	440.8	474.8	459.8
58	30/01/2003	11:25	29	-54	56.98	34	15.89	54	55	34	17.14	340	589.2	623.9	611.9
60 <sup>a</sup>	30/01/2014	16:50	30	-54	51.89	34	21.51	54	51.88	34	21.59				
63	31/01/2003	09:48	30	-53	47.85	35	51.15	53	48.95	35	48.34	124	678.8	719.1	699
65	31/01/2003	17:10	30	-53	56.56	35	55.76	53	58.38	35	57.61	211	252.5	276.1	268.8

<sup>a</sup> indicates sensor failure, haul excluded from analyses; <sup>b</sup> indicates short haul, but species presence/absence data used in analyses



## Appendix One

**Table 4** Complete catch record of all fish species from the 2003 deepwater survey at Shag Rocks and South Georgia

Order/Family/Species	Catch (kg)
<b>Rajidae</b>	
<i>Bathyraja meridionalis</i>	1.26
<i>Raja georgiana</i>	67.22
Rajidae spp. <sup>a</sup>	0.03
<b>Notacanthidae</b>	
Notacanthidae sp.	0.33
<b>Nemichthyidae</b>	
<i>Nemichthys curvirostratus</i> <sup>a</sup>	0.05
<b>Bathylagidae</b>	
<i>Bathylagus</i> spp. <sup>a</sup>	0.71
<b>Astronesthidae</b>	
<i>Astronesthidae</i> sp.	0.03
<b>Stomiidae</b>	
<i>Borostomias antarcticus</i> <sup>a</sup>	0.29
<i>Stomias boa</i> <sup>a</sup>	0.05
Stomiidae spp <sup>a</sup>	0.01
<b>Scopelarchidae</b>	
<i>Benthalbella macropinna</i> <sup>a</sup>	0.05
<i>Benthalbella elongate</i> <sup>a</sup>	0.10
Scopelarchidae sp. <sup>a</sup>	0.13
<b>Notosudidae</b>	
<i>Scopelosaurus hamiltoni</i> <sup>a</sup>	0.37
<b>Paralepididae</b>	
<i>Magnisudis prionosa</i> <sup>a</sup>	0.23
<i>Notolepis annulata</i> <sup>a</sup>	0.03
<i>Notolepis coatsi</i> <sup>a</sup>	0.10
<i>Notolepis rissoi</i> <sup>a</sup>	0.72
<b>Myctophidae</b>	
<i>Electrona antarctica</i> <sup>a</sup>	0.46
<i>Electrona carlsbergi</i> <sup>a</sup>	0.10
<i>Electrona</i> spp. <sup>a</sup>	0.22
<i>Gymnoscopelus bolini</i> <sup>a</sup>	16.18
<i>Gymnoscopelus braueri</i> <sup>a</sup>	0.60
<i>Gymnoscopelus hintonoides.</i> <sup>a</sup>	1.27
<i>Gymnoscopelus nicholsi</i> <sup>a</sup>	89.65
<i>Gymnoscopelus opisthopterus</i> <sup>a</sup>	3.76
<i>Lampanyctus achirus</i> <sup>a</sup>	0.05
Myctophidae spp. <sup>a</sup>	32.88
<i>Protomyctophum</i> spp. <sup>a</sup>	0.02
<b>Muraenolepididae</b>	
<i>Muraenolepis microps</i>	45.50
<b>Moridae</b>	
<i>Antimora rostrata</i>	192.41
<i>Halargyreus johnsonii</i>	28.29
<i>Lepidion</i> sp.	2.57

<b>Macrouridae</b>	
<i>Coryphenooides subserrulatus</i>	0.46
Macrouridae spp. <sup>a</sup>	7.44
<i>Macrourus caml</i>	44.52
<i>Macrourus carinatus</i>	32.01
<i>Macrourus holotrachys</i>	15.67
<b>Carapidae</b>	
<i>Echiodon cryomargarites</i>	0.03
<b>Melamphaidae</b>	
<i>Poromitra crassiceps</i>	0.57
<b>Liparidae</b>	
<i>Careproctus</i> sp.	0.02
Liparidae spp. <sup>a</sup>	4.80
<b>Zoarcidae</b>	
<i>Lycodapus antarcticus</i>	0.33
<i>Lycodapus pachysoma</i>	0.02
<i>Lycodapus</i> sp. <sup>a</sup>	0.05
<i>Melanostigma gelatinosum</i>	2.42
<i>Melanostigma</i> spp. <sup>a</sup>	0.45
<i>Seleniolycus</i> sp. <sup>a</sup>	0.05
Zoarcidae spp. <sup>a</sup>	0.20
<b>Nototheniidae</b>	
<i>Aethotaxis mitopteryx</i>	0.50
<i>Dissostichus eleginoides</i>	239.16
<i>Gobionotothen gibberifrons</i>	251.37
<i>Lepidonotothen larseni</i>	3.70
<i>Lepidonotothen nudifrons</i>	6.60
<i>Lepidonotothen squamifrons</i>	2580.10
<i>Notothenia rossii</i>	29.83
<i>Patagonotothen guntheri</i>	1864.71
<i>Pleuragramma antarcticum</i>	0.01
<i>Trematomus hansonii</i>	3.70
<b>Artedidraconidae</b>	
<i>Artedidraco mirus</i>	0.05
<b>Harpagiferidae</b>	
<i>Harpagifer</i> sp.	0.01
<b>Bathydraconidae</b>	
<i>Bathydraco joannae</i>	0.68
<i>Parachaenichthys georgianus</i>	2.22
<i>Psilodraco breviceps</i>	0.15
<b>Channichthyidae</b>	
<i>Chaenocephalus aceratus</i>	102.83
<i>Champscephalus gunnari</i>	972.79
<i>Pseudochaenichthys georgianus</i>	53.58
<b>Gempylidae</b>	

<i>Paradiplospinus gracilis</i> <sup>a</sup>	1.96
<b>Achiropsettidae</b>	
<i>Mancopsetta maculata antarctica</i>	6.51

<sup>a</sup> indicates species not included in analyses due to pelagic life history or insufficiently detailed identification




Figure 1




Figure 2




Figure 3




Figure 4




Figure 5






Figure 6a




Figure 6b




Figure 7