BRITISH GEOLOGICAL SURVEY Fort **McMurray** Observatory Monthly Magnetic Bulletin September 2015 15/09/FM

Fort McMurray

Flin Flon

Saskatoon L.

Vinnipeg





British Geological Survey

GREENLA

1. Introduction

The British Geological Survey (BGS) now operate or assist in the operation of six overseas geomagnetic observatories. The most recently established of these, Fort McMurray Observatory, is a joint venture between BGS and Sperry Drilling, Halliburton Group Canada in support of directional drilling programmes. The installation was carried out in December 2014 and the earliest data and data products are available from January 2015.

This bulletin is published to provide rapid access to the provisional geomagnetic observatory results. The information is freely available for personal, academic, educational and non-commercial research or use. Magnetic observatory data are presented as a series of plots of one-minute, hourly and daily values, followed by tabulations of monthly values. The operation of the observatory and presentation of data are described in the rest of this section.

Enquiries about the data should be addressed to:

Geomagnetism Team Earth Hazards and Systems British Geological Survey Murchison House, West Mains Road Edinburgh EH9 3LA Scotland, UK

 Tel:
 +44 (0) 131 667 1000

 Fax:
 +44 (0) 131 668 2683

 E-mail:
 enquiries@bgs.ac.uk

 Internet:
 www.geomag.bgs.ac.uk

2. Position

Fort McMurray Observatory is situated on Halliburton land, to the south of the town of Fort McMurray in Alberta, Canada. The observatory coordinates are:-

Geographic:	56° 35'53"N	248°41'06" E
Geomagnetic:	63 <i>° 31'41''N</i>	309 <i>° 32'49" E</i>
Height above m	386m	

The geographical co-ordinates and altitude were determined by a surveying company using a differential GPS and reference station. Data were provided in GPS co-ordinates to UTM 12 NAD83: Ellipsoid GRS1980, and converted to WGS84 by BGS. The geomagnetic co-ordinates are approximations, calculated using the 12th

generation International Geomagnetic Reference Field (IGRF) at epoch 2015.5. On-line access to models (including IGRF), charts and navigational data are available at

http://www.geomag.bgs.ac.uk/data_service/models _compass/home

3. The Observatory Operation

3.1 GDAS

The observatory operates under the control of the Geomagnetic Data Acquisition System (GDAS), which was developed by BGS staff, installed and became fully operational from December 2014. The data acquisition software, running on QNX operated computers, controls the data logging and the communications.

There are two sets of sensors used for making magnetic measurements. A tri-axial linear-core fluxgate magnetometer, manufactured by DTU Space at the Technical University of Denmark, is used to measure the variations in the horizontal (H) and vertical (Z) components of the field. The third sensor is oriented perpendicular to these, and measures variations, which are proportional to the changes in declination (D). Measurements are made at a rate of 1 Hz.

In addition to the fluxgate sensors, a GEM Systems GSM-90 Overhauser magnetometer makes measurements of the absolute total field intensity (*F*) every 5 seconds.

The raw unfiltered data are retrieved automatically via Internet connections to the BGS office in Edinburgh in near real-time. The fluxgate data are filtered to produce one-minute values using a 61point cosine filter and the total field intensity samples are filtered using a 13-point cosine filter.

3.2 Absolute Observations

The GDAS fluxgate magnetometers accurately measure variations in the components of the geomagnetic field, but not the absolute magnitudes. Two sets of absolute measurements of the field are made manually twice per month. A fluxgate sensor mounted on a theodolite is used to determine D and inclination (1); the GDAS PPM measurements, with a site difference correction applied, are used for F. The absolute observations are used in conjunction with the **GDAS** variometer measurements to produce a continuous record of the absolute values of the geomagnetic field elements as if they had been measured at the observatory reference pillar.

4. Observatory Results

The data presented in the bulletin are in the form of plots and tabulations described in the following sections.

4.1 Absolute Observations

The absolute observation measurements made during the month are tabulated. Also included are the corresponding baseline values, which are the differences between the absolute measurements and the variometer measurements of D, H and Z (in the sense absolute–variometer). These are also plotted (markers) along with the derived preliminary daily baseline values (line) throughout the year. Daily mean differences between the measured absolute F and the F computed from the baseline corrected H and Z values are plotted in the fourth panel (in the sense measured–derived). The bottom panel shows the daily mean temperature in the fluxgate chamber.

4.2 Summary magnetograms

Small-scale magnetograms are plotted which allow the month's data to be viewed at a glance. They are plotted 16 days to a page and show the one-minute variations in D, H and Z. The scales are shown on the right-hand side of the page. On disturbed days the scales are multiplied by a factor, which is indicated above the panel for that day. The variations are centred on the monthly mean value, shown on the left side of the page.

4.3 Magnetograms

The daily magnetograms are plotted using oneminute values of D, H and Z from the fluxgate sensors. The magnetograms are plotted to a variable scale; scale bars are shown to the right of each plot. The absolute level (the monthly mean value) is indicated on the left side of the plots.

4.4 Hourly Mean Value Plots

Hourly mean values of D, H and Z for the past 12 months are plotted in 27-day segments corresponding to the Bartels solar rotation number. Magnetic disturbances associated with active regions and/or coronal holes on the Sun may recur after 27 days: the same is true for geomagnetically

© NERC 2015. All rights reserved

Edinburgh

4.5 Daily and Monthly Mean Values

Daily mean values of D, H, Z and F are plotted throughout the year. In addition, a table of monthly mean values of all the geomagnetic elements is provided. These values depend on accurate specification of the fluxgate sensor baselines. It is anticipated that these provisional values will not be altered by more than a few nT or tenths of arcminutes before being made definitive at the end of the year.

5. Conditions of Use

The outputs presented in this bulletin are provided for personal, academic, educational, noncommercial research or other non-commercial use and are not for sale or distribution to third parties without written permission from BGS.


Reproduction of any part of this bulletin should be accompanied by the statement: 'Reproduced with the permission of the British Geological Survey ©NERC. All rights Reserved'. Publications making use of the data should include an acknowledgment statement of the form: 'The results presented in this paper rely on the data collected at Fort McMurray observatory, operated by Sperry Drilling, Halliburton Group Canada and the British Geological Survey.'


Commercial users can contact the geomagnetism team for information on the range of applications and services offered. Full contact details are available at www.geomag.bgs.ac.uk/contactus/staff


FORT MCMURRAY OBSERVATORY


ABSOLUTE OBSERVATIONS


	Declination			Inclination		Total Field		Horizontal Intensity		Vertical Intensity			
Date	Day Number	Time (UT)	Absolute (°)	Baseline (°)	Time (UT)	Absolute (°)	Site difference (nT)	Absolute corrected (nT)	Absolute (nT)	Baseline (nT)	Absolute (nT)	Baseline (nT)	Observer
22-Sep-15	265	21:06	14.7385	14.5500	21:10	77.6654	0.5	57764.8	12339.7	12323.0	56431.4	56487.2	ТМ
22-Sep-15	265	21:20	14.7100	14.5317	21:27	77.6601	0.5	57766.1	12345.2	12325.6	56431.5	56486.7	SP
22-Sep-15	265	21:39	14.7646	14.5833	21:45	77.6662	0.5	57761.4	12338.2	12322.7	56428.2	56487.3	TM
23-Sep-15	266	16:38	14.8779	14.5967	16:44	77.7042	0.5	57744.6	12297.3	12326.9	56420.1	56487.4	ТМ
23-Sep-15	266	16:54	14.8761	14.5833	17:02	77.7367	0.5	57740.0	12264.3	12327.5	56422.5	56487.3	SP
23-Sep-15	266	17:11	14.7408	14.5750	17:17	77.7529	0.5	57727.7	12245.7	12323.1	56414.0	56488.4	ТМ























































Fort McMurray Observatory: Declination (degrees)




Fort McMurray Observatory: Horizontal Intensity fhTŁ

Hourly Mean Values Plotted By Bartels Solar Rotation Number



Fort McMurray Observatory: Vertical Intensity (nT)



Monthly Mean Values for Fort McMurray Observatory 2015

Month	D	Н	Ι	X	Y	Ζ	F
January	14° 55.8´	12310 nT	77° 42.4´	11895 nT	3172 nT	56492 nT	57818 nT
February	**° ***´	***** nT	**° ***´	***** nT	***** nT	***** nT	***** nT
March	14° 54.3´	12300 nT	77° 42.8′	11886 nT	3164 nT	56473 nT	57796 nT
April	14° 52.6´	12318 nT	77° 41.6′	11905 nT	3162 nT	56466 nT	57794 nT
May	14° 51.4′	12328 nT	77° 40.8´	11916 nT	3161 nT	56449 nT	57780 nT
June	14° 50.1´	12315 nT	77° 41.4′	11905 nT	3153 nT	56438 nT	57766 nT
July	14° 48.3´	12330 nT	77° 40.5′	11920 nT	3151 nT	56434 nT	57765 nT
August September	14° 47.9′ 14° 47.2′	12304 nT 12306 nT	77° 40.3 77° 41.9′ 77° 41.7′	11920 nT 11896 nT 11898 nT	3143 nT 3141 nT	56425 nT 56419 nT	57751 nT 57746 nT

Note

i. The values shown here are provisional.