

## Article (refereed) - postprint

---

Smart, Simon Mark; Glanville, Helen Catherine; Blanes, Maria del Carmen; Mercado, Lina Maria; Emmett, Bridget Anne; Jones, David Leonard; Cosby, Bernard Jackson; Marrs, Robert Hunter; Butler, Adam; Marshall, Miles Ramsvik; Reinsch, Sabine; Herrero-Jáuregui, Cristina; Hodgson, John Gavin. 2017. **Leaf dry matter content is better at predicting above-ground net primary production than specific leaf area.** *Functional Ecology*, 31 (6). 1336-1344. [10.1111/1365-2435.12832](https://doi.org/10.1111/1365-2435.12832)

© 2017 The Authors. *Functional Ecology* © 2017 British Ecological Society

This version available <http://nora.nerc.ac.uk/515882/>

NERC has developed NORA to enable users to access research outputs wholly or partially funded by NERC. Copyright and other rights for material on this site are retained by the rights owners. Users should read the terms and conditions of use of this material at <http://nora.nerc.ac.uk/policies.html#access>

**This document is the author's final manuscript version of the journal article, incorporating any revisions agreed during the peer review process. There may be differences between this and the publisher's version. You are advised to consult the publisher's version if you wish to cite from this article.**

The definitive version is available at <http://onlinelibrary.wiley.com/>

Contact CEH NORA team at  
[noraceh@ceh.ac.uk](mailto:noraceh@ceh.ac.uk)

Received Date : 09-Sep-2016

Revised Date : 23-Nov-2016

Accepted Date : 08-Dec-2016

Article type : Standard Papers

Section: Ecosystems Ecology

Editor: Katie Field

**Leaf Dry Matter Content is better at predicting above-ground Net Primary Production than Specific Leaf Area**

Smart, Simon Mark<sup>1</sup>,

Glanville, Helen Catherine<sup>2</sup>,

Blanes, Maria del Carmen<sup>2</sup>,

Mercado, Lina Maria<sup>3,4</sup>,

Emmett, Bridget Anne<sup>5</sup>,

Jones, David Leonard<sup>2</sup>,

Cosby, Bernard Jackson<sup>5</sup>,

Marrs, Robert Hunter<sup>6</sup>,

Butler, Adam<sup>7</sup>,

Marshall, Miles Ramsvik<sup>3</sup>,

Reinsch, Sabine<sup>3</sup>,

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as doi: 10.1111/1365-2435.12832

This article is protected by copyright. All rights reserved.

Herrero-Jáuregui, Cristina<sup>8</sup>,

Hodgson, John Gavin<sup>9</sup>

1 Land Use Group, NERC Centre for Ecology & Hydrology, Library Avenue, Bailrigg LA1 4AP  
UK

2 School of Environment, Natural Resources & Geography, Bangor University, Bangor LL57  
2UW UK

3 College of Life and Environmental Sciences, Geography Department, University of Exeter,  
Rennes Drive, Exeter EX4 4RJ

4 NERC Centre for Ecology & Hydrology, Maclean Building, Benson Lane, Crowmarsh Gifford  
Wallingford, Oxfordshire, OX10 8BB4 UK

5 NERC Centre for Ecology & Hydrology, Environment Centre Wales, Deiniol Road, Bangor,  
Gwynedd, LL57 2UW UK

6 School of Environmental Sciences, University of Liverpool, Nicholson Building, Liverpool,  
L69 3GP UK

7 Biomathematics & Statistics Scotland, JCMB, The King's Buildings, Peter Guthrie Tait Road,  
Edinburgh EH9 3FD UK

8 Department of Ecology, Complutense University of Madrid, C/ José Antonio Novais 12  
28040 Madrid, España

9 Department of Animal and Plant Sciences, Alfred Denny Building, University of Sheffield, Western  
Bank, Sheffield S10 2TN UK

Author for correspondence:

Simon M. Smart

Email: [ssma@ceh.ac.uk](mailto:ssma@ceh.ac.uk)

**Running headline:** LDMC and SLA as predictors of primary production.

### Summary

1. Reliable modelling of above-ground Net Primary Production (aNPP) at fine resolution is a significant challenge. A promising avenue for improving process models is to include response and effect trait relationships. However, uncertainties remain over which leaf traits are correlated most strongly with aNPP.
2. We compared abundance-weighted values of two of the most widely used traits from the Leaf Economics Spectrum (Specific Leaf Area and Leaf Dry Matter Content) with measured aNPP across a temperate ecosystem gradient.
3. We found that Leaf Dry Matter Content (LDMC) as opposed to Specific Leaf Area (SLA) was the superior predictor of aNPP ( $R^2=0.55$ ).
4. Directly measured *in situ* trait values for the dominant species improved estimation of aNPP significantly. Introducing intra-specific trait variation by including the effect of

replicated trait values from published databases did not improve the estimation of aNPP.

5. Our results support the prospect of greater scientific understanding for less cost because LDMC is much easier to measure than SLA.

**Key-words:** Bayesian modelling, ecosystem, global change, measurement error, ecosystem function, intra-specific variation,

## Introduction

Net Primary Production (NPP), defined as the rate at which plants convert CO<sub>2</sub> and water into dry matter, is the basis for life on Earth and is a fundamental ecosystem function supporting food production, soil formation and climate stabilisation. An estimated 28.8% of global NPP (Haberl *et al.* 2007) is appropriated by humans as food, fibre and fuel with consumption often spatially far removed from the area of production (Erb *et al.* 2009).

Accurate prediction of NPP is therefore critical to ecological and economic assessments of the links between land-use change, human well-being and impacts on biodiversity and other ecosystem services (DeFries 2002; Haberl *et al.* 2007). NPP is, however, challenging to measure and predict accurately (Cramer *et al.* 1999; Scurlock *et al.* 2002; Jung *et al.* 2007). A way forward is to derive regionally applicable relationships between plant traits and NPP thereby providing empirical understanding that can potentially be built into global ecosystem models to improve their performance (Wright *et al.* 2006; Van Bodegom *et al.* 2012). New empirical predictions of NPP in terms of plant trait abundance also allow process models to be tested at fine resolution across a range of ecosystems.

Trait-based ecology has become a unifying strand in global change biology because the same sets of key plant traits respond to global change drivers while also driving subsequent effects on ecosystem function (Tateno & Chapin 1997; Suding *et al.* 2008; Reich 2014). We test the performance of two leaf traits – Leaf Dry Matter Content (LDMC) and Specific Leaf Area (SLA) – as predictors of above-ground NPP (aNPP) across a realistically wide productivity gradient using comprehensive measurements of aNPP comprising the full range of plant functional types that dominate temperate ecosystems. Our study seeks to resolve an outstanding question concerning the relative merits of each trait as a correlate of soil fertility and ecosystem productivity (Wilson *et al.* 1999; Hodgson *et al.* 2011). LDMC and SLA both correlate strongly with nutrient availability but it is not clear which of these is the best predictor of aNPP (Wilson *et al.* 1999; Ordoñez *et al.* 2009; 2010; Fortunel *et al.* 2009; Hodgson *et al.* 2011; Pakeman 2011). Given its repeatedly proven alignment with the soil available nutrients axis, SLA has become the pre-eminent predictive leaf trait (Reich 2014). However, the sensitivity of SLA to light availability means that it is not a reliable partial predictor of soil fertility as irradiance changes during succession. Since primary production reflects the availability of resources that include light and nutrients it could mean that SLA is actually a better predictor of aNPP. To test this relationship requires treating SLA as an effect trait rather than as a response trait where variation in abundance-weighted values are explained by abiotic factors (Hodgson *et al.* 2011).

Unlike SLA, LDMC varies independently of leaf thickness (Shipley 1995; Wilson *et al.* 1999; Roche *et al.* 2004) but is also strongly correlated with resource availability and with relative growth rate (Weiher *et al.* 1999; Garnier *et al.* 2004; Fortunel *et al.* 2009). LDMC has been recommended as a more reliable correlate of soil fertility at least in biomes not subject to

severe water limitation (Vendramini *et al.* 2002). Here we explore the role of SLA and LDMC as predictors of ecosystem function and ask which best predicts aNPP across ecosystems.

Since there has been a growing appreciation of the influence of within-species trait variation (Albert *et al.* 2010; Siefert *et al.* 2015) we also test whether including intra-specific trait variation improves the fitted relationship between traits and aNPP. We investigate the performance of each trait as a predictor of aNPP when species of low abundance are excluded and when plant species abundance-weighted trait values for the dominant species among habitats are based on database values or *in situ* measurements.

Our starting point was to compute abundance-weighted trait values based on published UK database values. This is the easiest method to apply for constructing trait-derived variables.

However, if locally measured trait-values differ appreciably from database means and correlate with aNPP then database-derived means will be a poorer predictor of local aNPP.

We tested the importance of intra-specific variation in two ways. First, we substituted mean database trait values for the dominant species in each sampling plot with *in situ* measurements of leaf traits for those species. The two most abundant species were selected to ensure adequate sampling of the species contributing the most biomass to each stand. Secondly, we introduced intra-specific trait variation via its effect on the variance of the abundance-weighted mean trait values. Thus, rather than employing one abundance-weighted mean trait value per sampling plot, a prior distribution of values was calculated based on repeated draws of trait values for each individual plant species. The distributions of trait values for each species were derived from readily accessible replicated database measurements. We then applied a Bayesian measurement error model that allows the observed values of aNPP to feedback onto the posterior estimates of the abundance-

weighted trait values potentially improving the fit between aNPP and trait-based explanatory variable. If successful, this would suggest that better use could be made of the variation in trait values that is readily accessible from databases, rather than just utilising trait means.

In summary we test the following hypotheses:

1. Abundance-weighted LDMC is a better predictor of aNPP than abundance-weighted SLA.
2. Estimation of aNPP is improved when trait values for the dominant plant species are based on *in situ* measurements rather than database averages.
3. Estimation of aNPP is improved when intra-specific trait variation based on replicated database values is included in the model.

## Materials and Methods

### Study region and sampling locations

Fifteen sites were located in the River Conwy catchment in north Wales, UK. The remaining two sites (limestone grassland and upland unimproved hay meadow) were located within the Ingleborough National Nature Reserve in North West England in the upper reaches of the Ribble catchment (Fig. 1; Table 1). The regional climate for all sites is temperate maritime (Peel, Finlayson & McMahon 2007). Annual precipitation lies between 1000 and 1300 mm at Ingleborough and between 600 and 4700 mm in the Conwy valley. Average daily minimum January temperature across the sites is in the range -1 to 3 °C and average daily maximum July temperatures from 17 to 21 °C (long term annual averages 1981-2010, <http://www.metoffice.gov.uk/public/weather/climate/#?region=uk>).



Above-ground NPP was measured in 49 vegetation sampling plots through 2013 and 2014. These plots were nested into 17 sites arranged along a productivity gradient from lowland grasslands intensively managed for agriculture through to montane heath. Within each site, an area of target habitat was selected as a roughly rectangular fraction of the wider habitat of interest. In enclosed land this rectangle was defined by field boundaries. In woodlands and unenclosed habitats a rectangular area was selected to encompass a large area (0.25-1 ha) of the target habitat, for example blanket bog, acid grassland and montane heath. Sampling locations within each site were then chosen at random. Together, these sites sample all common habitat and land-use types in Britain and thus were intended to represent the principal plant biodiversity and productivity gradients in NW Europe.

#### Measurement of above-ground Net Primary Production

Above-ground NPP ( $\text{g dry mass m}^{-2} \text{ yr}^{-1}$ ) was measured using a variety of methods according to the plant functional types present. These types comprised C3 graminoids (Poaceae, Junaceae, Cyperaceae), broad-leaved and needle-leaved trees, dwarf shrubs, forbs and bryophytes (Table 1). All plots were visited in early January at the start of each measurement year. Any green herbaceous material was removed by clipping to 1 cm vegetation height. Standing litter was, as far as possible, not removed nor disturbed. In sheep and cattle-grazed systems (grasslands and mires), livestock enclosures were installed and the vegetation cut twice throughout the growing season; first at estimated peak biomass and a second time to capture late summer and autumn regrowth. These two values were then summed. While this method does not overcome possible issues with negative and positive compensatory growth as a result of grazing, uncertainty over the importance and direction of these effects is great and no clearly superior method appears to exist that

accounts for these effects whilst also excluding grazers (McNaughton *et al.* 1996; Pontes *et al.* 2007).

The biomass fractions attributable to functional types within woodland and peatland ecosystems were measured using differing methods. In peatlands, growth of *Sphagnum* species was measured over two years using the cranked wire method (Clymo 1970; Kivimäki 2011). Peatland graminoids were measured by harvesting annual biomass accumulation in livestock enclosures over one year using the same methods applied to grazed grasslands.

In woodlands, different methods were used to measure annual production of trees. Leaf litter was collected using 20 randomly placed buckets (26cm in diameter) per 200m<sup>2</sup> sampling plot. These were installed in September before litterfall and visited and emptied every two to four weeks until no leaves were visible in the canopy. Annual woody mass increment was measured by combining tree-coring, DBH (tree diameter at 1.3 m height), wood density and tree height measurements. Herbaceous understorey growth was harvested in spring and summer after cutting back in January. Where present, annual production of the bryophyte layer was measured by harvesting the moss mat that had grown through coarse plastic meshes of known size pegged securely to the substrate in early January and harvested after one year.

Measurement of aNPP was carried out using plots of varying dimensions scaled to the size of the plant types present, but then expressed as production per m<sup>2</sup> across all vegetation types (Table 1). Full details of all the methods used for measurement of aNPP on each site are described in Supplementary Material.

## Plant species abundance

In each plot in which aNPP was measured, all vascular plant species and bryophytes were identified and cover was estimated in intervals of 5 % except for species recorded at  $\leq 1$  % cover which were given a value of 1. Percentage cover was based on horizontal leaf projection over the plot so that total cover over all species was allowed to exceed 100, for example, where the understorey comprised a bryophyte layer, a fern layer and a tree canopy. Only species recorded with  $\geq 5$  % cover were used in the calculation of mean abundance-weighted trait values.

## Plant traits

*In situ* measurements of SLA and LDMC were carried out by focussing on the dominant vascular plant species in each plot defined as the two species contributing maximum standing biomass in the year of sampling (Table 1). LDMC (g dry mass  $g^{-1}$  fresh mass) was measured by weighing fresh material consisting of 10 to several hundred mature but non-senescent leaves from different plants depending on leaf size. Leaves were weighed fresh, then dried for 24 hours at 80 °C, and weighed again. SLA ( $mm^2 mg^{-1}$  dry mass) was measured by sampling 10 leaves from different plants. Leaf area was calculated based on scanned photographs analysed using the Image J software v1.46r (<http://imagej.nih.gov/>). Dry weight was measured as for LDMC (Pérez-Harguindeguy *et al.* 2013).

Database values for SLA and LDMC for all vascular plant species encountered in the sample were extracted from LEDA (Kleyer *et al.* 2008) and ECPE (Grime *et al.* 2007). Only values for UK material were included except in four instances where German values were included because no UK data were available. These were *Carex bigelowii*, *C.nigra*, *Agrostis canina* and *Anthoxanthum odoratum*. None of these species were dominant in any of the sample plots. Out of a total pool of 136 vascular plant species recorded in the 49 aNPP plots all had database trait values. The ranges of trait values, including measured and database values, were 57.2 for SLA (4.81, *Picea sitchensis* to 62.1, *Oxalis acetosella*) and 0.45 for LDMC (0.08, *Stellaria media* to 0.53, *Sesleria caerulea*).

## Analysis

Mean abundance-weighted trait values ( $x_{jk}$ ) for SLA and LDMC were computed for each NPP sampling plot  $j$  within each site  $k$  as follows;

$$x_{jk} = \frac{\sum_i \tau_{ijk} p_{ijk}}{\sum_i p_{ijk}}$$

where ( $p_{ijk}$ ) was either the raw percentage cover or square-root transformed cover value for species  $i$  in each sample plot  $j$  within site  $k$  (e.g. Manning *et al.* 2015). The trait values ( $\tau_{ijk}$ )

for each species  $i$  in each sample plot  $j$  and within site  $k$  were based either on replicated *in situ* measurements on the two plant species with the highest cover in each plot, or mean values of each trait extracted from the databases described above.

Two values of the mean abundance-weighted trait (SLA or LDMC) were derived for each plot based on either trait values derived solely from UK databases or supplemented by *in situ* trait measurements for the dominant species in each plot where this value substituted for the database average for those species (Table 1). Abundance-weighted values for SLA and LDMC were used as covariates in regression models designed to test the three hypotheses by determining which model best predicted measured aNPP.

#### Statistical modelling

Model building was carried out using the 'lm' and 'lmer' functions in the lme4 R package (Bates *et al.* 2015). Initial data exploration and preparation followed the steps outlined in Zuur *et al.* (2010) and Crawley (2013). We identified outliers using the outlierTest function in the 'lm' R package. The boxcox function in the 'mass' R package was used to assess homogeneity of variance and the nature of any transformation required to aNPP.

Tests of hypotheses 1 to 3 were carried out by comparing models where each model was of the form,

$$y_{ik} = m_{ik} + \gamma_k + \epsilon_{ik}, \quad 1)$$

$$m_{jk} = a + b \cdot x_{jk}$$

2)

$$\gamma_k \sim N(0, \sigma_\gamma^2)$$

$$\varepsilon_{ik} \sim N(0, \sigma_\varepsilon^2).$$

Where  $y_{jk}$  was the natural log transformed aNPP for plot  $j$  in site  $k$ ,  $x_{jk}$  was the abundance-weighted trait variable and  $\gamma_k$  was a random intercept for each site  $k$ .

Model performance was evaluated by likelihood ratio test and the difference in AICc values between pairs of models. The AICc statistic was used in light of the small sample size (Burnham & Anderson 2002).

To test whether estimation of aNPP was improved by the inclusion of intra-specific trait-variation (Hypothesis 3), a Bayesian measurement-error model was constructed in OpenBUGS ver 3.2.2 rev 1063 (Lunn *et al.* 2013). We modelled the variation in abundance-weighted trait values in each plot by adjusting equation 2) to become,

$$m_{jk} = a + b \cdot z_{jk}$$

3)

$$x_{jk} \sim N(z_{jk}, \sigma_x^2).$$

Here, we now assume that the observed abundance-weighted mean  $x_{jk}$  is an imperfect measure of the true abundance-weighted mean  $z_{jk}$  with its variance being a function of the

distributions of species' trait values contributing to the abundance-weighted trait value for each aNPP plot. These distributions were derived from published replicated database measurements of the trait for each species present. The variance of each species trait value is, therefore, likely to be part measurement error and part ecologically meaningful intra-specific variation in the trait. Thus,  $\sigma_x^2$  conveys the variance in the trait-derived predictor of aNPP that is attributable to known variation in the trait for each contributing species in each plot. An estimate of  $\sigma_x^2$  was generated by first calculating the mean and standard deviation of the database measurements for each plant species which had replicate measurements in the database. Then 1000 random draws of trait values were made based on the parameters of each species' trait distribution. At each draw, a new dataset of abundance-weighted trait values was computed for each of the aNPP plots. An estimate of  $\sigma_x^2$  was then derived by drawing bootstrap samples of increasing size from this dataset until its value stabilised (Fig. S1). Note that this approach implicitly assumes that measurement errors are independent between species and plots. Since we did not derive the trait distributions from measured values from all the species populations within each plot, we cannot directly test this.

The fitted Bayesian measurement error model allows feedback from the aNPP data such that model fit can potentially be improved. Thus the posterior distribution of the slope  $b$  (Equation 3) is also a function of new updated posterior distributions for the abundance-weighted means that optimise the fit between these and aNPP. Without feedback, the effect of intra-specific variation on the abundance-weighted trait means would simply increase the uncertainty around the estimated slope. Measurement error models with feedback are common in pharmacokinetic studies (see Lunn *et al* (2009; 2013) for further details and discussion).

Hypothesis 3 was tested by comparing models with or without intra-specific variation (Equations 2 versus 3). We computed the marginal  $R^2$  (m) value of Nakagawa & Schielzeth (2014) for each model within our BUGS code. This quantifies the explanatory power of the fixed effects (abundance-weighted trait values) as a proportion of the sum of all the variance components; fixed effects plus random effects plus residual. Tests of the difference in  $R^2$  (m) between models were carried out by inspecting the 95 % credible interval (2.5 % and 97.5 % quantiles) of the distribution of differences between 1000 values of  $R^2$  (m) drawn randomly from the posterior distribution of the variable for each model to see whether or not it contained zero. This was achieved in an R script applied to the converged MCMC output for  $R^2$  (m).

The percentage variance attributable to the random effect of site was also calculated with and without the fixed trait effect. This firstly conveys the amount of variation in aNPP between versus within sites and then estimates the extent to which these differing components of variation in aNPP were explained by the abundance-weighted trait (Crawley 2013).

## Results

Initial data exploration showed that aNPP should be transformed to achieve normally distributed residuals and a linear response to abundance-weighted traits. The boxcox function (R package MASS) was applied, confirming that a natural log transformation was most appropriate. Models were also fitted with either untransformed, or square-root transformed plant species cover values in an attempt to reduce the influence of recording



error associated with small differences in % cover. Seven out of eight models based on square-root transformed cover had lower AICc values than the respective model with untransformed cover. In three cases, including the final best fitting model, the difference was greater than the rule-of-thumb value of 2 (Table S2). Thus all subsequent modelling was performed using abundance-weighted trait variables calculated from square-root transformed cover.

Across the 49 plots nested into 17 sites, measured aNPP ranged from 99 g dry mass m<sup>-2</sup> yr<sup>-1</sup> in montane heath to a maximum of 1481 g dry mass m<sup>-2</sup> yr<sup>-1</sup> in intensively-managed lowland improved grassland (Fig. 2). Overall, 91 % of the variation in aNPP occurred between sites.

AICc values for models based on abundance-weighted LDMC were all lower than for models including only SLA (LDMC: 25.6 for a model based on *in situ* trait measurements for the dominants and 30.7 for a model derived from database values only. SLA: 44.8 for a model based on *in situ* trait measurements for the dominants and 42.1 for a model derived from database values only) and differed significantly from these models based on likelihood ratio tests. Thus LDMC was the better trait for predicting aNPP and the best model included *in situ* measurement of LDMC on the dominant species. Hypotheses 1 and 2 were, therefore, supported.

When intra-specific variation in LDMC was included, the model with the highest R<sup>2</sup> (m) included *in situ* trait measurements and the effect of variation in LDMC derived from replicate values in the database. This model explained 55% of the variation in ln(aNPP) with a 95% credible interval of 0.34-0.71, but its R<sup>2</sup> (m) value was not significantly different from a model without database-derived intra-specific variation when their differences were

bootstrapped. The model with the highest  $R^2$  (m) explained 63 % of the within-site, between-plot variation and 34 % of the between-site variation.

## Discussion

### LDMC versus SLA?

We show that LDMC is a superior predictor of aNPP compared to SLA. Our result is novel since we tested SLA and LDMC as *effect* traits across a gradient comprising all major terrestrial ecosystems in the temperate zone. This contrasts with the large number of studies that have explored their role as *response* traits expressing inter- and intra-specific trait variation as a function of environmental gradients such as climate and soil conditions. While LDMC was the superior trait, low variance was explained. In particular, abundance-weighted LDMC only explained a relatively small proportion of the between-site variance that dominated the dataset. It is possible that other plant species-derived predictors could be usefully included in the analysis to increase explanatory power. Leaf traits exhibit differences between plant functional types that are linked to phylogenetically-conserved patterns of biomass allocation (Shiple 1995; Wilson *et al.* 1999; Wright *et al.* 2005; Poorter *et al.* 2012). Therefore, introducing proportional cover of each plant functional type might be worthwhile. However, such categorical units have reduced information content because they do not express continuous variation in plant properties that influence ecosystem function (Van Bodegom *et al.* 2012). This is especially critical for our study. Because we included a range of successional stages across sites and because our sites were located in the oceanic western edge of Europe, the most obvious additional axes of functional

variation across our dataset are plant height and bryophyte cover. In forest ecosystems, aNPP may poorly correlate with lower SLA or higher LDMC because lower production per mass of leaf is compensated by higher absolute foliage mass (Wright *et al.*, 2005; Garnier *et al.*, 2004). When we included abundance-weighted canopy height alongside abundance-weighted LDMC, it failed to explain significant variation in aNPP (see Supplementary Material; Text S1, Table S1, Fig. S3). Because a number of bryophyte genera, including *Sphagnum*, are capable of fixing atmospheric nitrogen (Cornelissen *et al.* 2007), the inclusion of bryophyte cover was also tested as an additional predictor alongside LDMC and SLA but this was also not significant (see Supplementary Material; Text S1, Table S1, Fig. S3). It is quite possible that the addition of climate variables could have explained further variation in aNPP. We did not explore this because (a) we expect considerable collinearity between climate and abundance-weighted trait means (e.g. Ordoñez *et al.* 2009) and (b), our principal aim was to explore the ability of each trait to explain variation in aNPP rather than to develop a full, empirical predictive model for aNPP. While an advantage of our study is in the breadth of ecosystem variation sampled, this also trades-off against our ability to measure and model ecosystem-specific factors and their interactions that are likely to have more fully explained observed aNPP (e.g. Minden & Kleyer 2015).

There are a number of possible reasons why LDMC outperformed SLA in our analysis. SLA exhibits a plastic response to irradiance via changes in leaf thickness such that values can vary significantly with canopy depth even on the same tree (Hollinger 1989). Thus thin shaded leaves have high SLA because they optimise light capture rather than being associated with high soil fertility and therefore higher aNPP (Hodgson *et al.* 2011). These erroneous SLA signals may well have contributed to the variation in published database

values and so to poorer performance of SLA in our analysis of database-derived means.

However, if intra-specific trait variation is partly an adaptive response to local conditions,

then one might have expected LDMC to perform less well because it appears to be

somewhat less plastic than SLA across environmental gradients (Siefert *et al.* 2014; but see

Roche *et al.* 2004). Our result is consistent with other evidence. In a study investigating the

response of leaf plant traits to cutting frequency and nitrogen supply among temperate

grass species, many of which also dominated our grassland samples, Pontes *et al.* (2007)

found that within-species, between treatment-variation in SLA and LDMC was around 14 %

and that LDMC but not SLA was correlated with aNPP. Even where significant intra-specific

variation has been observed, it has proved difficult to explain by abiotic factors (Ordoñez *et*

*al.* 2009; Laughlin *et al.* 2012) often ending up as residual variance rather than predicting

local coupling between trait values and environmental (Albert *et al.* 2010; Jung *et al.* 2010).

This suggests that we might not expect a major jump in predictive power by including intra-

specific variation alongside inter-specific variation especially when derived from replicated

database measurements rather than *in situ* plant populations.

#### Field measurements versus database values

Our results indeed showed that including *in situ* field measurements increased explanatory

power to a greater extent than introducing intra-specific variation via replicated database

values. *In situ* measurements ought to be a better physiological reflection of the

performance of the particular vegetation stand than database averages, and this was indeed

found to be the case. It is likely that the residual error associated with our best model was in

part attributable to low *in situ* trait measurement effort. For example Baroloto *et al.* (2010)

recommended sampling each species at least once in every plot. Even this level of effort may under-represent the variation that can occur in trait values between leaves on the same plant (Shipley, 1995), between individuals of the same species (Albert *et al.* 2010) and throughout the growing season (Pierce *et al.* 1994; Gunn *et al.* 1999; Jagodziński *et al.* 2016). Thus sampling a few individuals in a site at one point in time may lead to unrepresentative trait values poorly coupled to prevailing conditions. Evidently our level of *in situ* sampling effort was sufficient to improve model fit even though our best model still explained just over 55% of the variation in aNPP.

The extent to which *in situ* sampling should focus on capturing inter- or intra-specific variation depends upon the relative importance of each source of variation. Intra-specific variation appears to be greater in less species-rich ecosystems and towards the more extreme end of environmental gradients (Huslof *et al.* 2013; Siefert *et al.* 2015; Baroloto *et al.* 2010). In the absence of any *in situ* measurements, trait means must be calculated from existing databases. Cordlandwehr *et al.* (2013) showed that for less variable traits such as LDMC, database values could satisfactorily approximate ecosystem averages but would be less sensitive to between-patch variation within an ecosystem. Relying solely on database measurements may therefore only weakly capture trait-environment relationships (Manning *et al.* 2015). However, our results indicated that even when derived as database means, LDMC outperformed SLA in prediction of aNPP.

## Conclusions

Using finely-resolved plant trait measurements across a representative vegetation productivity gradient, we show that LDMC was the superior predictor of aNPP compared to SLA. Intra-specific variation, as expressed by *in situ* trait measurements of the two highest abundance species in each plot, led to improved estimation of aNPP but including trait variation as expressed in published database trait values did not. Thus, including database-derived intra-specific variation and allowing this to improve model fit is not an effective substitute for *in situ* trait measurements. However, since LDMC is much easier to measure than SLA, our results suggest that for prediction of aNPP, the burden of data collection can be reduced significantly, thereby offering the prospect of greater scientific understanding for less cost.

## Acknowledgements

We thank the following for arranging access to field sites: Colin Newlands, Andrew Roberts, Alex Turner, Isobelle Griffith, Mike McCabe, Alun Davies, Edward Ritchie, Eifion Jones and Edgar Davies, Thanks to Sue Ward, Amy Goodwin, Susan Jarvis, Laurence Jones, Chris Evans, David Cooper, Hannah Tobermann and Lucy Sheppard for assistance with vegetation sampling and to Simon Pierce, Ed Tipping and Peter Henrys for commenting on earlier versions of the manuscript. The work was funded by the UK Natural Environment Research Council Macronutrients Program.

## Author contributions

SMS, MM, BE, LM, DJ, JC planned and designed the research. SMS, HG, MM, LM, SR conducted fieldwork and analysed data. SMS, ADB carried out the statistical modelling. All authors wrote the manuscript.

## Data accessibility

Smart, S., Reinsch, S., Mercado, L., Blanes, M.C., Cosby, B.J., Glanville, H.C., Jones, D.L., Marshall, M.R. & Emmett, B.A. (2016). Plant structural measurements in North Wales and Northwest England 2013 and 2014. NERC Environmental Information Data Centre. <https://doi.org/10.5285/8899768c-cc5a-4885-a88b-c08374ee568e>.

## Supplementary Material

**Figure S1:** Standard deviation of abundance-weighted LDMC values versus bootstrap sample size.

**Fig S2:** Exploratory plots of abundance-weighted traits against  $\ln(aNPP)$ .

**Text S1:** Additional modelling of bryophyte cover and canopy height.

**Table S2:** Comparison of models derived from either square-root transformed or untransformed cover data for each plant species in each sampling plot.

**Text S2:** Field methods for measuring above-ground Net Primary Production

## References

- Albert, C.H., Thuiller, W., Yoccoz, N.G., Douzet, R., Aubert, S. & Lavorel, S. (2010) A multi-trait approach reveals the structure and the relative importance of intra- vs. interspecific variability in plant traits. *Functional Ecology*, **24**, 1192–1201.
- Baroloto, C., Paine, C.E.T., Patiño, S., Bonal, D., Hérault, B. & Chave, J. (2010) Functional trait variation and sampling strategies in species-rich plant communities. *Functional Ecology* **24**, 208-216.
- Bates, D., Maechler, M., Bolker, B. & Walker, S. (2015). Fitting Linear Mixed-Effects Models Using lme4. *Journal of Statistical Software*, **67**, 1-48
- Burnham, K.P. & Anderson, D.R. (2002) *Model Selection and Multimodel Inference: A Practical Information-Theoretic Approach*. Springer: New York, Berlin.
- Clymo, R.S. (1970) The growth of *Sphagnum*: Methods of measurement. *Journal of Ecology*, **58**, 13-49.
- Cordlandwehr, V., Meredith, R.L., Ozinga, W.A., Bekker, R.M., van Groenendael, J.M. & Bakker, J.P. (2013) Do plant traits retrieved from a database accurately predict on site? *Journal of Ecology*, **101**, 662-670.
- Cornelissen, J.H., Lang, S.I., Soudzilovskaia, N.A. & During, H.J. (2007) Comparative Cryptogam Ecology: A review of bryophyte and lichen traits that drive biogeochemistry. *Annals of Botany*, **99**, 987-1001.
- Cramer, W., Kicklighter, D.W., Bondeau, A., Moore, B., Churkina, G., Nemry, B., Ruimy, A. & Schloss, A.L. (1999) Comparing global models of terrestrial net primary productivity (NPP); overview and key results. *Global Change Biology* (Supplement 1) **5**: 1-15.
- Crawley, M.J. (2013) *The R Book*. 2<sup>nd</sup> edition. John Wiley & Sons Ltd: UK.



DeFries, R. (2002) Past and future sensitivity of primary production to human modification of the landscape. *Geophysical Research Letters*, **29**, 36-1–36-4.

Erb, K-H., Krausmann, F., Lucht, W. & Haberl, H. (2009) Embodied HANPP: Mapping the spatial disconnect between global biomass production and consumption. *Ecological Economics*, **69**, 328-334.

Fortunel, C., Garnier, E., Joffre, R., Kazakou, E., Quested, H., Grigulis, K., Lavorel, S., Ansquer, P., Castro, H., Cruz, P. *et al.* (2009) Leaf traits capture the effects of land use changes and climate on litter decomposability of grasslands across Europe. *Ecology*, **90**, 598-611.

Garnier, E., Cortez, J., Billes, G., Navas, M.L., Ruomet, C., Debussche, M., Laurent, G., Blanchard, A., Aubry, D., Bellmann, A., Neill, C. & Toussaint, J.P. (2004) Plant functional markers capture ecosystem properties during secondary succession. *Ecology*, **85**, 2630-2637.

Grime, J.P., Hodgson, J.G. & Hunt, R. (2007) *Comparative Plant Ecology*. 2nd edn. Dalbeattie, Scotland: Castlepoint Press.

Gunn, S., Farrar, J.F., Collis, B.E. & Nason, M. (1999) Specific Leaf Area in Barley: individual leaves versus whole plants. *New Phytologist*, **143**, 45-51.

Haberl, H., Erb, K-H., Krausmann, F., Gaube, V., Bondeau, A., Plutzer, C., Gingrich, S., Lucht W. & Fischer-Kowalski, M. (2007) Quantifying and mapping the human appropriation of net primary production in earth's terrestrial ecosystems. *Proceedings of the National Academy of Sciences of the USA*, **104**, 12942–12947.

Hill, M. O., Blackstock, T. H., Long, D. G. & Rothero, G. (2008) *A checklist and census catalogue of British and Irish bryophytes: updated 2008*. Middlewich: British Bryological Society.

Hodgson, J.G., Montserrat-Marti, G., Charles, M., Jones, G., Wilson, P., Shipley, B., Sharafi, M., Cerabolini, B.E.L., Cornelissen, J.H.C., Band, S.R., Bogard, A., Castro-Diez, P., Guerro-Campo, J., Palmer, C., Perez-Rontome, M.C., Carter, G., Hynd, A., Romo-Diez, A., Espuny, L.D. & Pla, F.R. (2011) Is leaf dry matter content a better predictor of soil fertility than specific leaf area? *Annals of Botany*, **108**, 1337-1345.

Hollinger, D.Y. (1989) Canopy organization and foliage photosynthetic capacity in a broad-leaved evergreen montane forest. *Functional Ecology*, **3**, 53-62.

Huslof, C.M., Violle, C., Spasojevic, M.J., McGill, B., Damschen, E., Harrison, S. & Enquist, B.J. (2013) Intra-specific and inter-specific variation in specific leaf area reveal the importance of abiotic and biotic drivers of species diversity across elevation and latitude. *Journal of Vegetation Science*, **24**, 921-931.

Jagodziński, A.M., Dyderski, M.K., Rawlik, K. & Kątna, B. (2016) Seasonal variability of biomass, total leaf area and specific leaf area of forest understorey herbs reflects their life strategies. *Forest Ecology and Management*, **374**, 71-81.

Jung, M., Vetter, M., Herold, M., Churkina, G., Reichstein, M., Zaehle, S., Ciais, P., Viovy, N., Bondeau, A., Chen Y *et al.* (2007) Uncertainties of modeling gross primary productivity over Europe: a systematic study on the effects of using different drivers and terrestrial biosphere models. *Global Biogeochemical Cycles*, **21**, (GB4021).

Jung, V., Violle, C., Mondy, C., Hoffmann, L., Muller, S. (2010) Intraspecific variability and trait-based community assembly. *Journal of Ecology*, **98**, 1134-1140.

Kivimäki, S.K. (2011) *Changes in carbon and nitrogen dynamics in Sphagnum capillifolium under enhanced nitrogen deposition*. PhD thesis. University of Edinburgh.

Kleyer, M., Bekker, R.M., Knevel, I.C., Bakker, J.P., Thompson, K., Sonnenschein, M., Poschlod, P., van Groenendael, J.M., Klimes, L., Klimesova, J. *et al.* (2008) The LEDA traitbase: a database of life-history traits of the NW European flora. *Journal of Ecology*, **96**, 1266-1274.

Laughlin, D.C., Chaitanya, J., van Bodegom, A.M., Bastow, Z.A. & Fulé, P.Z. (2012) A predictive model of community assembly that incorporates intraspecific trait variation. *Ecology Letters*, **15**, 1291-1299.

Lunn, D., Best, N., Spiegelhalter, D., Graham, G. & Neuenschwander, B. (2009) Combining MCMC with PKPD modelling. *Journal of Pharmacokinetics Pharmacodynamics*, **36**, 19-38.

Lunn, D., Jackson, C., Best, N., Thomas, A. & Spiegelhalter, D. (2013) *The BUGS book: A practical introduction to Bayesian analysis*. CRC Press, Taylor & Francis Group, Florida, USA, Pp.198-200.

Manning, P., de Vries, F.T., Tallowin, J.R.B., Smith, R., Mortimer, S.R., Pilgrim, E.S., Harrison, K.A., Wright, D.G., Quirk, H., Benson, J. *et al.* (2015) Simple measures of climate, soil properties and plant traits predict national-scale grassland soil carbon stocks. *Journal of Applied Ecology*, **52**, 1188-1196.

McNaughton, S.J., Milchunas, D.G. & Frank, D.A. (1996) How can net primary productivity be measured in grazing ecosystems? *Ecology*, **77**, 974-977.

Minden, V. & Kleyer, M. (2015) Ecosystem multifunctionality of coastal marshes is determined by key plant traits. *Journal of Vegetation Science* **26**, 651-662.

Nakagawa, S. & Schielzeth, H. (2014) A general and simple method for obtaining R<sup>2</sup> from generalized linear mixed effects models. *Method in Ecology and Evolution*, **4**, 133-142.

Ordoñez, J.C., van Bodegom, P.M., Witte, J.P.M., Wright, I.J., Reich, P.B. & Aerts, R. (2009) A global study of relationships between leaf traits, climate and soil measure of nutrient fertility. *Global Ecology and Biogeography*, **18**, 137-149.

Pakeman, R. (2011) Multivariate identification of plant functional response and effect traits in an agricultural landscape. *Ecology*, **92**, 1353-1365.

Peel, M.C., Finlayson, B.L. & McMahon, T.A. (2007) Updated world map of the Köppen-Geiger climate classification. *Hydrology & Earth System Sciences*, **11**, 1633–1644.

Pérez-Harguindeguy, N., Díaz, S., Garnier, E., Lavorel, S., Poorter, H., Jaureguiberry, P., Bret-Harte, M.S., Cornwell, W.K., Craine, J.M., Gurvich, D.E., Urcelay, C., Veneklaas, E.J., Reich, P.B., Poorter, L., Wright, I.J., Ray, P., Enrico, L., Pausas, J.G., de Vos, A.C., Buchmann, N., Funes, G., Quétier, F., Hodgson, J.G., Thompson, K., Morgan, H.D., ter Steege, H., van der Heijden, M.G.A., Sack, L., Blonder, B., Poschlod, P., Vaieretti, M.V., Conti, G., Staver, A.C., Aquino, S. & Cornelissen, J.H.C. (2013). New handbook for standardised measurement of plant functional traits worldwide. *Australian Journal of Botany* **61**, 167-234.

Pierce, L.L., Running, S.W. & Walker, J. (1994) Regional-scale relationships of Leaf Area Index to Specific Leaf Area and Leaf Nitrogen Content. *Ecological Applications*, **4**, 313-321.

Poorter, H., Niklas, K.J., Reich, P.B., Oleksyn, J., Poot, P. & Mommer, L. (2012) Biomass allocation to leaves, stems and roots: meta-analyses of interspecific variation and environmental control. *New Phytologist*, **193**, 30-50.

Pontes, L. da S., Soussana, J –F., Louault, F., Andueza, D. & Carrère P. (2007) Leaf traits affect the above-ground productivity and quality of pasture grasses. *Functional Ecology*, **21**, 844-853.

Reich, P.B. 2014. The world-wide ‘fast-slow’ plant economics spectrum: a traits manifesto. *Journal of Ecology*, **102**, 275-301.

Roche, P., Díaz-Burlinson, N. & Gachet, S. (2004) Congruency analysis of species ranking based on leaf traits: which traits are the more reliable? *Plant Ecology* **174**, 37-48.

Scurlock, J.M.O., Johnson, K. & Olson, R.J. (2002) Estimating net primary productivity from grassland biomass dynamics measurements. *Global Change Biology*, **8**, 736-753.

Siefert, A., Fridley, J.D. & Ritchie, M.E. (2014) Community functional responses to soil and climate at multiple spatial scales: when does intraspecific variation matter? *PLOS ONE*, **9**, e111189.

Siefert, A., Violle, C., Chalmandrier, L., Albert, C.H., Taudiere, A., Fajardo, A., Aarssen, L.W., Baraloto, C., Carlucci, M.B., Cianciaruso, M.V. *et al.* (2015) A global meta-analysis of the relative extent of intraspecific trait variation in plant communities. *Ecology Letters*, **18**, 1406-1419.

Shipley, B. (1995) Structured interspecific determinants of Specific Leaf Area in 34 species of herbaceous angiosperms. *Functional Ecology*, **9**, 312-319.

Stace, C. (1997) *New Flora of the British Isles*. 2<sup>nd</sup> edn. Cambridge University Press, Cambridge, UK.

Suding, K.N., Lavorel, S., Chapin, F.S., Cornelissen, J.H.C., Díaz, S., Garnier, E., Goldberg, D., Hooper, D.U., Jackson, S.T. & Navas, M.L. (2008) Scaling environmental change through the community-level; a trait-based response-and-effect framework for plants. *Global Change Biology*, **14**, 1125-1140.

Tateno, M. & Chapin III, F.S. (1997) The logic of carbon and nitrogen interactions in terrestrial ecosystems. *American Naturalist*, **149**, 723-744.

Accepted Article

Van Bodegom, P.M., Douma, J.C., Witte, J.P.M., Ordoñez, J.C., Bartholomeus, R.P. & Aerts, R. (2012) Going beyond limitations of plant functional types when predicting global ecosystem-atmosphere fluxes: exploring the merits of trait-based approaches. *Global Ecology & Biogeography*, **21**, 625-636.

Vendramini, F., Díaz, S., Gurvich, D.E., Wilson, P.J., Thompson, K. & Hodgson, J.G. (2002) Leaf traits as indicators of resource-use strategy in floras with succulent species. *New Phytologist*, **154**, 147-157.

Weihner, E., van der Werf, A., Thompson, K., Roderick, M., Garnier, E. & Eriksson, O. (1999) Challenging Theophrastus: A common core list of plant traits for functional ecology. *Journal of Vegetation Science*, **10**, 609-620.

Wilson, P.J., Thompson, K. & Hodgson, J.G. (1999) Specific leaf area and leaf dry matter content as alternative predictors of plant strategies. *New Phytologist* **143**, 155-162.


Wright, I.J., Reich, P.B., Cornelissen, J.H.C., Falster, D.S., Garnier, E., Hikosaka, K., Lamont, B.B., Lee, W., Veneklaas, E.J., Oleksyn, J. *et al.* (2005) Assessing the generality of global leaf-trait relationships. *New Phytologist*, **166**, 485-496.

Wright, J.P., Naeem, S., Hector, A., Lehman, C., Reich, P.B., Schmid, B. & Tilman, D. (2006) Conventional functional classification schemes underestimate the relationship with ecosystem functioning. *Ecology Letters*, **9**, 111-120.

Zuur, A.F., Ieno, E.N. & Elphick, C.S. (2010) A protocol for data exploration to avoid common statistical problems. *Methods in Ecology and Evolution*, **1**, 3-14.

Figure 1: Maps showing the sample sites in (a) North West England and (b) North Wales.

a)



b)




Figure 2: Measured above-ground NPP (aNPP) values across temperate ecosystem types sampled in 2013 and 2014. The median is shown as a black point. Boxes indicate the interquartile range and the whiskers the range of the measurements.






Figure 3: Best fitting model of  $\ln(\text{aNPP})$  predicted by cover-weighted Leaf Dry Matter Content incorporating the effect of database-derived intra-specific variation and including *in situ* trait measurements.  $R^2(m)=0.55$ .  $y=x$  line is shown.




Table 1: Details of study sites and plots in which aNPP was measured. Sampling methods are fully described in Supplementary Material. Nomenclature for vascular plants follows Stace (1997) and Hill *et al.* (2008) for bryophytes.

Site	Habitat type	Mean soil pH (0-15cm)	Management status	Dominant plant species	Number of plots (plot size)	aNPP methods
Beryl's Wood	Broadleaved woodland	4.62	Unmanaged	<i>Quercus petraea/robur</i> , <i>Fraxinus excelsior</i> , <i>Betula pendula</i>	2 (200m <sup>2</sup> tree canopy), 3 (1m <sup>2</sup> understorey)	Understorey biomass harvest.  Bryophyte mesh.  Litter buckets.  Annual woody increment from tree ring core, tree height & DBH.
Red Kite Wood	Broadleaved woodland	4.19	Unmanaged	<i>Acer pseudoplatanus</i>	1 (200m <sup>2</sup> tree canopy), 2 (1m <sup>2</sup> understorey)	As above.
Coed Dolgarrog	Broadleaved woodland	3.98	Unmanaged	<i>Quercus petraea/robur</i>	1 (200m <sup>2</sup> tree canopy), 2 (1m <sup>2</sup> understorey)	As above.
Glasgwm	Conifer plantation	4.2	30 year old Sitka subject to past thinning	<i>Picea sitchensis</i>	1 (200m <sup>2</sup> tree canopy), 2 (1m <sup>2</sup> understorey)	As above.

Nant-y-Coed	Improved grassland	5.68	Highly intensive cattle grazing. Cattle rotated fortnightly across heavily fertilized paddocks.	<i>Lolium perenne</i>	4 (1m <sup>2</sup> )	Two biomass harvests per year.
Blaen-y-Coed	Soligenous mire	4.56	Low intensity sheep grazing with periods unmanaged	<i>Molinia caerulea</i>	2 (1m <sup>2</sup> )	As above.
Migneint	Ombrogenous mire	3.82	Last burnt 30+ years ago. Very light sheep grazing.	<i>Sphagnum capillifolium</i> , <i>Eriophorum vaginatum</i>	16 cranked wires among 4 patches of <i>Sphagnum</i>	Wire length measurements over two years.
Nant-y-Brwyn	Ombrogenous mire	4.26	Last burnt 30+ years ago. Very light sheep grazing.	<i>Sphagnum fallax</i> , <i>Juncus effusus</i>	4 (1m <sup>2</sup> ) 6 cranked wires among 3 patches of <i>Sphagnum</i>	Two biomass harvests per year. Wire length measurements over two years.
Llyn Serw	Ombrogenous mire	3.82	Last burnt 30+ years ago. Very	<i>Calluna vulgaris</i>	3 (1m <sup>2</sup> in each of 5, 11, 30 year old <i>Calluna</i> )	Total biomass harvest for

			light sheep grazing.			growth curve construction.
Capel Curig VB	Soligenous mire	4.07	Very light sheep and cattle grazing.	<i>Molinia caerulea</i> , <i>Sphagnum papillosum</i>	4 (1m <sup>2</sup> ), 12 cranked wires among 4 patches	Two biomass harvests per year.  Wire length measurements over two years.
Capel Curig AG	Acid grassland	4.81	Light sheep and cattle grazing.	<i>Deschampsia flexuosa</i> , <i>Nardus stricta</i>	4 (1m <sup>2</sup> )	Two biomass harvests per year.
Carneddau	Montane heath	4.40	Light sheep grazing.	<i>Empetrum nigrum</i> , <i>Salix herbacea</i>	3 (0.25m <sup>2</sup> )	One biomass harvest per year.
Juniper Gill	Calcareous grassland	7.46	Wild deer and rabbit grazed.	<i>Sesleria caerulea</i>	2 (0.25m <sup>2</sup> )	Two biomass harvests per year.
Scar Close	Calcareous grassland	-	Wild deer and rabbit grazed.	<i>Sesleria caerulea</i> , <i>Pteridium aquilinum</i>	2 (0.25m <sup>2</sup> )	Two biomass harvests per year.
Colt Park	Unimproved grassland	5.07	Traditional hay meadow; cattle and sheep in	<i>Trifolium pratense</i> , <i>Anthoxanthum odoratum</i>	4 (0.25m <sup>2</sup> )	Two biomass harvests per year.

			spring then shut up for summer hay growth, then aftermath grazed.			
Hiraethlyn	Improved grassland	4.59	Intensive sheep grazing.	<i>Lolium perenne</i> , <i>Holcus lanatus</i>	4 (1m <sup>2</sup> )	Two biomass harvests per year.
Ysbyty-lfan IG	Improved grassland	5.67	Intensive cattle and sheep grazing.	<i>Lolium perenne</i> , <i>Poa trivialis</i>	4 (1m <sup>2</sup> )	Two biomass harvests per year.
Ysbyty-lfan SG	Semi-improved grassland	5.58	Intensive cattle and sheep grazing.	<i>Lolium perenne</i> , <i>Holcus lanatus</i>	4 (1m <sup>2</sup> )	Two biomass harvests per year.