

Climate change mitigation: potential benefits and pitfalls of enhanced rock weathering in tropical agriculture

David P. Edwards¹, Felix Lim¹, Rachael H. James², Christopher R. Pearce³, Julie Scholes¹, Robert P. Freckleton¹ and David J. Beerling¹

¹Department of Animal and Plant Sciences, University of Sheffield, Sheffield S10 2TN, UK

²Ocean and Earth Science, National Oceanography Centre Southampton, University of Southampton, Southampton, UK

³National Oceanography Centre, University of Southampton Waterfront Campus, European Way, Southampton, SO14 3ZH

Author for correspondence:

David P. Edwards

email: david.edwards@sheffield.ac.uk

Running head

Tropical farming and enhanced weathering

Keywords:

carbon dioxide removal, global temperature, negative emissions technologies, oil palm, agroecosystems and monoculture tree plantations, silicate weathering

Abstract

Restricting future global temperature increase to 2°C or less requires the adoption of Negative Emissions Technologies for carbon capture and storage. We review the potential for deployment of enhanced weathering (EW), via the application of crushed reactive silicate rocks (such as basalt), on over 680 million hectares of tropical agricultural and tree plantations to offset fossil fuel CO₂ emissions. Warm tropical climates and productive crops will substantially enhance weathering rates, with potential co-benefits including decreased soil acidification and increased phosphorus-supply promoting higher crop yields sparing forest for conservation, and reduced cultural eutrophication. Potential pitfalls include the impacts of mining operations on deforestation, producing the energy to crush and transport silicates, and the erosion of silicates into rivers and coral reefs that increase inorganic turbidity, sedimentation, and pH with unknown impacts for biodiversity. We identify nine priority research areas for untapping the potential of EW in the tropics, including effectiveness of tropical agriculture at EW for major crops in relation to particle sizes and soil types, impacts on human health, and effects on farmland, adjacent forest, and stream-water biodiversity.

45

1. Enhanced weathering as a negative emissions strategy

The 2015 Paris Agreement on climate change recognizes that restricting future temperature increases to 1.5-2°C requires deployment of unproven Negative Emissions Technologies (NETs) to remove CO₂ from the atmosphere. Currently, all proposed large-scale NETs have poorly developed feasibility, cost and acceptability [1] and few, if any, have had their impacts on ecosystem services or biodiversity considered [2].

Here we focus on the potential and consequences for the deployment of enhanced weathering (EW) on tropical agricultural lands by exploiting existing agricultural infrastructure. EW involves application of crushed reactive silicate rocks (particularly basalt and other mafic rocks) to vegetated landscapes to increase atmospheric CO₂ removal rates [3-5]. Natural rock weathering is regulated by climate and vegetation. CO₂ is removed by the chemical breakdown of calcium- and magnesium-rich silicate rocks and is accelerated by warm climates and vegetation rooting systems and their ubiquitous root-associating symbiotic fungi [6]. Weathered base cations and resulting bicarbonate in soils are flushed into rivers and delivered into the surface oceans, where CO₂ is stored either as dissolved inorganic carbon or permanently (on human timescales) as carbonate. Lower atmospheric CO₂ and an increased land-ocean flux of alkalinity generated by EW might help counteract ocean acidification [3, 5].

In this review, we briefly introduce why the tropics are likely to be particularly effective for EW and the kinds of tropical agricultural systems that could be used. We discuss the potential positives and pitfalls of tropical EW, both within the agroecosystems themselves and on wider-scales, and finish by providing a roadmap of critical outstanding research questions.

2. Why the tropics?

Silicate weathering rates depend on temperature, runoff and rate of physical erosion [7, 8]. Although warm and wet tropical conditions should theoretically enhance the rate of silicate rock weathering (Fig. 1a), natural rates are often very low [9] because lowland tropical environments are predominantly characterised by thick, mature soils that undergo little physical disturbance (Fig. 1). Primary minerals within these soil sequences have already been altered to weathering-resistant secondary minerals

depleted in the soluble cations (Ca^{2+} , Mg^{2+} , Na^+ , K^+) that support plant growth.

Furthermore, areas covered with thick layers of weathered soil prevent root access to fresh bedrock, and the roots themselves stabilise the soil surface reducing erosion and
80 lowering chemical weathering potential (Fig. 1b; [10]). Consequently, unlike other climate zones where the rate of silicate weathering is primarily controlled by kinetics, the rate of natural rock weathering in the tropics is limited by the supply of fresh mineral surfaces [7, 8].

Basalts are among the most susceptible silicate rocks to weathering (e.g., [11]).
85 Present-day CO_2 consumption from silicate weathering indicates that around 35% could be attributable to basaltic rocks, even though they constitute less than 5% of the continental area [12]. Amending tropical soils with freshly ground basalt could overcome issues associated with mineral supply and release the geochemical potential of the tropics for atmospheric CO_2 capture and storage (e.g., [5]; Fig. 1c). This will be
90 further enhanced by the secretion of organic acids and CO_2 during respiration by roots and acidification of the rhizosphere by root-associated mycorrhizal fungi [6]. Catchment-scale studies indicate that vegetation can increase weathering rates by five-fold or more compared to adjacent barren areas [6]. These considerations make the warm, highly productive tropics ideal for utilising EW as means of CO_2 removal.

95

3. Potential tropical agricultural systems for EW

We combine data from multiple sources to illustrate and compare the spatial extents and distribution of major land-use types across the tropics (Figure 2). Pan-tropically, over 676 million hectares (Mha) of land was under crop production in 2010 (Table S1),
100 indicating an extensive land area with potential for the large-scale application of EW. Tropical agriculture in each region is dominated by a few crops (Figure 2): Asia dominates production of rice, oil palm, seed cotton, coconut, and rubber; the Neotropics production of soybeans, sugar cane, and coffee; and Africa production of sorghum, millet, cowpeas, and cocoa. Given their extent and distribution, only twenty crops
105 accounted for 548 Mha (81%) of 2010 production (Table S1). Targeting these dominant crops for EW could maximise its effectiveness and efficiency. Additionally, substantial tree plantations of *Eucalyptus*, *Acacia*, etc. for paper-pulp and softwood exist in Brazil (7.3 Mha) and Indonesia (2.6 Mha) that might be utilized for EW (Figure 2). EW might

also have a role within forest restoration projects. Extensive tropical restoration
110 required for re-establishing lost biomass carbon sinks [13] might be deployed for EW to
further enhance carbon sequestration.

Crops (e.g., soybean, sugar cane, oil palm), tree and rubber plantations grown
intensively by large- to medium-scale agribusiness have the road and employment
infrastructural capacity required for spreading crushed silicates with many already
115 applying crushed limestone, as agricultural lime, and fertilizer [14]. By contrast, small-
scale farmers, especially those practicing shifting (slash-and-burn) agriculture, will
likely lack sufficient resources to apply crushed rocks. These practices make up a
substantial component of all tropical farming: shifting agriculture spans an estimated
258 Mha, with ~6-19% farmed annually; the remainder is naturally regenerating as
120 forest [15]. However, these systems are transitioning to more permanent and
mechanized farming with inputs, including via small-holders selling or leasing farmland
for monoculture conversion [16]. Further, improvements to road networks in such
areas aimed at reducing yield gaps [17] would aid the delivery of crushed silicates.
Thus, over time, much of these systems will probably become suitable for EW.

125

4. Potential positives

(a) Improved productivity and reduced CO₂ emissions from agriculture

Silicate rocks contain P, Mg, K, and Ca, which are limiting nutrients for plant growth,
thus their release via EW can fertilize crops [5]. There is a long history of amending soils
130 with ground silicate rocks to improve crop yields, especially in highly weathered
tropical soils in Africa and Brazil [18, 19]. For example, cocoa plants applied with basalt
(5 or 10 t ha⁻¹) had higher concentrations of K (1.4-fold), Mg (10-fold), and Ca (1.7-fold)
than untreated controls [20]; after 24 months, treated plants were 50% taller and 60%
thicker-stemmed than controls [20]. In many cases, silicate rocks are likely to be applied
135 in combination with fertiliser and/or manure. In Mauritius, addition of 60–250 t basalt
ha⁻¹, in combination with standard N, P, K fertilizer treatments, increased yields by 29%
over five successive crops and by 17% over three successive crops in two different sets
of replicated trials compared with plots receiving fertilizer only and no basalt addition
[21], indicating a positive interaction between basalt and fertilizer.

140 EW also releases silica into the soil and is taken up as silicic acid by major
tropical crops, including rice, oil palm, sugar cane, maize, and sorghum [3, 22, 23],
helping to confer resistance to economically important pests and diseases [3, 22, 23],
via mechanical cell wall strengthening (deposition of silicon within tissues) and defence
priming [24, 25]. Silicon also improves water-use efficiency by lowering leaf
145 transpiration rates, potentially increasing crop resilience to drought [3]. Application of
silicate rocks for EW might therefore contribute to improving food security in drought-
threatened areas and reduce the use and costs of pesticides.

Application of crushed basalt increases pH on highly weathered tropical soils
[20] and helps mitigate soil acidification in agricultural regions more generally [26] and
150 production constraints in crops established on acidic soils (e.g., heavy metal toxicity in
plants [20], including oil palm on drained peatlands in Southeast Asia [27]. EW effects
on soil pH broadly mirror those of liming agricultural soils to reduce acidification [28].
Substituting silicate EW for liming averts CO₂ emitted when lime reacts with soil water
and during its production [28].

155

(b) Land sparing

Expansion of tropical agricultural area continues at high rates (102 Mha from 2000-
2010; Table S1), mainly via deforestation (e.g., [27]). If application of silicate rocks
improves crop yields, food demand might be met on reduced land area, resulting in less
160 deforestation and/or more natural forest regeneration on abandoned marginal farms.
The Green Revolution in Asia and Latin America was land and greenhouse gas emissions
sparing [29], suggesting increased yields produced by EW could offer further land
savings. Absence of effective market regulation and land planning, however, may cause
perverse outcomes of higher-yielding, cheap tropical crops, including further
165 deforestation [30].

(c) Reduced risk of phytoplankton blooms in rivers and reefs

Fertilisers applied at high doses and incorrect times of year in tropical farmland are
frequently eroded, and deposited into rivers and nearby oceans, causing large
170 phytoplankton blooms [31], including toxic blue-green algae. Threat of eutrophication is

dependent on Si:N and Si:P ratios in run-off water [32, 33]. Cultural eutrophication occurs when high N and P but low Si causes algal blooms. EW of silicate rocks will likely generate high Si:P and Si:N ratios in run-off, increasing diatoms that remove nutrients from the water, preventing cultural eutrophication, and instead supporting diverse and productive food webs [33]. This could be a significant benefit for polluted riverine, reef, and oceanic ecosystems downstream of major areas of tropical agricultural production, while increased diatom production could increase CO₂ drawdown in the oceans [4, 34].

5. Potential pitfalls

180 (a) GHG emissions from grinding and transport

Global analyses indicate energy costs (i.e., CO₂ emissions) associated with mining, grinding and spreading rock dust could decrease efficiency of CO₂ sequestration by EW by 10-25%, depending on grain size [35]. However, this cost will likely decline as the world transitions to decarbonised energy sources. Increased transportation of crushed rock would increase NO_x emissions. In 16 Mha of oil palm plantations, which are high isoprene emitters, this could raise ground-level ozone (O₃) at harmful levels for plant and human health [36].

(b) Yield quality

190 Potentially toxic elements contained in some silicate minerals could become bioavailable under EW, reducing yields or accumulating in the food chain [3], with human health issues. In particular, high nickel and chromium content in olivine would be problematic in agriculture and in association with asbestos-related minerals in major mines [5]. EW with basalt appears the pragmatic choice for application in tropical agriculture to avoid unintended negative consequences [5]. The trade-off is that, theoretically at least, basalt is less effective than olivine for CO₂ capture (e.g., ~0.3 tCO₂ t⁻¹ basalt vs 0.8 tCO₂ t⁻¹ olivine [37]). Ancillary benefits of basalt for crop production, soil improvement and suppression of GHG emissions that are less likely to accrue from olivine and the lack of heavy metal toxicity would lower the practical barriers to take-up
200 by farmers in tropical agroecosystems.

(c) Biodiversity impacts within plantations and adjacent forest

Tropical farmland has wildlife that provides important ecosystem services for humans, including pollination and pest control. How these species will respond to silicate application is unknown. In particular, increasing pH could have negative consequences for species adapted to low pH soils, which are widespread in tropical regions, especially in peatlands. Forest edges are affected by environmental changes (e.g., increased wind, higher nutrient loads) that penetrate tens to hundreds of metres into forest interiors [38]. How far crushed silicates penetrates into forest from farmland and what the consequences would be for biodiversity adapted to nutrient-poor and acidic mature soils are uncertain. If consequences were negative, then this would be a major concern, given that 25% of the Amazon and Congo, and 91% Brazilian Atlantic forest is within 1 km of farmland edge [39].

(d) Reduced water quality in rivers and reefs

If unweathered silicates are washed into rivers, perhaps during intense tropical rainstorms, increased inorganic turbidity and sedimentation might follow, reducing reproduction and recruitment in river fish populations [40]. Higher sediment loads and inorganic turbidity cause coral mortality and reductions in reef diversity and depth limit [41]. There are thus potentially severe negative implications for local fisheries and conservation, although such losses would need to be weighed against any benefits gained from reduced organic turbidity (i.e., lower eutrophication, see section 4(c) above). Increased water pH might also negatively impact riverine plants and animals, especially in naturally acidic drainages (e.g., peatlands).

225

(e) Mining and infrastructural expansion

Although silicates are a waste product from mining and steel and iron production [42], if applied pan-tropically then new or larger mines could be required. For instance, rock application to 670 Mha of tropical cropland at 10 t ha⁻¹ yr⁻¹ would require 6.7 Pg of rock per year, and at 50 t ha⁻¹ yr⁻¹ would need 33.5 Pg annually [5]. By comparison, global coal production was 8.1 Pg in 2015 [43] and global aggregate production 40 Pg [44].

Mine creation is environmentally destructive, driving deforestation across the tropics and often occurring within or near to areas of high biodiversity value [45]. Development and expansion of road and rail infrastructure for mining can increase access to
235 biodiverse and remote ecosystems [45], which combined with employment opportunities, encourage population immigration, land clearing for agriculture and hunting [45].

6. Future directions and conclusions

240 We highlight nine major outstanding questions, indicating the need for further research on EW and clear protocols and regulations for any pan-tropical roll-out.

(1) How effective is tropical agriculture at enhanced rock weathering? Effectiveness of tropical agricultural systems at EW is a critical unknown and requires replicated pot experiments under field conditions for different key crops (Figure 2; Table S1), soil
245 types, application rates, and particle sizes. Resolving effective particle sizes that can be adopted in tropical agriculture will be critical because of the high energy costs associated with grinding rocks to fine particle sizes ($<10\ \mu\text{m}$ diameter) [35]. Once these questions have been addressed, field-scale trials are required to understand additional effects of catchment topography, drainage and soils on EW rates and evaluate
250 biogeochemical models. This information is critical for informing accurate spatial projections of pan-tropical carbon capture for EW in agriculture.

(2) What are the long-term effects of EW on farms and neighbouring forest? We need to quantify a range of processes at catchment scales before and after the application of silicate for multiple years (Shao et al. [46] added silicate (wollastonite) to the Hubbard
255 brook catchment and found effects lasting over a decade). These should include rates of weathering, as well as impacts on yield, sediment and chemical runoff into streams, and biodiversity within plantations. Application rates for crushed silicates required for carbon capture are uncertain (e.g., $\sim 10\text{-}50\ \text{t ha}^{-1}\ \text{yr}^{-1}$ [5]) and could be higher than current estimates. In practice, application rates would be optimized for crop type,
260 prevailing climate and soil, but will likely exceed those used for liming. On widespread highly weathered oxisols in the tropics, annual liming rates to obtain 90% of maximum yield (i.e., maximum economic rate) can reach $9\ \text{t ha}^{-1}$ for soybean, $8\ \text{t ha}^{-1}$ for corn, $6\ \text{t}$

ha⁻¹ for cotton and 3.8 t ha⁻¹ for sugarcane [47], with usual application rates for
Brazilian soy of ~4-6 t ha⁻¹ yr⁻¹ [14]. A key question is what happens to the
265 unweathered materials: if it accumulates in farmland or washes into rivers, then we
need to understand the implications for major biogeochemical processes and
biodiversity. Precision application methods might be necessary to optimize rates of
application and EW whilst minimizing any harmful biological effects.

Adopting farm catchments in proximity to natural forest will enable monitoring
270 of silicates penetration into adjacent forest, including if/how they affect plant growth,
interactions between species and biodiversity conservation value. If edge effects of EW
are severe, then research should identify which forest patches have sufficiently high
conservation value to require protection, and in those cases, silicates should only be
applied at a minimum distance from forest edge.

275 *(3) What is the effect of EW on tropical agriculture yields?* Using pot (1) and catchment-
scale (2) experiments, we need to investigate how crop yield is affected by EW and
investigate yield quality to determine the grades of silicate rocks that do not risk
bioaccumulation of toxic metals. Is the fertiliser effect sufficient to allow farmers to
reduce (or cease) application of commercially produced fertilisers? These data will
280 allow assessment of economic costs and benefits of EW to farmers, and determine when
and for which crops yield benefits are sufficient to promote adoption by agriculture.

Additional co-benefits of EW need to be understood given they might incentivise
widespread adoption. These include the benefits of increasing soil pH of widespread
highly weathered acidic tropical soils, increased plant resistance to pests, diseases and
285 drought. Each could reduce or remove the necessity for liming, pesticides and
fungicides, and increase crop yields with drought.

(4) How does EW affect hydrological cycles, rivers and coral reefs? By increasing plant
water-use efficiency EW might alter local hydrologic cycles, and this must be modelled
[3]. We also need to understand fluxes into rivers and coral reefs from treated
290 catchments to quantify likely effects on sedimentation, turbidity, pH, and enhanced Si:N
and Si:P ratios. This will identify the net balance between the potential positives of
reduced ocean acidification and cultural eutrophication versus the negatives of poorer
water quality. By sampling biodiversity within streams of catchment studies (2), any

295 local-scale impacts would provide an early warning system to larger river- or reef-scale impacts.

(5) How to minimise human health risks with silicate application? At small particle sizes, there are health risks for workers crushing or spreading silicates, including silicosis and other respiratory diseases [5]. Especially in areas where agriculture is not managed by agri-business, this would require a pan-tropical investment in education, safety
300 equipment and protocols. Additionally, application in tropical dry seasons could lead to large quantities of silicates being eroded by wind with potential issues for local population settlements.

(6) Can EW link with large-scale tropical reforestation programmes? As in (1), we need to understand optimal grain size and application of EW in large-scale reforestation
305 systems and how that affects growth and carbon sequestration across a range of tree species with differing mycorrhizal associations and soil types. We also need to understand whether it would be cost-efficient to apply EW to reforestation, given a lack of long-term manpower and transport networks, and impacts on biodiversity and ecosystem services.

(7) Will there be unintended mining and transport impacts of EW and how can they be prevented or mitigated? We need to understand the mass of silicate rock required for
310 tropic-wide application of EW and whether existing mines and infrastructure can meet this demand. If they cannot, then we must predict likely sources of silicates and resulting on and off-mine consequences for deforestation, biodiversity loss and
315 socioeconomic change. Investors in 'conservation mining' to reduce climate change via EW must then demand strict environmental standards to prevent such on and off mine impacts.

(8) Will the carbon savings from EW outweigh the carbon costs of producing and applying silicates? In (1) we highlight a need to understand the optimal particle size and
320 application quantities to maximise EW and thus CO₂ sequestration rates, plus CO₂ emissions savings from avoided liming. This needs to be balanced against the energy costs of mining, grinding, transport and spreading via a full life cycle assessment analysis across the tropics and different crop types. A related issue will likely be the
325 need to innovate and develop new high-efficiency low-carbon emitting grinding technologies, including adopting solar energy in tropical regions.

(9) *What role might carbon markets play in incentivising roll-out of EW?* We need to calculate the carbon market cost ($\text{\$t}^{-1} \text{CO}_2$) to subsidise silicate application across a range of crop types, and socioeconomic (e.g., labour cost) and geographic (distance to market, etc.) scenarios to make EW no net cost or profitable to farmers. This will entail
330 understanding and modelling the full range of economic costs and profits of EW,
combined with net carbon budgets from (8).

Conclusion

EW is a promising NET option that could deliver significant co-benefits to tropical
335 agriculture and coastal ocean ecosystems. However, major issues remain regarding the potential effectiveness of EW and the associated benefits and pitfalls of the related operation for tropical agroecosystems and natural habitats. If empirical evidence from field studies and carbon cycle modelling demonstrate a significant capacity of pan-tropical agroecosystems for net long-term carbon sequestration, then these benefits to
340 humanity will need balancing against negative impacts on biodiversity and ecosystem services.

Acknowledgements

345 We thank Paul Nelson and Michael Bird for generously sharing ideas, and gratefully acknowledge funding through a Leverhulme Research Centre Award (RC-2015-029). The authors declare no competing interests.

References

- 350 [1] Smith P *et al.* 2016 Biophysical and economic limits to negative CO₂ emissions. *Nat. Clim. Change* **6**, 42-50.
- [2] Williamson, P. 2016 Scrutinize CO₂ removal methods. *Nature* **530**, 153-155.
- [3] Hartmann J, West AJ, Renforth P, Kohler P, De La Rocha CL, Wolf-Gladrow DA, Durr HH, Scheffran J. 2013 Enhanced chemical weathering as a geoengineering strategy to
355 reduce atmospheric carbon dioxide, supply nutrients, and mitigate ocean acidification. *Revs Geophys.* **51**, 113-149.

- [4] Kohler P, Hartmann J, Wolf-Gladrow DA. 2010 Geoengineering potential of artificially enhanced silicate weathering of olivine. *Proc. Natl. Acad. Sci. USA* **107**, 20228-20233.
- 360 [5] Taylor LL, Quirk J, Thorley RMS, Kharecha PA, Hansen J, Ridgwell A, Lomas MR, Banwart SA, Beerling DJ. 2016 Enhanced weathering strategies for stabilizing climate and averting ocean acidification. *Nat. Clim. Change* **6**, 402-406.
- [6] Taylor LL, Leake JR, Quirk J, Hardy K, Banwart SA, Beerling DJ. 2009 Biological weathering and the long-term carbon cycle: integrating mycorrhizal evolution and
365 function into the current paradigm. *Geobiol.* **7**, 171-191.
- [7] West AJ, Galy A, Bickle M. 2005 Tectonic and climatic controls on silicate weathering. *Earth Planet. Sci. Lett.* **235**, 211-228.
- [8] Maher K, Chamberlain CP. 2014 Hydrologic regulation of chemical weathering and the geologic carbon cycle. *Science* **343**, 1502-1504.
- 370 [9] Stallard RF, Edmond JM. 1983 Geochemistry of the Amazon 2. The influence of geology and weathering environment on the dissolved load. *J Geophys. Res.-Oc. Atm.* **88**, 9671-9688.
- [10] Behrens R, Bouchez J, Schuessler JA, Dultz S, Hewawasam T, von Blanckenburg F. 2015 Mineralogical transformations set slow weathering rates in low-porosity
375 metamorphic bedrock on mountain slopes in a tropical climate. *Chem. Geol.* **411**, 283-298.
- [11] Gislason SR, Oelkers EH. 2003 Mechanism, rates, and consequences of basaltic glass dissolution: II. An experimental study of the dissolution rates of basaltic glass as a function of pH and temperature. *Geochim. Cosmochim. Acta* **67**, 3817-3832.
- 380 [12] Dessert C, Dupre B, Gaillardet J, Francois LM, Allegre CJ. 2003 Basalt weathering laws and the impact of basalt weathering on the global carbon cycle. *Chem. Geol.* **202**, 257-273.
- [13] Chazdon RL. 2014 *Second growth*. Chicago, MI: Chicago University Press.
- [14] Clay J. 2004 *World agriculture and the environment: A commodity-by-commodity
385 guide to impacts and practices*. Washington DC: Island Press
- [15] Silva JMN, Carreiras JMB, Rosa I, Pereira JMC. 2011 Greenhouse gas emissions from shifting cultivation in the tropics, including uncertainty and sensitivity analysis. *J Geophys. Res.-Atm.* **116**, D20304
- [16] van Vliet N *et al.* 2012 Trends, drivers and impacts of changes in swidden
390 cultivation in tropical forest-agriculture frontiers: A global assessment. *Glob. Environ. Change* **22**, 418-429.
- [17] Laurance WF *et al.* 2014 A global strategy for road building. *Nature* **513**, 229-232
- [18] Leonardos OH, Fyfe WS, Kronberg BI. 1987 The use of ground rocks in laterite systems - an improvement to the use of conventional soluble fertilizers. *Chem. Geol.* **60**,
395 361-370.
- [19] Van Straaten, P. 2006 Farming with rocks and minerals: challenges and opportunities. *An. Acad. Bras. Ciênc.* **78**, 731-747.

- 400 [20] Anda M, Shamshuddin J, Fauziah CI. 2013 Increasing negative charge and nutrient contents of a highly weathered soil using basalt and rice husk to promote cocoa growth under field conditions. *Soil Tillage Res.* **132**, 1-11.
- [21] de Villiers OD. 1961 Soil rejuvenation with crushed basalt in Mauritius. *Int. Sugar J.* **63**, 363-364.
- 405 [22] Keeping MG, Miles N, Sewpersad C. 2014 Silicon reduces impact of plant nitrogen in promoting stalk borer (*Eldana saccharina*) but not sugarcane thrips (*Fulmekiola serrata*) infestations in sugarcane. *Front. Plant Sci.* **5**, 289
- [23] Najihah NI, Hanafi MM, Idris A, Hakim MA. 2015 Silicon treatment in oil palms confers resistance to basal stem rot disease caused by *Ganoderma boninense*. *Crop Prot.* **67**, 151-159.
- 410 [24] Liang Y, Nikolic M, Bélanger R, Gong H, Song A. 2015 Silicon in agriculture: from theory to practice. *Springer Netherlands*.
- [25] Ye M *et al.* 2013 Priming of jasmonate-mediated antiherbivore defense responses in rice by silicon. *Proc. Natl. Acad. Sci. USA* **110**, E3631-E3639.
- 415 [26] Guo JH, Liu XJ, Zhang Y, Shen JL, Han WX, Zhang WF, Christie P, Goulding KWT, Vitousek PM, Zhang FS. 2010 Significant acidification in major Chinese croplands. *Science* **327**, 1008-1010.
- [27] Wilcove DS, Giam X, Edwards DP, Fisher B, Koh LP. 2013 Navjot's nightmare revisited: logging, agriculture, and biodiversity in Southeast Asia. *Trends Ecol Evol* **28**, 531-540.
- 420 [28] ten Berge HFM, van der Meer HG, Steenhuizen JW, Goedhart PW, Knops P, Verhagen J. 2012 Olivine weathering in soil, and its effects on growth and nutrient uptake in ryegrass (*Lolium perenne* L.): a pot experiment. *PLOS ONE* **7**, e42098
- [29] Hertel TW, Ramankutty N, Baldos ULC. 2014 Global market integration increases likelihood that a future African Green Revolution could increase crop land use and CO2 emissions. *Proc. Natl. Acad. Sci. USA* **111**, 13799-13804.
- 425 [30] Carrasco LR, Larrosa C, Milner-Gulland EJ, Edwards DP. 2014 A double-edged sword for tropical forests. *Science* **346**, 38-40.
- [31] Beman JM, Arrigo KR, Matson PA. 2005 Agricultural runoff fuels large phytoplankton blooms in vulnerable areas of the ocean. *Nature* **434**, 211-214.
- 430 [32] Sommer U, Stibor H, Katechakis A, Sommer F, Hansen T. 2002 Pelagic food web configurations at different levels of nutrient richness and their implications for the ratio fish production: primary production. *Hydrobiologia* **484**, 11-20.
- [33] Kiran MT, Bhaskar MV, Tiwari A. 2016 Phycoremediation of eutrophic lakes using diatom algae. In *Lake sciences and climate change* (ed. MN Rashed), pp. 103-115. *InTech*.
- 435 [34] Carey JC, Fulweiler RW. 2016 Human appropriation of biogenic silicon - the increasing role of agriculture. *Func. Ecol.* **30**, 1331-1339.
- [35] Moosdorf N, Renforth P, Hartmann J. 2014 Carbon dioxide efficiency of terrestrial enhanced weathering. *Environ. Sci. Technol.* **48**, 4809-4816.

- 440 [36] Hewitt CN *et al.* 2009 Nitrogen management is essential to prevent tropical oil palm plantations from causing ground-level ozone pollution. *Proc. Natl. Acad. Sci. USA* **106**, 18447-18451.
- [37] Renforth P. 2012 The potential of enhanced weathering in the UK. *Int. J. Greenh. Gas Control* **10**, 229-243.
- [38] Laurance WF, Lovejoy TE, Vasconcelos HL, Bruna EM, Didham RK, Stouffer PC, Gascon C, Bierregaard RO, Laurance SG, Sampaio E. 2002 Ecosystem decay of Amazonian forest fragments: A 22-year investigation. *Conserv. Biol.* **16**, 605-618.
- 445 [39] Haddad NM *et al.* 2015 Habitat fragmentation and its lasting impact on Earth's ecosystems. *Science Adv.* **1**, e1500052.
- [40] Kemp P, Sear D, Collins A, Naden P, Jones I. 2011 The impacts of fine sediment on riverine fish. *Hydrol. Process.* **25**, 1800-1821.
- 450 [41] Fabricius, K.E. 2005 Effects of terrestrial runoff on the ecology of corals and coral reefs: review and synthesis. *Marine Poll. Bull.* **50**, 125-146.
- [42] Renforth P, Washbourne CL, Taylder J, Manning DAC. 2011 Silicate Production and Availability for Mineral Carbonation. *Environ. Sci. Technol.* **45**, 2035-2041.
- [43] WCA. 2016 *Coal mining*. World Coal Association, <http://www.worldcoal.org/coal/coal-mining> accessed August 2016.
- 455 [44] UEPG. 2016 *GAIN – Global Aggregates Information Network*. European Aggregates Association, <http://www.uepg.eu/media-room/links/gain-global-aggregates-information-network> accessed August 2016.
- [45] Edwards DP, Sloan S, Weng L, Dirks P, Sayer J, Laurance WF. 2014 Mining and the African Environment. *Conserv. Lett.* **7**, 302-311.
- 460 [46] Shao S, Driscoll CT, Johnson CE, Fahey TJ, Battles JJ, Blum JD. 2016 Long-term responses in soil solution and stream-water chemistry at Hubbard Brook after experimental addition of wollastonite. *Environ. Chem.* **13**, 528-540.
- [47] Fageria NK, Baligar VC. 2008 Ameliorating soil acidity of tropical oxisols by liming for sustainable crop production. In *Advances in Agronomy, Vol 99* (ed. DL Sparks), pp. 345-399.
- 465 [48] Monfreda C, Ramankutty N, Foley JA. 2008 Farming the planet: 2. Geographic distribution of crop areas, yields, physiological types, and net primary production in the year 2000. *Glob. Biogeochem. Cycles* **22**, GB1022
- 470 [49] Ramankutty N, Evan AT, Monfreda C, Foley JA. 2008 Farming the planet: 1. Geographic distribution of global agricultural lands in the year 2000. *Global Biogeochemical Cycles* **22**, GB1003
- [50] Transparent World. 2015 *Tree plantations*. <http://www.globalforestwatch.org> accessed August 2016.
- 475 [51] Bontemps S, Defourny P, Bogaert EV, Arino O, Kalogirou V, Perez JR. 2011 *GLOBCOVER 2009-Products description and validation report*. Technical Report for ESA GlobCover project: UCLouvain & ESA Team





Figure 1. Schematic illustration of enhanced rock weathering for CO₂ removal in the
 480 tropics. Relative CO₂ consumption rates are graphed for eight hypothetical tropical
 rivers draining (a) highlands with limited vegetation and thin/absent soil profiles, (b)
 lowlands with thick, mature, weathering-resistant soils, and (c) lowlands dressed with
 reactive ground basalt.



485

490

495

Figure 2. Extent of the 16 most frequently cultivated tropical crops, plus pasture, wood-fibre/timber plantations, rubber plantations and natural forest in (a) Latin America, (b) Africa and (c) Asia-Pacific. Crop and pasture distribution in 2000 (data averaged across 1997–2003) was obtained from [48] and [49]. We displayed the dominant crop of each cell (i.e., with the highest proportion; [48]), provided harvest area exceeded 10% of the cell. Likewise, pastures were displayed if they occupied an area exceeding 10% of the cell, although any crop present (>10% area) was displayed over pasture. Information on the distribution of timber and wood fibre plantations were only obtained for five countries: Brazil, Cambodia, Indonesia, Malaysia and Peru via [50]. Information on the extent of forests across the tropics was obtained for 2009 [51], and includes all forest types. Each habitat was mapped at a resolution of 5 by 5 arcminutes (approximately 10 km x 10 km along the equator).

Climate change mitigation: potential benefits and pitfalls of enhanced rock weathering in tropical agriculture

505 **David P. Edwards¹, Felix Lim¹, Rachael H. James², Christopher R. Pearce³, Julie Scholes¹, Robert P. Freckleton¹ and David J. Beerling¹**

¹Department of Animal and Plant Sciences, University of Sheffield, Sheffield S10 2TN, UK

510 ²Ocean and Earth Science, National Oceanography Centre Southampton, University of Southampton, Southampton, UK

³National Oceanography Centre, University of Southampton Waterfront Campus, European Way, Southampton, SO14 3ZH

515

Author for correspondence:

David P. Edwards

email: david.edwards@sheffield.ac.uk

520 **Table S1.** Year 2000 and 2010 area in millions of hectares (Mha) of tropical crops, as
per FAOSTAT [1]. Area for each crop is given in total and split between the Neotropics,
tropical Africa, and tropical Asia and Oceania (i.e., excluding Australia). Crops are listed
from largest to smallest total areal extent in 2010. Areas for 2000 were obtained by
averaging across 1997-2003, and areas for 2010 by averaging across 2007-2013
525 (averaging across multiple years removes the problem of missing cells). Total areas of
the major 20 crops and of all 163 tropical crops are shown.

Crop Name	Total		Neotropics		Africa		Asia	
	2000	2010	2000	2010	2000	2010	2000	2010
1 Rice, paddy	109.6	118.3	5.8	5.5	7.0	9.4	96.8	103.4
2 Maize	59.5	73.5	24.0	26.7	20.6	28.3	14.9	18.4
3 Soybeans	24.8	40.5	16.4	28.1	0.8	1.1	7.6	11.3
4 Sorghum	36.1	36.8	3.1	3.2	22.8	26.0	10.2	7.6
5 Wheat	32.0	35.6	3.1	4.1	1.6	2.3	27.3	29.2
6 Millet	32.4	31.0	<0.05	<0.05	19.7	20.2	12.7	10.9
7 Beans, dry	21.4	25.6	6.9	6.3	4.6	6.7	9.9	12.6
8 Cassava	16.6	20.4	2.4	2.6	11.0	14.1	3.2	3.7
9 Sugar cane	15.7	20.2	8.2	11.7	0.8	1.0	6.6	7.5
10 Vegetables Primary	15.6	19.5	2.1	2.4	4.2	5.6	9.3	11.5
11 Groundnuts, with shell	17.2	18.9	0.3	0.3	8.7	11.3	8.2	7.3
12 Seed cotton	14.6	16.6	1.3	1.5	4.4	3.9	8.9	11.2
13 Oil, palm fruit	10.1	16.1	0.5	0.8	4.1	4.6	5.5	10.7
14 Forage products	11.7	14.0	2.3	2.3	0.1	0.2	9.2	11.5
15 Coconuts	10.4	11.4	0.6	0.7	0.9	1.1	8.9	9.6
16 Cow peas, dry	9.3	11.3	0.1	0.1	9.2	11.1	0.1	0.2
17 Coffee, green	10.2	10.3	5.7	5.6	2.3	2.2	2.2	2.5
18 Cocoa, beans	7.0	9.6	1.5	1.6	4.7	6.2	0.9	1.8
19 Chick peas	7.3	9.2	0.1	0.1	0.4	0.4	6.8	8.6
20 Rubber, natural	6.9	8.7	0.1	0.2	0.6	0.7	6.2	7.8
Subtotal (20 crops)	468.4	547.7	84.6	103.9	128.5	156.5	255.3	287.3
21 Sesame seed	5.6	7.3	0.2	0.3	2.5	3.6	2.8	3.5
22 Vegetables, fresh nes	6.2	7.1	0.4	0.5	1.8	2.2	4.0	4.4
23 Rapeseed	6.1	6.6	<0.05	0.1	<0.05	<0.05	6.1	6.4
24 Yams	4.0	5.7	0.1	0.2	3.9	5.5	<0.05	<0.05
25 Plantains	5.0	5.3	0.9	0.9	4.0	4.3	<0.05	0.1
26 Pigeon peas	4.3	5.3	<0.05	0.1	0.5	0.7	3.8	4.5
27 Cashew nuts, with shell	3.2	5.2	0.6	0.8	1.1	2.5	1.5	1.9
28 Potatoes	3.4	4.7	0.9	0.9	0.9	1.4	1.6	2.4
29 Bananas	3.8	4.4	1.2	1.2	1.3	1.5	1.3	1.8
30 Sweet potatoes	3.7	4.4	0.2	0.2	2.5	3.4	0.9	0.7
31 Mangoes, mangosteens, guavas	2.9	4.2	0.4	0.5	0.4	0.6	2.1	3.2
32 Pulses, nes	3.7	3.9	<0.05	<0.05	1.5	1.7	2.2	2.2

33	Sunflower seed	2.7	3.4	0.3	0.5	0.4	1.1	1.9	1.9
34	Fruit, fresh nes	2.4	3.4	0.2	0.3	0.7	0.7	1.4	2.4
35	Cereals, nes	2.5	3.3	<0.05	<0.05	2.4	3.2	0.1	0.1
36	Barley	2.6	2.6	0.8	0.7	1.0	1.1	0.8	0.7
37	Oranges	2.1	2.3	1.5	1.4	0.1	0.2	0.4	0.7
38	Onions, dry	1.1	2.1	0.2	0.2	0.2	0.5	0.7	1.4
39	Fruit, tropical fresh nes	1.4	2.0	0.1	0.2	0.1	0.2	1.1	1.6
40	Tobacco, unmanufactured	1.7	1.9	0.5	0.5	0.4	0.6	0.8	0.8
41	Lentils	1.7	1.8	<0.05	<0.05	0.1	0.1	1.6	1.6
42	Tomatoes	1.2	1.8	0.3	0.3	0.4	0.7	0.5	0.8
43	Chillies and peppers, dry	1.7	1.7	<0.05	0.1	0.5	0.5	1.2	1.2
44	Jute & Jute-like Fibres	1.6	1.5	<0.05	<0.05	<0.05	<0.05	1.6	1.5
45	Peas, dry	1.3	1.5	0.1	0.1	0.4	0.6	0.8	0.8
46	Castor oil seed	1.1	1.4	0.2	0.2	0.1	0.2	0.8	1.0
47	Jute	1.4	1.4	<0.05	<0.05	<0.05	<0.05	1.4	1.4
48	Taro (cocoyam)	1.3	1.3	<0.05	<0.05	1.2	1.2	0.1	0.1
49	Tea	1.0	1.3	<0.05	<0.05	0.2	0.3	0.8	1.0
50	Oilseeds nes	1.2	1.2	NA	NA	0.6	0.7	0.6	0.5
51	Okra	0.8	1.2	<0.05	<0.05	0.4	0.7	0.4	0.5
52	Roots and tubers, nes	0.9	1.2	0.1	0.1	0.7	0.9	0.1	0.1
53	Fruit, citrus nes	0.9	1.1	<0.05	0.1	0.9	0.9	<0.05	0.1
54	Melonseed	0.9	0.9	<0.05	<0.05	0.9	0.9	NA	NA
55	Pumpkins, squash and gourds	0.7	0.9	0.1	0.1	0.1	0.2	0.4	0.6
56	Pineapples	0.7	0.8	0.2	0.2	0.2	0.3	0.3	0.3
57	Areca nuts	0.5	0.8	NA	NA	NA	NA	0.5	0.8
58	Forage and silage, grasses nes	0.8	0.8	0.4	0.5	NA	NA	0.3	0.3
59	Spices, nes	0.7	0.8	<0.05	<0.05	<0.05	<0.05	0.6	0.7
60	Broad beans, horse beans, dry	0.6	0.8	0.2	0.2	0.4	0.6	NA	NA
61	Eggplants (aubergines)	0.6	0.7	<0.05	<0.05	<0.05	<0.05	0.5	0.7
62	Cabbages and other brassicas	0.5	0.7	<0.05	<0.05	0.1	0.2	0.4	0.5
63	Cashewapple	0.7	0.7	0.6	0.6	0.1	0.1	NA	NA
64	Chillies and peppers, green	0.6	0.7	0.2	0.2	0.3	0.2	0.2	0.3
65	Maize, green	0.8	0.7	0.1	0.1	0.6	0.4	0.1	0.2
66	Anise, badian, fennel, coriander	0.4	0.6	<0.05	<0.05	<0.05	<0.05	0.4	0.6
67	Lemons and limes	0.4	0.6	0.2	0.3	<0.05	<0.05	0.2	0.3
68	Beans, green	0.7	0.6	<0.05	<0.05	<0.05	<0.05	0.6	0.5
69	Forage and silage, alfalfa	0.4	0.6	0.4	0.6	NA	NA	NA	NA
70	Linseed	0.8	0.6	<0.05	<0.05	0.1	0.1	0.7	0.4
71	Fonio	0.4	0.5	NA	NA	0.4	0.5	NA	NA
72	Cauliflowers and broccoli	0.3	0.5	<0.05	0.1	<0.05	<0.05	0.3	0.4
73	Pepper (piper spp.)	0.4	0.5	<0.05	<0.05	<0.05	<0.05	0.3	0.4
74	Karite nuts (sheanuts)	0.4	0.5	NA	NA	0.4	0.5	NA	NA
75	Peas, green	0.4	0.5	0.1	0.1	<0.05	<0.05	0.3	0.4
76	Kola nuts	0.4	0.5	NA	NA	0.4	0.5	NA	NA
77	Forage and silage, maize	0.3	0.5	0.3	0.5	NA	NA	NA	NA
78	Watermelons	0.3	0.4	0.2	0.2	<0.05	0.1	0.1	0.2

79	Nuts, nes	0.4	0.4	0.1	0.1	0.1	0.1	0.2	0.2
80	Apples	0.4	0.4	0.1	0.1	<0.05	<0.05	0.2	0.3
81	Sisal	0.3	0.4	0.3	0.3	0.1	0.1	<0.05	<0.05
82	Papayas	0.3	0.4	0.1	0.1	0.1	0.1	0.1	0.2
83	Cloves	0.5	0.4	NA	NA	0.1	0.1	0.4	0.3
84	Safflower seed	0.6	0.4	0.1	0.1	<0.05	<0.05	0.4	0.3
85	Cucumbers and gherkins	0.3	0.3	<0.05	<0.05	0.1	0.2	0.1	0.1
86	Garlic	0.2	0.3	<0.05	<0.05	<0.05	<0.05	0.2	0.3
87	Fibre crops nes	0.3	0.3	NA	NA	0.3	0.3	<0.05	<0.05
88	Avocados	0.2	0.3	0.2	0.2	<0.05	0.1	<0.05	<0.05
89	Oats	0.4	0.3	0.3	0.3	0.1	<0.05	NA	NA
90	Nutmeg, mace and cardamoms	0.2	0.3	0.1	0.1	<0.05	<0.05	0.1	0.2
91	Tea nes	0.2	0.3	<0.05	<0.05	0.2	0.3	NA	NA
92	Grapes	0.2	0.3	0.1	0.1	<0.05	<0.05	<0.05	0.1
93	Ginger	0.3	0.2	<0.05	<0.05	0.2	0.1	0.1	0.2
94	Forage and silage, sorghum	0.2	0.2	0.2	0.2	NA	NA	NA	NA
95	Bambara beans	0.1	0.2	NA	NA	0.1	0.2	NA	NA
96	Lettuce and chicory	0.2	0.2	<0.05	<0.05	<0.05	<0.05	0.1	0.2
97	Tangerines, mandarins, etc.	0.2	0.2	0.1	0.1	<0.05	<0.05	0.1	<0.05
98	Melons, other (inc.cantaloupes)	0.2	0.2	0.1	0.1	<0.05	<0.05	<0.05	0.1
99	Kapok fruit	0.2	0.2	NA	NA	NA	NA	0.2	0.2
100	Grapefruit (inc. pomelos)	0.1	0.2	0.1	0.1	<0.05	<0.05	0.1	0.1
101	Bastfibres, other	0.3	0.2	0.0	<0.05	<0.05	<0.05	0.2	0.1
102	Vetches	0.1	0.2	0.0	<0.05	0.1	0.2	NA	NA
103	Carrots and turnips	0.1	0.2	0.1	0.1	<0.05	<0.05	0.1	0.1
104	Manila fibre (abaca)	0.1	0.2	<0.05	<0.05	<0.05	<0.05	0.1	0.1
105	Cinnamon (canella)	0.1	0.2	<0.05	<0.05	<0.05	<0.05	0.1	0.2
106	Peaches and nectarines	0.1	0.1	0.1	0.1	<0.05	<0.05	<0.05	<0.05
107	Quinoa	0.1	0.1	0.1	0.1	NA	NA	NA	NA
108	Sugar crops, nes	0.1	0.1	0.0	0.0	NA	NA	0.1	0.1
109	Walnuts, with shell	0.1	0.1	<0.05	0.1	NA	NA	<0.05	<0.05
110	Maté	0.1	0.1	0.1	0.1	NA	NA	NA	NA
111	Canary seed	<0.05	0.1	<0.05	<0.05	NA	NA	<0.05	0.1
112	Mustard seed	<0.05	0.1	<0.05	<0.05	<0.05	<0.05	<0.05	0.1
113	Vegetables, leguminous nes	0.1	0.1	0.1	0.1	<0.05	<0.05	<0.05	<0.05
114	Vanilla	<0.05	0.1	<0.05	<0.05	<0.05	0.1	<0.05	<0.05
115	Onions, shallots, green	0.1	0.1	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05
116	Dates	<0.05	0.1	<0.05	<0.05	<0.05	0.1	NA	NA
117	Leeks, other alliaceous vegetables	<0.05	0.1	<0.05	<0.05	<0.05	<0.05	<0.05	0.1
118	Asparagus	<0.05	0.1	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05
119	Spinach	<0.05	0.1	<0.05	<0.05	<0.05	<0.05	<0.05	0.1
120	Buckwheat	<0.05	0.1	<0.05	<0.05	<0.05	<0.05	NA	NA
121	Forage and silage, rye grass	<0.05	0.1	<0.05	0.1	NA	NA	NA	NA
122	Triticale	<0.05	0.1	<0.05	0.1	NA	NA	NA	NA
123	Agave fibres nes	0.1	0.1	<0.05	<0.05	NA	NA	<0.05	<0.05

124	Berries nes	<0.05	0.1	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05
125	Pears	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05
126	Plums and sloes	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05
127	Yautia (cocoyam)	<0.05	<0.05	<0.05	<0.05	NA	NA	NA	NA
128	Chestnut	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	NA	NA
129	String beans	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05
130	Vegetables and roots fodder	<0.05	<0.05	NA	NA	<0.05	<0.05	<0.05	<0.05
131	Olives	<0.05	<0.05	<0.05	<0.05	NA	NA	NA	NA
132	Hops	<0.05	<0.05	NA	NA	<0.05	<0.05	NA	NA
133	Pyrethrum, dried	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05
134	Tung nuts	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	NA	NA
135	Lupins	<0.05	<0.05	<0.05	<0.05	NA	NA	NA	NA
136	Strawberries	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05
137	Brazil nuts, with shell	<0.05	<0.05	<0.05	<0.05	NA	NA	NA	NA
138	Figs	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05
139	Persimmons	<0.05	<0.05	<0.05	<0.05	NA	NA	NA	NA
140	Forage and silage, clover	<0.05	<0.05	<0.05	<0.05	NA	NA	NA	NA
141	Artichokes	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	NA	NA
142	Cassava leaves	<0.05	<0.05	NA	NA	<0.05	<0.05	<0.05	<0.05
143	Apricots	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05
144	Almonds, with shell	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	NA	NA
145	Rye	<0.05	<0.05	<0.05	<0.05	NA	NA	0.0	0.0
146	Mixed Grasses and Legumes	<0.05	<0.05	<0.05	<0.05	NA	NA	NA	NA
147	Cherries	<0.05	<0.05	<0.05	<0.05	NA	NA	<0.05	<0.05
148	Fruit, stone nes	<0.05	<0.05	NA	NA	<0.05	<0.05	<0.05	<0.05
149	Ramie	<0.05	<0.05	<0.05	<0.05	NA	NA	<0.05	<0.05
150	Sugar beet	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	NA	NA
151	Pistachios	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	NA	NA
152	Quinces	<0.05	<0.05	<0.05	<0.05	NA	NA	NA	NA
153	Raspberries	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	NA	NA
154	Turnips for fodder	<0.05	<0.05	<0.05	<0.05	NA	NA	NA	NA
155	Chicory roots	<0.05	<0.05	<0.05	<0.05	NA	NA	<0.05	<0.05
156	Blueberries	<0.05	<0.05	<0.05	<0.05	NA	NA	NA	NA
157	Mushrooms and truffles	<0.05	<0.05	NA	NA	NA	NA	<0.05	<0.05
158	Cherries, sour	<0.05	<0.05	<0.05	<0.05	NA	NA	NA	NA
159	Jojoba seed	<0.05	<0.05	<0.05	<0.05	NA	NA	NA	NA
160	Beets for fodder	<0.05	<0.05	<0.05	<0.05	NA	NA	NA	NA
161	Hazelnuts, with shell	0.0	<0.05	NA	NA	0.0	<0.05	NA	NA
162	Carobs	<0.05	<0.05	<0.05	<0.05	NA	NA	NA	NA
163	Swedes for fodder	<0.05	<0.05	<0.05	<0.05	NA	NA	NA	NA
Total		573.6	676.2	99.1	120.4	165.2	204.6	307.9	349.7

Reference

530 [1] FAO. 2015. Food and agriculture organization of the United Nations Statistics division. Available from <http://faostat.fao.org/>. Assessed Aug 2016.